

7. Literaturverzeichnis

- (1) Stanek G. et al.: European Lyme borreliosis. Ann. NY Acad. Sci. 539 (1988), 274-82
- (2) Wilske B., Steinhilber R. et al.: Lyme-Borreliose in Süddeutschland. Dtsch. med. Wschr. 112 (1987), 1730-1736
- (3) Krause A., Burmester G.: Lyme- Borreliose. Thieme Verlag Stuttgart-New York, 1999
- (4) Kramer M.D. et al.: Symptomatik und Diagnostik der Lyme-Borreliose, Dtsch. med. Wschr. 118 (1993), 423-427
- (5) Merkblatt des Robert-Koch-Institutes: Merkblatt für Ärzte, Lyme-Borreliose-Erkennung und Verhütung. Robert Koch-Institut, 1999
- (6) Thalhammer F., Hollenstein U., et al.: Infektionen, Diagnostik und Therapie der Lyme-Borreliose in der Praxis. <http://www.ahc-net.at>, Forum Dr.Med 13/99
- (7) Merkblatt des Robert-Koch-Institutes: Ratgeber Infektionskrankheiten. Robert Koch-Institut, Fachgebiet 23 Infektionsepidemiologie, 1999
- (8) Hülße Ch., Herrmann H., et al.: Untersuchungen zur Verbreitung von *Borrelia burgdorferi* in Mecklenburg-Vorpommern. Hyg. Med. 20 (1995), 345-350
- (9) Hassler D., Zöllner L., et al.: Lyme-Borreliose in einem europäischen Endemiegebiet, Antikörper-prävalenz und klinisches Spektrum. Dtsch. med. Wschr. 117 (1992), 767-774
- (10) Kahl O., Schmidt K., et al.: Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* Ticks in Berlin (West) Zbl. Bakt. Hyg. A. 270 (1989), 434-440
- (11) Gupta S.K., Schönberg A., et al.: Prevalence of ticks in relation to their role as vector of *Borrelia burgdorferi* under autochthon conditions. Applied Parasitology 36 (1995), 97-106
- (12) Schönberg A., Loser C.: Vorkommen von Borrelien im Berliner Raum. Durch Zecken übertragene Erkrankungen, FSME und Lyme-Borreliose, Potsdamer Symposium, Weller-Verlag 1994, 84-93
- (13) Cimmino M., Granström M., et al.: European Lyme Borreliosis. Clinical Spektrum. Zbl. Bakt. 287 (1998), 248-252
- (14) O'Connell S., Granström M., et al.: Epidemiology of European Lyme Borreliosis. Zbl. Bakt. 287 (1998), 229-240

- (15) Rath P.-M., Ibershoff A., et al.: Seroprevalence of Lyme Borreliosis in Forestry Workers from Brandenburg, Germany. *Eur. J. Clin. Microbiol. Infect. Dis.*, (15) 1996, 372-377
- (16) Maiwald M., Petney T.N., et al.: Untersuchungen zur natürlichen Epidemiologie der Lyme-Borreliose anlässlich des gehäuften Auftretens von Erkrankungen in einem Vorort einer nord-badischen Gemeinde. *Gesundh.-Wes.*, 57 (1995), 419-425
- (17) Kaiser R., Kern A., et al.: Prevalence of Antibody to *Borrelia burgdorferi* and Tick-borne Encephalitis Virus in an Endemic Region in Southern Germany. *Zbl. Bakt.* 286 (1997), 534-541
- (18) Stanek G. et al.: Lyme borreliosis and emerging tick-borne diseases in Europe. *Wien. Klin. Wochenschr.* 110/24 (1998), 847-849
- (19) Burgdorfer W., Barbour A.G., et al.: Lyme disease- a tick-borne spirochetosis? *Science* 216 (1982), 1317-1319
- (20) Franz J.K., Krause A. : Lyme disease (Lyme borreliosis), *Best Practice & Research Clinical Rheumatology* 17/2 (2003), 241-264
- (21) Steere A.C.: Lyme disease. *The New England Journal of Medicine.* 345 (2001), 115-125
- (22) Nadelmann R.B. et al.: Prophylaxis with single-dose doxycycline for the prevention of Lyme disease after an *Ixodes scapularis* tick bite. *N. Engl. J. Med.* 345 (2001), 79-84
- (23) Heininger U., Zimmermann T., et al.: Zeckenstich und Lyme-Borreliose. Eine epidemiologische Untersuchung im Raum Erlangen. *Monatsschr. Kinderheilkd.* 141 (1993), 874-877
- (24) Paul H., Gerth H.J., et al.: Infectiousness for humans of *Ixodes ricinus* containing *Borrelia burgdorferi*. *Zbl. Bakt. Hyg. A* 263 (1986), 473-476
- (25) Assous M.V., Postic D. et al.: Western blot analysis of sera from Lyme borreliosis patients according to the genomic species of *Borrelia* strains used as antigens. *Eur. J. Clin. Microbiol. Infect. Dis.* 12 (1993), 261-268
- (26) Canica M.M., Nato F. et al.: Monoclonal antibodies for identification of *Borrelia afzelii* sp nov associated with late cutaneous manifestation of Lyme borreliosis. *Scand. J. Infect. Dis.* 25 (1993), 441-448
- (27) van Dam A.P., Kuiper H. et al.: Different genospecies of *Borrelia burgdorferi* are associated with distinct clinical manifestation of Lyme borreliosis. *Clin. Infect. Dis.* 17 (1993), 708-717

- (28) Mather T.N., Nicholson M.C. et al.: Entomologic index for human risk of Lyme disease. *J. Clin. Microbiol.* 144 (11) (1996), 1066-1069
- (29) Maupin G.O., Fish D. et al.: Landscape ecology of Lyme disease in a residential area of Westchester County, New York. *Am. J. Epidemiol.* 133 (1991), 1105-1113
- (30) Berger B.W., Johnson R.C. et al.: Cultivation of *Borrelia burgdorferi* from erythema migrans lesions and perilesional skin. *J. Clin. Microbiol.* 30 (1992), 359-361
- (31) Gustafson R.: Epidemiological studies of Lyme borreliosis and tick-borne encephalitis. *Scand. J. Infect. Dis. Suppl.* 92 (1994), 1-63
- (32) Strl F.: Lyme borreliosis in Slovenia. *Zbl. Bakteriol.* 289 (1999), 643-652
- (33) Huppertz H.I., Bohme M. et al.: Incidence of Lyme borreliosis in the Würzburg region of Germany. *Eur. J. Clin. Microbiol. Infect. Dis.* 18 (10) (1999), 697-703
- (34) Williams C.L., Strobino B. et al.: Lyme disease in childhood: clinical and epidemiologic feature: ninety cases. *Pediatr. Infect. Dis. J.* 9 (1990), 10-4
- (35) Hobusch D., Christen H.J. et al.: Diagnosis and therapy of Lyme borreliosis in children. *Practic guideline of the German Society for Pediatric. Infectious Disease. Klin. Pädiatr.* 211 (2) (1999), 70-4
- (36) Christen H.J.: Lyme neuroborreliosis in children. *Ann. Med.* 28 (3) (1996), 235-40
- (37) Aberer E., Kehldorfer M. et al.: The outcome of Lyme borreliosis in children. *Wien. Klin. Wochenschr.* 111 (22-23) (1999), 941-9
- (38) Niemann G., Koksal M.A. et al.: Facial palsy and Lyme borreliosis: long-term follow-up of children with antibioticly untreated „idiopathic” facial palsy. *Klin. Pädiatr.* 209 (3) (1997), 95-9
- (39) Christen H.J., Hanefeld F. et al.: Epidemiology and clinical manifestations of Lyme borreliosis in childhood. A prospectiv multicentre study with special regard in neuroborreliosis. *Acta. Pädiatr. Sppl.* 386 (1993), 1-75
- (40) Saint Girons I., Gern L. et al.: Identification of *Borrelia burgdorferi* sensu lato species in Europe. *Zbl. Bakteriol.* 287 (3) (1998), 190-5
- (41) Maiwald M., Oehme R. et al.: Transmission risk of *Borrelia burgdorferi* sensu lato from *Ixodes ricinus* ticks to humans in southwest Germany. *Epidemiol. Infect.* 121 (1) (1998), 103-8
- (42) Lünemann J.D., Krause A.: Heterogenität von *Borrelia burgdorferi*: ätiopathogenetische Relevanz und klinische Implikationen. *Zeitschr. Rheumatol.* 62 (2) (2003), 148-154
- (43) Hubalek Z., Halouzka J.: Distribution of *Borrelia* sensu lato genomic group in Europe, a review. *Eur. J. Epidemiol.* 13 (8) (1997), 951-7

- (44) Stanek G. et al.: European Union Concerted Action on Risk Assessment in Lyme Borreliosis: clinical case definitions for Lyme Borreliosis. *Wien. Klin. Wschr.* 108 (1996), 741-747
- (45) Hunfeld K.P., Stanek G. et al.: Lessons from the German Proficiency Testing Program 1999-2001. A preliminary report. *Wien. Klin. Wschr.* 114 (13-14) (2002), 591-600
- (46) Wilske B.: Microbiological diagnosis in Lyme borreliosis. *Int. J. Med. Microbiol.* 291 (Suppl 33) (2002), 114-119
- (47) Reimer B., Erbas B. et al.: Seroprevalence of *Borrelia* infection in occupational tick-exposed people in Bavaria (Germany). *Int. J. Med. Microbiol.* 291 (Suppl.33) (2002), 215
- (48) Fitzner J., Ammon A. et al.: Risk factors in Lyme borreliosis: a German case-control study. *Int. J. Med. Microbiol.* 291 (Suppl. 33) (2002), 220
- (49) Kampen H., Rotzel D.C. et al.: Substantial rise in the prevalence of Lyme borreliosis spirochetes in a region of western Germany over a 10-year period. *Appl. Environ. Microbiol.* 70 (3) (2004), 1576-82
- (50) Oehme R., Hartelt K. et al.: Foci of tick-borne diseases in southwest Germany. *Int. J. Med. Microbiol.* 291 (Suppl. 33) (2002), 22-29
- (51) Hubalek Z., Halouzka J.: Prevalence rates of *Borrelia burgdorferi* sensu lato in host-seeking *Ixodes ricinus* ticks in Europe. *Parasitol. Res.* 84 (1998), 167-172
- (52) Klein J.D., Eppes S.C. et al.: Environmental and life-style risk factors for Lyme disease in children. *Clin. Pediatr.(Phila.)* 35 (7) (1996), 359- 363
- (53) Robertson J., Guy E. et al.: A european multicenter study of immunoblotting in serodiagnosis of lyme borreliosis. *J. Clin. Microbiol.* 38 (6) (2000), 2097- 2102
- (54) Priem S., Rittig M.G. et al.: An optimized PCR leads to rapid and highly sensitive dedection of *Borrelia burgdorferi* in patients with Lyme borreliosis. *J. Clin. Microbiol.* 35 (3) (1997), 685- 690
- (55) Maiwald M., Stockinger C. et al.: Evaluation of the detection of *Borrelia burgdorferi* DNA in urin samples by Polymerase chain reaction. *Infection.* 23 (3) (1995), 173- 179
- (56) Franz J.K., Priem S. et al.: Studies on the pathogenesis and treatment of Lyme arthritis. *Wien. Klin. Wochenschr.* 111 (22-23) (1999), 981- 984
- (57) Lunemann J.D., Zarmas S. et al.: Rapid typing of *Borrelia burgdorferi* sensu lato species in specimens from patients with different manifestations of Lyme borreliosis. *J. Clin. Microbiol.* 39 (3) (2001), 1130- 1133

- (58) Pleyer U., Priem S. et al.: Detection of *Borrelia burgdorferi* DNA in urin of patients with ocular Lyme borreliosis. *Br. J. Ophthalmol.* 85 (5) (2001), 552- 555
- (59) Sigal L.H.: Lyme disease and the Lyme disease vaccines. *Bull. Rheumat. Dis.* 48 (4) (1999), 1-4
- (60) Steere A.C., Sikand V.K. et al.: Vaccination against Lyme disease with rekombinant *Borrelia burgdorferi* outer-surface lipoprotein A with adjuvant. Lyme Disease Vaccine Study Group. *N. Engl. J. Med.* 339 (4) (1998), 209-215
- (61) Nocton J.J., Dressler F. et al.: Detection of *Borrelia burgdorferi* DNA by Polymerase chain reaction in synovial fluid from patients with Lyme arthritis. *N. Engl. J. Med.* 330 (4) (1994), 229- 234
- (62) Nocton J.J., Bloom B. J. et al.: Detection of *Borrelia burgdorferi* DNA by polymerase chain reaction in cerebrospinal fluid in Lyme neuroborreliosis. *J. Infect. Dis.* 174 (3) (1996), 623- 627
- (63) Steere A.C., Green J. et al.: Treatment of Lyme disease. *Zbl. Bakteriол. Mikrobiol. Hyg.* 263 (3) (1987), 352-356
- (64) Wormser G.P., Ramanathan R. et al.: Duration of antibiotic therapy for early Lyme disease. A randomized, double-blind, placebo-controlled trial. *Ann. Intern. Med.* 138 (9) (2003), 697-704
- (65) Steere A.C., Gross D. et al.: Autoimmune mechanisms in antibiotic treatment-resistant lyme arthritis. *J. Autoimmun.* 16 (3) (2001), 263- 268
- (66) Drouin E.E., Glickstein L.J. et al.: Molecular characterization of the OspA (161-175) T cell epitope associated with treatment-resistant Lyme arthritis: differences among the three pathogenic species of *Borrelia burgdorferi sensu lato*. *J. Autoimmun.* 23 (3) (2004), 281-292
- (67) Priem S., Burmester G. et al.: Detection of *Borrelia burgdorferi* by polymerase chain re-action in synovial membrane, but not in synovial fluid in patients with persisting Lyme arthritis after antibiotic therapy. *Ann. Rheum. Dis.* 57 (2) (1998), 118-121
- (68) Wang G., van Dam A.P. et al.: Molecular typing of *Borrelia burgdorferi sensu lato*: taxonomic, epidemio-logical and clinical implications. *Clin. Microbiol. Rev.* 12 (4) (1999), 633-653
- (69) Berglund J., Eitrem R. et al.: An epidemiologic study of Lyme disease in southern Sweden. *N. Eng. J. Med.* 333 (20) (1995), 1319-1327
- (70) Breier F.H., Aberer E. et al.: Isolation of *Borrelia afzelii* from circumscribed scleroderma. *Br. J. Dermatol.* 140 (5) (1999), 925- 930

- (71) Steere A.C., Bartenhagen N.H. et al.: Clinical manifestations of Lyme disease. *Zbl. Bakteriolog. Mikrobiolog. Hyg.* 263 (1-2) (1986), 201-205
- (72) Steere A.C., Batsford W.P. et al.: Lyme carditis: cardiac abnormalities of Lyme disease. *Ann. Intern. Med.* 93 (1) (1980), 8-16
- (73) Kaplan R.F., Jones-Woodward L. et al.: Neuropsychological deficits in Lyme disease patients with and without other evidence of central nervous system pathology. *Appl. Neuropsychol.* 6 (1) (1999), 3-11
- (74) Steere A.C.: Musculoskeletal manifestations of Lyme disease. *Am. J. Med.* 98 (4A) (1995), 44-48
- (75) Logigian E.L., Kaplan R.F., Steere A.C.: Chronic neurologic manifestations of Lyme disease. *N. Engl. J. Med.* 323 (21) (1990), 1438-1444
- (76) Steere A.C., Schoen R.T., Taylor E.: The clinical evolution of Lyme arthritis. *Ann. Intern. Med.* 107 (5) (1987), 725-31
- (77) Kornmehl E.W., Lesser R.L. et al.: Bilateral keratitis in Lyme disease. *Ophthalmology* 96 (8) (1989), 1194-1197
- (78) Wormser G.P., Nadelmann R.B. et al.: Practise guideline for the treatment of Lyme disease. *Infect. Dis. Soc. Am.* 31 Suppl.1 (2000), 1-14
- (79) Huppertz H.I., Karch H. et al.: Lyme arthritis in European children and adolescents. The Pediatric Rheumatology Collaborative Group. *Arthritis. Rheum.* 38 (3) (1995), 361-368
- (80) Smith R.P., Schoen R.T. et al.: Clinical characteristics and treatment outcome of early Lyme disease in patients with microbiologically confirmed erythema migrans. *Ann. Intern. Med.* 136 (6) (2002), 421-428
- (81) Steere A.C., Taylor E. et al.: The overdiagnosis of Lyme disease. *JAMA* 269 (14) (1993), 1812-1816
- (82) Shadick N.A., Phillips C.B. et al.: The long-term clinical outcomes of Lyme disease. A population-based retrospective cohort study. *Ann. Intern. Med.* 121 (8) (1994), 560-567
- (83) Nowakowski J., Nadelmann R.B. et al.: Long-term follow-up of patients with culture-confirmed Lyme disease. *Am. J. Med.* 115 (2) (2003), 91-96
- (84) Steere A.C., Green J. et al.: Treatment of Lyme disease. *Zbl. Bakteriolog. Mikrobiolog. Hyg.* 263 (3) (1987), 352-356
- (85) Logigian E.L., Kaplan R.F., Steere A.C.: Successful treatment of Lyme encephalopathy with intravenous ceftriaxon. *J. Infect. Dis.* 180 (2) (1999), 377-383

- (86) Steere A.C., Levine R.E., Molloy P.J. et al.: Treatment of Lyme arthritis. *Arthritis Rheum.* 37 (6) (1994), 878-888
- (87) Kalish R.A., Mc Hugh G. et al.: Persistence of immunoglobulin M or immunoglobulin G antibody responses to *Borrelia burgdorferi* 10-20 years after active Lyme disease. *Clin. Infect. Dis.* 33 (6) (2001), 780-785
- (88) Steere A.C., Berardi V.P. et al.: Evaluation of the intrathecal antibody response to *Borrelia burgdorferi* as a diagnostic test for Lyme neuroborreliosis. *J. Infect. Dis.* 161 (6) (1990), 1203-1209
- (89) Steere A.C., Coburn J. et al.: The emergence of Lyme disease. *J. Clin. Invest.* 113 (8) (2004), 1093-1101
- (90) Bannwarth A.: Zur Klinik und Pathogenese der "chronischen lymphozytären Meningitis". *Archiv für Psychiatrie und Nervenkrankheiten.* 117 (1944), 161-185
- (91) Garin C., Bujadoux A.: Paralysis par le tiques. *J. Med. Lyon* 71 (1922), 765-767
- (92) Herxheimer K., Hartmann K.: Über die Acrodermatitis chronica atrophicans. *Arch. Dermatol. (Berl)*, 61 (1902), 57-76
- (93) Afzelius A.: Verhandlungen der dermatologischen Gesellschaft zu Stockholm. Sitzung vom 28. Oktober 1909. *Arch. Dermatol. Syph.* 101 (1910), 403-404
- (94) Robert-Koch-Institut: Zur Lyme-Borreliose im Land Brandenburg. Erfahrungen nach langjähriger Surveillance. *Epid. Bull.* 20 (2005), 173-178
- (95) Talaska T., Bätzing-Feigenbaum J. et al.: Untersuchungen zu Seroprävalenz und Klinik von Lyme-Borreliose und „Humaner Granulozytärer Ehrlichiose“ bei Waldarbeitern und Förstern in Brandenburg und Berlin. *RKI: Epid. Bull.* 19 (2001), 109-110
- (96) Robert-Koch-Institut: Risikofaktoren für Lyme-Borreliose: Ergebnisse einer Studie in einem Brandenburger Landkreis. *Epid. Bull.* 16 (2001), 147-149
- (97) Robert-Koch-Institut: Zur Situation der wichtigen Infektionskrankheiten in Deutschland. Erkrankungen an Lyme-Borreliose in den sechs östlichen Bundesländern 2002-2003. *Epid. Bull.* 28 (2004), 219-222
- (98) Orloski K. A. et al. : Surveillance for Lyme disease-United States 1992-1998. *MMWR* 49 (2000), 1-11
- (99) Darai G., Sonntag H.G.(Hrsg): *Lexikon der Infektionskrankheiten des Menschen*, Heidelberg 2003, S.74-75
- (100) Robert-Koch-Institut: *Ratgeber Infektionskrankheiten*, Berlin 2001