

6. Literatur

- Acheson, A., Sunshine, J.L. und Rutishauser, U. (1991):
NCAM polysialic acid can regulate both cell-cell and cell-substrate interactions.
J Cell Biol, 114: 143-53.
- Albelda, S.M., Smith, C.W. und Ward, P.A. (1994):
Adhesion molecules and inflammatory injury.
FASEB J, 8: 504-12.
- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K. und Watson, J. (1997):
Molekularbiologie der Zelle.
VHC Verlagsgesellschaft mbH, Weinheim.
- Angata, K., Suzuki, M. und Fukuda, M. (1998):
Differential and cooperative polysialylation of the neural cell adhesion molecule by two polysialyltransferases, PST and STX [In Process Citation].
J Biol Chem, 273: 28524-32.
- Aplin, A.E., Howe, A., Alahari, S.K. und Juliano, R.L. (1998):
Signal transduction and signal modulation by cell adhesion receptors: the role of integrins, cadherins, immunoglobulin-cell adhesion molecules, and selectins.
Pharmacol Rev, 50: 197-263.
- Arimura, N., Inagaki, N., Chihara, K., Menager, C., Nakamura, N., Amano, M., Iwamatsu, A., Goshima, Y. und Kaibuchi, K. (2000):
Phosphorylation of collapsin response mediator protein-2 by Rho-kinase. Evidence for two separate signalling pathways for growth cone collapse.
J Biol Chem, 275: 23973-80.
- Atkins, A.R., Osborne, M.J., Lashuel, H.A., Edelman, G.M., Wright, P.E., Cunningham, B.A. und Dyson, H.J. (1999):
Association between the first two immunoglobulin-like domains of the neural cell adhesion molecule N-CAM.
FEBS Lett, 451: 162-8.
- Avraham, H., Park, S.Y., Schinkmann, K. und Avraham, S. (2000):
RAFTK/Pyk2-mediated cellular signalling.
Cell Signal, 12: 123-33.
- Avraham, S., London, R., Fu, Y., Ota, S., Hiregowdara, D., Li, J., Jiang, S., Pasztor, L.M., White, R.A., Groopman, J.E. und et al. (1995):
Identification and characterization of a novel related adhesion focal tyrosine kinase (RAFTK) from megakaryocytes and brain.
J Biol Chem, 270: 27742-51.
- Baldwin, T.J., Fazeli, M.S., Doherty, P. und Walsh, F.S. (1996):
Elucidation of the molecular actions of NCAM and structurally related cell adhesion molecules.
J Cell Biochem, 61: 502-13.
- Barthels, D., Vopper, G., Boned, A., Cremer, H. und Wille, W. (1992):
High Degree of NCAM Diversity Generated by Alternative RNA Splicing in Brain and Muscle.
Eur J Neurosci, 4: 327-337.

- Beavis, R.C. und Chait, B.T. (1996):
Matrix-assisted laser desorption ionization mass-spectrometry of proteins.
Methods Enzymol, 270: 519-51.
- Beggs, H.E., Baragona, S.C., Hemperly, J.J. und Maness, P.F. (1997):
NCAM140 interacts with the focal adhesion kinase p125(fak) and the SRC- related tyrosine kinase p59(fyn).
J Biol Chem, 272: 8310-9.
- Beggs, H.E., Soriano, P. und Maness, P.F. (1994):
NCAM-dependent neurite outgrowth is inhibited in neurons from Fyn-minus mice.
J Cell Biol, 127: 825-33.
- Birnboim, H.C. und Doly, J. (1979):
A rapid alkaline extraction procedure for screening recombinant plasmid DNA.
Nucleic Acids Res, 7: 1513-23.
- Blaukat, A., Ivankovic-Dikic, I., Gronroos, E., Dolfi, F., Tokiwa, G., Vuori, K. und Dikic, I. (1999):
Adaptor proteins Grb2 and Crk couple Pyk2 with activation of specific mitogen- activated protein kinase cascades.
J Biol Chem, 274: 14893-901.
- Bollen, M. (2001):
Combinatorial control of protein phosphatase-1.
Trends Biochem Sci, 26: 426-31.
- Bourtchuladze, R., Frenguelli, B., Blendy, J., Cioffi, D., Schutz, G. und Silva, A.J. (1994):
Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive element-binding protein.
Cell, 79: 59-68.
- Bradford, M.M. (1976):
A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.
Anal Biochem, 72: 248-54.
- Brennan, C.M., Gallouzi, I.E. und Steitz, J.A. (2000):
Protein ligands to HuR modulate its interaction with target mRNAs in vivo.
J Cell Biol, 151: 1-14.
- Brittis, P.A., Silver, J., Walsh, F.S. und Doherty, P. (1996):
Fibroblast Growth Factor Receptor Function Is Required for the Orderly Projection of Ganglion Cell Axons in the Developing Mammalian Retina.
Mol Cell Neurosci, 8: 120-8.
- Brooks, J.E. (1987):
Properties and uses of restriction endonucleases.
Methods Enzymol, 152: 113-29.
- Brummendorf, T. und Rathjen, F.G. (1995):
Cell adhesion molecules 1: immunoglobulin superfamily.
Protein Profile, 2: 963-1108.
- Bruses, J.L. und Rutishauser, U. (2001):
Roles, regulation, and mechanism of polysialic acid function during neural development.
Biochimie, 83: 635-43.

- Buchner, J. und Rudolph, R. (1991):
Routes to active proteins from transformed microorganisms.
Curr Opin Biotechnol, 2: 532-8.
- Byk, T., Dobranksy, T., Cifuentes-Diaz, C. und Sobel, A. (1996):
Identification and molecular characterization of Unc-33-like phosphoprotein (Ulip), a putative mammalian homolog of the axonal guidance-associated unc-33 gene product.
J Neurosci, 16: 688-701.
- Cary, L.A., Han, D.C., Polte, T.R., Hanks, S.K. und Guan, J.L. (1998):
Identification of p130Cas as a mediator of focal adhesion kinase-promoted cell migration.
J Cell Biol, 140: 211-21.
- Cavallaro, U., Niedermeyer, J., Fuxa, M. und Christofori, G. (2001):
N-CAM modulates tumour-cell adhesion to matrix by inducing FGF-receptor signalling.
Nat Cell Biol, 3: 650-7.
- Chen, H.C., Chan, P.C., Tang, M.J., Cheng, C.H. und Chang, T.J. (1998):
Tyrosine phosphorylation of focal adhesion kinase stimulated by hepatocyte growth factor leads to mitogen-activated protein kinase activation.
J Biol Chem, 273: 25777-82.
- Choi, J., Krushel, L.A. und Crossin, K.L. (2001):
NF-kappaB activation by N-CAM and cytokines in astrocytes is regulated by multiple protein kinases and redox modulation.
Glia, 33: 45-56.
- Chou, D.K., Ilyas, A.A., Evans, J.E., Costello, C., Quarles, R.H. und Jungalwala, F.B. (1986):
Structure of sulfated glucuronyl glycolipids in the nervous system reacting with HNK-1 antibody and some IgM paraproteins in neuropathy.
J Biol Chem, 261: 11717-25.
- Cole, G.J. und Akeson, R. (1989):
Identification of a heparin binding domain of the neural cell adhesion molecule N-CAM using synthetic peptides.
Neuron, 2: 1157-65.
- Cowin, P. und Burke, B. (1996):
Cytoskeleton-membrane interactions.
Curr Opin Cell Biol, 8: 56-65.
- Cremer, H., Chazal, G., Carleton, A., Goridis, C., Vincent, J.D. und Lledo, P.M. (1998):
Long-term but not short-term plasticity at mossy fiber synapses is impaired in neural cell adhesion molecule-deficient mice.
Proc Natl Acad Sci U S A, 95: 13242-7.
- Cremer, H., Chazal, G., Goridis, C. und Represa, A. (1997):
NCAM is essential for axonal growth and fasciculation in the hippocampus.
Mol Cell Neurosci, 8: 323-35.
- Cremer, H., Lange, R., Christoph, A., Plomann, M., Vopper, G., Roes, J., Brown, R., Baldwin, S., Kraemer, P., Scheff, S. und et al. (1994a):
Inactivation of the N-CAM gene in mice results in size reduction of the olfactory bulb and deficits in spatial learning.
Nature, 367: 455-9.

- Cremer, H., Lange, R., Christoph, A., Plomann, M., Vopper, G., Roes, J., Brown, R., Baldwin, S., Kraemer, P., Scheff, S. und et al. (1994b):
Inactivation of the N-CAM gene in mice results in size reduction of the olfactory bulb and deficits in spatial learning.
Nature, 367: 455-9.
- Crossin, K.L., Tai, M.H., Krushel, L.A., Mauro, V.P. und Edelman, G.M. (1997):
Glucocorticoid receptor pathways are involved in the inhibition of astrocyte proliferation.
Proc Natl Acad Sci U S A, 94: 2687-92.
- Cunningham, B.A., Hemperly, J.J., Murray, B.A., Prediger, E.A., Brackenbury, R. und Edelman, G.M. (1987):
Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing.
Science, 236: 799-806.
- Dagert, M. und Ehrlich, S.D. (1979):
Prolonged incubation in calcium chloride improves the competence of Escherichia coli cells.
Gene, 6: 23-8.
- Dai, R.M. und Li, C.C. (2001):
Valosin-containing protein is a multi-ubiquitin chain-targeting factor required in ubiquitin-proteasome degradation.
Nat Cell Biol, 3: 740-4.
- Dalby, K.N., Morrice, N., Caudwell, F.B., Avruch, J. und Cohen, P. (1998):
Identification of regulatory phosphorylation sites in mitogen-activated protein kinase (MAPK)-activated protein kinase-1a/p90rsk that are inducible by MAPK.
J Biol Chem, 273: 1496-505.
- Daniloff, J.K., Levi, G., Grumet, M., Rieger, F. und Edelman, G.M. (1986):
Altered expression of neuronal cell adhesion molecules induced by nerve injury and repair.
J Cell Biol, 103: 929-45.
- Deak, M., Clifton, A.D., Lucocq, L.M. und Alessi, D.R. (1998):
Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38, and may mediate activation of CREB.
Embo J, 17: 4426-41.
- Dickson, G., Gower, H.J., Barton, C.H., Prentice, H.M., Elsom, V.L., Moore, S.E., Cox, R.D., Quinn, C., Putt, W. und Walsh, F.S. (1987):
Human muscle neural cell adhesion molecule (N-CAM): identification of a muscle-specific sequence in the extracellular domain.
Cell, 50: 1119-30.
- Dietrich, J., Kastrup, J., Nielsen, B.L., Odum, N. und Geisler, C. (1997):
Regulation and function of the CD3gamma DxxxLL motif: a binding site for adaptor protein-1 and adaptor protein-2 in vitro.
J Cell Biol, 138: 271-81.
- Ditlevsen, D.K., Kohler, L.B., Pedersen, M.V., Risell, M., Kolkova, K., Meyer, M., Berezin, V. und Bock, E. (2003):
The role of phosphatidylinositol 3-kinase in neural cell adhesion molecule-mediated neuronal differentiation and survival.
J Neurochem, 84: 546-56.

Divecha, N. und Irvine, R.F. (1995):

Phospholipid signaling.

Cell, 80: 269-78.

Doherty, P., Ashton, S.V., Moore, S.E. und Walsh, F.S. (1991a):

Morphoregulatory activities of NCAM and N-cadherin can be accounted for by G protein-dependent activation of L- and N-type neuronal Ca²⁺ channels.

Cell, 67: 21-33.

Doherty, P., Ashton, S.V., Moore, S.E. und Walsh, F.S. (1991b):

Morphoregulatory activities of NCAM and N-cadherin can be accounted for by G protein-dependent activation of L- and N-type neuronal Ca²⁺ channels.

Cell, 67: 21 - 33.

Doherty, P., Barton, C.H., Dickson, G., Seaton, P., Rowett, L.H., Moore, S.E., Gower, H.J. und Walsh, F.S. (1989):

Neuronal process outgrowth of human sensory neurons on monolayers of cells transfected with cDNAs for five human N-CAM isoforms.

J. Cell Biol., 109: 789-798.

Doherty, P., Cohen, J. und Walsh, F.S. (1990):

Neurite outgrowth in response to transfected N-CAM changes during development and is modulated by polysialic acid.

Neuron, 5: 209-19.

Doherty, P., Rimon, G., Mann, D.A. und Walsh, F.S. (1992a):

Alternative splicing of the cytoplasmic domain of neural cell adhesion molecule alters its ability to act as a substrate for neurite outgrowth.

J Neurochem, 58: 2338-41.

Doherty, P., Skaper, S.D., Moore, S.E., Leon, A. und Walsh, F.S. (1992b):

A developmentally regulated switch in neuronal responsiveness to NCAM and N-cadherin in the rat hippocampus.

Development, 115: 885-92.

Doherty, P. und Walsh, F.S. (1992):

Cell adhesion molecules, second messengers and axonal growth.

Curr Opin Neurobiol, 2: 595-601.

Doherty, P. und Walsh, F.S. (1996):

CAM-FGF Receptor Interactions: A Model for Axonal Growth.

Mol Cell Neurosci, 8: 99-111.

Dolfi, F., Garcia-Guzman, M., Ojaniemi, M., Nakamura, H., Matsuda, M. und Vuori, K. (1998):

The adaptor protein Crk connects multiple cellular stimuli to the JNK signaling pathway.

Proc Natl Acad Sci U S A, 95: 15394-9.

Doyle, E., Nolan, P.M., Bell, R. und Regan, C.M. (1992):

Hippocampal NCAM180 transiently increases sialylation during the acquisition and consolidation of a passive avoidance response in the adult rat.

J Neurosci Res, 31: 513-23.

Dzhandzhugazyan, K. und Bock, E. (1997):

Demonstration of an extracellular ATP-binding site in NCAM: functional implications of nucleotide binding.

Biochemistry, 36: 15381-95.

Eckhardt, M., Mühlenhoff, M., Bethe, A., Koopman, J., Frosch, M. und Gerardy-Schahn, R. (1995):

Molecular characterization of eukaryotic polysialyltransferase-1.

Nature, 373: 715-8.

Edelman, G.M. (1985):

Cell adhesion and the molecular processes of morphogenesis.

Ann. Rev. Biochem., 54: 135-169.

Erbe, D.V., Watson, S.R., Presta, L.G., Wolitzky, B.A., Foxall, C., Brandley, B.K. und Lasky, L.A. (1993):

P- and E-selectin use common sites for carbohydrate ligand recognition and cell adhesion.

J Cell Biol, 120: 1227-35.

Falasca, M., Logan, S.K., Lehto, V.P., Baccante, G., Lemmon, M.A. und Schlessinger, J. (1998):

Activation of phospholipase C gamma by PI 3-kinase-induced PH domain-mediated membrane targeting.

Embo J, 17: 414-22.

Foxall, C., Watson, S.R., Dowbenko, D., Fennie, C., Lasky, L.A., Kiso, M., Hasegawa, A., Asa, D. und Brandley, B.K. (1992):

The three members of the selectin receptor family recognize a common carbohydrate epitope, the sialyl Lewis(x) oligosaccharide.

J Cell Biol, 117: 895-902.

Friedlander, D.R., Milev, P., Karthikeyan, L., Margolis, R.K., Margolis, R.U. und Grumet, M. (1994):

The neuronal chondroitin sulfate proteoglycan neurocan binds to the neural cell adhesion molecules Ng-CAM/L1/NILE and N-CAM, and inhibits neuronal adhesion and neurite outgrowth.

J Cell Biol, 125: 669-80.

Geschwind, D.H., Kelly, G.M., Fryer, H., Feeser-Bhatt, H. und Hockfield, S. (1996):

Identification and characterization of novel developmentally regulated proteins in rat spinal cord.

Brain Res Dev Brain Res, 97: 62-75.

Goridis, C. und Brunet, J.F. (1992):

NCAM: structural diversity, function and regulation of expression.

Semin Cell Biol, 3: 189-97.

Goridis, C., Deagostini-Bazin, H., Hirn, M., Hirsch, M.R., Rougon, G., Sadoul, R., Langley, O.K., Gombos, G. und Finne, J. (1983):

Neural surface antigens during nervous system development.

Cold Spring Harb Symp Quant Biol, 48 Pt 2: 527-37.

Goshima, Y., Nakamura, F., Strittmatter, P. und Strittmatter, S.M. (1995):

Collapsin-induced growth cone collapse mediated by an intracellular protein related to UNC-33.

Nature, 376: 509-14.

Greene, L.A. und Tischler, A.S. (1976):

Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor.

Proc Natl Acad Sci U S A, 73: 2424-8.

- Hagag, N., Halegoua, S. und Viola, M. (1986):
Inhibition of growth factor-induced differentiation of PC12 cells by microinjection of antibody to ras p21.
Nature, 319: 680-2.
- Hall, H., Williams, E.J., Moore, S.E., Walsh, F.S., Prochiantz, A. und Doherty, P. (1996):
Inhibition of FGF-stimulated phosphatidylinositol hydrolysis and neurite outgrowth by a cell-membrane permeable phosphopeptide.
Curr Biol, 6: 580-7.
- Heasley, L.E. und Johnson, G.L. (1992):
The beta-PDGF receptor induces neuronal differentiation of PC12 cells.
Mol Biol Cell, 3: 545-53.
- Hoffman, S. und Edelman, G.M. (1987):
A proteoglycan with HNK-1 antigenic determinants is a neuron-associated ligand for cytactin.
Proc Natl Acad Sci U S A, 84: 2523-7.
- Horstkorte, R., Leßner, N., Gerardy-Schahn, R., Lucka, L., Danker, K. und Reutter, W. (1999):
Expression of the polysialyltransferase ST8SiaIV: polysialylation interferes with adhesion of PC12 cells in vitro.
Exp Cell Res, 246: 122-8.
- Horstkorte, R., Schachner, M., Magyar, J.P., Vorherr, T. und Schmitz, B. (1993):
The fourth immunoglobulin-like domain of NCAM contains a carbohydrate recognition domain for oligomannosidic glycans implicated in association with L1 and neurite outgrowth.
J Cell Biol, 121: 1409-21.
- Hu, H., Tomasiewicz, H., Magnuson, T. und Rutishauser, U. (1996):
The role of polysialic acid in migration of olfactory bulb interneuron precursors in the subventricular zone.
Neuron, 16: 735-43.
- Hynes, R.O. und Lander, A.D. (1992):
Contact and adhesive specificities in the associations, migrations, and targeting of cells and axons.
Cell, 68: 303-22.
- Jessen, U., Novitskaya, V., Pedersen, N., Serup, P., Berezin, V. und Bock, E. (2001):
The transcription factors CREB and c-Fos play key roles in NCAM-mediated neuritogenesis in PC12-E2 cells.
J Neurochem, 79: 1149-60.
- Johnson, G.L. und Lapadat, R. (2002):
Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases.
Science, 298: 1911-2.
- Jones, F.S., Holst, B.D., Minowa, O., De Robertis, E.M. und Edelman, G.M. (1993):
Binding and transcriptional activation of the promoter for the neural cell adhesion molecule by HoxC6 (Hox-3.3).
Proc Natl Acad Sci U S A, 90: 6557-61.

Jones, F.S., Prediger, E.A., Bittner, D.A., De Robertis, E.M. und Edelman, G.M. (1992):
Cell adhesion molecules as targets for Hox genes: neural cell adhesion molecule promoter activity is modulated by cotransfection with Hox-2.5 and -2.4.
Proc Natl Acad Sci U S A, 89: 2086-90.

Juliano, R.L. (2002):
Signal transduction by cell adhesion receptors and the cytoskeleton: functions of integrins, cadherins, selectins, and immunoglobulin-superfamily members.
Annu Rev Pharmacol Toxicol, 42: 283-323.

Kamiguchi, H., Long, K.E., Pendergast, M., Schaefer, A.W., Rapoport, I., Kirchhausen, T. und Lemmon, V. (1998):
The neural cell adhesion molecule L1 interacts with the AP-2 adaptor and is endocytosed via the clathrin-mediated pathway.
J Neurosci, 18: 5311-21.

Katayose, Y., Li, M., Al-Murrani, S.W., Shenolikar, S. und Damuni, Z. (2000):
Protein phosphatase 2A inhibitors, I(1)(PP2A) and I(2)(PP2A), associate with and modify the substrate specificity of protein phosphatase 1.
J Biol Chem, 275: 9209-14.

Keilhauer, G., Faissner, A. und Schachner, M. (1985):
Differential inhibition of neurone-neurone, neurone-astrocyte and astrocyte-astrocyte adhesion by L1, L2 and N-CAM antibodies.
Nature, 316: 728-30.

Kessels, M.M. und Qualmann, B. (2002):
Syndapins integrate N-WASP in receptor-mediated endocytosis.
Embo J, 21: 6083-94.

Kirchhausen, T., Bonifacino, J.S. und Riezman, H. (1997):
Linking cargo to vesicle formation: receptor tail interactions with coat proteins.
Curr Opin Cell Biol, 9: 488-95.

Kiselyov, V.V., Berezin, V., Maar, T.E., Soroka, V., Edvardsen, K., Schousboe, A. und Bock, E. (1997a):
The first immunoglobulin-like neural cell adhesion molecule (NCAM) domain is involved in double-reciprocal interaction with the second immunoglobulin-like NCAM domain and in heparin binding.
J Biol Chem, 272: 10125-34.

Kiselyov, V.V., Berezin, V., Maar, T.E., Soroka, V., Edvardsen, K., Schousboe, A. und Bock, E. (1997b):
The first immunoglobulin-like neural cell adhesion molecule (NCAM) domain is involved in double-reciprocal interaction with the second immunoglobulin-like NCAM domain and in heparin binding.
J Biol Chem, 272: 10125-34.

Kiselyov, V.V., Skladchikova, G., Hinsby, A.M., Jensen, P.H., Kulahin, N., Soroka, V., Pedersen, N., Tsetlin, V., Poulsen, F.M., Berezin, V. und Bock, E. (2003):
Structural basis for a direct interaction between FGFR1 and NCAM and evidence for a regulatory role of ATP.
Structure (Camb), 11: 691-701.

Kishimoto, A., Takai, Y., Mori, T., Kikkawa, U. und Nishizuka, Y. (1980):
Activation of calcium and phospholipid-dependent protein kinase by diacylglycerol, its possible relation to phosphatidylinositol turnover.
J Biol Chem, 255: 2273-6.

Kiss, J.Z., Troncoso, E., Djebbara, Z., Vutskits, L. und Muller, D. (2001):
The role of neural cell adhesion molecules in plasticity and repair.
Brain Res Brain Res Rev, 36: 175-84.

Kiss, J.Z., Wang, C., Olive, S., Rougon, G., Lang, J., Baetens, D., Harry, D. und Pralong, W.F. (1994):
Activity-dependent mobilization of the adhesion molecule polysialic NCAM to the cell surface of neurons and endocrine cells.
Embo J, 13: 5284-92.

Klausner, R.D., Van Renswoude, J., Ashwell, G., Kempf, C., Schechter, A.N., Dean, A. und Bridges, K.R. (1983):
Receptor-mediated endocytosis of transferrin in K562 cells.
J Biol Chem, 258: 4715-24.

Kolkova, K., Novitskaya, V., Pedersen, N., Berezin, V. und Bock, E. (2000a):
Neural cell adhesion molecule-stimulated neurite outgrowth depends on activation of protein kinase C and the Ras-mitogen-activated protein kinase pathway.
J Neurosci, 20: 2238-46.

Kolkova, K., Pedersen, N., Berezin, V. und Bock, E. (2000b):
Identification of an amino acid sequence motif in the cytoplasmic domain of the NCAM-140 kDa isoform essential for its neuritogenic activity.
J Neurochem, 75: 1274-82.

Kozma, R., Sarner, S., Ahmed, S. und Lim, L. (1997):
Rho family GTPases and neuronal growth cone remodelling: relationship between increased complexity induced by Cdc42Hs, Rac1, and acetylcholine and collapse induced by RhoA and lysophosphatidic acid.
Mol Cell Biol, 17: 1201-11.

Krog, L., Olsen, M., Dalseg, A.M., Roth, J. und Bock, E. (1992):
Characterization of soluble neural cell adhesion molecule in rat brain, CSF, and plasma.
J Neurochem, 59: 838-47.

Krushel, L.A., Tai, M.H., Cunningham, B.A., Edelman, G.M. und Crossin, K.L. (1998):
Neural cell adhesion molecule (N-CAM) domains and intracellular signaling pathways involved in the inhibition of astrocyte proliferation.
Proc Natl Acad Sci U S A, 95: 2592-6.

Laemmli, U.K. (1970):
Cleavage of structural proteins during the assembly of the head of bacteriophage T4.
Nature, 227: 680-5.

Leshchyns'ka, I., Sytnyk, V., Morrow, J.S. und Schachner, M. (2003):
Neural cell adhesion molecule (NCAM) association with PKC β 2 via betaI spectrin is implicated in NCAM-mediated neurite outgrowth.
J Cell Biol, 161: 625-39.

Leung, T., Chen, X.Q., Manser, E. und Lim, L. (1996):
The p160 RhoA-binding kinase ROK alpha is a member of a kinase family and is involved in the reorganization of the cytoskeleton.
Mol Cell Biol, 16: 5313-27.

Leung, T., Manser, E., Tan, L. und Lim, L. (1995):
A novel serine/threonine kinase binding the Ras-related RhoA GTPase which translocates the kinase to peripheral membranes.
J Biol Chem, 270: 29051-4.

- Li, M., Makkinje, A. und Damuni, Z. (1996):
Molecular identification of I1PP2A, a novel potent heat-stable inhibitor protein of protein phosphatase 2A.
Biochemistry, 35: 6998-7002.
- Little, E.B., Crossin, K.L., Krushel, L.A., Edelman, G.M. und Cunningham, B.A. (2001):
A short segment within the cytoplasmic domain of the neural cell adhesion molecule (N-CAM) is essential for N-CAM-induced NF-kappa B activity in astrocytes.
Proc Natl Acad Sci U S A, 98: 2238-43.
- Little, E.B., Edelman, G.M. und Cunningham, B.A. (1998):
Palmitoylation of the cytoplasmic domain of the neural cell adhesion molecule N-CAM serves as an anchor to cellular membranes.
Cell Adhes Commun, 6: 415-30.
- Liu, L., Haines, S., Shew, R. und Akeson, R.A. (1993):
Axon growth is enhanced by NCAM lacking the VASE exon when expressed in either the growth substrate or the growing axon.
J Neurosci Res, 35: 327-45.
- Luo, L. (2000):
Rho GTPases in neuronal morphogenesis.
Nat Rev Neurosci, 1: 173-80.
- Maccioni, R.B. und Cambiazo, V. (1995):
Role of microtubule-associated proteins in the control of microtubule assembly.
Physiol Rev, 75: 835-64.
- Main, A.L., Harvey, T.S., Baron, M., Boyd, J. und Campbell, I.D. (1992):
The three-dimensional structure of the tenth type III module of fibronectin: an insight into RGD-mediated interactions.
Cell, 71: 671-8.
- Marais, R. und Marshall, C.J. (1996):
Control of the ERK MAP kinase cascade by Ras and Raf.
Cancer Surv, 27: 101-25.
- Matsuoka, K., Taoka, M., Satozawa, N., Nakayama, H., Ichimura, T., Takahashi, N., Yamakuni, T., Song, S.Y. und Isobe, T. (1994):
A nuclear factor containing the leucine-rich repeats expressed in murine cerebellar neurons.
Proc Natl Acad Sci U S A, 91: 9670-4.
- Meiri, K.F., Saffell, J.L., Walsh, F.S. und Doherty, P. (1998):
Neurite outgrowth stimulated by neural cell adhesion molecules requires growth-associated protein-43 (GAP-43) function and is associated with GAP-43 phosphorylation in growth cones.
J Neurosci, 18: 10429-37.
- Minana, R., Duran, J.M., Tomas, M., Renau-Piqueras, J. und Guerri, C. (2001):
Neural cell adhesion molecule is endocytosed via a clathrin-dependent pathway.
Eur J Neurosci, 13: 749-56.
- Minturn, J.E., Fryer, H.J., Geschwind, D.H. und Hockfield, S. (1995):
TOAD-64, a gene expressed early in neuronal differentiation in the rat, is related to unc-33, a C. elegans gene involved in axon outgrowth.
J Neurosci, 15: 6757-66.

- Mohammadi, M., Froum, S., Hamby, J.M., Schroeder, M.C., Panek, R.L., Lu, G.H., Eliseenkova, A.V., Green, D., Schlessinger, J. und Hubbard, S.R. (1998):
Crystal structure of an angiogenesis inhibitor bound to the FGF receptor tyrosine kinase domain.
Embo J, 17: 5896-904.
- Muller, D., Djebbara-Hannas, Z., Jourdain, P., Vutskits, L., Durbec, P., Rougon, G. und Kiss, J.Z. (2000):
Brain-derived neurotrophic factor restores long-term potentiation in polysialic acid-neural cell adhesion molecule-deficient hippocampus.
Proc Natl Acad Sci U S A, 97: 4315-20.
- Muller, D., Wang, C., Skibo, G., Toni, N., Cremer, H., Calaora, V., Rougon, G. und Kiss, J.Z. (1996):
PSA-NCAM is required for activity-induced synaptic plasticity.
Neuron, 17: 413-22.
- Mullis, K.B. und Falloona, F.A. (1987):
Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction.
Meth. Enzymol, 155: 335-50.
- Nakayama, J., Fukuda, M.N., Fredette, B., Ranscht, B. und Fukuda, M. (1995):
Expression cloning of a human polysialyltransferase that forms the polysialylated neural cell adhesion molecule present in embryonic brain.
Proc Natl Acad Sci U S A, 92: 7031-5.
- Nelson, R.W., Bates, P.A. und Rutishauser, U. (1995):
Protein determinants for specific polysialylation of the neural cell adhesion molecule.
J Biol Chem, 270: 17171-9.
- Niethammer, P., Delling, M., Sytnyk, V., Dityatev, A., Fukami, K. und Schachner, M. (2002):
Cosignaling of NCAM via lipid rafts and the FGF receptor is required for neuritogenesis.
J Cell Biol, 157: 521-32.
- Nishiki, T., Narumiya, S., Morii, N., Yamamoto, M., Fujiwara, M., Kamata, Y., Sakaguchi, G. und Kozaki, S. (1990):
ADP-ribosylation of the rho/rac proteins induces growth inhibition, neurite outgrowth and acetylcholine esterase in cultured PC-12 cells.
Biochem Biophys Res Commun, 167: 265-72.
- Nusser, N., Gosmanova, E., Zheng, Y. und Tigyi, G. (2002):
Nerve growth factor signals through TrkA, phosphatidylinositol 3-kinase, and Rac1 to inactivate RhoA during the initiation of neuronal differentiation of PC12 cells.
J Biol Chem, 277: 35840-6.
- Olsen, M., Krog, L., Edvardsen, K., Skovgaard, L.T. und Bock, E. (1993):
Intact transmembrane isoforms of the neural cell adhesion molecule are released from the plasma membrane.
Biochem J, 295: 833-40.
- Olsen, M., Zuber, C., Roth, J., Linnemann, D. und Bock, E. (1995):
The ability to re-express polysialylated NCAM in soleus muscle after denervation is reduced in aged rats compared to young adult rats.
Int J Dev Neurosci, 13: 97-104.

- Ono, K., Tomasiewicz, H., Magnuson, T. und Rutishauser, U. (1994):
N-CAM mutation inhibits tangential neuronal migration and is phenocopied by enzymatic removal of polysialic acid.
Neuron, 13: 595-609.
- Opal, P., Garcia, J.J., Propst, F., Matilla, A., Orr, H.T. und Zoghbi, H.Y. (2003):
Mapmodulin/leucine-rich acidic nuclear protein binds the light chain of microtubule-associated protein 1B and modulates neuritogenesis.
J Biol Chem, 278: 34691-9.
- Orsulic, S. und Peifer, M. (1996):
Cell-cell signalling: Wingless lands at last.
Curr Biol, 6: 1363-7.
- Oudega, M., Touri, F., Deenen, M.G., Riederer, B.M. und Marani, E. (1998):
Microtubule-associated protein 1a is involved in the early development.
Neurosci Lett, 246: 81-4.
- Owens, G.C., Edelman, G.M. und Cunningham, B.A. (1987):
Organization of the neural cell adhesion molecule (N-CAM) gene: alternative exon usage as the basis for different membrane-associated domains.
Proc Natl Acad Sci U S A, 84: 294-8.
- Pappin, D.J. (2003):
Peptide mass fingerprinting using MALDI-TOF mass spectrometry.
Methods Mol Biol, 211: 211-9.
- Paratcha, G., Ledda, F. und Ibanez, C.F. (2003):
The neural cell adhesion molecule NCAM is an alternative signaling receptor for GDNF family ligands.
Cell, 113: 867-79.
- Park, T.U., Lucka, L., Reutter, W. und Horstkorte, R. (1997):
Turnover studies of the neural cell adhesion molecule NCAM: degradation of NCAM in PC12 cells depends on the presence of NGF.
Biochem Biophys Res Commun, 234: 686-9.
- Peck, D. und Walsh, F.S. (1993):
Differential effects of over-expressed neural cell adhesion molecule isoforms on myoblast fusion.
J Cell Biol, 123: 1587-95.
- Perkins, D.N., Pappin, D.J., Creasy, D.M. und Cottrell, J.S. (1999):
Probability-based protein identification by searching sequence databases using mass spectrometry data.
Electrophoresis, 20: 3551-67.
- Persohn, E., Pollerberg, G.E. und Schachner, M. (1989):
Immunoelectron-microscopic localization of the 180 kD component of the neural cell adhesion molecule N-CAM in postsynaptic membranes.
J Comp Neurol, 288: 92-100.
- Pleasure, I.T., Black, M.M. und Keen, J.H. (1993):
Valosin-containing protein, VCP, is a ubiquitous clathrin-binding protein.
Nature, 365: 459-62.

- Pollerberg, E.G., Sadoul, R., Goridis, C. und Schachner, M. (1985):
Selective expression of the 180-kD component of the neural cell adhesion molecule N-CAM during development.
J Cell Biol, 101: 1921-9.
- Pollerberg, G.E., Burridge, K., Krebs, K.E., Goodman, S.R. und Schachner, M. (1987):
The 180-kD component of the neural cell adhesion molecule N-CAM is involved in a cell-cell contacts and cytoskeleton-membrane interactions.
Cell Tissue Res, 250: 227-36.
- Pollerberg, G.E., Nolte, C. und Schachner, M. (1990):
Accumulation of N-CAM 180 at Contact Sites Between Neuroblastoma Cells and Latex Beads Coated with Extracellular Matrix Molecules.
Eur J Neurosci, 2: 879-887.
- Pollerberg, G.E., Schachner, M. und Davoust, J. (1986):
Differentiation state-dependent surface mobilities of two forms of the neural cell adhesion molecule.
Nature, 324: 462-5.
- Qualmann, B. und Kelly, R.B. (2000):
Syndapin isoforms participate in receptor-mediated endocytosis and actin organization.
J Cell Biol, 148: 1047-62.
- Qualmann, B., Roos, J., DiGregorio, P.J. und Kelly, R.B. (1999):
Syndapin I, a synaptic dynamin-binding protein that associates with the neural Wiskott-Aldrich syndrome protein.
Mol Biol Cell, 10: 501-13.
- Quinn, C.C., Gray, G.E. und Hockfield, S. (1999):
A family of proteins implicated in axon guidance and outgrowth.
J Neurobiol, 41: 158-64.
- Ranheim, T.S., Edelman, G.M. und Cunningham, B.A. (1996)
Homophilic adhesion mediated by the neural cell adhesion molecule involves multiple immunoglobulin domains.
Proc Natl Acad Sci USA, 93: 4071-5
- Rao, Y., Zhao, X. und Siu, C.H. (1994)
Mechanism of homophilic binding mediated by the neural cell adhesion molecule NCAM . Evidence for isologous interaction.
J Biol Chem, 269: 27540-8.
- Rhee, S.G. und Bae, Y.S. (1997):
Regulation of phosphoinositide-specific phospholipase C isozymes.
J Biol Chem, 272: 15045-8.
- Ridley, A.J. und Hall, A. (1992):
The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors.
Cell, 70: 389-99.
- Ridley, A.J., Paterson, H.F., Johnston, C.L., Diekmann, D. und Hall, A. (1992):
The small GTP-binding protein rac regulates growth factor-induced membrane ruffling.
Cell, 70: 401-10.

- Roberson, E.D., English, J.D., Adams, J.P., Selcher, J.C., Kondratick, C. und Sweatt, J.D. (1999):
The mitogen-activated protein kinase cascade couples PKA and PKC to cAMP response element binding protein phosphorylation in area CA1 of hippocampus.
J Neurosci, 19: 4337-48.
- Ronn, L.C., Bock, E., Linnemann, D. und Jahnsen, H. (1995):
NCAM-antibodies modulate induction of long-term potentiation in rat hippocampal CA1.
Brain Res, 677: 145-51.
- Ronn, L.C., Doherty, P., Holm, A., Berezin, V. und Bock, E. (2000):
Neurite outgrowth induced by a synthetic peptide ligand of neural cell adhesion molecule requires fibroblast growth factor receptor activation.
J Neurochem, 75: 665-71.
- Ronn, L.C., Olsen, M., Ostergaard, S., Kiselyov, V., Berezin, V., Mortensen, M.T., Lerche, M.H., Jensen, P.H., Soroka, V., Saffell, J.L., Doherty, P., Poulsen, F.M., Bock, E., Holm, A. und Saffells, J.L. (1999):
Identification of a neuritogenic ligand of the neural cell adhesion molecule using a combinatorial library of synthetic peptides.
Nat Biotechnol, 17: 1000-5.
- Rutishauser, U., Acheson, A., Hall, A.K., Mann, D.M. und Sunshine, J. (1988):
The neural cell adhesion molecule (NCAM) as a regulator of cell-cell interactions.
Science, 240: 53-7.
- Saffell, J.L., Doherty, P., Tiveron, M.C., Morris, R.J. und Walsh, F.S. (1995):
NCAM requires a cytoplasmic domain to function as a neurite outgrowth-promoting neuronal receptor.
Mol Cell Neurosci, 6: 521-31.
- Saffell, J.L., Walsh, F.S. und Doherty, P. (1994):
Expression of NCAM containing VASE in neurons can account for a developmental loss in their neurite outgrowth response to NCAM in a cellular substratum.
J Cell Biol, 125: 427-36.
- Saffell, J.L., Williams, E.J., Mason, I.J., Walsh, F.S. und Doherty, P. (1997):
Expression of a dominant negative FGF receptor inhibits axonal growth and FGF receptor phosphorylation stimulated by CAMs.
Neuron, 18: 231-42.
- Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A. und Arnheim, N. (1985):
Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia.
Science, 230: 1350-4.
- Sakai, R., Iwamatsu, A., Hirano, N., Ogawa, S., Tanaka, T., Nishida, J., Yazaki, Y. und Hirai, H. (1994):
Characterization, partial purification, and peptide sequencing of p130, the main phosphoprotein associated with v-Crk oncogene product.
J Biol Chem, 269: 32740-6.
- Sambrook, J., Fritsch, E.F. und Maniatis, T. (1989):
Molecular cloning: A laboratory manual.
Cold Spring Harbour Laboratory Press, Cold Spring Harbour, New York.

- Sanger, F., Nicklen, S. und Coulson, A.R. (1977):
DNA sequencing with chain-terminating inhibitors.
Proc Natl Acad Sci U S A, 74: 5463-7.
- Sano, M. und Kitajima, S. (1998):
Activation of mitogen-activated protein kinases is not required for the extension of neurites from PC12D cells triggered by nerve growth factor.
Brain Res, 785: 299-308.
- Santoni, M.-J., Barthels, D., Barbas, J.A., Hirsch, M.-R., Steinmetz, M., Goridis, C. und Wille, W. (1987):
Analysis of cDNA clones that code for the transmembrane form of the mouse neural cell adhesion molecule (N-CAM) and are generated by alternative RNA splicing.
Nucl. Acids Res., 15: 8621-8641.
- Scheidegger, E.P., Sternberg, L.R., Roth, J. und Lowe, J.B. (1995):
A human STX cDNA confers polysialic acid expression in mammalian cells.
J Biol Chem, 270: 22685-8.
- Schlaepfer, D.D., Broome, M.A. und Hunter, T. (1997):
Fibronectin-stimulated signaling from a focal adhesion kinase-c-Src complex: involvement of the Grb2, p130cas, and Nck adaptor proteins.
Mol Cell Biol, 17: 1702-13.
- Schlaepfer, D.D., Hanks, S.K., Hunter, T. und van der Geer, P. (1994):
Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase.
Nature, 372: 786-91.
- Schlaepfer, D.D., Hauck, C.R. und Sieg, D.J. (1999):
Signaling through focal adhesion kinase.
Prog Biophys Mol Biol, 71: 435-78.
- Schmid, R.S., Graff, R.D., Schaller, M.D., Chen, S., Schachner, M., Hemperly, J.J. und Maness, P.F. (1999):
NCAM stimulates the Ras-MAPK pathway and CREB phosphorylation in neuronal cells.
J Neurobiol, 38: 542-58.
- Schmidt, C., Kunemund, V., Wintergerst, E.S., Schmitz, B. und Schachner, M. (1996):
CD9 of mouse brain is implicated in neurite outgrowth and cell migration in vitro and is associated with the alpha 6/beta 1 integrin and the neural adhesion molecule L1.
J Neurosci Res, 43: 12-31.
- Schuch, U., Lohse, M.J. und Schachner, M. (1989):
Neural cell adhesion molecules influence second messenger systems.
Neuron, 3: 13-20.
- Seki, T. und Rutishauser, U. (1998):
Removal of polysialic acid-neural cell adhesion molecule induces aberrant mossy fiber innervation and ectopic synaptogenesis in the hippocampus.
J Neurosci, 18: 3757-66.
- Shin, M.H., Lee, E.G., Lee, S.H., Lee, Y.S. und Son, H. (2002):
Neural cell adhesion molecule (NCAM) promotes the differentiation of hippocampal precursor cells to a neuronal lineage, especially to a glutamatergic neural cell type.
Exp Mol Med, 34: 401-10.

Skladchikova, G., Ronn, L.C., Berezin, V. und Bock, E. (1999):
Extracellular adenosine triphosphate affects neural cell adhesion molecule (NCAM)-mediated cell adhesion and neurite outgrowth.
J Neurosci Res, 57: 207-18.

Small, S.J. und Akeson, R. (1990):
Expression of the unique NCAM VASE exon is independently regulated in distinct tissues during development.
J Cell Biol, 111: 2089-96.

Small, S.J., Shull, G.E., Santoni, M.J. und Akeson, R. (1987):
Identification of a cDNA clone that contains the complete coding sequence for a 140-kD rat NCAM polypeptide.
J Cell Biol, 105: 2335-45.

Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J. und Klenk, D.C. (1985):
Measurement of protein using bicinchoninic acid [published erratum appears in Anal Biochem 1987 May 15;163(1):279].
Anal Biochem, 150: 76-85.

Son, H., Seuk Kim, J., Mogg Kim, J., Lee, S.H. und Lee, Y.S. (2002):
Reciprocal actions of NCAM and tPA via a Ras-dependent MAPK activation in rat hippocampal neurons.
Biochem Biophys Res Commun, 298: 262-8.

Sorkin, B.C., Hoffman, S., Edelman, G.M. und Cunningham, B.A. (1984):
Sulfation and phosphorylation of the neural cell adhesion molecule, N- CAM.
Science, 225: 1476-8.

Storms, S.D., Jensen, J.J., Yaghmai, D. und Murray, B.A. (1994):
Multiple mechanisms of N2A and CHO cell adhesion to NCAM purified from chick embryonic brain and retina.
Exp Cell Res, 214: 100-12.

Sweatt, J.D. (2001):
The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory.
J Neurochem, 76: 1-10.

Takeichi, M. (1995):
Morphogenetic roles of classic cadherins.
Curr Opin Cell Biol, 7: 619-27.

Tedder, T.F., Steeber, D.A., Chen, A. und Engel, P. (1995):
The selectins: vascular adhesion molecules.
FASEB J, 9: 866-73.

Thiery, J.P., Brackenbury, R., Rutishauser, U. und Edelman, G.M. (1977):
Adhesion among neural cells of the chick embryo. II. Purification and characterization of a cell adhesion molecule from neural retina.
J Biol Chem, 252: 6841-5.

Thomaidou, D., Coquillat, D., Meintanis, S., Noda, M., Rougon, G. und Matsas, R. (2001):
Soluble forms of NCAM and F3 neuronal cell adhesion molecules promote Schwann cell migration: identification of protein tyrosine phosphatases zeta/beta as the putative F3 receptors on Schwann cells.
J Neurochem, 78: 767-78.

- Tomasiewicz, H., Ono, K., Yee, D., Thompson, C., Goridis, C., Rutishauser, U. und Magnuson, T. (1993):
Genetic deletion of a neural cell adhesion molecule variant (N-CAM-180) produces distinct defects in the central nervous system.
Neuron, 11: 1163-74.
- Tonissen, K.F. und Krieg, P.A. (1993):
Two neural-cell adhesion molecule (NCAM)-encoding genes in Xenopus laevis are expressed during development and in adult tissues.
Gene, 127: 243-7.
- Ueda, Y., Hirai, S., Osada, S., Suzuki, A., Mizuno, K. und Ohno, S. (1996):
Protein kinase C activates the MEK-ERK pathway in a manner independent of Ras and dependent on Raf.
J Biol Chem, 271: 23512-9.
- Uehata, M., Ishizaki, T., Satoh, H., Ono, T., Kawahara, T., Morishita, T., Tamakawa, H., Yamagami, K., Inui, J., Maekawa, M. und Narumiya, S. (1997):
Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension.
Nature, 389: 990-4.
- Ulitzur, N., Rancano, C. und Pfeffer, S.R. (1997):
Biochemical characterization of mapmodulin, a protein that binds microtubule-associated proteins.
J Biol Chem, 272: 30577-82.
- Vaughn, D.E. und Bjorkman, P.J. (1996):
The (Greek) key to structures of neural adhesion molecules.
Neuron, 16: 261-73.
- Vestweber, D. und Blanks, J.E. (1999):
Mechanisms that regulate the function of the selectins and their ligands.
Physiol Rev, 79: 181-213.
- Vossler, M.R., Yao, H., York, R.D., Pan, M.G., Rim, C.S. und Stork, P.J. (1997):
cAMP activates MAP kinase and Elk-1 through a B-Raf- and Rap1-dependent pathway.
Cell, 89: 73-82.
- Vutskits, L., Djebbara-Hannas, Z., Zhang, H., Paccaud, J.P., Durbec, P., Rougon, G., Muller, D. und Kiss, J.Z. (2001):
PSA-NCAM modulates BDNF-dependent survival and differentiation of cortical neurons.
Eur J Neurosci, 13: 1391-402.
- Walsh, F.S., Meiri, K. und Doherty, P. (1997):
Cell signalling and CAM-mediated neurite outgrowth.
Soc Gen Physiol Ser, 52: 221-6.
- Wary, K.K., Mariotti, A., Zurzolo, C. und Giancotti, F.G. (1998):
A requirement for caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth.
Cell, 94: 625-34.
- Weiss, B., Live, T.R. und Richardson, C.C. (1968):
Enzymatic breakage and joining of deoxyribonucleic acid. V. End group labeling and analysis of deoxyribonucleic acid containing single stranded breaks.
J Biol Chem, 243: 4530-42.

- Whitmarsh, A.J., Shore, P., Sharrocks, A.D. und Davis, R.J. (1995):
Integration of MAP kinase signal transduction pathways at the serum response element.
Science, 269: 403-7.
- Williams, A.F. und Barclay, A.N. (1988):
The immunoglobulin superfamily--domains for cell surface recognition.
Annu Rev Immunol, 6: 381-405.
- Williams, E.J., Furness, J., Walsh, F.S. und Doherty, P. (1994a):
Activation of the FGF receptor underlies neurite outgrowth stimulated by L1, N-CAM, and N-cadherin.
Neuron, 13: 583-94.
- Williams, E.J., Furness, J., Walsh, F.S. und Doherty, P. (1994b):
Characterisation of the second messenger pathway underlying neurite outgrowth stimulated by FGF.
Development, 120: 1685-93.
- Williams, E.J., Mittal, B., Walsh, F.S. und Doherty, P. (1995):
A Ca²⁺/calmodulin kinase inhibitor, KN-62, inhibits neurite outgrowth stimulated by CAMs and FGF.
Mol Cell Neurosci, 6: 69-79.
- Williams, E.J., Walsh, F.S. und Doherty, P. (1994c):
The production of arachidonic acid can account for calcium channel activation in the second messenger pathway underlying neurite outgrowth stimulated by NCAM, N-cadherin, and L1.
J Neurochem, 62: 1231-4.
- Yoshida, Y., Kojima, N., Kurosawa, N., Hamamoto, T. und Tsuji, S. (1995):
Molecular cloning of Sia alpha 2,3Gal beta 1,4GlcNAc alpha 2,8- sialyltransferase from mouse brain.
J Biol Chem, 270: 14628-33.
- Zalk, R. und Shoshan-Barmatz, V. (2003):
ATP-binding sites in brain p97/VCP (valosin-containing protein), a multifunctional AAA ATPase.
Biochem J, 374: 473-80.
- Zanassi, P., Paolillo, M., Feliciello, A., Avvedimento, E.V., Gallo, V. und Schinelli, S. (2001):
cAMP-dependent protein kinase induces cAMP-response element-binding protein phosphorylation via an intracellular calcium release/ERK-dependent pathway in striatal neurons.
J Biol Chem, 276: 11487-95.
- Zhou, F.Q., Zhong, J. und Snider, W.D. (2003):
Extracellular crosstalk: when GDNF meets N-CAM.
Cell, 113: 814-5.