

6. Literatur

1. Akimatsu T, Gute DC, Korthuis RJ. Leukocyte adhesion induced by inhibition of nitric oxide production in skeletal muscle. *J Appl Physiol* 1995;78: 1725-1732
2. Allen DM, Chen LE, Seaber AV, Urbaniak JR. Pathophysiology and related studies of the no reflow phenomenon in skeletal muscle. *Clin Orthop* 1995;314: 122-133
3. Aruoma OI, Halliwell B, Hoey BM, Butler J. The antioxidant action of N-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide, and hypochlorous acid. *Free Radic Biol Med* 1989;6: 593-597
4. Bagge U, Blixt A, Strid KG. The initiation of post-capillary margination of leukocytes: studies in vitro on the influence of erythrocyte concentration and flow velocity. *Int J Microcirc Clin Exp* 1983;2: 215-227
5. Baltzer AWA, Lattermann C, Whalen JD, Wooley P, Weiss K, Grimm M, Ghivizzani SC, Robbins PD, Evans CH. Genetic enhancement of fracture repair: healing of an experimental segmental defect by adenoviral transfer of the BMP-2 gene. *Gene Ther* 2000;7: 734-739
6. Barlow Y, Willoughby J. Pathophysiology of soft tissue repair. *Br Med Bull* 1992;48: 698-711
7. Behrens FF. Fractures with soft tissue injuries. In: Browner D, Jupiter JB, Levine AM, Trafton PG. *Skeletal trauma*. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo: W.B. Saunders Company 1998
8. Bombardier C, Reicin A, Shapiro D, Burgos-Vargas R, Davis B, Day R, Ferraz MB, Hawkey CJ, Hochberg MC, Kvien TK, Schnitzer TJ; VIGOR Study Group. Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. VIGOR Study Group. *N Engl J Med* 2000;343: 1520-1528
9. Bonnarens F, Einhorn TA. Production of a standard closed fracture in laboratory animal bone. *J Orthop Res* 1984;2: 97-101
10. Botha AJ, Moore EE. Base-deficit after major trauma directly relates to neutrophil CD 11b expression: a proposed mechanism of shock-induced organ injury. *Intensive Care Med* 1997;23: 504-509

11. Brookes M, Elkin AC, Harrison RG. A new concept of capillary circulation in bone cortex. *Lancet* 1961;1: 1078
12. Brooks PM. Nonsteroidal antiinflammatory drugs-differences and similarities. *N Engl J Med* 1991;325: 747
13. Buckwalter JA. Activity vs. rest in the treatment of bone, soft tissue and joint injuries. *Iowa Orthop J* 1995;15: 29-42
14. Buckwalter JA. Pharmacological treatment of soft-tissue injuries. *J Bone Joint Surg Am* 1995;77: 1902-1914
15. Buckwalter JA. Should bone, soft-tissue, and joint injuries be treated with rest or activity? [editorial]. *J Orthop Res* 1995;13: 155-156
16. Buckwalter JA, Woo SL. Drug treatment of soft tissue injuries efficacy and tissue effects. *Iowa Orthop J* 1993;13: 40-48
17. Bulger EM, Maier RV. Antioxidants in critical illness. *Arch Surg* 2001;136: 1201-1207
18. Byrd HS, Cierny G, Tebbetts JB. The management of open tibial fractures with associated soft tissue loss: external pin fixation with early flap coverage. *Plast Reconstr Surg* 1981;68: 73-79
19. Carden DL, Korthuis RJ. Mechanisms of postischemic vascular dysfunction in skeletal muscle: implications for therapeutic intervention. *Microcirc Endothelium Lymphatics* 1989;5: 277-298
20. Carden DL, Smith KJ, Korthuis RJ. Neutrophil-mediated microvascular dysfunction in postischemic canine skeletal muscle: Role of granulocyte adherence. *Circulation Research* 1990;66: 1436-1444
21. Carlos TM, Harlan JM. Leukocyte-endothelial adhesion molecules. *Blood* 1994;84: 2068-2101
22. Cernak I, O'Connor C, Vink R. Inhibition of cyclooxygenase-1 by nimesulfide improves cognitive outcome more than motor outcomes following diffuse traumatic brain injury in rats. *Exp Brain Res* 2002;147: 193-199
23. Chen LE, Seaber AV, Nasser RM, Stamler JS, Urbaniak JR. Effects of S-nitroso-N-acetylcysteine on contractile function of reperfused skeletal muscle. *Am J Physiol* 1998;274: R822-829
24. Chiba Y, Muraoka R, Ihaya A, Morioka K, Sasaki M, Uesaka T. Leukocyte depletion and prevention of reperfusion injury during cardiopulmonary bypass: a clinical study. *Cardiovasc Surg* 1993;1: 350-356

25. Chien S. Role of sympathetic nervous system in hemorrhage. *Physiol Rev* 1967;47: 214-228
26. Claudi BF, Oedekoven G. "Biologische Osteosynthesen". *Chirurg* 1991;62: 367-377
27. Clayton JM, Hayes AC, Barnes RW. Tissue pressure and perfusion in the compartment syndrome. *J Surg Res* 1977;22: 333-339
28. Crisco JJ, Jokl P, Heinen GT, Connell MD, Panjabi MM. A muscle contusion injury model. Biomechanics, physiology, and histology. *Am J Sports Med* 1994;22: 702-710
29. Cuzzocrea S, Mazzon E, Costantino G, Serraino I, De Sarro A, Caputi AP. Effects of n-acetylcysteine in a rat model of ischemia and reperfusion injury. *Cardiovasc Res* 2000;47: 537-548
30. Cuzzocrea S, Mazzon E, Dugo L, Serraino I, Ciccolo A, Centorrino T, De Sarro A, Caputi AP. Protective effects of n-acetylcysteine on lung injury and red blood cell modification induced by carrageenan in the rat. *FASEB J* 2001;15: 1187-1200
31. Deal DN, Tipton J, Rosencrance E, Curl WW, Smith TL. Ice reduces edema. *J Bone Joint Surg (Am)* 2002;84-A: 1573-1578
32. Dervin GF. Skeletal fixation of grade IIIB tibial fractures. The potential of metaanalysis. *Clin Orthop* 1996;332: 10-15
33. Dixon CE, Clifton GL, Lighthall JW, Yaghmai AA, Hayes RL. A controlled cortical impact model of traumatic brain injury in the rat. *J Neurosci Methods* 1991;39: 253-262
34. Dorow C, Markgraf E. Therapy of soft tissue injuries--biological strategies. *Zentralbl Chir* 1997;122: 962-969
35. Dunne JB, Davenport M, Williams R, Tredger JM. Evidence that S-adenosylmethionine and N-acetylcysteine reduce injury from sequential cold and warm ischemia in the isolated perfused rat liver. *Transplantation* 1994;57: 1161-1168
36. Echtermeier V. Kompartmentsyndrom. Prinzipien der Therapie. *Unfallchirurg* 1991;94: 225-230
37. Einhorn TA. The cell and molecular biology of fracture healing. *Clin Orthop* 1998;355 (Suppl.): S7-21

38. Elder CL, Dahners LE, Weinhold PS. A cyclooxygenase-2 inhibitor impairs ligament healing in the rat. *Am J Sports Med* 2001;29: 801-805.
39. Esterhai JL, Jr., Queenan J. Management of soft tissue wounds associated with type III open fractures. *Orthop Clin North Am* 1991;22: 427-432!
40. Fisher BD, Baracos VE, Reid DC. Effect of systemic inhibition of prostaglandin synthesis on muscle protein balance after trauma in the rat. *Can J Physiol Pharmacol* 1991;69: 831-836
41. Flaherty JT, Weisfeldt ML. Reperfusion Injury. *Free Radical Biol Med* 1988;5: 409-419
42. Freitelson J, Kulenovic E, Beck DJ. Endogenous norepinephrine regulates blood flow to the intact rat tibia. *J Orthop Res* 2002;20: 391-396
43. Fukuzawa K, Emre S, Senyuz O, Acarli K, Schwartz ME, Miller CM. N-acetylcysteine ameliorates reperfusion injury after warm hepatic ischemia. *Transplantation* 1995;59: 6-9
44. Gautier E, Ganz R. Die biologische Plattenosteosynthese. *Zentralbl Chir* 1994;119: 564-572
45. Gierer P, Mittlmeier T, Bordel R, Schaser KD, Gradi G, Vollmar B. Selective cyclooxygenase-2 inhibition reverses microcirculatory sequelae of closed soft tissue trauma in an animal model. *J Bone Joint Surg [Am]* 2005;87: 153-160
46. Gierer P, Vollmar B, Schaser KD, Andreas C, Gradi G, Mittlmeier T. Efficiency of small-volume resuscitation in restoration of disturbed skeletal muscle microcirculation after soft tissue trauma and haemorrhagic shock. *Langenbecks Arch Surg* 2004;389: 40-45
47. Goodman JC, Cherian L, Bryan RM, Jr., Robertson CS, Cherian L, Robertson CS, Contant CF, Jr., Bryan RM, Jr. Lateral cortical impact injury in rats: pathologic effects of varying cortical compression and impact velocity. Lateral cortical impact injury in rats: cerebrovascular effects of varying depth of cortical deformation and impact velocity. *J Neurotrauma* 1994;11: 587-597
48. Granger DN, Kubes P. The microcirculation and inflammation: modulation of leukocyte-endothelial cell adhesion. *J Leukoc Biol* 1994;55: 662-675
49. Gullo A, Berlot G. Ingredients of organ dysfunction or failure. *World J Surg* 1997;20: 430-436

50. Gustilo RB, Anderson JT. Prevention of infection in the treatment of 1025 open fractures of long bones: retrospective and prospective analyses. *J Bone Joint Surg [Am]* 1984;58: 453-458
51. Gustilo RB, Mendoza RM. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma* 1984;24: 742-746
52. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures. *J Trauma* 1984;24: 742-746
53. Gustilo RB, Merkow RL, Templeman D. The management of open fractures. *J Bone Joint Surgery [Am]* 1990;72: 299-304
54. Haas NP, Schütz M, Mauch C, Hoffmann R, Sudkamp NP. Management of ipsilateral fractures of the femur shaft and proximal femur--therapy overview and current management. *Zentralbl Chir* 1995;120: 856-861
55. Hansen M, Voltmer W, Rommens PM. [The extraarticular proximal tibial fractures]. *Unfallchirurg* 2002;105: 858-872.
56. Hansen ST. Soft-tissue injuries in the lower extremity. In: Orthopedic trauma protocols. 1993
57. Harris AG, Skalak TC. Effects of leukocyte capillary plugging in skeletal muscle ischemia-reperfusion injury. *Am J Physiol* 1996;271: H2653-2660
58. Harrison PM, Wendon JA, Gimson AE, Alexander GJ, Williams R. Improvement by acetylcysteine of hemodynamics and oxygen transport in fulminant hepatic failure. *N Engl J Med* 1991;324: 1852-1857
59. Hayes AA, Bower GD, Pitstock KL. Chronic (exertional) compartment syndrome of the legs diagnosed with thallous chloride scintigraphy. *J Nucl Med* 1995;36: 1618-1624.
60. Henley MB, Chapman JR, Agel J, Harvey EJ, Whorton AM, Swiontkowski MF. Treatment of II, IIIA, and IIIB open fractures of the tibial shaft: a prospective comparison of unreamed interlocking intramedullary nails and half-pin fixators. *J Orthop Trauma* 1998;12: 1-7
61. Heppenstall RB. Compartment Syndrome: Pathophysiology, Diagnosis, and Treatment. *Techniques in Orthopaedics* 1997;12: 92-108
62. Hietaniemi K Peltonen J, Paavolainen P. An experimental model for producing non-unions in rat. *Injury* 1995;26: 681-686

63. Hoffmann JE, Schmidmaier G, Duda G, Raschke M. Ein neues standardisiertes Frakturmodell an der Ratte. In: Hrsg. P. Hertel. Hefte zur Zeitschrift "Der Unfallchirurg." Berlin, Heidelberg, New York: Springer Verlag 1999
64. Ince C Coremans J, Bruining HA. In vivo NADH Autofluorescence. *Adv Exp Med Biol* 1992;317: 277-296
65. Ito Y, Katagiri H, Ishii K, Kakita A, Hayashi I, Majima M. Effects of selective cyclooxygenase inhibitors on ischemia/reperfusion-induced hepatic microcirculatory dysfunction in mice. *Eur Surg Res* 2003;35: 408-416
66. Janssen GH, Tangelder GJ, Oude Egbrink MG, Reneman RS. Spontaneous leukocyte rolling in venules in untraumatized skin of conscious and anesthetized animals. *Am J Physiol* 1994;267: H1199-1204
67. Jarvinen M, Sorvari T. Healing of a crush injury in rat striated muscle. 1. Description and testing of a new method of inducing a standard injury to the calf muscles. *Acta Pathol Microbiol Scand A* 1975;83: 259-265
68. Jensen T, Kharazmi A, Schiotz PO, Nielsen H, Stenvang Pedersen S, Stafanger G, Koch C, Hoiby N. Effect of oral N-acetylcysteine administration on human blood neutrophil and monocyte function. *APMIS* 1988;96: 62-67
69. Johansen KJ, Daines M, Howey T. Objective criteria accurately predict amputation following lower extremity trauma. *J Trauma* 1990;30: 568
70. Joussen AM, Poulaki V, Mitsiades N, Kirchhof B, Koizumi K, Dohmen S, Adamis AP. Nonsteroidal anti-inflammatory drugs prevent early diabetic retinopathy via TNF-alpha suppression. *FASEB J* 2002;16: 438-440
71. Kellett J. Acute soft tissue injuries - a review of the literature. *Med Sci Sports Exerc* 1986;18: 489-500
72. Kharazmi A, Nielsen H, Schiotz PO. N-acetylcysteine inhibits human neutrophil and monocyte chemotaxis and oxidative metabolism. *Int J Immunopharmacol* 1988;10: 39-46
73. Kiaer T, Kristensen KD. Intracompartmental pressure, PO₂, PCO₂ and blood flow in the human skeletal muscle. *Arch Orthop Trauma Surg* 1988;107: 114-116
74. Kirkpatrick CJ, Bittinger F, Klein CL, Hauptmann S, Klosterhalfen, B. The role of the microcirculation in multiple organ dysfunction syndrome (MODS): a review and perspective. *Virchows Arch* 1996;427: 461-476

75. Kobczik L, Reid M, Bredt DS, Stamler JS. Nitric oxide in skeletal muscle. *Nature* 1994;372: 546-548
76. Korthuis RJ, Grisham MB, Granger DN. Leukocyte depletion attenuates vascular injury in postischemic skeletal muscle. *Am J Physiol* 1988;254: H823-827
77. Köstler W Strohm P, Südkamp NP. Acute compartment syndrome of the limb. *Injury* 2004;35: 1221-1227.
78. Kowalski MJ, Schemitsch EH, Kregor PJ, Senft D, Swiontkowski MF. Effect of periosteal stripping on cortical bone perfusion: A laser-Doppler study in sheep. *Calcif Tissue Int* 1996;59: 24-26
79. Kramer GC, Wade CE, Prough DS. Hypertonic saline dextran: efficacy and regulatory approval [editorial]. *Acta Anaesthesiol Scand* 1998;42: 141-144
80. Kregor P, Senft D, Parvin D. Cortical bone perfusion in plated fractured sheep tibiae. *J Orthop Res* 1995;13: 715-724
81. Kreimeier U, Christ F, Frey L, Habler O, Thiel M, Welte M, Zwissler B, Peter K. Small-volume resuscitation for hypovolemic shock. Concept, experimental and clinical results. *Anaesthetist* 1997;46: 309-328
82. Kreimeier U, Messmer K. Hemodilution in clinical surgery: state of the art 1996. *World J Surg* 1996;20: 1208-1217
83. Krettek C. Fracture and soft tissue damage. *Chirurg* 1998;69: 684-700
84. Kubes P, Suzuki M, Granger D. Nitric oxide: an endogenous modulator of leukocyte adhesion. *Proc Natl Acad Sci USA* 1991;88: 4651-4655
85. Kunkel EJ, Jung U, Bullard DC. Absence of trauma-induced leukocyte rolling in mice deficient for both P-selectin and intercellular adhesion molecule. *J Exp Med* 1996;183: 57-65
86. Kunkel EJ, Ley K. Distinct phenotype of E-selectin-deficient mice. E-selectin is required for slow leukocyte rolling in vivo. *Circ Res* 1996;79: 1196-1204
87. Kuziel WA, Morgan SJ, Dawson TC, Griffin S, Smithies O, Ley K, Maeda N. Severe reduction in leukocyte adhesion and monocyte extravasation in mice deficient in CC chemokine receptor 2. *Proc Natl Acad Sci USA* 1997;94: 12053-12058
88. Landry PS, Marino AA, Sadasivan KK, Albright JA. Effect of soft-tissue trauma on the early periosteal response of bone to injury. *J Trauma* 2000;48: 479-483

89. Lefer AM. Attenuation of myocardial ischemia-reperfusion injury with nitric oxide replacement therapy. *Ann Thorac Surg* 1995;60: 847-851
90. Lefer AM, Lefer DJ. The role of nitric oxide and cell adhesion molecules on the microcirculation in ischaemia-reperfusion. *Cardiovasc Res* 1996;32: 743-751
91. Lefer AM, Ma XL. Cytokines and growth factors in endothelial dysfunction. *Crit Care Med* 1993;21: S9-14
92. Lefer AM, Ma XL, Weyrich A, Lefer DJ. Endothelial dysfunction and neutrophil adherence as critical events in the development of reperfusion injury. *Agents Actions Suppl* 1993;41: 127-135
93. Lehr HA, Leunig M, Menger MD, Nolte D, Messmer K. Dorsal skinfold chamber technique for intravital microscopy in nude mice. *Am J Pathol* 1993;143: 1055-1062
94. Leunig M, Yuan F, Gerweck LE, Berk DA, Jain RK. Quantitative analysis of angiogenesis and growth of bone: effect of indometacin exposure in a combined in vitro-in vivo approach. *Res Exp Med* 1995;195: 275-288
95. LeVan TAL, Levin SL. Principles of soft tissue handling. *Techniques in Orthopedics* 1995;10: 94-104
96. Levin LS, Condit DP. Combined injuries--soft tissue management. *Clin Orthop* 1996;327: 172-181
97. Levin SL. Debridement. *Techniques in Orthopedics* 1995;10: 104-108
98. Levin SL. Personality of soft tissue injury. *Techniques in Orthopedics* 1995;10: 65-73
99. Levin SL. The reconstructive ladder. *Techniques in Orthopedics* 1995;10: 88-94
100. Lighthall JW, Dixon CE, Anderson TE. Experimental models of brain injury. *J Neurotrauma* 1989;6: 83-97
101. Liu K, Chen LE, Seaber AV, Urbaniak JR. S-nitroso-N-acetylcysteine protects skeletal muscle against reperfusion injury. *Microsurgery* 1998;18: 299-305
102. Luh EH, Shackford SR, Shatos MA, Pietropaoli JA. The effects of hyperosmolarity on the viability and function of endothelial cells. *J Surg Res* 1996;60: 122-128

103. Maningas PA, DeGuzman LR, Tillman FJ, Hinson CS, Priegnitz KJ, Volk KA, Bellamy RF. Small-volume infusion of 7.5% NaCl in 6% Dextran 70 for the treatment of severe hemorrhagic shock in swine. *Ann Emerg Med* 1986;15: 1131-1137
104. Mazzoni MC, Borgstrom P, Arfors KE, Intaglietta M. Dynamic fluid redistribution in hyperosmotic resuscitation of hypovolemic hemorrhage. *Am J Physiol* 1988;255: H629-637
105. Mazzoni MC, Borgstrom P, Intaglietta M, Arfors KE. Capillary narrowing in hemorrhagic shock is rectified by hyperosmotic saline-dextran reinfusion. *Circ Shock* 1990;31: 407-418
106. Mazzoni MC, Borgstrom P, Warnke KC, Skalak TC, Intaglietta M, Arfors KE. Mechanisms and implications of capillary endothelial swelling and luminal narrowing in low-flow ischemias. *Int J Microcirc Clin Exp* 1995;15: 265-270
107. McCord JM. Oxygen-derived free radicals in postischemic tissue injury. *N Engl J Med* 1985;312: 159-163
108. McGeachie JK, Grounds MD. Initiation and duration of muscle precursor replication after mild and severe injury to skeletal muscle of mice. An autoradiographic study. *Cell Tissue Res* 1987;248: 125-130
109. Menger M, Vollmar B. Adhesion molecules as determinants of disease: from molecular biology to surgical research. *Br J Surg* 1996;83: 588-601
110. Menger MD. Microcirculatory disturbances secondary to ischemia-reperfusion. *Transplant Proc* 1995;27: 2863-2865
111. Menger MD, Kerger H, Geisweid A, Leu AJ, Hecht R, Nolte D, Messmer K. Leukocyte-endothelium interaction in the microvasculature of postischemic striated muscle. *Adv Exp Med Biol* 1994;361: 541-545
112. Menger MD, Lehr HA. Scope and perspectives of intravital microscopy--bridge over from in vitro to in vivo. *Immunol Today* 1993;14: 519-522
113. Menger MD, Pelikan S, Steiner D, Messmer K. Microvascular ischemia-reperfusion injury in striated muscle: significance of "reflow paradox". *Am J Physiol* 1992;263: H1901-1906
114. Menger MD, Rucker M, Vollmar B. Capillary dysfunction in striated muscle ischemia/reperfusion: on the mechanisms of capillary "no-reflow". *Shock* 1997;8: 2-7

115. Menger MD, Steiner D, Messmer K. Microvascular ischemia-reperfusion injury in striated muscle: significance of "no reflow". Am J Physiol 1992;263: H1892-1900
116. Menger MD, Thierjung C, Hammersen F, Messmer K. Dextran vs. hydroxyethylstarch in inhibition of postischemic leukocyte adherence in striated muscle. Circ Shock 1993;41: 248-255
117. Menger MD, Vollmar B. In vivo analysis of microvascular reperfusion injury in striated muscle and skin. Microsurgery 1994;15: 383-389
118. Menth-Chiari WA, Curl, WW, Rosenrance E, Smith TL. Contusion of skeletal muscle increases leukocyte-endothelial cell interactions: an intravital microscopic study in rats. J Trauma 1998;45: 709-714
119. Mercer NS, Moss AL. Soft tissue management of compound leg fractures: a 5-year experience. J R Coll Surg Edinb 1988;33: 263-266
120. Miles W. Soft tissue trauma. Hand Clin 1986;2: 33-43
121. Mitchell D, Tyml K. Nitric oxide release in rat skeletal muscle capillary. Am J Physiol 1996;270: H1696-1703
122. Mittlmeier T Khodadadyan C, Haas NP. Grundlagen der Frakturbehandlung. In: Mutschler W, Haas N (Hrsg.) Praxis der Unfallchirurgie 1998;55-98
123. Mittlmeier T, Schaser KD, Kroppenstedt S, Raschke M, Peine R, Menger M, Vollmar B, Haas N. Microvascular response to closed soft tissue injury. Trans Orthop Res Soc 1997;44: 317
124. Mittlmeier T Vollmar B, Menger MD, Schewior L, Raschke M, Bail HJ, Haas NP, Schaser KD. Small volume resuscitation reduces microvascular dysfunction following closed soft tissue trauma. Trans Orthop Res Soc 1999;46: 257
125. Mittlmeier T Vollmar B, Raschke M, Menger MD, Schewior L, Schaser KD. Small volume hypertonic hydroxyethylstarch reduces acute microvascular dysfunction following closed soft tissue trauma. J Bone Joint Surg [Br] 2002;85-B: 126-132
126. Morgan SJ, Moore MW, Cacalano G, Ley K. Reduced leukocyte adhesion response and absence of slow leukocyte rolling in interleukin-8 receptor-deficient mice. Microvasc Res 1997;54: 188-191
127. Morisaki H, Suematsu M, Wakabayashi Y, Moro Oka S, Fukushima K, Ishimura Y, Takeda J. Leukocyte-endothelium interaction in the rat

- mesenteric microcirculation during halothane or sevoflurane anesthesia. Anesthesiology 1997;87: 591-598
128. Myers PR, Minor RL, Jr., Guerra R, Jr., Bates JN, Harrison DG. Vasorelaxant properties of the endothelium-derived relaxing factor more closely resemble S-nitrosocysteine than nitric oxide. Nature 1990;345: 161-163
129. Nakayama M, Uchimura K, Zhu R, Nagayama T, Rose ME, Stetler RA, Isakson PC, Chen J, Graham SH. Cyclooxygenase-2 inhibition prevents delayed death of CA1 hippocampal neurons following global ischemia. Proc Natl Acad Sci 1998;95: 10954-10959
130. Nishina K, Akamatsu H, Mikawa K, Shiga M, Maekawa N, Obara H, Niwa Y. The inhibitory effects of thiopental, midazolam, and ketamine on human neutrophil functions. Anesth Analg 1998;86: 159-165
131. Nolte D, Bayer M, Lehr HA, Becker M, Krombach F, Kreimeier U, Messmer K. Attenuation of postischemic microvascular disturbances in striated muscle by hyperosmolar saline dextran. Am J Physiol 1992;263: H1411-1416
132. Nolte D, Menger MD, Messmer K. Microcirculatory models of ischaemia-reperfusion in skin and striated muscle. Int J Microcirc Clin Exp 1995;15 Suppl 1: 9-16
133. Noveck RJ, Laurent A, Kuss M, Talwalker S. Parecoxib sodium does not impair platelet function in healthy elderly and non-elderly individuals. Clin Drug Invest 2001;21: 465-476
134. Oestern HJ, Tscherne H. Pathophysiology and classification of soft tissue damage in fractures. Orthopade 1983;12: 2-8
135. Osborn L. Leukocyte adhesion to endothelium in inflammation. Cell 1990;62: 3-6
136. Owens S, Edwards P, Miles K, Jenner J, Allen M. Chronic compartment syndrome affecting the lower limb: MIBI perfusion imaging as an alternative to pressure monitoring: two case reports. Br J Sports Med 1999;33: 49-51.
137. Pascual JL, Ferri L, Seely AJ, Campisi G, Chaudhury P, Giannias B, Evans DC, Razek T, Michel RP, Christou NV. Hypertonic saline resuscitation of hemorrhagic shock diminishes neutrophil rolling and adherence to endothelium and reduces in vivo vascular leakage. Ann Surg 2002;236: 634-642

138. Pemberton M, Anderson G, Barker J. In vivo microscopy of microcirculatory injury in skeletal muscle following ischemia/reperfusion. *Microsurgery* 1994;15 (6): 374-382
139. Perler BA, Tohmeh AG, Bulkley GB. Inhibition of the compartment syndrome by the ablation of free radical-mediated reperfusion injury. *Surgery* 1990;108: 40-47
140. Rhinelander FW, Phillips RS, Steel WM, Beer JC. Microangiography in bone healing. Displaced closed fractures. *J Bone Joint Surg (Am)* 1968; 50: 643-662
141. Rhinelander FW. Some aspects of the microcirculation of healing bone. *Clin Orthop* 1965;400: 12-16
142. Rhinelander FW. Tibial blood supply in relation to fracture healing. *Clin Orthop* 1974;105: 34-81
143. Richards RR, Schemitsch EH. Effect of muscle flap coverage on the repair of devascularized tibia. An experimental investigation in dog. *J Orthop Res* 1989;7: 550-558
144. Rodgers CM, Ketch LL. Management of associated soft-tissue injury. *Surg Clin North Am* 1988;68: 823-835
145. Rose S, Marzi I. Pathophysiology of polytrauma. *Zentralbl Chir* 1996;121: 896-913
146. Rubinstein I, Abassi Z, Coleman R, Winaver J, Better OS. Involvement of nitric oxide system in experimental muscle crush injury. *J Clin Invest* 1998;101: 1325-1333
147. Ruffmann R, Wendel A. GSH rescue by N-acetylcysteine. *Klin Wochenschr* 1991;69: 857-862
148. Salyer JL, Bohnsack JF, Knape WA, Shigeoka AO, Ashwood ER, Hill HR. Mechanisms of tumor necrosis factor-alpha alteration of PMN adhesion and migration. *Am J Pathol* 1990;136: 831-841
149. Schandelmaier P, Stephan C, Krettek C, Tscherne H. Distale Femurfrakturen. *Der Unfallchirurg* 2000;103: 428-436
150. Schaser KD, Vollmar B, Kroppenstedt S, Schewior L, Raschke M, Menger MD, Lübbe A, Haas NP, Mittlmeier T. In vivo analysis of microcirculation following closed soft tissue injury. *J Orthop Res* 1999;17: 678-685

151. Schaser KD, Vollmar B, Menger MD, Schewior L, Haas NP, Duda G, Mittlmeier T. Nitric oxide Donor N-acetylcysteine ameliorates skeletal muscle microcirculation following closed soft tissue injury. *Trans Orthop Res Soc* 1999;324
152. Schaser KD, Zhang L, Haas NP, Mittlmeier T, Duda G, Bail HJ. Temporal profile of microvascular disturbances in rat tibial periosteum following closed soft tissue trauma. *Langenbecks Arch Surg* 2003;388: 323-340
153. Schaser KD, Bail H, Schewior L, Stover JF, Melcher I, Haas NP, Mittlmeier T. Acute effects of N-acetylcysteine on skeletal muscle microcirculation following closed soft tissue trauma in rats. *J Orthop Res* 2005;23: 231-241
154. Schaser KD, Zhang L, Mittlmeier T, Ostapowicz D, Schmidmaier G, Duda G, Haas NP, Bail HJ. Effect of soft tissue damage on fracture healing: intravitalmicroscopic and biomechanical investigations in rats. *Trans Orthop Res Soc* 2003;0120
155. Schaser KD, Zhang L, Mittlmeier T, Ostapowicz D, Schmidmaier G, Duda G, Haas NP, Bail HJ. Einfluss des traumatischen Weichteilschadens auf die Frakturheilung: intravitalmikroskopische und biomechanische Untersuchungen an der Ratte. *Chirurgisches Forum (Fritz-Lindner Preisträgersitzung)* 2003;32: 9-11
156. Schewior L, Schaser KD, Vollmar B, Duda G, Hoffmann J, Mittlmeier T. Protektion der Mikrozirkulation nach schwerem geschlossenem Weichteiltrauma durch Infusion hyperosmolarer-hyperonkotischer Lösung. *Hefte zur "Der Unfallchirurg"* 1998;272: 678-679
157. Shrier I, Magder S. Pressure-flow relationships in in vitro model of compartment syndrome. *J Appl Physiol* 1995;79: 214-221
158. Silverstein FE, Faich G, Goldstein JL, Simon LS, Pincus T, Whelton A, Makuch R, Eisen G, Agrawal NM, Stenson WF, Burr AM, Zhao WW, Kent JD, Lefkowith JB, Verburg KM, Geis GS. Gastrointestinal toxicity with celecoxib vs nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis: the CLASS study: A randomized controlled trial. *Celecoxib Long-term Arthritis Safety Study.* *JAMA* 2000;284: 1247-1255
159. Skjeldal S, Svindland A, Hvaal K, Kase T, Reikeras O, Nordsletten L. Severe hindlimb ischemia causes periosteal proliferation in the rat tibia. *Acta Orthop Scand* 1997;68: 593-597

160. Smail N, Messiah A, Edouard A, Descorts Declere A, Duranteau J, Vigue B, Mimoz O, Samii K. Role of systemic inflammatory response syndrome and infection in the occurrence of early multiple organ dysfunction syndrome following severe trauma. *Intensive Care Med* 1995;21: 813-816
161. Smith TL, Curl WW, Smith BP, Holden MB, Wise T, Marr A, Koman LA. New skeletal muscle model for the longitudinal study of alterations in microcirculation following contusion and cryotherapy. *Microsurgery* 1993;14: 487-493
162. Song AM, Bhagat L, Singh VP, Van Acker GG, Steer ML, Saluja AK. Inhibition of cyclooxygenase-2 ameliorates the severity of pancreatitis and associated lung injury. *Am J Physiol (Gastrointest Liver Physiol)* 2002;283: G 1166-1174
163. Stamler J, Mendelsohn ME, Amarante P, Smick D, Andon N, Davies PF, Cooke JP, Loscalzo J. N-acetylcysteine potentiates platelet inhibition by endothelium-derived relaxing factor. *Circ Res* 1989;65: 789-795
164. Steinbauer M, Harris AG, Leiderer R, Abels C, Messmer K. Impact of dextran on microvascular disturbances and tissue injury following ischemia/reperfusion in striated muscle. *Shock* 1998;9: 345-351
165. Steinbauer M, Harris AG, Messmer K. Effects of dextran on microvascular ischemia-reperfusion injury in striated muscle. *Am J Physiol* 1997;272: H1710-1716
166. Stichtenoth D, Frolich JC. The second generation of COX-2 inhibitors: what advantages do the newest offer? *Drugs* 2003;63: 33-45
167. Südkamp NP, Barbey N, Veuskens A, Tempka A, Haas NP, Hoffmann R, Tscherne H. The incidence of osteitis in open fracturees: an analysis of 948 open fractures (review of the hannover expierience). *J Orthop Trauma* 1993;7: 473-482
168. Südkamp NP Haas NP, Tempka A, Barbey N, Veuskens A, Kirchhoff A, Tscherne H. Indications and incidence of spongiosa transplantation in open fractures: analysis of 470 open fractures. *Aktuelle Traumatol* 1993;23: 169-177
169. Sun LL, Ruff P, Austin B. Early up-regulation of intercellular adhesion molecule-1 and vascular cell adhesion molecule-1 expression in rats with hemorrhagic shock and resuscitation. *Shock* 1999;11: 416-422

170. Sunose Y, Takeyoshi I, Tsutsumi H, Ohwada S, Oriuchi N, Matsumoto K, Morishita Y. Effect of a cyclooxygenase-2 inhibitor, FK3311, in a canine lung transplantation model. *J Am Coll Surg* 2001;192: 54-62
171. Svindland AD, Nordsletten L, Reikeras O, Skjeldal S. Periostal response to transient ischemia. *Acta Orthop Scand* 1995;66: 468-472
172. Szczesny G, Nolte D, Veihlmann A, Messmer K. A new chamber technique for intravital microscopic observations in the soft tissue layers of mouse hindleg. *J Trauma* 2000;46: 1108-1115
173. Taut FJ, Schmidt H, Zapletal CM, Thies JC, Grube C, Motsch J, Klar E, Martin E. N-acetylcysteine induces shedding of selectins from liver and intestine during orthotopic liver transplantation. *Clin Exp Immunol* 2001;124: 337-341
174. Thiel M, Buessecker F, Chouker A. Effect of hypertonic saline on the expression of adhesion molecules of human polymorphonuclear leukocytes. *Shock* 1997;7 (Suppl): 44 (Abstract)
175. Thorlacius H, Vollmar B, Westermann S, Torkvist L, Menger MD. Effects of local cooling on microvascular hemodynamics and leukocyte adhesion in the striated muscle of hamsters. *J Trauma* 1998;45: 715-719
176. Tornetta P, Bergmann M, Watnik N, Berkowitz G, Steuer J. Treatment of grade IIIb open fractures. A prospective randomised comparison of external fixation and non-reamed nailing. *J Bone Joint Surg [Br]* 1994;76: 13-19
177. Trease L, van Every B, Bennell K, Brukner P, Rynderman J, Baldey A, Turlakow A, Kelly MJ. A prospective blinded evaluation of exercise thallium-201 SPECT in patients with suspected chronic exertional compartment syndrome of the leg. *Eur J Nucl Med* 2001;28: 688-695.
178. Tscherne H, Oestern HJ. [A new classification of soft-tissue damage in open and closed fractures (author's transl)]. *Unfallheilkunde* 1982;85: 111-115
179. Turnipseed WD, Hurschler C, Vanderby R, Jr. The effects of elevated compartment pressure on tibial arteriovenous flow and relationship of mechanical and biochemical characteristics of fascia to genesis of chronic anterior compartment syndrome. *J Vasc Surg* 1995;21: 810-816; discussion 816-817.

180. Tym K, Budreau CH. A new preparation of rat extensor digitorum longus muscle for intravital investigation of the microcirculation. *Int J Microcirc Clin Exp* 1991;10: 335-343
181. Utvag SE, Grundnes O, Rindal DB, Reikeras O. Influence of extensive muscle injury on fracture healing in rat tibia. *J Orthop Trauma* 2003;17: 430-435
182. Utvag SE, Grundnes O, Reikeras O. Early muscle-periosteal lesion inhibits fracture healing in rats. *Acta Orthop Scand* 1999;70: 62-66
183. Utvag SE, Grundnes O, Reikeras O. Effects of lesion between bone, periosteum and muscle on fracture healing in rats. *Acta Orthop Scand* 1998;69: 177-180
184. Utvag SE, Iversen K, Grundnes O, Reikeras O. Poor muscle coverage delays fracture healing in rats. *Acta Orthop Scand* 2002;73: 471-474
185. van der Laan L, Kapitein PJ, Oyen WJ, Verhofstad AA, Hendriks T, Goris RJ. A novel animal model to evaluate oxygen derived free radical damage in soft tissue. *Free Radic Res* 1997;26: 363-372
186. van der Laan L, Oyen WJ, Verhofstad AA, Tan EC, ter Laak HJ, Gabreels Festen A, Hendriks T, Goris RJ. Soft tissue repair capacity after oxygen-derived free radical-induced damage in one hindlimb of the rat. *J Surg Res* 1997;72: 60-69
187. Vane JR, Bakhle YS, Botting RM. Cyclooxygenases 1 and 2. *Annu Rev Pharmacol Toxicol* 1998;38: 97-120
188. Vassar MJ, Perry CA, Holcroft JW. Analysis of potential risks associated with 7.5% sodium chloride resuscitation of traumatic shock [published erratum appears in Arch Surg 1991 Jan;126(1):43]. *Arch Surg* 1990;125: 1309-1315
189. Vollmar B, Lang G, Menger MD, Messmer K. Hypertonic hydroxyethyl starch restores hepatic microvascular perfusion in hemorrhagic shock. *Am J Physiol* 1994;266: H1927-1934
190. Vollmar B, Preissler G, Menger MD. Hemorrhagic hypotension induces arteriolar vasomotion and intermittent capillary perfusion in rat pancreas. *Am J Physiol* 1994;267: H1936-1940
191. Waydhas C, Kanz KG, Ruchholtz S, Nast Kolb D. [Algorithms in trauma management]. *Unfallchirurg* 1997;100: 913-921

192. Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med* 1989;320: 365-376
193. Westermann S, Vollmar B, Menger MD. [In vivo analysis of the microcirculation of striated skin muscle during induction of a compartment syndrome] *Langenbecks Arch Chir Suppl Kongressbd* 1997;114: 485-488
194. Whiteside LA, Lesker, P.A. The effects of extraperiosteal and subperiosteal dissection. I. On blood flow in muscle. II. On fracture healing. *J Bone Joint Surg (Am)* 1978;60: 23-30
195. Wichmann MW, Ayala, A, Chaudry IH. Severe depression of host immune functions following closed-bone fracture, soft tissue trauma, and hemorrhagic shock. *Crit Care Med* 1998;26: 1372-1378
196. Wiley JP, Short WB, Wiseman DA, Miller SD. Ultrasound catheter placement for deep posterior compartment pressure measurements in chronic compartment syndrome. *Am J Sports Med* 1990;18: 74-79.
197. Winet H. The role of microvasculature in normal and perturbed bone healing as revealed by intravital microscopy. *Bone* 1996;19: 39S-57S
198. Wolfard A, Csaszar, J, Gera L, Petri A, Simonka JA., Balogh A, Boros M. Endothelin-a receptor antagonist treatment improves the periosteal microcirculation after hindlimb ischemia and reperfusion in the rat. *Microcirculation* 2002;9: 471-476
199. Zhang L, Bail H, Mittlmeier T, Haas NP, Schaser KD. Immediate microcirculatory derangements in skeletal muscle and periosteum following closed tibial fracture. *J Trauma* 2003;54: 979-985