

6. LITERATURVERZEICHNIS

Akaike, T.: Neuronal organization of the vestibulospinal system in the cat. *Brain Res.*, 259:217-227 (1983)

Altman, J. and Bayer, S.A.: Time of origin and distribution of a new cell type in the rat cerebellar cortex. *Exp. Brain Res.*, 29:265-274 (1977)

Anden, N.E., Fuxe, K. and Ungerstedt, U.: Monoamine pathways to the cerebellum and cerebral cortex. *Experientia*, 23:838-839 (1967)

Andre, P., Fascetti, F., Pompeiano, O. and White, S.R.: The muscarinic agonist, bethanechol, enhances GABA-induced inhibition of Purkinje cells in the cerebellar cortex. *Brain Res.*, 637:1-9 (1994)

Anelli, R. and Mugnaini, E.: Enrichment of unipolar brush cell-like neurons in primary rat cerebellar cultures. *Anat. Embryol. (Berl.)*, 203:283-292 (2001)

Anniko, M., Sjöström, B. and Webster, D.: The effects of auditory deprivation on morphological maturation of the ventral cochlear nucleus. *Arch. Otorhinolaryngol.*, 246:43-47 (1989)

Arnhold, S., Andressen, C., Angelov, D.N., Vajna, R., Volsen, S.G., Hescheler, J. and Addicks, K.: Embryonic stem-cell derived neurones express a maturation dependent pattern of voltage-gated calcium channels and calcium-binding proteins. *Int. J. Dev. Neurosci.*, 18:201-212 (2000)

Babinski, J.: Sur le rôle du cervelet dans les actes volitionnels nécessitant une succession rapide de mouvements. *Rev. Neurol.*, 21:1-3 (1902)

Baird, R.A., Desmadryl, G., Fernández, C. and Goldberg, J.M.: The vestibular nerve of the chinchilla. II. Relation between afferent response properties and peripheral innervation patterns in the semicircular canals. *J. Neurophysiol.*, 60:182-203 (1988)

Bartles, J.R.: Parallel actin bundles and their multiple actin-bundling proteins. *Curr. Opin. Cell Biol.*, 12:72-78 (2000)

Bartles, J.R., Wierda, A. and Zheng, L.: Identification and characterization of espin, an actin-binding protein localized to the F-actin-rich junctional plaques of Sertoli cell ectoplasmic specializations. *J. Cell Sci.*, 109:1229-1239 (1996)

Bartles, J.R., Zheng, L., Li, A., Wierda, A. and Chen, B.: Small espin: a third actin-bundling protein and potential forked protein ortholog in brush border microvilli. *J. Cell Biol.*, 143:107-119 (1998)

Bäurle, J.: Cerebelläre und cortico-nukleäre Kompensation, Untersuchungen an cerebellären Mutanten. *Institut für Physiologie der Freien Universität Berlin* (2001)

Bäurle, J. and Grüsser-Cornehls, U.: Differential number of glycine- and GABA-immunopositive neurons and terminals in the deep cerebellar nuclei of normal and Purkinje cell degeneration mutant mice. *J. Comp. Neurol.*, 382:443-458 (1997)

LITERATURVERZEICHNIS

- Bäurle, J. and Guldin, W.: Unbiased number of ganglion neurons in the mouse. *Neurosci. Lett.*, 246:89-92 (1998a)
- Bäurle, J. and Guldin, W.: Vestibular ganglion neurons survive the loss of their cerebellar targets. *NeuroReport*, 9:4119-4122 (1998b)
- Bäurle, J., Helmchen, C. and Grüsser-Cornehls, U.: Diverse effects of Purkinje cell loss on deep cerebellar and vestibular nuclei neurons in Purkinje cell degeneration mutant mice: a possible compensatory mechanism. *J. Comp. Neurol.*, 384:580-596 (1997)
- Bäurle, J., Hoshi, M. and Grüsser-Cornehls, U.: Dependence of parvalbumin expression on Purkinje cell input in the deep cerebellar nuclei. *J. Comp. Neurol.*, 392:499-514 (1998a)
- Bäurle, J., Kleine, J., Grüsser, O.J. and Guldin, W.: Co-localisation of glycine and calbindin D-28K in the vestibular ganglion of the rat. *NeuroReport*, 8:2443-2447 (1997)
- Bäurle, J., Vogten, H. and Grüsser-Cornehls, U.: Course and targets of the Calbindin D-28k subpopulation of primary vestibular afferents. *J. Comp. Neurol.*, 402:111-128 (1998b)
- Berghaus, A., Rettinger, G. and Böhme, G.: Hals-Nasen-Ohren-Heilkunde, 7.1.2.: Peripher-vestibuläre Erkrankungen. Hippokrates-Verlag, Stuttgart, pp 195-196 (1996)
- Billing-Marczak, K. and Kuźnicki, J.: Calretinin - sensor or buffer - function still unclear. *Pol. J. Pharmacol.*, 51:173-178 (1999)
- Blatt, G.J. and Eisenmann, L.M.: A qualitative and quantitative light microscopic study of the inferior olfactory complex of normal, reeler, and weaver mutant mice. *J. Comp. Neurol.*, 232:117-128 (1985)
- Bouilleret, V., Schwaller, B., Schurmans, S., Celio, M.R. and Fritschy, J.M. : Neurodegenerative and morphogenic changes in a mouse model of temporal lobe epilepsy do not depend on the expression of the calcium-binding proteins parvalbumin, calbindin, or calretinin. *Neuroscience*, 97:47-58 (2000)
- Brichta, A.M. and Peterson, E.H.: Functional architecture of vestibular primary afferents from the posterior semicircular canal of a turtle, *Pseudemys (Trachemys) scripta elegans*. *J. Comp. Neurol.*, 344:481-507 (1994)
- Brodal, A.: Experimentelle Untersuchungen über die olivo-cerebellare Lokalisation. *Z. Neurol. Psychiat.*, 1-153 (1940)
- Brodal, A.: Afferent cerebellar connections. In: J. Jansen and A. Brodal (eds.), Aspects of cerebellar anatomy. *Johan Grundt Tanum*, Oslo (1954)
- Brodal, A.: Anatomy of the vestibular nuclei and their connections. In: Autrum, H., Jung, R., Loewenstein, W.R., MacKay, D.M. and Teuber, H.L. (eds.), *Handbook of Sensory Physiology*. Vol. VI/1, Springer: Berlin, pp 240-352 (1974)
- Brodal, A.: Neurological Anatomy. In Relation to Clinical Medicine, *Oxford University Press*: Oxford, New York, 3rd edition (1981)
- Brodal, A.: The vestibular nuclei in the macaque monkey. *J. Comp. Neurol.*, 227:252-266 (1984)

LITERATURVERZEICHNIS

- Brodal, A. and Pompeiano, O.: The vestibular nuclei in the cat. *J. Anat.*, 91:435-454 (1957)
- Brodal, A., Taber, E. and Walberg, F.: The raphe nuclei of the brainstem in the cat. II. Efferent projections. *J. Comp. Neurol.*, 114:239-260 (1960)
- Brooks, V.B. and Thach, W.T.: Cerebellar control of posture and movement. In: V.B. Brooks (ed.), *Handbook of Physiology*. Section 1: The Nervous System, Vol. II., Motor Control, Part 2, American Physiological Society: Bethesda, Md, pp 877-946 (1981)
- Büttner, U. and Buettner, U.W.: Parietal cortex (2v) neuronal activity in the alert monkey during natural vestibular and optokinetic stimulation. *Brain Res.*, 153:392-397 (1978)
- Büttner, U., Henn V., and Oswald, H.P.: Vestibular-related neuronal activity in the thalamus of the alert monkey during sinusoidal rotation in the dark. *Exp. Brain Res.*, 30:435-444 (1977)
- Büttner-Ennever, J.A.: Patterns of connectivity in the vestibular nuclei. *Ann. N.Y. Acad. Sci.*, 656:363-378 (1992)
- Büttner-Ennever, J.A.: A review of otolith pathway to brainstem and cerebellum. *Ann. N.Y. Acad. Sci.*, 871:51-64 (1999)
- Cable, J. and Steel, K.P.: Combined cochleo-saccular and neuroepithelial abnormalities in the Varitint-waddler-J (Va^J) mouse. *Hear. Res.*, 123:125-136 (1998)
- Caddy, K.W.T. and Biscoe, T.J.: Preliminary observations on the cerebellum in the mutant mouse Lurcher. *Brain Res.*, 91:276-280 (1975)
- Caddy, K.W.T. and Biscoe, T.J.: Structural and quantitative studies on the normal C3H and Lurcher mutant mouse. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 287:167-201 (1979)
- Carpenter, M.B.: Vestibular nuclei: afferent and efferent projections. *Prog. Brain Res.*, 76:5-15 (1988)
- Cass, S.P., Davidson, P. and Goshgarian, H.: Survival of the vestibular nerve after labyrinthectomy in the cat. *Otolaryngol. Head Neck Surg.*, 101:459-465 (1989)
- Caviness, Jr., V.S. and Rakic, P.: Mechanisms of cortical development: a view from mutations in mice. *Ann. Rev. Neurosci.*, 1:297-326 (1978)
- Celio, M.R.: Calbindin D-28K and Parvalbumin in the rat nervous system. *Neuroscience*, 35:375-475 (1990)
- Chan-Palay, V.: *Cerebellar Dentate Nucleus: Organisation, Cytology and Transmitters*. Springer: Berlin (1977)
- Chen, B., Li, A., Wang, D., Wang, M., Zheng, L. and Bartles, J.R.: Espin contains an additional actin-binding site in its N terminus and is major actin-bundling protein of the Sertoli cell-spermatid ectoplasmic specialization junctional plaque. *Mol. Biol. Cell*, 10:4327-4339 (1999)
- Chimento, T.C. and Ross, M.D.: Evidence of a Sensory Processing Unit in the Mammalian Macula. *Ann. N.Y. Acad. Sci.*, 781:196-212 (1996)

LITERATURVERZEICHNIS

- Cloudman, A.M. and Bunker, L.E.: The varitint-waddler mouse. *J. Hered.*, 36:258-263 (1945)
- Coggeshall, R.E. and Lekan, H.A.: Methods for determining numbers of cells and synapses: a case for more uniform standards of review. *J. Comp. Neurol.*, 364:6-15 (1996)
- Cohen, B.: Vestibular and Oculomotor Physiology: Internal Meeting of the Bárány Society, *Ann. N.Y. Acad. Sci.*, 374:1,892 (1981)
- Cools, A.R.: Asymmetrical spotting and direction of circling in the varitint-waddler mouse. *J. Hered.*, 63:161-171 (1972a)
- Cools, A.R.: Neurochemical correlates of the waltzing-shaker syndrome in the Varitint-waddler mouse. *Psychopharmacologia*, 24:384-396 (1972b)
- Cowling, K., Robbins, R.J., Haigh, G.R., Teed, S.K. and Dawson, W.D.: Coat color genetics of Peromyscus: IV. Variable white, a new dominant mutation in the deer mouse. *J. Hered.*, 85:48-52 (1994)
- Crepel, F. and Audinat, E.: Excitatory amino acid receptors of cerebellar Purkinje cell: development and plasticity. *Prog. Biophys. Mol. Biol.*, 55:31-46 (1991)
- De Jager, P.L. and Heintz, N.: The Lurcher mutation and ionotropic glutamate receptors: contributions to programmed neuronal death *in vivo*. *Brain Pathol.*, 8:795-807 (1998)
- Dechesne, C.J., Thomasset, M., Brehier, A. and Sans, A.: Calbindin (CaBP 28 kDa) localization in the peripheral vestibular system of various vertebrates. *Hear. Res.*, 33:273-278 (1988)
- Dechesne, C.J., Winsky, L., Kim, H.N., Goping, G., Vu, T.D., Wenthold, R.J. and Jacobowitz, D.M.: Identification and ultrastructural localization of a calretinin-like calcium binding protein (protein 10) in the guinea pig and rat inner ear. *Brain Res.*, 560:139-148 (1991)
- Demêmes, D., Raymond, J., Atger, P., Grill, C., Winsky, L. and Dechesne, C.J.: Identification of neuron subpopulations in the rat vestibular ganglion by Calbindin-D 28K, calretinin and neurofilament proteins immunoreactivity. *Brain Res.*, 582:168-172 (1992)
- Deol, M.S.: The anomalies of the labyrinth of the mutants varitint-waddler, shaker-2 and jerker in the mouse. *J. Genet.*, 52:562-588 (1954)
- Deol, M.S.: The anatomy and development of the mutants pirouette, shaker-1 and waltzer in the mouse. *Proc. R. Soc. Lond. B. Biol. Sci.*, 145:206-213 (1956)
- Desmadryl, G., Chambard, J.-M., Valmier, J. and Sans, A.: Multiple voltage-dependent calcium currents in acutely isolated mouse vestibular neurons. *Neuroscience*, 78:511-522 (1997)
- Desmadryl, G. and Dechesne, C.J.: Calretinin immunoreactivity in chinchilla and guinea pig vestibular end organs characterizes the calyx unit subpopulation. *Exp. Brain. Res.*, 89:105-108 (1992)
- Diño, M.R., Perachio, A.A. and Mugnaini, E.: Cerebellar unipolar brush cells are targets of primary vestibular afferents: an experimental study in the gerbil. *Exp. Brain Res.*, 140:162-170 (2001)

LITERATURVERZEICHNIS

- Di Palma, F., Belyantseva, I.A., Kim, H.J., Vogt, T.F., Kachar, B. and Noben-Trauth, K.: Mutations in *Mcoln3* associated with deafness and pigmentation defects in varitint-waddler (*Va*) mice. *Proc. Natl. Acad. Sci. USA*, 99:14994-14999 (2002)
- Dow, R.S. and Moruzzi, G.: *The Physiology and Pathology of the Cerebellum*, University of Minnesota Press: Minneapolis (1958)
- Dow, R.S.: The fiber connections of the posterior parts of the cerebellum in the rat and cat. *J. Comp. Neurol.*, 63:527-548 (1936)
- Eatock, R.A., Rüschi, A., Lysakowski, A. and Saeki, M.: Hair cells in mammalian utricles. *Otolaryngol. Head Neck Surg.*, 119, 172-181 (1998)
- Eccles, J.C., Llinás, R. and Sasaki, K.: The excitatory synaptic action of climbing fibres on the Purkinje cells of the cerebellum, *J. Physiol. (Lond.)*, 182:268-296 (1966)
- Edmonds, B., Reyes, R., Schwaller, B. and Roberts, W.M.: Calretinin modifies presynaptic calcium signaling in frog saccular hair cells. *Nat. Neurosci.*, 3:786-790 (2000)
- El-Amraoui, A., Sahly, I., Picaud, S., Sahel, J., Abitbol, M. and Petit, C.: Human Usher IB/mouse *shaker-1*: the retinal phenotype discrepancy explained by the presence/absence of myosin VIIA in the photoreceptor cells. *Hum. Mol. Genet.*, 5:1171-1178 (1996)
- Engström, H., Ades, H.W. and Hawkins, Jr., J.E.: Structure and functions of the sensory hair of the inner ear. *J. Acoust. Soc. Am.*, 34:1356 (1962)
- Ernfors, P., Lee, K.F. and Jaenisch, R.: Mice lacking brain-derived neurotrophic factor develop with sensory deficits. *Nature*, 368:147-150 (1994)
- Fernández, C., Baird, R.A. and Goldberg, J.M.: The vestibular nerve of the chinchilla. I. Peripheral innervation patterns in the horizontal and superior semicircular canals. *J. Neurophysiol.*, 60:167-181 (1988)
- Fernández, C., Goldberg, J.M. and Baird, R.A.: The vestibular nerve of the chinchilla. III. Peripheral innervation patterns utricular macula. *J. Neurophysiol.*, 63:767-790 (1990)
- Fernández, C., Lysakowski, A. and Goldberg, J.M.: Hair-cell counts and afferent innervation patterns in the cristae ampullares of the squirrel monkey with a comparison to the chinchilla. *J. Neurophysiol.*, 73:1253-1269 (1995)
- Fernandez-Gonzalez, A., La Spada, A.R., Treadaway, J., Higdon, J.C., Harris, B.S., Sidman, R.L., Morgan, J.I. and Zuo, J.: *Purkinje cell degeneration (pcd)* phenotypes caused by mutations in the axotomy-induced gene, *Nna1*. *Science*, 295:1904-1906 (2002)
- Floureens, P.: *Recherches expérimentales sur les propriétés et les fonctions du système nerveux dans les animaux vertébrés*. Edition 1. Crevot: Paris. (1824)
- Gabbott, P.L.A., Somogyi, J., Stewart, M.G. and Hámori, J.: GABA-immunoreactive neurons in the rat cerebellum: a light and electron microscope study. *J. Comp. Neurol.*, 251:474-490 (1986)

LITERATURVERZEICHNIS

- Garthwaite, J. and Brodbelt, A.R.: Glutamate as the principal mossy fibre transmitter in rat cerebellum: pharmacological evidence. *Eur. J. Neurosci.*, 2:177-180 (1989)
- German, D.C., Manaye, K.F., Sonsalla, P.K. and Brooks, B.A.: Midbrain dopaminergic cell loss in Parkinson's disease and MPTP-induced parkinsonism: sparing of calbindin-D28K-containing cells. *Ann. N.Y. Acad. Sci.*, 648:42-62 (1992)
- Gerrits, N.M.: Vestibular nuclear complex. In: G. Paxinos (ed.), *The Human Nervous System*. Academic Press: San Diego, New York, Boston, London, Sydney, Tokyo, Toronto, pp 863-888 (1990)
- Ghetti, B., Alyea, C., Hillman, D. and McBride, W.: PCD (Purkinje cell degeneration) mutant mouse: morphometric and neurochemical analyses of the cerebellum after Purkinje cell (Pc) loss. In: *Proceedings of the IXth International Congress of Neuropathology 69*, Egermann: Vienna (1977)
- Ghetti, B., Alyea, C.J. and Muller, J.: Studies on the Purkinje cell degeneration (pcd) mutant: primary; pathology and transneuronal changes. *J. Neuropath. Exp. Neurol.*, 37:617 (1978)
- Ghetti, B., Norton, J. and Triarhou, L.C.: Nerve cell atrophy and loss in the inferior olfactory complex of "Purkinje cell degeneration" mutant mice. *J. Comp. Neurol.*, 260:409-422 (1987)
- Ghetti, B., Truex, L., Sawyer, B., Strada, S. and Schmidt, M.: Exaggerated cyclic AMP accumulation and glial cell reaction in the cerebellum during Purkinje cell degeneration in pcd mutant mice. *J. Neurosci. Res.*, 6:789-801 (1981)
- Gibson, F., Walsh, J., Mburu, P., Varela, A., Brown, K.A., Antonio, M., Beisel, K.W., Steel, K.P. and Brown, S.D.M.: A type VII myosin encoded by the mouse deafness gene shaker-1. *Nature*, 374:62-64 (1995)
- Gilman, S., Bloedel, J.R. and Lechtenberg, R.: Disorders of the cerebellum. Contemporary neurology series. F.A. Davis Company: Philadelphia (1981)
- Goldberg, J.M. and Fernández, C.: Physiology of peripheral neurons innervating semicircular canals of the squirrel monkey. 3. Variations among units in their discharge properties. *J. Neurophysiol.*, 34:676-684 (1971)
- Goldberg, J.M. and Fernández, C.: Conduction times and background discharge of vestibular afferents. *Brain Res.*, 122:545-550 (1977)
- Goldberg, J.M., Lysakowski, A. and Fernández, C.: Morphophysiological and ultrastructural studies in the mammalian cristae ampullares. *Hear. Res.*, 49:89-102 (1990)
- Goldowitz, D.: Interspecific mouse chimeras to study weaver mutant gene action and cell lineage in the CNS. *Soc. Neurosci. Abstr.*, 12:1583 (1986)
- Goldowitz, D. and Mullen, R.J.: Granule cell as a site of gene action in the weaver mouse cerebellum: evidence from heterozygous mutant chimeras. *J. Neurosci.*, 2:1474-1485 (1982)
- Green, M.C.: Genetic variants and strains of the laboratory mouse. Gustav Fischer: Stuttgart (1981)

LITERATURVERZEICHNIS

Grüneberg, H., Burnett, J.B. and Snell, G.D.: The origin of the jerker, a new gene mutation of the house mouse, and linkage studies made with it. *Proc. Natl. Acad. Sci. U.S.A.*, 27:562-565 (1941)

Grüsser-Cornehls, U.: Compensatory mechanisms at the level of the vestibular nuclei following postnatal degeneration of specific cerebellar cell classes and ablation of the cerebellum in mutant mice. In: H. Flohr (ed.), *Postlesion Neural Plasticity*. Springer: Berlin, Heidelberg, pp 431-442 (1988)

Grüsser-Cornehls, U. and Bäurle, J.: Mutant mice as a model for cerebellar ataxia. *Prog. Neurobiol.*, 63:489-540 (2001)

Guldin, W.O. and Grüsser, O.J.: Is there a vestibular cortex? *Trends Neurosci.*, 21:254-259 (1998)

Gundersen, H.J.G.: Stereology of arbitrary particles. *J. Microsc.*, 143:3-45 (1986)

Gundersen, H.J.G., Bagger, P., Bendtsen, T.F., Evans, S.M., Korbo, L., Marcussen, N., Møller, A., Nielsen, K., Nyengaard, J.R., Pakkenberg, B., Sørensen, F.B., Vesterby, A. and West, M.J.: The new stereological tools: disector, fractionator, nucleator and point sampled intercepts and their use in pathological research and diagnosis. *APMIS*, 96:857-881 (1988b)

Gundersen, H.J.G., Bendtsen, T.F., Korbo, L., Marcussen, N., Møller, A., Nielsen, K., Nyengaard, J.R., Pakkenberg, B., Sørensen, F.B., Vesterby, A. and West, M.J.: Some new, simple and efficient stereological methods and their use in pathological research and diagnosis. *APMIS*, 96:379-394 (1988a)

Hallgren, B.: Retinitis pigmentosa combined with congenital deafness, with vestibulo-cerebellar ataxia and mental abnormality in a proportion of cases: a clinical and genetic statistic study. *Acta Psychiat. Scand.*, 34:5-101 (1959)

Hammer, J.A. III.: The structure and function of unconventional myosins: a review. *J. Muscle Res. Cell Motil.*, 15:1-10 (1994)

Harding, A.E.: Friedreich's ataxia: a clinical and genetic study of 90 families with an analysis of early diagnostic criteria and intrafamilial clustering of clinical features. *Brain*, 104:589-620 (1981)

Hasson, T. and Mooseker, M.S.: Molecular motors, membrane movements and physiology: emerging roles for myosins. *Curr. Opin. Cell Biol.*, 7:587-594 (1995)

Hatten, M.E., Liem, R.K.H. and Mason, C.A.: Weaver mouse cerebellar granule cells fail to migrate on wild-type astroglial processes in vitro. *J. Neurosci.*, 6:2676-2683 (1986)

Hauglie-Hanssen, E.: Intrinsic neuronal organization of the vestibular nuclear complex in the cat: a Golgi study. *Ergeb. Anat. Entw. Gesch.*, 40:1-15 (1968)

Heizmann, C.W.: Calcium signaling in the brain. *Acta Neurobiol. Exp. (Warsz.)*, 53:15-23 (1993)

Henn, V.: The correlation between motion sensation, nystagmus and activity in the vestibular nerve and nuclei. In: V. Honrubia, M.A.B. Brazier (eds.), *Nystagmus and Vertigo: Clinical Approaches to the Patient with Dizziness*. Academic Press: New York, pp 115-124 (1982)

Henn, V.: E. Mach on the analysis of motion sensation. *Human Neurobiol.*, 3:145-148 (1984)

LITERATURVERZEICHNIS

- Henn, V.: Motion Sense. In: G. Adelmann (ed.), *Encyclopedia of Neuroscience*. Birkhäuser: Boston, Basel, Stuttgart, pp 680-681 (1987)
- Herrup, K. and Trenkner, E.: Regional differences in cytoarchitecture of the weaver cerebellum suggest a new model for weaver gene action. *Neuroscience*, 23:871-885 (1987)
- Highstein, S.M. and McCrea, R.A.: The anatomy of the vestibular nuclei. *Rev. Oculomot. Res.*, 2:177-202 (1988)
- Hökfelt, T. and Fuxe, K.: Cerebellar monoamine nerve terminals, a new type of afferent fibres to the cortex cerebelli. *Exp. Brain Res.*, 9:63-72 (1969)
- Holmes, G.: The symptoms of acute cerebellar injuries due to gunshot injuries. *Brain*, 40:461-535 (1917)
- Holmes, G.: The clinical symptoms of cerebellar disease and their interpretation. The Croonian Lectures I. *Lancet*, 1:1177-1182 (1922a)
- Holmes, G.: The clinical symptoms of cerebellar disease and their interpretation. The Croonian Lectures II. *Lancet*, 1:1231-1237 (1922b)
- Holmes, G.: The clinical symptoms of cerebellar disease and their interpretation. The Croonian Lectures III. *Lancet*, 2:59-65 (1922c)
- Holmes, G.: The clinical symptoms of cerebellar disease and their interpretation. The Croonian Lectures IV. *Lancet*, 2:111-115 (1922d)
- Holmes, G.: The cerebellum of man. *Brain*, 62:1-30 (1939)
- Hopf, H.C., Deuschl, G., Diener, H.C. and Reichmann, H.: Schwindel: Ein multisensorisches Syndrom, In: Neurologie in Praxis und Klinik, Band I, 3. Auflage, Thieme: Stuttgart, New York, pp 285-311 (1999)
- Hubbard, M.J. and McHugh, N.J.: Calbindin28kDa and calbindin30kDa (calretinin) are substantially localised in the particulate fraction of rat brain. *FEBS Lett.*, 374:333-337 (1995)
- Inanobe, A., Yoshimoto, Y., Horio, Y., Morishige, K.I., Hibino, H., Matsumoto, S., Tokunaga, Y., Maeda, T., Hata, Y., Takai, Y. and Kurachi, Y.: Characterization of G-protein-gated K⁺ channels composed of Kir3.2 subunits in dopaminergic neurons of the substantia nigra. *J. Neurosci.*, 19:1006-1017 (1999)
- Ito, M.: The Cerebellum and Neural Control. Raven Press: New York (1984)
- Ito, M. and Yoshida, M.: The cerebellar-evoked monosynaptic inhibition of Deiters' neurones. *Experientia* (Basel), 20:515-516 (1964)
- Ito, M. and Yoshida, M.: The origin of cerebellar-induced inhibition of Deiters' neurones. I. Monosynaptic initiation of the inhibitory postsynaptic potentials. *Exp. Brain. Res.*, 2:330-349 (1965)
- Ivry, R.B. and Keele, S.W.: Timing functions of the cerebellum. *J. Cogn. Neurosci.*, 1:136-152 (1989)

LITERATURVERZEICHNIS

Jansen, J. and Brodal, A.: Experimental studies on the intrinsic fibers of the cerebellum. II. The cortico-nuclear projection. *J. Comp. Neurol.*, 73:267-321 (1940)

Jansen, J. and Brodal, A.: Experimental studies on the intrinsic fibers of the cerebellum. III. The cortico-nuclear projection in the rabbit and the monkey. *Norske Vid. Akad., Avh. I, Math. Nat. Kl.*, 3:1-50 (1942)

Jarolimek, W., Bäurle, J. and Misgeld, U.: Pore mutation in a G protein-gated inwardly rectifying K⁺ channel subunit causes loss of K⁺-dependent inhibition in *weaver* hippocampus. *J. Neurosci.*, 18:4001-4007 (1998)

Jarolimek, W., Bäurle, J. and Misgeld, U.: Impaired inhibition of epileptiform activity by baclofen, but not by adenosine in the *weaver* hippocampus. *Neuropharmacology*, 39:246-253 (2000)

Jensen, P., Surmeier, J. and Goldowitz, D.: Rescue of cerebellar granule cells from death in *weaver* NR1 double mutants. *J. Neurosci.*, 19:7991-7998 (1999)

Kanowski, S., Fischhof, P., Hiersemzel, R., Röhmel, J. and Kern, U.: Wirksamkeitsnachweis von Nootropika am Beispiel von Nimodipin - ein Beitrag zur Entwicklung geeigneter klinischer Prüfmodelle. *Z. Gerontopsych. Psychiat.*, 1:35-44 (1988)

Karschin, C., Dißmann, E., Stühmer, W. and Karschin, A.: GIRK(1-3) and GIRK(1-4) inwardly rectifying K⁺ channel mRNAs are differentially expressed in the adult rat brain. *J. Neurosci.*, 16:3559-3570 (1996)

Keller, A.D., Roy, R.S. and Chase, W.P.: Exstirpation of the neocerebellar cortex without eliciting so-called cerebellar signs. *Am. J. Physiol.*, 118:720-733 (1937)

Kevetter, G.A. and Leonard, R.B.: Calretinin stains the most irregular firing vestibular primary afferents. *Proc. Bárány Soc. Meet.*, S7/PHY19 (1996)

Kevetter, G.A. and Leonard, R.B.: Molecular probes of the vestibular nerve. II. Characterization of neurons in Scarpa's ganglion to determine separate populations within the nerve. *Brain Res.*, 928:18-29 (2002)

Kevetter, G.A., Shoher, A.D., and Leonard, R.B.: Calyx- and bouton-only afferents comprise forty per cent of the vestibular nerve in gerbil. *Soc. Neurosci. Abstract*, 24:650.14 (1998)

Kikuchi, K. and Hilding, D.: The development of the organ of Corti in the mouse. *Acta Otolaryngol.*, 60:207-222 (1965)

Kobayashi, T., Ikeda, K., Ichikawa, T., Abe, S., Togashi, S. and Kumanishi, T.: Molecular cloning of a mouse G-protein-activated K⁺ channel (mGIRK1) and distinct distributions of three GIRK (GIRK1, 2 and 3) mRNAs in mouse brain. *Biochem. Biophys. Res. Commun.*, 208:1166-1173 (1995)

Kölliker, A.: Der feinere Bau des verlängerten Markes. *Anat. Anz.*, 6:427-431 (1891)

Kotchabhakdi, N. and Walberg, F.: Cerebellar afferent projections from the vestibular nuclei in the cat: an experimental study with the method of retrograde axonal transport of horseradish peroxidase. *Exp. Brain Res.*, 31:591-604 (1978)

LITERATURVERZEICHNIS

Kuźnicki, J., Isaacs, K.R. and Jacobowitz, D.M.: The expression of calretinin in transfected PC12 cells provides no protection against Ca(2+)-overload or trophic factor deprivation. *Biochim. Biophys. Acta*, 1313:194-200 (1996)

Kuźnicki, J., Strauss, K.I. and Jacobowitz, D.M.: Conformational changes and calcium binding by calretinin and its recombinant fragments containing different sets of EF hand motifs. *Biochemistry*, 34:15389-15394 (1995b)

Kuźnicki, J., Wang, T.L., Martin, B.M., Winsky, L. and Jacobowitz, D.M.: Localization of Ca(2+)-dependent conformational changes of calretinin by limited tryptic proteolysis. *Biochem. J.*, 308:607-612 (1995a)

Lainé, J. and Axelrad, H.: Morphology of the Golgi-impregnated Lugano cell in the rat cerebellar cortex: a reappraisal with a description of its axon. *J. Comp. Neurol.*, 375:618-640 (1996)

Landis, S.C. and Mullen, R.J.: The development and degeneration of Purkinje cells in *pcd* mutant mice. *J. Comp. Neurol.*, 177:125-143 (1978)

Lane, P.W.: Weaver, wv, recessive. In: R.L. Sidman, M.C. Green, S.H. Appel (eds.): Catalog of the Neurological Mutations of the Mouse. *Harvard University Press*: Cambridge, MA, pp 66-67 (1965)

LaVail, M.M., Blanks, J.C. and Mullen, R.J.: Retinal degeneration in the *pcd* mutant mouse. I. Light microscopic and autoradiographic analysis. *J. Comp. Neurol.*, 212:217-230 (1982)

Liao, Y.J., Jan, Y.N. and Jan, L.Y.: Heteromultimerization of G-protein-gated inwardly rectifying K⁺ channel proteins GIRK1 and GIRK2 and their altered expression in *weaver* brain. *J. Neurosci.*, 16:7137-7150 (1996)

Lim, D.J.: Functional structure of the organ of Corti: a review. *Hear. Res.*, 22:117-146 (1986)

Liss, B., Neu, A. and Roeper, J.: The weaver mouse *gain-of-function* phenotype of dopaminergic midbrain neurons is determined by coactivation of wvGIRK2 and K-ATP channels. *J. Neurosci.*, 19:8839-8848 (1999)

Llinás, R., Hillman, D.E. and Precht, W.: Neuronal circuit reorganization in mammalian agranular cerebellar cortex. *J. Neurobiol.*, 4:69-94 (1973)

Lord, E.M. and Gates, W.H.: Shaker, a new mutation in the house mouse. *Am. Nat.*, 63:435-442 (1929)

Luciani, L.: *Il cervelletto: Nuovi studi di fisiologia normale et pathologica*. Le Monnier: Florence. (1891)

Lugaro, E.: Sulle conessioni tra gli elementi nervosi della corteccia cerebellare con considerazioni generali sul significato fisiologico dei rapporti tra gli elementi nervosi. *Riv. Sper. Freniat.*, 20:297-331 (1894)

Lüscher, C., Jan, L.Y., Stoffel, M., Malenka, R.C. and Nicoll, R.A.: G protein-coupled inwardly rectifying K⁺ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron*, 19:687-695 (1997)

LITERATURVERZEICHNIS

Mach, E.: *Grundlinien der Lehre von den Bewegungsempfindungen*. Engelmann: Leipzig (Reprint 1967), Bonset: Amsterdam (1875)

Magnin, M. and Putkonen, P.T.S.: A new vestibular thalamic area: electrophysiological study of the thalamic reticular nucleus of the ventral lateral geniculate complex of the cat. *Exp. Brain Res.*, 32:91-104 (1978)

Makowiec, R.L., Cha, J.J., Penney, J.B. and Young, A.B.: Cerebellar excitatory amino acid binding sites in normal, granuloprival and Purkinje cell-deficient mice. *Neuroscience*, 42:671-681 (1991)

Mannen, H.: Arborisations dendritique. Étude topographique et quantitative dans le noyaux vestibulaire du chat. *Arch. Ital. Biol.*, 103:197-219 (1965)

Mehler, W.R. and Rubertone, J.A.: Anatomy of the vestibular nucleus complex. In: G. Paxinos (ed.), The rat nervous system. Vol. 2. Hindbrain and spinal cord, *Academic Press*: Sydney, pp 185-219 (1985)

Mikaelian, D.O. and Ruben, R.J.: Hearing degeneration in shaker-1 mouse. *Arch. Otolaryngol. Head Neck Surg.*, 80:418-430 (1964)

Mikaelian, D.O., Alford, B.R. and Ruben, R.J.: Cochlear potentials and VIII nerve action potentials in normal and genetically deaf mice. *Ann. Otol. Rhinol. Laryngol.*, 74:146-157 (1965)

Midtgård, J.: Stellate cell inhibition of Purkinje cells in the turtle cerebellum in vitro. *J. Physiol.*, 457:355-367 (1992)

Morris, R.J., Beech, J.N. and Heizmann, C.W.: Two distinct phases and mechanisms of axonal growth shown by primary vestibular fibres in the brain, demonstrated by parvalbumin immunohistochemistry. *Neuroscience*, 27:571-596 (1988)

Mugnaini, E., Floris, A. and Wright-Goss, M.: Extraordinary synapses of the unipolar brush cell: an electron microscopic study in the rat cerebellum. *Synapse*, 16:284-311 (1994)

Mullen, R.J.: Genetic dissection of the CNS with mutant-normal mouse and its chimeras. In: W.H. Cowan and J.A. Ferrendell (eds.), Neuroscience Symposia, Vol. 2, *Society for Neuroscience*, Bethesda: Maryland, pp 47-65 (1977)

Mullen, R.J., Eicher, E.M. and Sidman, R.L.: Purkinje cell degeneration, a new neurological mutation in the mouse. *Proc. Natl. Acad. Sci. U.S.A.*, 73:208-212 (1976)

Mumenthaler, M.: Schwindel, Gleichgewichtsstörungen und Nystagmus. In: Neurologische Differentialdiagnostik, 4. Auflage, *Thieme*: Stuttgart, New York, pp 270-278 (1997)

Murer, G., Adelbrecht, C., Lauritzen, I., Lesage, F., Lazdunski, M., Agid, Y. and Raisman-Vozari, R.: An immunocytochemical study on the distribution of two G-protein-gated inward rectifier potassium channels (GIRK2 and GIRK4) in the adult rat brain. *Neuroscience*, 80:345-357 (1997)

O'Gorman, S. and Sidman, R.L.: Degeneration of thalamic neurons in "Purkinje cell degeneration" mutant mice. I. Distribution of neurons loss. *J. Comp. Neurol.*, 234:277-297 (1985)

LITERATURVERZEICHNIS

- Ottersen, O.P. and Storm-Mathisen, J.: Glutamate- and GABA-containing neurons in the mouse and rabbit brain, as demonstrated with a new immunocytochemical technique. *J. Comp. Neurol.*, 229:374-392 (1984)
- Palay, S.L. and Chan-Palay, V.: Cerebellar Cortex. Cytology and Organisation. *Springer*: Berlin, Heidelberg, New York (1974)
- Patil, N., Cox, D.R., Bhat, D., Faham, M., Myers, R.M. and Peterson, A.S.: A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation. *Nat. Genet.*, 11:126-129 (1995)
- Peters, J.: Mouse gene list. *Mouse Genome*, 86:27-80 (1990)
- Peterson, B.W. and Coulter, J.D.: A new long spinal projection from the vestibular nuclei in the cat. *Brain Res.*, 122:351-356 (1977)
- Phillips, R.J.S.: ‘Lurcher‘, a new gene in linkage group XI of the house mouse. *J. Genet.*, 57:35-42 (1960)
- Pickles, J.O. and Corey, D.P.: Mechanoelectrical transduction by hair cells. *Trends Neurosci.*, 15:254-259 (1992)
- Purkinje, J.E.: Neueste Untersuchungen aus der Nerven- und Hirnanatomie. In: K. Sternberg und J.V. von Krombholtz (eds.), *Bericht über die Versammlung deutscher Naturforscher und Aerzte in Prag im September*, pp 177-180 (1837)
- Qizilbash, N., Lopez-Arrieta, J.M. and Birks, J.: Nimodipine for primary degenerative, mixed and vascular dementia (Cochrane Review). In: *The Cochrane Library*, Issue 1 (1999)
- Rakic, P. and Sidman, R.L.: Sequence of developmental abnormalities leading to granule cell deficit in cerebellar cortex of weaver mutant mice. *J. Comp. Neurol.*, 152:103-132 (1973a)
- Rakic, P. and Sidman, R.L.: Organization of cerebellar cortex secondary to deficit of granule cells in weaver mutant mice. *J. Comp. Neurol.*, 153:133-162 (1973b)
- Ramón y Cajal, S.: *Histologie du Système Nerveux de l'Homme et des Vertébrés*, Tome I. Maloine: Paris (1909)
- Raymond, J., Dechesne, C.J., Desmadryl, G., and Demêmes, D.: Different calcium-binding proteins identify subpopulations of vestibular ganglion neurons in the rat. *Acta Otolaryngol. Suppl.*, 503:114-118 (1993)
- Rezai, Z. and Yoon, C.H.: Abnormal rate of granule cell migration in the cerebellum of “Weaver“ mutant mice. *Dev. Biol.*, 29:17-26 (1972)
- Roberts, W.M., Howard, J. and Hudspeth, A.J.: Hair cells: transduction, tuning, and transmission in the inner ear. *Annu. Rev. Cell Biol.*, 4:63-92 (1988)

LITERATURVERZEICHNIS

Roffler-Tarlov, S., Beart, P.M., O'Gorman, S., and Sidman, R.L.: Neurochemical and morphological consequences of axon terminal degeneration in cerebellar deep nuclei of mice with inherited Purkinje cell degeneration. *Brain Res.*, 168:75-95 (1979)

Roffler-Tarlov, S. and Graybiel, A.M.: Expression of the weaver gene in dopamine-containing neural systems is dose-dependent and affects both striatal and nonstriatal regions. *J. Neurosci.*, 6:3319-3330 (1986)

Rogers, J.H.: Calretinin: a gene for a novel calcium-binding protein expressed principally in neurons. *J. Cell Biol.*, 105:1343-1353 (1987)

Rogers, J.H.: Immunoreactivity for calretinin and other calcium-binding proteins in cerebellum. *Neuroscience*, 31:711-721 (1989)

Sahly, I., El-Amraoui, A., Abitbol, M., Petit, C. and Dufier, J.-L.: Expression of myosin VIIA during mouse embryogenesis. *Anat. Embryol. (Berl.)*, 196:159-170 (1997)

Scheibel, M.E. and Scheibel, A.B.: Observations on the intracortical relations of the climbing fibres of the cerebellum. *J. Comp. Neurol.*, 101:733-763 (1954)

Schiffmann, S.N., Cheron, G., Lohof, A., d'Alcantara, P., Meyer, M., Parmentier, M. and Schurmans, S.: Impaired motor coordination and Purkinje cell excitability in mice lacking calretinin. *Proc. Natl. Acad. Sci. USA*, 96:5257-5262 (1999)

Schmidt, M.J., Sawyer, B.D., Perry, K.W., Fuller, R.W., Foreman, M.M. and Ghetti, B.: Dopamine deficiency in the weaver mutant mouse. *J. Neurosci.*, 2:376-380 (1982)

Schurmans, S., Schiffmann, S.N., Gurden, H., Lemaire, M., Lipp, H.-P., Schwam, V., Pochet, R., Imperato, A., Böhme, G.A. and Parmentier, M.: Impaired long-term potentiation induction in dentate gyrus of calretinin-deficient mice. *Proc. Natl. Acad. Sci. USA*, 94:10415-10420 (1997)

Schwarz, I.E. and Schwarz, D.W.: The primary vestibular projection to the cerebellar cortex in the pigeon (*Columba livia*). *J. Comp. Neurol.*, 216:438-444 (1983)

Seguela, P., Geffard, M., Bujis, R.M. and le Moal, M.: Antibodies against gamma-aminobutyric acid: specificity studies and immunocytochemical results. *Proc. Natl. Acad. Sci. U.S.A.*, 81:3888-3892 (1984)

Sekerková, G., Loomis, P.A., Changyaleket, B., Zheng, L., Etyan, R., Chen, B., Mugnaini, E. and Bartles, J.R.: Novel Espin Actin-Bundling Proteins Are Localized to Purkinje Cell Dendritic Spines and Bind the Src Homology 3 Adapter Protein Insulin Receptor Substrate p53. *J. Neurosci.*, 23:1310-1319 (2003)

Sekiguchi, M., Nowakowski, R.S., Nagato, Y., Tanaka, O., Guo, H., Madoka, M. and Abe, H.: Morphological abnormalities in the hippocampus of the weaver mutant mouse. *Brain Res.*, 696:262-267 (1995)

Self, T., Mahony, M., Fleming, J., Walsh, J., Brown, S.D.M. and Steel, K.P.: Shaker-1 mutations reveal roles for myosin VIIA in both development and function of cochlear hair cells. *Development*, 125:557-566 (1998)

LITERATURVERZEICHNIS

- Shnerson, A., Lenoir, M., Van de Water, T.R. and Pujol, R.: The pattern of sensorineural degeneration in the cochlea of the deaf shaker-1 mouse: ultrastructural observations. *Brain Res.*, 285:305-315 (1983)
- Shojaeian, H., Delhaye-Bouchaud, N. and Mariani, J.: Stability of inferior olivary neurons in rodents. I. Moderate cell loss in adult Purkinje cell degeneration mutant mouse. *Brain Res.*, 466:211-218 (1988)
- Sidman, R.L., Green, M.C. and Appel, S.H.: Catalog of the neurological mutants of the mouse. *Harvard University Press*: Cambridge, Massachusetts (1965)
- Sirkin, D.W., Precht, W. and Courjon, J.H.: Initial, rapid phase of recovery from unilateral vestibular lesion in rat not dependent on survival of central portion of vestibular nerve. *Brain Res.*, 302:245-256 (1984)
- Sjöström, B. and Anniko, M.: Morphologically specific vestibular hair cell degeneration in the jerker mutant mouse. *Eur. Arch. Otorhinolaryngol.*, 247:51-55 (1990a)
- Sjöström, B. and Anniko, M.: Variability in genetically induced age-related impairment of auditory brainstem response thresholds. *Acta Otolaryngol.*, 109:353-360 (1990b)
- Sjöström, B. and Anniko, M.: Cochlear structure and function in a recessive type of genetically induced inner ear degeneration. *ORL J. Otorhinolaryngol. Relat. Spec.*, 54:220-228 (1992a)
- Sjöström, B. and Anniko, M.: Genetically induced inner ear degeneration. A structural and functional study. *Acta Otolaryngol. Suppl.*, 493:141-146 (1992b)
- Sloviter, R.S., Sollas, A.L., Barbaro, N.M. and Laxer, K.D.: Calcium-binding protein (Calbindin-D28K) and parvalbumin immunocytochemistry in the normal and epileptic human hippocampus. *J. Comp. Neurol.*, 308:381-396 (1991)
- Smeyne, R.J. and Goldowitz, D.: Development and death of external granular layer cells in the weaver mouse cerebellum: a quantitative study. *J. Neurosci.*, 9:1808-1820 (1989)
- Smeyne, R.J. and Goldowitz, D.: Postnatal development of the wild-type and weaver cerebellum after embryonic administration of propylthiouracil (PTU). *Dev. Brain Res.*, 54:282-286 (1990)
- Smith, R.J.H., Berlin, C.I., Hejtmancik, J.F., Keats, B.J.B., Kimberling, W.J., Lewis, R.A., Möller, C.G., Pelias, M.Z. and Tranebjærg, L.: Clinical diagnosis of the Usher syndromes. *Am. J. Med. Genet.*, 50:32-38 (1994)
- Smolyaninov, V.V.: Some special features of organization of the cerebellar cortex. In: I.M. Gelfand, V.S., V.S. Gurfinkel, S.V. Fomin and M.L. Tsetlin (eds.), *Models of the Structural-Functional Organization of Certain Biological Systems*. MIT Press: Cambridge, pp 250-423 (1971)
- Sotelo, C.: Mutant mice and the formation of cerebellar circuitry. *Trends Neurosci.*, 3:33-36 (1980)
- Sotelo, C.: Synaptic remodeling in agranular cerebella. In: S.L. Palay, V. Chan-Palay (eds.), *The Cerebellum – New Vistas*. Springer: Berlin, Heidelberg, New York, pp 50-68 (1982)

LITERATURVERZEICHNIS

- Sotelo, C. and Alvarado-Mallart, R. M.: Integration of grafted Purkinje cell into the host of cerebellar circuitry in Purkinje cell degeneration mutant mouse. *Prog. Brain Res.*, 18:141-154 (1988)
- Sotelo, C. and Alvarado-Mallart, R.M.: The reconstruction of cerebellar circuits. *Trends Neurosci.*, 14: 350-355 (1991)
- Sotelo, C. and Changeaux, J.P.: Bergmann fibers and granular cell migration in the cerebellum of homozygous weaver mutant mouse. *Brain Res.*, 77:484-491 (1974)
- Steel, K.P.: Inherited hearing defects in mice. *Annu. Rev. Genet.*, 29:675-701 (1995)
- Steel, K.P.: Varitint-waddler: a double whammy for hearing. *Proc. Natl. Acad. Sci. USA*, 99:14613-14615 (2002)
- Steel, K.P. and Bock, G.R.: Cochlear dysfunction in the jerker mouse. *Behav. Neurosci.*, 97: 381-391 (1983)
- Stein, R.B. and Lee, R.G.: Tremor and Clonus. In: J.M. Brookhart, V.B. Mountcastle, V.B. Brooks (eds.): *Handbook of Physiology*, Section 1: The Nervous System, Vol. II, Motor Control, Part 1. *American Physiological Society*: Bethesda, pp 325-344 (1981)
- Sterio, D.C.: The unbiased estimation of number and sizes of arbitrary particles using the disector. *J. Microsc.*, 134:127-136 (1984)
- Sun, J.C., van Alphen, A.M., Wagenaar, M., Huygen, P., Hoogenraad, C.C., Hasson, T., Koekkoek, S.K.E., Bohne, B.A. and de Zeeuw, C.I.: Origin of vestibular dysfunction in usher syndrome type 1B. *Neurobiol. Dis.*, 8:69-77 (2001)
- Swisher, D.A., and Wilson, D.B.: Cerebellar histogenesis in the lurcher (*Lc*) mutant mouse. *J. Comp. Neurol.*, 173:205-217 (1977)
- Thach, W.T.: Ataxia, cerebellar. In: G. Adelmann (ed.), *Encyclopedia of Neuroscience*. *Birkhäuser*: Boston, Basel, Stuttgart, pp 83-86 (1987)
- Tilney, L.G., Tilney, M.S., and DeRosier, D.J.: Actin filaments, stereocilia and hair cells: how cells count and measure. *Annu. Rev. Cell Biol.*, 8:257-274 (1992)
- Tilney, M.S., Tilney, L.G., Stephens, R.E., Merte, C., Drenckhahn, D., Contanche, D.A. and Bretscher, A.: Preliminary characterization of the stereocilia and cuticular plate of hair cells in the chick cochlea. *J. Cell Biol.*, 109:1711-1723 (1989)
- Todorov, P.T., Hardisty, R.E. and Brown, S.D.M.: Myosin VIIA is specifically associated with calmodulin and microtubule-associated protein-2B (MAP-2B). *Biochem. J.*, 354:267-274 (2001)
- Trepel, M.: Mikroskopische Anatomie der Kleinhirnrinde. In: *Neuroanatomie: Struktur und Funktion*. *Urban & Fischer*: München, Stuttgart, Jena, Lübeck, Ulm, p144 (1999)
- Triarhou, L.C. and Ghetti, B.: The dendritic dopamine projection of the substantia nigra: phenotypic denominator of weaver gene action in hetero- and homozygosity. *Brain Res.*, 501:373-381 (1989)

LITERATURVERZEICHNIS

- Triarhou, L.C. and Ghetti, B.: Stabilization of neurone number in the inferior olfactory complex of aged 'Purkinje cell degeneration' mutant mice. *Acta Neuropathol. (Berl.)*, 81:597-602 (1991)
- Triarhou, L.C., Norton J., Alyea, C. and Ghetti, B.: A quantitative study of the granule cells in the Purkinje cell degeneration mutant. *Ann. Neurol.*, 18:146/P108 (1985)
- Triarhou, L.C., Norton, J. and Ghetti, B.: Anterograde transsynaptic degeneration in the deep cerebellar nuclei of Purkinje cell degeneration (*pcd*) mutant mice. *Exp. Brain Res.*, 66:577-588 (1987)
- Tsuboi, K., Kimber, T.A. and Shults, C.W.: Calretinin-containing axons and neurons are resistant to an intrastriatal 6-hydroxydopamine lesion. *Brain Res.*, 866:55-64 (2000)
- Udovichenko, I.P., Gibbs, D. and Williams, D.S.: Actin-based motor properties of native myosin VIIa. *J. Cell Sci.*, 115:445-450 (2002)
- Usher, C.R.: On the inheritance of retinitis pigmentosa with notes of cases. *R. Lond. Ophthalmol. Hosp. Rep.*, 19:130-236 (1913/14)
- Vincent, P., Armstrong, C.M. and Marty, A.: Inhibitory synaptic currents in rat cerebellar Purkinje cells: modulation by postsynaptic depolarization. *J. Physiol.*, 456:453-471 (1992)
- Voogd, J., Gerrits, N.M. and Marani, E.: Cerebellum. In: G. Paxinos (ed.), The rat nervous system. Vol. 2. Hindbrain and spinal cord, *Academic Press*: Sydney, pp 251-291 (1985)
- Voogd, J., Gerrits, N.M. and Ruigrok, T.J.: Organization of the vestibulocerebellum. *Ann. N.Y. Acad. Sci.*, 781:553-579 (1996)
- Wassem, M., Simons, J., Tappaz, M.L. and Sotelo, C.: Non-Purkinje cell GABAergic innervation of the deep cerebellar nuclei: a quantitative immunocytochemical study in C57BL and in Purkinje cell degeneration mutant mice. *Brain Res.*, 399:125-135 (1986)
- Wei, J., Dlouhy, S.R., Bayer, S., Piva, R., Verina, T., Wang, Y., Feng, Y., Dupree, B., Hodes, M.E. and Ghetti, B.: In situ hybridization analysis of *Girk2* expression in the developing central nervous system in normal and *weaver* mice. *J. Neuropathol. Exp. Neurol.*, 56:762-771 (1997)
- Weil, D., Blanchard, S., Kaplan, J., Guilford, P., Gibson, F., Walsh, J., Mburu, P., Varela, A., Levilliers, J., Weston, M.D., Kelley, P.M., Kimberling, W.J., Wagenaar, M., Levi-Acobas, F., Larget-Piet, D., Munnich, A., Steel, K.P., Brown, S.D.M. and Petit, C.: Defective myosin VIIA gene responsible for Usher syndrome type 1B. *Nature*, 374:60-61 (1995)
- Wersäll, J.: Studies of the structure and innervation of the sensory epithelium of the cristae ampullares in the guinea pig. *Acta Otolaryngol. (Stockh.)*, Suppl. 126:1-85 (1956)
- West, M.J.: New stereological methods for counting neurons. *Neurobiol. Aging*, 14:275-285 (1993)
- West, M.J.: Advances in the study of age-related neuron loss. *Sem. Neurosci.*, 6:403-411 (1994)
- Wetts, H. and Herrup, K.: Interaction of granule, Purkinje and inferior olfactory neurons in lurcher chimeric mice. II. Granule cell death. *Brain Res.*, 250:358-362 (1982)

LITERATURVERZEICHNIS

Wilson, V.J., Peterson, B.W., Fukushima, K., Hirai, N. and Uchino, Y.: Analysis of vestibulocollic reflexes by sinusoidal polarization of vestibular afferent fibers. *J. Neurophysiol.*, 42:331-346 (1979)

Yagi, T., Simpson, N.E. and Markham, C.H.: The relationship of conduction velocity to other physiological properties of the cat's horizontal canal neurons. *Exp. Brain Res.*, 30:587-600 (1977)

Zheng, L., Sekerkova, G., Vranich, K., Tilney, L.G., Mugnaini, E. and Bartles, J.R.: The deaf jerker mouse has a mutation in the gene encoding the espin actin-bundling proteins of hair cell stereocilia and lacks espins. *Cell*, 102:377-385 (2000)

Zuo, J., De Jager, P.L., Takashi, K.A., Jiang, W., Linden, D.J. and Heintz, N.: Neurodegeneration in Lurcher mice caused by mutation in $\delta 2$ glutamate receptor gene. *Nature*, 388:769-773 (1997)