

8. Literatur

Augustine PC. Cell: sporozoite interactions and invasion by apicomplexan parasites of the genus *Eimeria*. *Int J Parasitol*. 2001 Jan;31(1):1-8. Review.

Bruno S, Duschak VG, Ledesma B, Ferella M, Andersson B, Guarnera EA, Angel SO. Identification and characterization of serine proteinase inhibitors from *Neospora caninum*. *Mol Biochem Parasitol*. 2004 Jul;136(1):101-7.

Candolfi E, Hunter CA, Remington JS. Mitogen- and antigen-specific proliferation of T cells in murine toxoplasmosis is inhibited by reactive nitrogen intermediates. *Infect Immun*. 1994 May;62(5):1995-2001.

Cappello M, Hawdon JM, Jones BF, Kennedy WP, Hotez PJ. *Ancylostoma caninum* anticoagulant peptide: cloning by PCR and expression of soluble, active protein in *E. coli*. *Mol Biochem Parasitol*. 1996 Sep;80(1):113-7.

Carter P, Wells JA. Dissecting the catalytic triad of a serine protease. *Nature*. 1988 Apr 7;332(6164):564-8.

Channon JY, Kasper LH. *Toxoplasma gondii*-induced immune suppression by human peripheral blood monocytes: role of gamma interferon. *Infect Immun*. 1996 Apr;64(4):1181-9.

Chu D, Bungiro RD, Ibanez M, Harrison LM, Campodonico E, Jones BF, Mieszczanek J, Kuzmic P, Cappello M. Molecular characterization of *Ancylostoma ceylanicum* Kunitz-type serine protease inhibitor: evidence for a role in hookworm-associated growth delay. *Infect Immun*. 2004 Apr;72(4):2214-21.

Conseil V, Soete M, Dubremetz JF. Serine protease inhibitors block invasion of host cells by *Toxoplasma gondii*. *Antimicrob Agents Chemother*. 1999 Jun;43(6):1358-61.

Corvera CU, Dery O, McConalogue K, Bohm SK, Khitin LM, Caughey GH, Payan DG, Bunnett NW. Free in PMC Mast cell tryptase regulates rat colonic myocytes through proteinase-activated receptor 2. *J Clin Invest*. 1997 Sep 15;100(6):1383-93.

Dobrowolski JM, Sibley LD. *Toxoplasma* invasion of mammalian cells is powered by the actin cytoskeleton of the parasite. *Cell*. 1996 Mar 22;84(6):933-9.

Foey AD, Feldmann M, Brennan FM. CD40 ligation induces macrophage IL-10 and TNF-alpha production: differential use of the PI3K and p42/44 MAPK-pathways. *Cytokine*. 2001 Nov 21;16(4):131-42.

van Gent D, Sharp P, Morgan K, Kalsheker N. Serpins: structure, function and molecular evolution. *Int J Biochem Cell Biol*. 2003 Nov;35(11):1536-47. Review.

Gomez-Escobar N, Gregory WF, Maizels RM. Identification of tgh-2, a filarial nematode homolog of *Caenorhabditis elegans* daf-7 and human transforming growth factor beta,

expressed in microfilarial and adult stages of *Brugia malayi*. *Infect Immun*. 2000 Nov;68(11):6402-10.

Goodridge HS, Wilson EH, Harnett W, Campbell CC, Harnett MM, Liew FY. Free Full Text Modulation of macrophage cytokine production by ES-62, a secreted product of the filarial nematode *Acanthocheilonema viteae*. *J Immunol*. 2001 Jul 15;167(2):940-5.

Gu L, Tseng S, Horner RM, Tam C, Loda M, Rollins BJ. Control of TH2 polarization by the chemokine monocyte chemoattractant protein-1. *Nature*. 2000 Mar 23;404(6776):407-11.

Habeck M. Vaccinating against ticks. *Drug Discov Today*. 2002 Jun 1;7(11):586-8. No abstract available.

Hall W, Hall E. *Guidebook to Cytokines and Their Receptors*. Oxford University Press, Oxford, New York, Tokyo. 1994: 83-5.

Hamblin AS. *Cytokines and Cytokine Receptors*. Oxford University Press. New York. 1993, 2nd edition: 14-8, 33-4.

Hartmann S, Kyewski B, Sonnenburg B, Lucius R. A filarial cysteine protease inhibitor down-regulates T cell proliferation and enhances interleukin-10 production. *Eur J Immunol*. 1997 Sep;27(9):2253-60.

Huang DR, Wang J, Kivisakk P, Rollins BJ, Ransohoff RM. Absence of monocyte chemoattractant protein 1 in mice leads to decreased local macrophage recruitment and antigen-specific T helper cell type 1 immune response in experimental autoimmune encephalomyelitis. *J Exp Med*. 2001 Mar 19;193(6):713-26.

Huang X, Zhao N, Zhang K. Abstract Extracellular enzymes serving as virulence factors in nematophagous fungi involved in infection of the host. *Res Microbiol*. 2004 Dec;155(10):811-6. Review.

Huntington JA, Read RJ, Carrell RW. Structure of a serpin-protease complex shows inhibition by deformation. *Nature*. 2000 Oct 19;407(6806):923-6.

Irving JA, Pike RN, Lesk AM, Whisstock JC. Phylogeny of the serpin superfamily: implications of patterns of amino acid conservation for structure and function. *Genome Res*. 2000 Dec;10(12):1845-64.

Irving JA, Askew DJ, Whisstock JC. Computational analysis of evolution and conservation in a protein superfamily. *Methods*. 2004 Feb;32(2):73-92.

Jiang H, Kanost MR. Characterization and functional analysis of 12 naturally occurring reactive site variants of serpin-1 from *Manduca sexta*. *J Biol Chem*. 1997 Jan 10;272(2):1082-7.

Kaslik G, Patthy A, Balint M, Graf L. Trypsin complexed with alpha 1-proteinase inhibitor has an increased structural flexibility. *FEBS Lett*. 1995 Aug 21;370(3):179-83.

Kim JY, Park SC, Kim MH, Lim HT, Park Y, Hahm KS. Antimicrobial activity studies on a trypsin-chymotrypsin protease inhibitor obtained from potato. *Biochem Biophys Res Commun*. 2005 May 13;330(3):921-7.

Kim K, Weiss LM. Abstract *Toxoplasma gondii*: the model apicomplexan. *Int J Parasitol*. 2004 Mar 9;34(3):423-32. Review.

Komiyama T, Ray CA, Pickup DJ, Howard AD, Thornberry NA, Peterson EP, Salvesen G. Inhibition of interleukin-1 beta converting enzyme by the cowpox virus serpin CrmA. An example of cross-class inhibition. *J Biol Chem*. 1994 Jul 29;269(30):19331-7.

Lindh JG, Botero-Kleiven S, Arboleda JJ, Wahlgren M. A protease inhibitor associated with the surface of *Toxoplasma gondii*. *Mol Biochem Parasitol*. 2001 Sep 3;116(2):137-45.

Loke P, MacDonald AS, Robb A, Maizels RM, Allen JE. Alternatively activated macrophages induced by nematode infection inhibit proliferation via cell-to-cell contact. *Eur J Immunol*. 2000 Sep;30(9):2669-78.

Lomas DA, Carrell RW. Serpinopathies and the conformational dementias. *Nat Rev Genet*. 2002 Oct;3(10):759-68. Review

Maizels RM, Yazdanbakhsh M. Abstract Immune regulation by helminth parasites: cellular and molecular mechanisms. *Nat Rev Immunol*. 2003 Sep;3(9):733-44. Review.

Macfarlane SR, Seatter MJ, Kanke T, Hunter GD, Plevin R. Proteinase-activated receptors. *Pharmacol Rev*. 2001 Jun;53(2):245-82. Review.

Maizels RM, Gomez-Escobar N, Gregory WF, Murray J, Zang X. Immune evasion genes from filarial nematodes. *Int J Parasitol*. 2001 Jul;31(9):889-98. Review.

Maizels RM, Balic A, Gomez-Escobar N, Nair M, Taylor MD, Allen JE. Helminth parasites--masters of regulation. *Immunol Rev*. 2004 Oct;201:89-116. Review.

Manoury B, Gregory WF, Maizels RM, Watts C. Bm-CPI-2, a cystatin homolog secreted by the filarial parasite *Brugia malayi*, inhibits class II MHC-restricted antigen processing. *Curr Biol*. 2001 Mar 20;11(6):447-51.

Markwardt F, Hirudin as alternative anticoagulant--a historical review. *Semin Thromb Hemost*. 2002 Oct;28(5):405-14. Review.

Menssen HD, Brandt N, Leben R, Muller F, Thiel E, Melber K. Measurement of hematological, clinical chemistry, and infection parameters from hirudinized blood collected in universal blood sampling tubes. *Semin Thromb Hemost*. 2001 Aug;27(4):349-56.

Milstone AM, Harrison LM, Bungiro RD, Kuzmic P, Cappello M. A broad spectrum Kunitz type serine protease inhibitor secreted by the hookworm *Ancylostoma ceylanicum*. *J Biol Chem*. 2000 Sep 22;275(38):29391-9.

Morris MT, Coppin A, Tomavo S, Carruthers VB. Functional analysis of *Toxoplasma gondii* protease inhibitor 1. *J Biol Chem*. 2002 Nov 22;277(47):45259-66. Epub 2002 Sep 11.

- Morris MT, Carruthers VB. Identification and partial characterization of a second Kazal inhibitor in *Toxoplasma gondii*. *Mol Biochem Parasitol*. 2003 Apr 25;128(1):119-22.
- Mule F, Baffi MC, Cerra MC. Dual effect mediated by protease-activated receptors on the mechanical activity of rat colon. *Br J Pharmacol*. 2002 Jun;136(3):367-74.
- Muleng A, Sugino M, Nakajim M, Sugimoto C, Onuma M. Tick-Encoded serine proteinase inhibitors (serpins); potential target antigens for tick vaccine development. *J Vet Med Sci*. 2001 Oct;63(10):1063-9. Review.
- Nguyen TT, Qasim MA, Morris S, Lu CC, Hill D, Laskowski M Jr, Sakanari JA. Expression and characterization of elastase inhibitors from the ascarid nematodes *Anisakis simplex* and *Ascaris suum*. *Mol Biochem Parasitol*. 1999 Jul 30;102(1):79-89.
- Nufer O, Corbett M, Walz A. Amino-terminal processing of chemokine ENA-78 regulates biological activity. *Biochemistry*. 1999 Jan 12;38(2):636-42.
- Owen CA, Campbell EJ. The cell biology of leukocyte-mediated proteolysis. *J Leukoc Biol*. 1999 Feb;65(2):137-50. Review.
- Pak SC, Kumar V, Tsu C, Luke CJ, Askew YS, Askew DJ, Mills DR, Bromme D, Silverman GA. SRP-2 is a cross-class inhibitor that participates in postembryonic development of the nematode *Caenorhabditis elegans*: initial characterization of the clade L serpins. *J Biol Chem*. 2004 Apr 9;279(15):15448-59. Epub 2004 Jan 22.
- Pastrana DV, Raghavan N, FitzGerald P, Eisinger SW, Metz C, Bucala R, Schleimer RP, Bickel C, Scott AL. Filarial nematode parasites secrete a homologue of the human cytokine macrophage migration inhibitory factor. *Infect Immun*. 1998 Dec;66(12):5955-63.
- Petit F, Bertagnoli S, Gelfi J, Fassy F, Boucraut-Baralon C, Milon A. Characterization of a myxoma virus-encoded serpin-like protein with activity against interleukin-1 beta-converting enzyme. *J Virol*. 1996 Sep;70(9):5860-6.
- Pfaff AW, Schulz-Key H, Soboslay PT, Taylor DW, MacLennan K, Hoffmann WH. *Litomosoides sigmodontis* cystatin acts as an immunomodulator during experimental filariasis. *Int J Parasitol*. 2002 Feb;32(2):171-8.
- Plotnick MI, Rubin H, Schechter NM. The effects of reactive site location on the inhibitory properties of the serpin alpha(1)-antichymotrypsin. *J Biol Chem*. 2002 Aug 16;277(33):29927-35. Epub 2002 Jun 7.
- Potempa J, Korzus E, Travis J. The serpin superfamily of proteinase inhibitors: structure, function, and regulation. *J Biol Chem*. 1994 Jun 10;269(23):15957-60. Review. No abstract available.
- Pszenny V, Angel SO, Duschak VG, Paulino M, Ledesma B, Yabo MI, Guarnera E, Ruiz AM, Bontempi EJ. Molecular cloning, sequencing and expression of a serine proteinase inhibitor gene from *Toxoplasma gondii*. *Mol Biochem Parasitol*. 2000 Apr 15;107(2):241-9.

Pszenny V, Ledesma BE, Matrajt M, Duschak VG, Bontempi EJ, Dubremetz JF, Angel SO. Subcellular localization and post-secretory targeting of TgPI, a serine proteinase inhibitor from *Toxoplasma gondii*. *Mol Biochem Parasitol*. 2002 May;121(2):283-6.

Quan LT, Caputo A, Bleackley RC, Pickup DJ, Salvesen GS. Granzyme B is inhibited by the cowpox virus serpin cytokine response modifier A. *J Biol Chem*. 1995 May 5;270(18):10377-9.

Rhoads ML, Fetterer RH, Hill DE, Urban JF Jr. *Trichuris suis*: a secretory chymotrypsin/elastase inhibitor with potential as an immunomodulator. *Exp Parasitol*. 2000 May;95(1):36-44.

Riahi Y, Siman-Tov R, Ankri S. Molecular cloning, expression and characterization of a serine proteinase inhibitor gene from *Entamoeba histolytica*. *Mol Biochem Parasitol*. 2004 Feb;133(2):153-62.

Roggwiler E, Betoulle ME, Blisnick T, Braun Breton C. A role for erythrocyte band 3 degradation by the parasite gp76 serine protease in the formation of the parasitophorous vacuole during invasion of erythrocytes by *Plasmodium falciparum*. *Mol Biochem Parasitol*. 1996 Nov 12;82(1):13-24.

Sant'Anna Azzolini S, Sasaki SD, Torquato RJ, Andreotti R, Andreotti E, Tanaka AS. Abstract *Rhipicephalus sanguineus* trypsin inhibitors present in the tick larvae: isolation, characterization, and partial primary structure determination. *Arch Biochem Biophys*. 2003 Sep 15;417(2):176-82.

Schierack P, Lucius R, Sonnenburg B, Schilling K, Hartmann S. Parasite-specific immunomodulatory functions of filarial cystatin. *Infect Immun*. 2003 May;71(5):2422-9.

Shaw MK, Roos DS, Tilney LG. Abstract Cysteine and serine protease inhibitors block intracellular development and disrupt the secretory pathway of *Toxoplasma gondii*. *Microbes Infect*. 2002 Feb;4(2):119-32.

Silverman GA, Bird PI, Carrell RW, Church FC, Coughlin PB, Gettins PG, Irving JA, Lomas DA, Luke CJ, Moyer RW, Pemberton PA, Remold-O'Donnell E, Salvesen GS, Travis J, Whisstock JC. The serpins are an expanding superfamily of structurally similar but functionally diverse proteins. Evolution, mechanism of inhibition, novel functions, and a revised nomenclature. *J Biol Chem*. 2001 Sep 7;276(36):33293-6. Epub 2001 Jul 2. Review.

Simonet G, Claeys I, Broeck JV. Abstract Structural and functional properties of a novel serine protease inhibiting peptide family in arthropods. *Comp Biochem Physiol B Biochem Mol Biol*. 2002 May;132(1):247-55. Review.

Stanley P, Stein PE. BmSPN2, a serpin secreted by the filarial nematode *Brugia malayi*, does not inhibit human neutrophil proteinases but plays a noninhibitory role. *Biochemistry*. 2003 May 27;42(20):6241-8.

Steinhoff M, Buddenkotte J, Shpacovitch V, Rattenholl A, Moormann C, Vergnolle N, Luger TA, Hollenberg MD. Proteinase-activated receptors: transducers of proteinase-mediated signaling in inflammation and immune response. *Endocr Rev*. 2005 Feb;26(1):1-43. Review.

Stratikos E, Gettins PG. Formation of the covalent serpin-proteinase complex involves translocation of the proteinase by more than 70 Å and full insertion of the reactive center loop into beta-sheet A. *Proc Natl Acad Sci U S A*. 1999 Apr 27;96(9):4808-13.

Stryer D. *Biochemistry*. WH Freeman. New York. 1995, 4th edition: 222-7, 254-8.

Tanaka AS, Andreotti R, Gomes A, Torquato RJ, Sampaio MU, Sampaio CA. A double headed serine proteinase inhibitor--human plasma kallikrein and elastase inhibitor--from *Boophilus microplus* larvae. *Immunopharmacology*. 1999 Dec;45(1-3):171-7.

Taylor MD, LeGoff L, Harris A, Malone E, Allen JE, Maizels RM. Removal of regulatory T cell activity reverses hyporesponsiveness and leads to filarial parasite clearance in vivo. *J Immunol*. 2005 Apr 15;174(8):4924-33.

Trimnell AR, Hails RS, Nuttall PA. Dual action ectoparasite vaccine targeting 'exposed' and 'concealed' antigens. *Vaccine*. 2002 Oct 4;20(29-30):3560-8.

Uehara A, Muramoto K, Takada H, Sugawara S. Neutrophil serine proteinases activate human nonepithelial cells to produce inflammatory cytokines through protease-activated receptor 2. *J Immunol*. 2003 Jun 1;170(11):5690-6.

Vergnolle N, Derian CK, D'Andrea MR, Steinhoff M, Andrade-Gordon P. Characterization of thrombin-induced leukocyte rolling and adherence: a potential proinflammatory role for proteinase-activated receptor-4. *J Immunol*. 2002 Aug 1;169(3):1467-73.

Whisstock JC, Pike RN, Jin L, Skinner R, Pei XY, Carrell RW, Lesk AM. Conformational changes in serpins: II. The mechanism of activation of antithrombin by heparinidagger. *J Mol Biol*. 2000 Sep 1;301(5):1287-305.

Ye S, Goldsmith EJ. Serpins and other covalent protease inhibitors. *Curr Opin Struct Biol*. 2001 Dec;11(6):740-5. Review.

Yenbutr P, Scott AL. Molecular cloning of a serine proteinase inhibitor from *Brugia malayi*. *Infect Immun*. 1995 May;63(5):1745-53.

Yoo BC, Aoki K, Xiang Y, Campbell LR, Hull RJ, Xoconostle-Cazares B, Monzer J, Lee JY, Ullman DE, Lucas WJ. Characterization of cucurbita maxima phloem serpin-1 (CmPS-1). A developmentally regulated elastase inhibitor. *J Biol Chem*. 2000 Nov 10;275(45):35122-8.

Zang X, Yazdanbakhsh M, Jiang H, Kanost MR, Maizels RM. A novel serpin expressed by blood-borne microfilariae of the parasitic nematode *Brugia malayi* inhibits human neutrophil serine proteinases. *Blood*. 1999 Aug 15;94(4):1418-28.

Zang X, Atmadja AK, Gray P, Allen JE, Gray CA, Lawrence RA, Yazdanbakhsh M, Maizels RM. The serpin secreted by *Brugia malayi* microfilariae, Bm-SPN-2, elicits strong, but short-lived, immune responses in mice and humans. *J Immunol*. 2000 Nov 1;165(9):5161-9.

Zang X, Maizels RM. Serine proteinase inhibitors from nematodes and the arms race between host and pathogen. *Trends Biochem Sci*. 2001 Mar;26(3):191-7. Review.