

THE USE OF ANTISENSE OLIGONUCLEOTIDES TO STUDY MCM FUNCTION DURING THE INITIATION OF DNA REPLICATION

Inaugural Dissertation zur Erlangung
des Doktorgrades der Naturwissenschaften

eingereicht am
Fachbereich Biologie der Freien Universität Berlin

vorgelegt von **Pierre Charles Debs**
aus Brooklyn, New York

Berlin 2004

1. Gutachter: Prof. Dr. Thomas Blankenstein

2. Gutachter: Prof. Dr. Volker Erdmann

Tag der Disputation: 02 November 2004

"For Angelika, Lulia and MonkeyBoy"

| | |
|--|-----------|
| Introduction | 7 |
| The initiation of DNA replication..... | 7 |
| Components of the Pre-Replicative Complex | 12 |
| Eukaryotic MCM proteins..... | 16 |
| Goal of this thesis | 19 |
| Methods | 21 |
| Culturing primary human fibroblasts | 21 |
| Serum starvation of primary human fibroblasts | 22 |
| Preparation of antisense oligonucleotides and transfection lipids..... | 22 |
| Tranfection of primary human fibroblasts with GeneBloks..... | 23 |
| Analysis of mRNA levels by Real-time rtPCR | 24 |
| Preparation of whole cell extracts for immunoblotting..... | 25 |
| Preparation of fractionated cell extracts for immunoblotting | 25 |
| Polyacrylamide Gel Electrophoresis (PAGE)..... | 26 |
| Transfer of proteins to PVDF membranes | 26 |
| Immunoblotting | 27 |
| Chemiluminescence | 27 |
| Coomassie staining and destaining of PVDF membranes..... | 27 |
| Preparation of primary human fibroblasts for flow cytometry | 28 |
| Flow cytometry..... | 28 |
| BrdU incorporation and detection | 29 |
| Detection of MPM2 reactivity by flow cytometry..... | 30 |
| Hoechst staining of DNA for microscopy | 30 |
| Materials..... | 31 |
| Computers | 35 |
| Programs | 35 |
| Machines, laboratory equipment and chemicals | 36 |
| Mcm4 cDNA | 37 |
| Real-time rtPCR amplicons | 38 |
| Phosphorothioated antisense oligonucleotides..... | 40 |

| | |
|--|------------|
| Results | 43 |
| Testing of amplicons for Real-time rtPCR..... | 43 |
| Design and production of mcm4 Geneblocs | 45 |
| Screening of MCM4 Geneblocs in IMR90 cells | 45 |
| Time course of Mcm4 depletion in IMR90 cells transfected with GB22174 | 51 |
| MCM4 protein synthesis is blocked in IMR90 cells transfected with GB 22174 | 54 |
| Other MCM proteins are degraded in IMR90 cells after MCM4 depletion..... | 54 |
| The subcellular localization of MCM proteins in IMR90 | 60 |
| The subcellular localization of MCM is altered in IMR90 cells depleted of MCM4 | 60 |
| Cellular proliferation of IMR90 cells is drastically reduced in the absence of MCM | 66 |
| The cell cycle distribution of MCM4 depleted IMR90 cells is drastically altered | 69 |
| IMR90 cells transfected with GB22174 fail to synthesize DNA..... | 73 |
| IMR90 cells depleted of MCM4 accumulate in G2 | 76 |
| Hoechst staining of IMR90 cells transfected with GB22174 | 77 |
| Discussion | 80 |
| Blocking mcm4 gene expression with antisense oligonucleotides | 82 |
| Blocking MCM4 synthesis destabilizes MCM complex subunits | 85 |
| The synthesis of MCM proteins is blocked in the absence of mcm4 gene expression | 87 |
| Chromatin Bound MCM proteins are solubilized in response to MCM4 degradation..... | 88 |
| Inactivation of the MCM complex stops proliferation and prevents completion of S phase | 90 |
| MCM proteins in human cells: future prospects | 94 |
| Summary | 96 |
| Zussamenfassung..... | 97 |
| Abbreviations | 98 |
| References | 99 |
| Eidesstaatliche Erklärung | 110 |

| | |
|------------------------------|------------|
| Lebenslauf | 111 |
| Acknowledgements..... | 115 |