

## Literaturverzeichnis

- (1) Klein S, Kinney J, Jeejeebhoy K, Alpers D, Hellerstein M, Murray M, Twomey P: Nutrition support in clinical practice: Review of published data and recommendations for future research directions. *American Journal of Clinical Nutrition* 1997; 66: 683-706
- (2) Keller U: Malnutrition im Spital. *Schweizer Medizinische Wochenschrift* 1996; 126, Suppl. 79: 9S-13S
- (3) Pirlich M, Schwenk A, Müller MJ, Ockenga J, Schmidt S, Schütz T et al. DGEM-Leitlinie Enterale Ernährung: Ernährungsstatus. *Akt. Ernähr. Med* 2003; 28(S1): 10-25
- (4) Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr* 2003; 22(4): 415-421
- (5) Detsky AS, McLaughlin JR, Baker JP, Johnston N; Whittaker S, Mendelsohn RA, Jeejeebhoy KN: What is subjective global assessment of nutritional status? *JPEN* 1987; 11: 8-13
- (6) Mensink GMB. Daten des Bundesgesundheitsveys 1998. Personal communication, Robert Koch Institut, Berlin, Germany
- (7) Norman K, Lochs H, Pirlich M. Malnutrition als prognostischer Faktor. *Chir Gastroenterol* 2004; 20:175-180
- (8) Coats KG, Morgan SL, Bartolucci AA, Weinsier RL. Hospital-associated malnutrition: a reevaluation 12 years later. *J AM Dietetic Ass* 1993; 93: 27-33
- (9) Mc Whirter JP & Pennington CR: Incidence and recognition of malnutrition in hospital. *British Medical Journal* 1994; 308: 945-948
- (10) Gariballa SE, Parker SG, Taub N, Castleden CM: Influence of nutritional status on clinical outcome after acute stroke. *American Journal of Clinical Nutrition* 1998; 68: 275-81

- (11) Martineau J, Bauer JD, Isenring E, Cohen S: Malnutrition determined by the patient-generated subjective global assessment is associated with poor outcomes in acute stroke patients. *Clinical Nutrition* 2005; 24: 1073-1077
- (12) Bistrain BR, Blackburn GL, Hallowell E, et al: Protein status of general surgical patients. *JAMA* 1974; 230: 858
- (13) Pham NV, Cox-Reijven PLM, Greve JW, Soeters PB: Application of subjective global assessment as a screening tool for malnutrition in surgical patients in Vietnam. *Clinical Nutrition* 2006; 25: 102-108
- (14) Hill GL, Blackett RL, Pickford I, et al: Malnutrition in surgical patients; an unrecognized problem. *Lancet* 1977; 1: 689
- (15) Bruun LI, Bosaeus I, Bergstad I, Nygaard K: Prevalence of malnutrition in surgical patients: evaluation of nutritional support and documentation. *Clinical Nutrition* 1999; 18(3):141-147
- (16) Tan YS, Nambiar R, Yo SL: Prevalence of protein calorie malnutrition in general surgical patients. *Annals Academy of Medicine* 1992; Vol.21 No.3: 334-337
- (17) Pirlich M, Luhmann N, Schütz T, Plauth M, Lochs H: Mangelernährung bei Klinikpatienten: Diagnostik und klinische Bedeutung. *Aktuelle Ernährungsmedizin* 1999; 24: 260-266
- (18) Waitzberg DL, Caiaffa WT, Correia MITD: Hospital Malnutrition: the Brazilian National Survey (IBRANUTRI): a study of 4000 patients. *Nutrition* 2001; 17: 573-580
- (19) Persson C, Sjöden PO, Glimelius B: The Swedish version of the patient generated subjective global assessment of nutritional status: gastrointestinal vs. urological cancers. *Clinical Nutrition* 1999; 18: 71-77
- (20) Cederholm T, Jägren C, Hellström K: Outcome of protein-energy malnutrition in elderly medical patients. *Am J Med* 1995; 98: 67-74
- (21) Sacks GS, Dearman K, Replogle WH, Cora VL, Meeks M, Canada T: Use of subjective global assessment to identify nutrition-associated complications and death in geriatric long-term care facility residents. *J Am Coll Nutr* 2000; 19: 570-577

- (22) Pirlich M, Lochs H: Nutrition in the elderly. *Best Pract Res Clin Gastroenterol* 2001; 15: 869-884
- (23) Robinson G, Goldstein M, Levine GM: Impact of nutritional status on DRG length of stay. *JPEN* 1987; 11: 49-51
- (24) Kyle UG, Kossovsky MP, Karsegard VL, Pichard C: Comparison of tools for nutritional assessment and screening at hospital admission: A population study. *Clinical Nutrition*, im Druck
- (25) Naber TNJ, Schermer T, de Bree A, Nusteling K, et al: Prevalence of malnutrition in nonsurgical hospitalized patients and its association with disease complications. *American Journal of Clinical nutrition* 1997; 66: 1232-9
- (26) Nakamura K, Moriyama Y, Kariyazono H et al. Influence of preoperative nutritional state on inflammatory response after surgery. *Nutrition* 1999; 15: 834-41
- (27) Tucker HN, Miguel SG: Cost containment through nutritional intervention. *Nutrition review* 1996; 54: 111-121
- (28) Odelli C, Burgess D, Bateman L, Hughes A, Ackland S, Gillies J, Collins CE: Nutrition support improves patient outcomes, treatment tolerance and admission characteristics in oesophageal cancer. *Clinical Oncology* 2005; 17: 639-645
- (29) Lawson JA, Lazarus R, Kelly JJ: Prevalence and prognostic significance of malnutrition in chronic renal insufficiency. *J Ren Nutr* 2001; 11: 16-22
- (30) Buzby GP, Knox LS, Crosby LO et al.: Study protocol: a randomized clinical trial of total parenteral nutrition in malnourished surgical patients. *Am J Clin Nutr* 1988; 47: 357-365
- (31) Durnin JVGA, Womersley J: Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 1974; 32: 77-97
- (32) Frisancho AR: New norms of upper limb fat and muscle areas for assessment of nutritional status. *Am J Clin Nutr* 1981; 34: 2540-2545
- (33) Kushner RF: Bioelectrical impedance analysis: a review of principles and applications. *J Am Coll Nutr* 1992; 11: 199-209

- (34) Kushner RF, Gudivaka R, Schoeller DA. Clinical characteristics influencing bioelektrical impedance analysis measurements. *Am J Clin Nutr Suppl.* 1996; 64: 423S-427S
- (35) Kushner RF, Schoeller DA: Estimation of total body water with bioelectrical impedance analysis. *Am J Clin Nutr* 1986; 44: 417-424
- (36) Pirlich M, Schütz T, Spachos T, Ertl S, Weiß M-L, Lochs H, Plauth M: Bioelectrical impedance analysis is a useful bedside technique to assess malnutrition in cirrhotic patients with and without ascites. *Hepatology* 2000; 32: 1208-1215
- (37) Süttmann U, Ockenga J, Selberg O, Hoogestraat L, Deicher H, Müller MJ: Incidence and prognostic value of malnutrition and wasting in human immunodeficiency virus-infected outpatients: *J Acquir Immune Defic Syndr Hum Retrovirol* 1995; 8: 239-246
- (38) Pirlich M, Plauth M, Lochs H: Bioelektrische Impedanzanalyse: Fehlerquellen und methodische Grenzen bei der klinischen Anwendung zur Analyse der Körperzusammensetzung. *Akt. Ernährungsmed.* 1999; 24: 81-90
- (39) Selberg O: The adjunctive value of routine biochemistry in nutritional assessment of hospitalized patients. *Clinical Nutrition* 2001; 20: 477-85.
- (40) Ott M, Lembcke B, Fischer H, et al: Early changes of body composition in human immunodeficiency virus-infected patients: tetrapolar body impedance analysis indicates significant malnutrition. *Am J Clin Nutr* 1993, 57:15-9
- (41) Bistran BR, Blackburn GL, Vitale J, et al: Prevalence of malnutrition in general medical patients. *JAMA* 1976; 235: 1567
- (42) Weinsier RL, Humber EM, Krumdieck CL, et al: Hospital malnutrition: a prospective evaluation of general medical patients during the course of hospitalization. *American Journal of Clinical Nutrition* 1979; 32: 418
- (43) Mullen J, Gertner M, Buzby P, Goodhart G, Rosato E: Implication of malnutrition in the surgical patient. *Arch Surg* 1979; 114: 121-125

- (44) Pirlich M, Schütz T, Kemps M, Luhmann N, Burmester GR, Baumann G, Plauth M, Lübke HJ, Lochs H: Prevalence of malnutrition in hospitalized medical patients: Impact of underlying disease. *Dig Dis* 2003; 21: 245-251
- (45) Gurski RR, Schirmer CC, Rosa AR, Bretano L: Nutritional assessment in patients with squamous cell carcinoma of the esophagus. *Hepatogastroenterology* 2003; 50(54): 1943-7
- (46) Belghiti J, Langonnet F, Boursstyn E, Fekete F: Surgical implications of malnutrition and immunodeficiency in patients with carcinoma of the oesophagus. *Br J Surg* 1983; 70: 339-341
- (47) Detsky A, Baker J, O'Rourke K, et al: Predicting nutrition-associated complications for patients undergoing gastrointestinal surgery. *Journal of Parenteral and Enteral Nutrition* 1987; 11: 440-446
- (48) Andreyev HJN, Norman AR, Oates J, Cunningham D: Why do patients with weight loss have a worse outcome when undergoing chemotherapy for gastrointestinal malignancies? *Eur J Cancer* 1998; Vol 34, No 4: 503-509
- (49) Müller MJ, Selberg O, Lautz HU, et al: Tumorkachexie: Pathophysiologische Grundlagen und ernährungsmedizinische Aufgabe. *Akt. Ernähr.-Med.* 1991; 16: 1-6
- (50) Figueiredo FA, Dickson ER, et al: Utility of standard parameters in detecting body cell mass depletion in patients with end-stage liver disease. *Liver Transplantation* 2000;5,5: 575-581
- (51) Abdullah MS, Wild G, et al: Cytokines and the malnutrition of chronic renal failure. *Miner Electrolyte Metab* 1997; 23: 237-242
- (52) Enia G, Sicuso C, Alati G, Zoccali C: Subjective global assessment of nutrition in dialysis patients. *Nephrol Dial Transplant* 1993; 8: 1094-1098
- (53) Arora NS, Rochester DF: Respiratory muscle strength and maximal voluntary ventilation in undernourished patients. *Am Rev Respir Dis* 1982; 126: 5
- (54) Anker SD, Ponikowski P, Varney S et al. Wasting as independent risk factor for mortality in chronic heart failure. *Lancet* 1997; 349: 1050-3
- (55) Grimble RF: Fasting in healthy individuals and adaptation to undernutrition during chronic disease. *Curr Opin Clin Nutr Metab Care* 1998; 1: 369-374

- (56) Geerling BJ, Badart-Smook A, Stockbrügger RW, Brummer RJM: Comprehensive nutritional status in patients with longstanding Crohn disease currently in remission. *Am J Clin Nutr* 1998; 67: 919-26
- (57) Bistrain BR: Role of the systemic inflammatory response syndrome in the development of protein-calorie malnutrition in ESRD. *American Journal of Kidney Diseases*, 1998; 32, 6(Suppl 4); S113-S117
- (58) Volkert D: Ernährungsprobleme in der Geriatrie – Mangelernährung bei geriatrischen Patienten. *Aktuelle Ernährungsmedizin* 1996; 21: 200-202
- (59) Tierney AJ. Undernutrition and elderly hospital patients: a review. *Journal of Advanced Nursing* 1996; 23: 228-246
- (60) Volkert D, Kruse W, Oster P, Schlierf G. Malnutrition in geriatric patients: diagnosis and prognostic significance of nutritional parameters. *Ann. Nutr. Metab* 1992; 36: 97-112.
- (61) White H, Pieper C, Schmader K. The association in weight change in Alzheimers disease with severity of disease and mortality: a longitudinal analysis. *J Am Geriatr Soc* 1998; 46: 1223-7
- (62) DeWys WD, Begg C, Lavin PT, et al. Prognostic effect of weight loss prior to chemotherapy in cancer patients. *Am J Med* 1980; 69: 491-7
- (63) Pirlich M, Schütz T, Kemps M, Luhmann N, Minko N, Lübke HJ, Rossnagel K, Willich SN, Lochs H. Social risk factors for hospital malnutrition. *Nutrition* 2005; 20: 295-300
- (64) Ezzati M, Lopez AD, Rodgers A, Hoorn SV, Murray CJL et al. Selected major risk factors and global and regional burden of disease. *Lancet* 2002; 360: 1347-60
- (65) Langnase K, Muller MJ. Nutrition and health in an adult urban homeless population in Germany. *Public Health Nutr* 2001; 4: 805-811
- (66) Reuben DB, Greendale GA, Harrison GG. Nutrition screening in older persons. *J Am Geriatr Soc* 1995; 43: 415-25
- (67) Pirlich M, Schütz T, Gastell S, Lochs H: Malnutrition affects long-term prognosis in hospitalized patients. *Gastroenterology* 2002; 122(suppl 1): A 363
- (68) Kyle UG, Pirlich M, Schuetz T, Lochs H, Pichard C. Is nutritional depletion by Nutritional Risk Index (NRI) associated with increased length of hospital stay (LOS)? *J Parent Enter Nutri* 2004; 28: 99-104

- (69) Barton AD, Beigg CL, MacDonald IA, Allison SP. High food wastage and low nutritional intakes in hospital patients. *Clin Nutr* 2000;19(6): 445-449
- (70) Barton AD, Beigg CL, MacDonald IA, Allison SP. A recipe for improving food intaks in elderly hospitalized patients. *Clin Nutr* 2000; 19(6): 445-449
- (71) Tremblay A, Bandi V. Impact of body mass index on outcomes following critical care. *Chest* 2003; 123: 1202-1207
- (72) Nowbar S, Burkart KM, Gonzales R, Fedorowicz A, Gozansky WS, Gaudio JC, Taylor MRG, Zwillich CW. Obesity-associated hypoventilation in hospitalized patients: prevalence, effects and outcome. *Am J Med.* 2004; 116:1-7
- (73) Nightingale JMD, Reeves J. Knowledge about the assessment and management of undernutrition: a pilot questionnaire in a UK teaching hospital. *Clin Nutr* 1999; 18(1): 23-27
- (74) Reilly JJ, Hull SF, Albert N, Waller A, Bringardener S. Economic impact of malnutrition: a model system for hospitalized patients. *JPEN* 1988; 12: 371-376
- (75) Edington J, Boorman J, Durrant ER et al. Prevalence of malnutrition on admission to four hospitals in England. *Clin Nutr* 2000; 19(3): 191-195
- (76) Committee of Experts on Nutrition FsaCP. Food and nutritional care in hospitals: how to prevent undernutrition. Strasburg: Council of Europe Publishing, 2002