

6 LITERATURVERZEICHNIS

- 1 Ammann P, Bourrin S, Bonjour JP, et al. Protein undernutrition-induced bone loss is associated with decreased IGF-I levels and oestrogen deficiency. *Bone Miner Res*, 2000. 15(4): S. 683-669.
- 2 An HS, Simpson JM, Stepany J. Comparison between allograft plus demineralized bone matrix versus autograft in anterior cervical spine fusion: A prospective multicenter study. *Spine*, 1995. 20(20): S. 2211-2216.
- 3 Aronson N, Filtzer DL, Bagan M. Anterior cervical fusion by the Smith-Robinson approach. *Neurosurg*, 1968. 29: S. 397-404.
- 4 Aspenberg P, Turek T. BMP-2 for intramuscular bone induction. Effect of squirrel monkeys is dependent on implantation site. *Acta Orthop Scand*, 1996. 67: S. 3-6.
- 5 Bagby G. Arthrodesis by a distraction-compression method using a stainless steel implant. *Orthopedics*, 1988. 11: S. 931-934.
- 6 Bailey AW, Badgley CE. Stabilization of the cervical spine by anterior fusion. *J Bone Joint Surg*, 1960. 42: S. 565-594.
- 7 Beck L, Amento E, Xu Y, Deguzman, et al. TGF- beta 1 induces bone closure of skull defects-Temporal dynamics of bone formation in defects exposed to rh TGF-beta 1. *Bone Min Res*, 1993. 8: S. 753-761.
- 8 Beck LS, Ammann AJ, Aufdemorte TB, et al. In vivo induction of bone by recombinant human transforming growth factor beta 1. *Bone Miner Res*, 1991. 6: S. 961- 968.
- 9 Blumenthal SL, Gill K. Can lumbar spine radiographs accurately determine fusion in postoperative patients? Correlation of routine radiographs with a second surgical look at lumbar fusions. *Spine*, 1993. 18: S 1186-1189.
- 10 Boden SD, Martin GJ Jr, Horton WC, et al. Laparoscopic anterior spinal arthrodesis with rh BMP-2 in a titanium interbody threaded cage. *Spinal Disord*, 1998. 11: S. 95-101.

- 11 Boden SD, Martin GJ Jr, Morona MA, et al. Posterolateral lumbar intertransverse process spine arthrodesis with recombinant human bone morphogenetic protein 2/ hydroxyapatite-tricalcium phosphate after laminectomy in the nonhuman primate. *Spine*, 1999. 24: S. 1179- 1185.
- 12 Boden SD, Zdeblick TA, Sandhu HS, et al. The use of rhBMP-2 in interbody fusion cages. Definitive evidence of osteoinduction in humans: a preliminary report. *Spine*, 2000. 25: S. 376- 381.
- 13 Bohlmann HH, Emery SE, Goodfellow DB, et al. Robinson anterior cervical discectomy and arthrodesis for cervical radiculopathy. *Bone Joint Surg*, 1993. 75A: S. 1298- 1307.
- 14 Bostworth DM. Technics of spinal fusion: Pseudarthrosis and method of repair. AAOS Instructional Course Lecture, 1948. 5: S. 295-313.
- 15 Brill T. Prä- und postoperative klinische Untersuchungen am Versuchstier. *Vet. Med. Diss. München* 1991.
- 16 Brodke DS, Zdeblick TA. Modified Smith- Robinson procedure for anterior cervical discectomy and fusion. *Spine*, 1992. 17(10): S. 427- 430.
- 17 Brown MD, Malinin TI, Brown MD. A röntgenographic evaluation of frozen allografts versus autografts in anterior cervical fusion. *Clin Orthop*, 1976. 119: S. 231- 236.
- 18 Brown A. Slosar P. Reynolds J. et al. Paired BAK proximity cages versus standard BAK: subsidence and clinical failures. North American Spine Society. 1999, (Oct): S.20-24.
- 19 Burkus J, Kenneh MD, Dorchak JD, et al. Radiographic assessment of interbody fusion using recombinant human bone morphogenetic protein type 2. *Spine*, 2003. 28(4): S. 372-377.
- 20 Burt DW. Evolutionary grouping of the transforming growth factor-beta superfamily. *Biochem Biophys Res Commun*. 1992, 184: S. 590-595.
- 21 Chipault A. Chirurgie Operatoire du Systeme Nerveux Reuff et Circe Ed. Paris, 1895.

- 22 Cloward RW. The anterior approach for removal of ruptured cervical discs. Neurosurg. 1958. 15: S. 602-617.
- 23 Cunningham BW, Kanayama M, Parker LM, et al. Osteogenic protein versus autologous interbody arthrodesis in the sheep thoracic spine. A comparative endoscopic study using the Bagby and Kuslich interbody fusion device. Spine, 1999. 24: S 509-518.
- 24 David SM, Gruber HE, Mayer RA Jr, et al. Lumbar spinal fusing using recombinant human bone morphogenetic protein in the canine. A comparison of three dosages and two carriers. Spine, 1999. 24: S.1973-1979.
- 25 De Palma AF, Rothmann LH, Lewinnek GE, et al. Anterior interbody fusion for severe cervical disc degeneration. Surg Gynecol Obstet, 1972. 134: S. 755-758.
- 26 Dereymaker A, Chosez JP, Kennes R. Les Traitment Chirurgical de la Discopathie Cervicale. Clin Orthop. 1965. 40: S. 113-122.
- 27 Eitel F, Seiler H, Schweiberer L. Morphologic examination of animalexperiment results: comparison with regeneration of the human bone-structure. J Research methods (author transl), Unfallheilkunde 1981. 84(6): S 250-254.
- 28 Emery SE, Bolesta MJ, Banks MA, et al. Robinson anterior cervical fusion: Comparison of the standard and modified techniques. Spine, 1994. 19: S. 660- 663.
- 29 Fischgrund JS, James SB, Chabot MC, et al. Augmentation of autograft using rhBMP-2 and different carrier media in the canine spinal fusion model. Spinal Disord, 1997. 10: S. 467- 472.
- 30 Fujimoto A, Tanizawa T, Nishida S Yamamoto N, et al. Local effects of transforming growth factor- beta 1 on rat calvaria: changes depending on the dose and the injection site. Bone Miner Metab, 1999. 17: S. 11-17.

- 31 Gerhart TN, Kirker- Head CA, Kriz MJ, et al. Healing segmental femoral defects in sheep using recombinant human bone morphogenetic protein. Clin Orthop, 1993. 293: S. 317-326.
- 32 Harda BE. Wiring of the vertebrae as a means of immobilization in fractures and Pott's disease. Medical Times an Register 22. 423, Reprint: Clin Orthop, 1975. 112: S 4
- 33 Hecht BP, Fischgrund JS, Herkowitz HN, et al. The use of recombinant human bone morphogenetic protein-2 (rhBMP-2) to promote spinal fusion in a nonhuman primate anterior interbody fusion model. Spine, 1999. S. 629-636.
- 34 Hermann R, Schmidmaier G, Markl B, et al. Antithrombogenic coating of stents using a biodegradable drug delivery technology. Thromb Haemost, 1999. 82: S. 51-57.
- 35 Hock J, Centrella M, Canalis E. Insulin like growth faktor I has independent effcts on bone matrix formation and cell replication. Endocrinology, 1988. 122: S. 254- 260.
- 36 Hoshi K, Amizuka N, Sakou T, et al. Fibroblasts of spinal ligaments pathologically differentiate into chondrocytes induced by recombinant human bone morphogenetic protein-2: morphological examinations for ossification of spinal ligaments. Bone, 1997. 21: S.155-162.
- 37 Irving JT, Ph. D. M.D., LeBolt SA, et al: Ectopic bone formation and aging. Clin Orthop, 1981. 154: S. 249-253.
- 38 Isgaard J, Nilson A, Lindahl A, et al. Effects of local administration of GH and IGF-I on longitudinal bone growth in rats. Physiol, 1986. 250: S. 367-372.
- 39 Ishidou Y, Kitajima I, Obama H, et al. Enhanced expression of type I bone morphogenetic proteins during bone formation. J Bone Miner Res, 1995. 10: S.1651-1659.

- 40 Itoh H, Ebara S, Kamimura M, et al. Experimental spinal fusion with use of recombinant human bone morphogenetic protein 2. Spine, 1999. 24: S.1402-1405.
- 41 Jost B, Cripton PA, Lund T, et al. Compressive strength of interbody cages in the lumbar spine: the effect of cage shape, posterior instrumentation and bone density. Eur Spine J, 1998. 7: S 132-141.
- 42 Joyce M, Jingushi S, Bolander M. Transforming growth factor in the regulation of fracture repair. Orthop Clin. 1990, 21A: S.199-209.
- 43 Kandziora F, Bail H, Schmidmaier G, et al. Bone morphogenetic protein-2 by a poly-(D,L-lactide) coated interbody cage: in vivo results of a new carrier for growth factors. J Neurosurg Spine, 2002. 97(1): S 40-48.
- 44 Kandziora F, Kerschbaumer F, Mittlmeier T. Stage related surgery for cervical spine instability in rheumatoid arthritis. Euro Spine, 1999. 8: S. 371-381.
- 45 Kandziora F, Pflugmacher R., Scholz M, et al. [Experimental fusion of the sheep cervical spine. Part I: Effect of cage design on interbody fusion]. Chirurg, 2002. 73: S 909-917.
- 46 Kandziora F, Pflugmacher R, Scholz M, et al. Comparison between sheep and human cervical spines: an anatomic, radiographic, bone mineral density, and biomechanical study. Spine, 2001. 26: S.1028-1037.
- 47 Kandziora F, Pflugmacher R, Scholz M, et al. Comparison between BMP-2 and combined IGF-I/TGF- β 1 application in a sheep cervical spine fusion model. Europ Spine J, 2002. 11: S. 482-493.
- 48 Kandziora F, Schmidmaier G, Bail H, et al. IGF-I and TGF- β 1 application by a poly-(D,L-lactide) coated interbody cage promotes fusion in the sheep cervical spine. Spine, 2002. 27: S, 1710-1723.
- 49 Kawamura M, Urist MR. Human Fibrin in a physiologic delivery System for bone morphogenetic protein. Clin. Orthop, 1988. 235: S. 301-310.

- 50 Kon T, Yamazaki M, Tagawa M, et al. Bone morphogenetic protein- 2 stimulates differentiation of cultured spinal ligament cells from patients with ossification of the posterior longitudinal ligament. *Calcif Tissue Int*, 1997. 60: S. 291- 296.
- 51 Kummer B. Biomechanical problems of upright posture. *Anat Anz*, 1992, 174: S 33-39.
- 52 Kuslich SD, Ulstrom CL, Griffith SL, et al. The Bagby and Kuslich method of lumbar interbody fusion. History, techniques, and 2- year follow -up results of a United States prospective, multicenter trial. *Spine*. 1998. 23: S.1267-1278.
- 53 Lacroix P. Organisers of the growth of bone. *J Bone Joint Surg*. 1947, 29: S. 292-296.
- 54 Larsen JM, Rimoldi RL, Capen DA, et al. Assessment of pseudarthrosis in pedicle screw fusion: a prospective study comparing plain radiographs, flexion/extension radiographs, CT scanning, and bone scintigraphy with operative findings. *J Spinal Disord*, 1996. 9: S. 117-120.
- 55 Laurencin C, Lane JM. Poly (lactide acid) AND Poly (glycolid acid): Orthopedic surgery applications. In: Bone formation and repair. Brighton C, Frielaender G, Lane MJ. (eds) Rosemont. American Academy of Orthopedic Surgeons, 1994. AQ1: S. 325-339.
- 56 Levander G. A study of bone regeneration. *Surg Gynec Obstetr*, 1938. 67: S. 705-717.
- 57 Lind M. Growth factor stimulation on bone healing: Effects on osteoblasts, osteotomies, and implants fixations. *Acta Orthop Scand Suppl*, 1998. 283: S. 2-37.
- 58 Lind M, Schumacker B, Soballe K, et al. Transforming growth factor- beta enhances fracture healing in rabbit tibiae. *Acta Orthop Scand*, 1993. 64: S. 553-556.

- 59 Lund T, Oxland TR, Jost B, et al. Interbody cage stabilisation in the lumbar spine: biomechanical evaluation of cage design, postreior instrumentation and bone density. *Joint Surg Br*, 1998. 80: S. 351-359.
- 60 Martin GJ Jr, Boden SD, Marone MA, et al. Posterolateral intertransverse process spinal arthrodesis with rhBMP-2 in a nonhuman primate: importand lessons learned regarding dose, carrier, and safety. *Spinal Disord*, 1999. 12: S.179-186.
- 61 McAfee PC. Interbody fusion cages in reconstructive operations on the spine. *J Bone Joint Surg Am*, 1999. 81: S. 859-880.
- 62 McAfee PC, Boden SD, Brantigan JW, et al. Symposium: a critical discrepancy-a criteria of successful arthrodesis following interbody spinal fusions. *Spine*, 2001. 26: S. 320-334.
- 63 McAffe PC, Regan JJ, Farey ID, et al. The biomechanical and histomorphometric properties of anterior lumbar fusions: a canine model. *J Spinal Disord*, 1988. S 101-110.
- 64 McCarthy TL, Centrella M, Canalis E. Insulin- like growth factor (IGF) and bone. *Connect tissue Res*, 1989. 20: S. 277-282.
- 65 Meyer RA Jr, Gruber HE, Howard BA, et al. Safety of recombinant human bone morphogenetic protein-2 after spinal laminectomy in the dog. *Spine*, 1999. 24: S.747-754.
- 66 Mimatsu K, Kishi S, Hashizume Y. Experimental chronic compressions on the spinal cord of the rabbit by ectopic bone formation in the ligamentum flavum with bone morphogenetic protein. *Spinal Cord*, 1997. 35: S. 740-746.
- 67 Mohan S. Insulin like growth factor binding proteins in bone cell regulation. *Growth Regul*. 1993, 3: S.67-70.
- 68 Mohan S, Baylink D. Bone growth factors. *Clin Orthop Rel Res*, 1991. 263: S. 30-48.

- 69 Mosdal C. Cervical osteochondrosis and disc herniation: Eighteen years use of interbody fusion by Clowards technique in 755 cases. *Acta Neurochir*, 1984. 70: S. 207-225.
- 70 Nakamura K, Kurokawa T, Kato T, et al. Lokal applikation of basic fibroblast growth factor into the bone increases bone mass at the applied site in rabbits. *Arch of Orthop and Trauma Surg*, 1996. 115: S. 344-346.
- 71 Newton PO, Cardelia MJ, Farnsworth CL, et al. A biomechanical comparison of open and thoracoscopic anterior spinal release in a goat model. *Spine*, 1998. 23: S. 530- 536.
- 72 Nielson H, Isgaard J, Lindahl A, et al. Effects of unilateral arterial infinfusion of GH and IGF-I on tibial longitudinal bone growth in hypophysectomized rats. *Calcif tissue Int*, 1987. 40: S. 91-96.
- 73 Oikarinen J. Experimental spine fusion with decalcified bone matrix and deep- frozen allogenic bone in rabbits. *Clin Orthop*, 1982. 273: S. 210-218.
- 74 Pettersson K, Hildingsson T, Fagerlund MB. Disc pathology after whiplash injury. *Spine*, 1997. 22: S.283-288.
- 75 Pfeilschifter J, Oechsner M, Naumann A, et al. Stimulation of bone matrix apposition in vitro by local growth factors: a comparison between Insulin-like growth factor I, platetderived growth factor and transforming growth factor beta. *Endokrinology*, 1990. 127: S. 69- 75.
- 76 Porchet F, Jaques B. Unusual complications at iliac crest bone graft donor site: experience with two cases. *Neurosurgery*, 1996. 39: S. 856-859.
- 77 Raschke M, Bail M, Kolbeck S, et al. Recombinant growth hormone accelerates bone regenerate consolidation in distraction osteogenesis. *Chirurg*, 2000. 71(9): S. 1009-1015.
- 78 Reddi AH. Age- depending decline in extracellular matrix- induced local bone differentiation. *Isr J med Sci*, 1985. 21: S. 312-313.

- 79 Reddi AH, Wietroub S, Muthukumaran N. Biologic principles of bone induction. Orthopedics clinics of North America, 1987. 18: S. 207-218.
- 80 Riley LH, Robinson RA, Johnson KA, et al. The results of anterior interbody fusion of the cervical spine. Neurosurg, 1969. 30: S.127-133.
- 81 Rish BL, Mc Fadden JT, Penix JO. Anterior cervical fusion using homologous bone grafts: A comparative study. Surg Neurol, 1975. 5: S. 119-121.
- 82 Roberts A, Sporn M, Bolander M. Transforming growth factor- beta and the initiation of chondrogenesis in the rat femur. Cell Biol, 1990. 110: S. 2195- 2207.
- 83 Robey PG, Young MF, Flanders KC, et al. Osteoblasts synthesize and respond to transforming growth factor-type β (TGF- β) in vitro. J Cell Biol, 1987. 105: S. 457-463.
- 84 Robinson R, Smith G. Antero- lateral cervicel disc removal and interbody fusion for cervical disc syndrome. Bull John Hopkins Hosp, 1955. 96: S. 223-224.
- 85 Robinson RA, Walker AE, Ferlic DC, et al. The results of anterior interbody fusion of the cervical spine. Bone Joint Surg, 1962. 44: S.1569- 1587.
- 86 Saitoh H, Takata T, Nikai H, et al. Effects of polylactid acid on osteoinduction of demineralized bone: preliminary study of the usefulness of polylactid acid as a carrier of bone morphogenetic protein. Oral Rehabil, 1994. 21: S. 431-438.
- 87 Sampath TK, Rashka KE, Doctor JS, et al. Drosophila TGF- β superfamily proteins induce enchondral bone formation in mammals. Proc Natl Acad Sci USA, 1993. 90: S. 6004-6008.
- 88 Sandhu HS, Kanim LE, Kabo JM, Toth et al. Effective dosis of recombinant human bone morphogenetic protein-2 in experimental spine fusion. Spine, 1996. 21: S. 2115- 2122.

- 89 Sandhu HS, Toth JM, Diwan A, et al. Histologic evaluation of the efficacy of rhBMP-2 compared with autograft bone in sheep spinal anterior interbody fusion. Spine, 2002. 27(6): S. 567-575.
- 90 Santos E, Russel KM, Fraser RD. Radiologic assessment of interbody fusion using carbon fiber cages. Spine, 2003. 28: S. 997-1001.
- 91 Savolainen S, Usenius JP, Hernesniemi J. Iliac crest versus artificial bone grafts in 250 cervical fusions. Acta Neurochir, 1994. 129: S. 54-57.
- 92 Schimandle JH, Boden SD, Hutton WC. Experimental spinal fusion with recombinant human bone morphogenetic protein- 2. Spine, 1995. 20: S. 1326- 1337.
- 93 Schmidmaier G, Wildemann B, Sternberger A et al. Biodegradable poly (D,L-lactide) coating of implants for continuous release of growth factors. Biomed Mat Res, 2001. 58: S. 449- 455.
- 94 Schmidmaier G, Wildemann B, Sternberger A, et al. Local application of growth factors (insulin-like growth factor-1 and transforming growth factor-beta 1) from a biodegradable poly (D,L-lactide) coating of osteosynthetic implants accelerates fracture healing in rats. Bone, 2001. 28: S. 341-350.
- 95 Schnaa CL, Freese A, Weil RJ, et al. Analysis of harvest morbidity and radiographic outcome using autograft for anterior cervical fusion. Spine, 1997. 22: S. 2222- 2227.
- 96 Schult E K, Clark CR, Goel VK. Kinematics of the cervical spine following discectomy and stabilization. Spine, 1989. 14: S. 1116-1121.
- 97 Schumacher B, Albrechtsen J, Keller J, et al. Periostal insulin- like growth factor I and bone formation- Changes during tibial lengthening in rabbits. Acta Orthop Scand, 1998. 67: S. 237-241.
- 98 Shah RR, Mohammed S, Saifuddin A, et al. Comparison of plain radiographs with CT scan to evaluate interbody fusion following the use of titanium interbody cages and transpedicular Instrumentation. Eur Spine J, 2003. 12: S. 378-385.

- 99 Simmers BN, Eisenstein SM. Donor site pain from ilium: A complication of lumbar spine fusion. Bone Joint Surg, 1989. 71: S. 667-680.
- 100 Sorensen TS, Sorensen AI, Merser S. Rapid release of gentamicin from collagen sponge. In vitro comparison plastic beads. Acta Orthop Scand, 1990. 61: S. 353-356.
- 101 Steinbrech DS, Behara BJ, Rowe NM, et al. Gene expression of TGF-beta, TGF-beta receptor, and extracellular matrix proteins during membranous bone healing in rats. Plast Reconstr Surg, 2000. 105: S. 2028-2038.
- 102 Takaoka K, Koezuka H. Teleopeptide-depleted bovine skin collagen as a carrier for bone morphogenetic protein. Orthop Res, 1991. 9: S. 902-907.
- 103 Takaoka K, Ono K, Amitani K, et al. Solubilization and concentration of a bone-inducing substance from a murine osteosarcoma. Clin. Orthop. Rel. Res, 1980. 148: S. 247-280.
- 104 Tencer AF, Hampton D, Eddy S. Biomechanical properties of threaded inserts for lumbar interbody spinal fusion. Spine, 1995. 20: S. 2408-2414.
- 105 Terrell TG, Working PK, Chow CP, et al. Pathology of recombinant human growth factor-beta 1 in rats and rabbits. Int Rev Exp Pathol, 1993. 34B: S. 43-67.
- 106 Thaller S, Hoyt J, Tesluk H, et al. The effect of insulin like growth factor- 1 on calvarial sutures in a Sprague- Dawley rat. Craniofac Surg, 1993. 4: S. 35-39.
- 107 Thielemann FW, Feller AM, Schmidt K. Defektersatz mit Osteogeninhaltiger Gelantine (OCG). Orthop, 1984. 122(6): S. 843-847.
- 108 Trippel S, Coutts R, Einhorn T, et al. Growth factors as therapeutic agents. Bone Joint Surg, 1996. 78A: S.1272-1286.
- 109 Tscherne H, Illgner A. Die ventrale interkorporelle Spondylodese der Halswirbelsäule. Operat Orthop Traumatolog, 1991. 3: S.147-157.

- 110 Tullberg T. Failure of a carbon fiber Implantat. A case report. Spine, 1998. 23: S.1804-1806.
- 111 Ulrich C, Woersdoerfer O, Klaff R, et al. Biomechanics of fixation systems to the cervical spine. Spine, 1991. 16: S. 4- 9.
- 112 Urist MR. Bone formation by autoinduction. Science, 1965. 159: S. 893-899.
- 113 Urist MR. Heterotopic bone formation. Fundamental and clinical bone physiology. JB Lippincott, Philadelphia USA, 1980. S.369-393.
- 114 Urist MR, Huo YK, Brownell AG, et al. Purification of bovine bone morphogenetic protein by hydroxyapatite chromatography. Proc Nat Acad Sci, 1984. 81: S. 371-375.
- 115 Urist MR, Lietze A, Dawson E. Beta- Tricalcium phosphat delivery system for bone morphogenetic protein. Clin. Orthop, 1984. 187: S. 277-280.
- 116 Urist MR, Mikulski A, Lietze A. Solubilized and insolubilized bone morphogenetic protein. Proc Natl Acad Sci USA, 1979. 76: S.1828-1832.
- 117 Wang EA, Rosen V, D'Alessandro JS. Recombinant human bone morphogenetic protein induces bone formation. Proc Natl Acad Sci USA, 1990. 87: S. 2220-2224.
- 118 Weiner BK, Fraser RD. Spine update. Lumbar interbody cages. Spine, 1998. 23: S. 634-640.
- 119 White AA, Southwick WO, Deponte RJ, et al. Relief of pain by anterior cervical- spine fusion for spondylodesis. A report of sixty- five patients. Bone Joint Surg (Am), 1973. 23: S. 634-640.
- 120 Wilke HJ, Kettler A, et al. Anatomy of the sheep spine and its comparison to the human spine. Anat Rec, 1997. 247(4): S. 542-555.
- 121 Wilke HJ, Kettler A, Claes LE. Are sheep spines a valid model for human spines ? Spine, 1997. 22: S. 2365-2374.

- 122 Wissing H, Stürmer KM, Breidenstein G. Die Wertigkeit verschiedener Versuchstierspezies für experimentelle Untersuchungen am Knochen. Hefte zur Unfallheikunde, 1990. 212: S. 479-488.
- 123 Wilton P. Treatment with recombinant human insulin-like growth factor I of children with growth hormone receptor deficiency. Acta Paediatr Suppl, 1992. 383: S.137-142.
- 124 Yamada Y, Harada A, Hosoi T, et al. Association of transforming growth factor beta I genotype with therapeutic response to active vitamin D for postmenopausal osteoporosis. Bone Miner Res, 2000. 15: S. 415- 420.
- 125 Yasko AW, Lane JM, Fellinger EJ, et al. The healing of segmental bone defects, induced by recombinant human bone morphogenetic protein (rhBMP-2). Bone Joint Surg (Am), 1992. 74: S. 659-670.
- 126 ZdeblickTA, Cooke ME, Kunz DN, Wilson et al. Anterior cervical discectomy and fusion using a porous hydroxyapatite bone graft substitute. Spine, 1994. 19: S. 2348-2357.
- 127 Zdeblick TA, Cooke ME, Wilson D, et al. Anterior cervical discectomy, fusion and plating: A comparative animal study. Spine, 1993. 18: S.1974-1983.
- 128 Zdeblick TA, Ganajem AJ, Rapoff AJ, et al. Cervical interbody fusion cages. An animal model with and without bone morphogenetic protein. Spine, 1998. 23: S. 758-765.
- 129 Zdeblick TA, Wilson D, Cooke ME, et al. Anterior cervical discectomy and fusion: A comparisons of techniques in an animal model. Spine, 1992. 17 (Suppl): S. 418- 426.
- 130 Zegzula HD, Buck DC, Brekke J, et al. Bone formation with use of rhBMP-2 (recombinant human bone morphogenetic protein-2). Bone Joint Surg Am, 1997. 79: S.1778-1790.