

8. Literaturverzeichnis

1. Wasinger, V. C. et al. Progress with gene-product mapping of the Mollicutes: Mycoplasma genitalium. *Electrophoresis* 16, 1090-1094 (1995).
2. Kahn, P. From genome to proteome: looking at a cell's proteins. *Science* 270, 369-370 (1995).
3. Fleischmann, R. D. et al. Whole-genome random sequencing and assembly of Haemophilus influenzae Rd. *Science* 269, 496-512 (1995).
4. Goffeau, A. et al. Life with 6000 genes. *Science* 274, 546, 563-567 (1996).
5. Consortium, C. e. S. Genome sequence of the nematode C-elegans: A platform for investigating biology. *Science*. 282, 2012-2018 (1998).
6. Adams, M. D. et al. The genome sequence of Drosophila melanogaster. *Science* 287, 2185-2195 (2000).
7. Venter, J. C. et al. The sequence of the human genome. *Science* 291, 1304-51 (2001).
8. Lander, E. S. et al. Initial sequencing and analysis of the human genome. *Nature* 409, 860-921 (2001).
9. Initiative, T. A. G. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408, 796-815 (2000).
10. Goff, S. A. et al. A draft sequence of the rice genome (*Oryza sativa* L. ssp japonica). *Science* 296, 92-100 (2002).
11. Yu, J. et al. A draft sequence of the rice genome (*Oryza sativa* L. ssp indica). *Science* 296, 79-92 (2002).
12. Anderson, N. G. & Anderson, N. L. Photo/essay. The human protein index. *JAMA* 246, 2620-2621 (1981).
13. Anderson, N. G. & Anderson, L. The Human Protein Index. *Clin. Chem.* 28, 739-748 (1982).
14. Cohen, S. N., Chang, A. C., Boyer, H. W. & Helling, R. B. Construction of biologically functional bacterial plasmids in vitro. *Proc Natl Acad Sci U S A* 70, 3240-3244 (1973).
15. Baltimore, D. RNA-dependent DNA polymerase in virions of RNA tumour viruses. *Nature* 226, 1209-1211 (1970).
16. Temin, H. M. & Mizutani, S. RNA-dependent DNA polymerase in virions of Rous sarcoma virus. *Nature* 226, 1211-3 (1970).
17. Bank, A., Terada, M., Metafora, S., Dow, L. & Marks, P. A. In vitro synthesis of DNA components of human genes for globins. *Nat New Biol* 235, 167-9 (1972).
18. Ross, J., Aviv, H., Scolnick, E. & Leder, P. In vitro synthesis of DNA complementary to purified rabbit globin mRNA (RNA-dependent DNA polymerase-reticulocyte-hemoglobin-density gradient centrifugation-oligo(dT) primer). *Proc Natl Acad Sci U S A* 69, 264-8 (1972).
19. Verma, I. M., Temple, G. F., Fan, H. & Baltimore, D. In vitro synthesis of DNA complementary to rabbit reticulocyte 10S RNA. *Nat New Biol* 235, 163-7 (1972).
20. Rougeon, F., Kourilsky, P. & Mach, B. Insertion of a rabbit beta-globin gene sequence into an E. coli plasmid. *Nucleic Acids Res* 2, 2365-78 (1975).
21. Rougeon, F. & Mach, B. Stepwise biosynthesis in vitro of globin genes from globin mRNA by DNA polymerase of avian myeloblastosis virus. *Proc Natl Acad Sci U S A* 73, 3418-22 (1976).
22. Rabbitts, T. H. Bacterial cloning of plasmids carrying copies of rabbit globin messenger RNA. *Nature* 260, 221-5 (1976).
23. Maniatis, T., Kee, S. G., Efstratiadis, A. & Kafatos, F. C. Amplification and characterization of a beta-globin gene synthesized in vitro. *Cell* 8, 163-82 (1976).
24. Alwine, J. C. et al. Detection of specific RNAs or specific fragments of DNA by fractionation in gels and transfer to diazobenzyloxymethyl paper. *Methods Enzymol* 68, 220-42 (1979).
25. Saiki, R. K. et al. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230, 1350-4 (1985).
26. Mullis, K. B. & Faloona, F. A. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol* 155, 335-50 (1987).
27. Dunne, P. W., Wang, S. W., Ashizawa, T., Perryman, M. B. & Epstein, H. F. cDNA surveying of specific tissue expression of human chromosome 19 sequences. *Genomics* 14, 263-9 (1992).

28. Maier, E., Roest Crolius, H. & Lehrach, H. Hybridisation techniques on gridded high density DNA and in situ colony filters based on fluorescence detection. *Nucleic Acids Res* 22, 3423-4 (1994).
29. Schena, M., Shalon, D., Davis, R. W. & Brown, P. O. Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270, 467-470 (1995).
30. Lockhart, D. J. et al. Expression monitoring by hybridization to high-density oligonucleotide arrays. *Nat Biotechnol* 14, 1675-80. (1996).
31. Lashkari, D. A. et al. Yeast microarrays for genome wide parallel genetic and gene expression analysis. *Proc Natl Acad Sci USA* 94, 13057-13062 (1997).
32. Anderson, L. & Seilhamer, J. A comparison of selected mRNA and protein abundances in human liver. *Electrophoresis*, 533-577 (1997).
33. Haynes, P. A., Gygi, S. P., Figeys, D. & Aebersold, R. Proteome analysis: biological assay or data archive? *Electrophoresis* 19, 1862-71 (1998).
34. Gygi, S. P., Rochon, Y., Franza, B. R. & Aebersold, R. Correlation between protein and mRNA abundance in yeast. *Mol Cell Biol* 19, 1720-1730 (1999).
35. Futcher, B., Latter, G. I., Monardo, P., McLaughlin, C. S. & Garrels, J. I. A sampling of the yeast proteome. *Mol Cell Biol* 19, 7357-68. (1999).
36. Wilkins, M. R. et al. Progress with proteome projects: why all proteins expressed by a genome should be identified and how to do it. [Review]. *Biotechnology & Genetic Engineering Reviews* 13, 19-50 (1996).
37. Smithies, O. & Poulik, M. 2-Dimensional electrophoresis of serum proteins. *Nature* 177, 1033 (1956).
38. Raymond, S. & Weintraub, L. Acrylamide gel as a supporting medium for zone electrophoresis. *Science* 130, 711-713 (1959).
39. Shapiro, A. L., Vinuela, E. & Maizel, J. V. Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels. *Biochemical & Biophysical Research Communications* 28, 815-20 (1967).
40. Laemmli, U. K. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-55 (1970).
41. Svensson, H. Isoelectric fractionation, analysis, and characterisation of ampholytes in natural pH gra-dients. I. The differential equation of solute concentra-tions at a steady state and its solution for simple cases. *Acta Chemica Scandinavica* 15, 325-341 (1961).
42. Svensson, H. Isoelectric fractionation, analysis, and characterisation of ampholytes in natural pH gradients. II. Buffering capacity and conductance of isotonic ampholytes. *Acta Chemica Scandinavica* 16, 456-466 (1962).
43. Vesterberg, O. & Svensson, H. Isoelectric fractionation, analysis, and characterization of ampholytes in natural pH gradients. IV. Further studies on the resolving power in connection with separation of myoglobins. *Acta Chemica Scandinavica* 20, 820-34. (1966).
44. Vesterberg, O. Synthesis and isoelectric fraction of carrier ampholytes. *Acta Chemica Scandinavia* 23, 2653-2668 (1969).
45. Klose, J. Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. A novel approach to testing for induced point mutations in mammals. *Humangenetik*, 231-243 (1975).
46. Scheele, G. A. Two-dimensional gel analysis of soluble proteins. Charaterization of guinea pig exocrine pancreatic proteins. *J Biol Chem* 250, 5375-85. (1975).
47. O'Farrell, P. H. High resolution two-dimensional electrophoresis of proteins. *J Biol Chem* 250, 4007-4021 (1975).
48. Klose, J. & Kobalz, U. Two-dimensional electrophoresis of proteins: an updated protocol and implications for a functional analysis of the genome. *Electrophoresis* 16, 1034-1059 (1995).
49. Merrill, C. R., Goldman, D., Sedman, S. A. & Ebert, M. H. Ultrasensitive stain for proteins in polyacrylamide gels shows regional variation in cerebrospinal fluid proteins. *Science* 211, 1437-1438 (1981).
50. Heukeshoven, J. & Dernick, R. in *Elektrophorese Forum* 92-97 (München, 1983).
51. Heukeshoven, H. & Dernick, R. Simplified method for silver staining of proteins in polyacrylamide gels and the mechanism of silver staining. *Electrophoresis* 6, 103-112 (1985).

52. Diezel, W., Kopperschläger, G. & Hofmann, E. An improved procedure for protein staining in polyacrylamide gels with a new type of coomassie brilliant blue. *Anal Biochem* 48, 617-620 (1972).
53. Neuhoff, V., Stamm, R. & Eibl, H. in *Electrophoresis* 427-448 (1985).
54. Neuhoff, V., Arold, N., Taube, D. & Ehrhardt, W. Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis* 9, 255-262 (1988).
55. Fernandez-Patron, C., Castellanos-Serra, L. & Rodriguez, P. Reverse staining of sodium dodecyl sulfate polyacrylamide gels by imidazole-zinc salts: Sensitive detection of unmodified proteins. *BioTechniques* 12, 564-573 (1992).
56. Ortiz, M. L. et al. Imidazole-SDS-Zn reverse staining of proteins in gels containing or not SDS and microsequence of individual unmodified electroblotted proteins. *FEBS Letters* 296, 300-4 (1992).
57. Patton, W. F. A thousand points of light: The application of fluorescence detection technologies to two-dimensional gel electrophoresis and proteomics [Review]. *Electrophoresis* 21, 1123-1144 (2000).
58. Anderson, N. L., Taylor, J., Scandora, A. E., Coulter, B. P. & Anderson, N. G. The TYCHO system for computerized analysis of two-dimensional gel protein mapping data. *Clin. Chem.* 27, 1807-1820 (1981).
59. Celis, J. E. et al. The Human Keratinocyte Two-Dimensional Gel Protein Database (Update 1995) - Mapping Components of Signal Transduction Pathways. *Electrophoresis*. 16, 2177-2240 (1995).
60. Edman, P. Method for determination of the amino acid sequence in peptides. *Acta Chemica Scandinavica* 4, 283-293 (1950).
61. Edman, P. & Begg, G. A protein sequenator. *European Journal of Biochemistry* 1, 80-91. (1967).
62. Peng, J. & Gygi, S. P. Proteomics: the move to mixtures. *J Mass Spectrom* 36, 1083-91 (2001).
63. Cameron, A. E. & Eggers, D. F. An Ion Velocitron. *Review of Scientific Instruments* 19, 605-607 (1948).
64. Wiley, W. C. & McLaren, I. H. Time-of-Flight Mass Spectrometer with Improved Resolution. *Review of Scientific Instruments* 26, 1150-1157 (1955).
65. Paul, W. & Steinwedel, H. Ein Neues Massenspektrometer Ohne Magnetfeld. *Zeitschrift Fur Naturforschung Section a-a Journal of Physical Sciences* 8, 448-450 (1953).
66. Paul, W. & Raether, M. Das Elektrische Massenfilter. *Zeitschrift Fur Physik* 140, 262-273 (1955).
67. Karas, M. & Hillenkamp, F. Influence of the wavelength in high-irradiance ultraviolet-laser desorption Mass-Spectrometry of organic-molecules. *Anal Chem* 57, 2935-2939 (1985).
68. Karas, M., Bachmann, D., Bahr, U. & Hillenkamp, F. Matrix-assisted ultraviolet laser desorption of non-volatile compounds. *International Journal of Mass Spectrometry and Ion Processes* 78, 53-68 (1987).
69. Karas, M. & Hillenkamp, F. Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *60*, 2299-2301 (1988).
70. Tanaka, K. et al. Protein and polymer analysis up to m/z 100 000 by laser ionization Time-of-Flight Mass Spectrometry. *Rapid Communications in Mass Spectrometry* 2, 151-153 (1988).
71. Hillenkamp, F. & Karas, M. Mass spectrometry of peptides and proteins by matrix-assisted ultraviolet laser desorption/ionization. *Methods Enzymol* 193, 280-95 (1990).
72. Hillenkamp, F., Karas, M., Beavis, R. C. & Chait, B. T. Matrix-assisted laser desorption/ionization mass spectrometry of biopolymers. *Anal Chem* 63, 1193A-1203A (1991).
73. Fenn, J. B., Mann, M., Meng, C. K., Wong, S. F. & Whitehouse, C. M. Electrospray ionization for mass spectrometry of large biomolecules. *Science* 246, 64-71 (1989).
74. Vertes, A., Gijbels, R. & Levine, R. D. Homogeneous Bottleneck Model of Matrix-Assisted Ultraviolet- Laser Desorption of Large Molecules. *Rapid Communications in Mass Spectrometry* 4, 228-233 (1990).

75. Vertes, A. & Levine, R. D. Sublimation Versus Fragmentation in Matrix-Assisted Laser Desorption. *Chemical Physics Letters* 171, 284-290 (1990).
76. Karas, M., Gluckmann, M. & Schafer, J. Ionization in matrix-assisted laser desorption/ionization: singly charged molecular ions are the lucky survivors. *Journal of Mass Spectrometry* 35, 1-12 (2000).
77. Zenobi, R. & Knochenmuss, R. Ion formation in MALDI mass spectrometry. *Mass Spectrom Rev* 17, 337-366 (1998).
78. Dole, M., Mack, L. L. & Hines, R. L. Molecular Beams of Macroions. *Journal of Chemical Physics* 49, 2240-& (1968).
79. Iribarne, J. V. & Thomson, B. A. Evaporation of Small Ions from Charged Droplets. *Journal of Chemical Physics* 64, 2287-2294 (1976).
80. Henzel, W. J. et al. Identifying proteins from two-dimensional gels by molecular mass searching of peptide fragments in protein sequence databases. *Proc Natl Acad Sci U S A* 90, 5011-5015 (1993).
81. Mann, M., Horjrup, P. & Roepstorff, p. Use of mass spectrometric molecular weight information to identify proteins in sequence database. *Biological Mass Spectrometry* 22, 338-345 (1993).
82. Pappin, D. J. C., Hojrup, P. & Bleasby, A. J. Rapid identification of proteins by peptide-mass fingerprinting. *Current Biology* 3, 327-332 (1993).
83. James, P., Quadroni, M., Carafoli, E. & Gonnet, G. Protein identification by mass profile fingerprinting. *Biochemical & Biophysical Research Communications* 195, 58-64 (1993).
84. Yates, J. R., 3rd, Speicher, S., Griffin, P. R. & Hunkapiller, T. Peptide mass maps: a highly informative approach to protein identification. *Anal Biochem* 214, 397-408 (1993).
85. Jensen, O. N., Podtelejnikov, A. & Mann, M. Delayed extraction improves specificity in database searches by matrix-assisted laser desorption/ionization peptide maps. *Rapid Commun Mass Spectrom* 10, 1371-8 (1996).
86. Morris, H. R. et al. High Sensitivity Collisionally-Activated Decomposition Tandem Mass Spectrometry On a Novel Quadrupole/Orthogonal-Acceleration Time-of-Flight Mass Spectrometer. *Rapid Communications in Mass Spectrometry*. 10, 889-896 (1996).
87. Krutchinsky, A. N., Zhang, W. & Chait, B. T. Rapidly switchable matrix-assisted laser desorption/ionization and electrospray quadrupole-time-of-flight mass spectrometry for protein identification. *J Am Soc Mass Spectrom* 11, 493-504. (2000).
88. Shevchenko, A., Loboda, A., Ens, W. & Standing, K. G. MALDI quadrupole time-of-flight mass spectrometry: a powerful tool for proteomic research. *Anal Chem* 72, 2132-41. (2000).
89. Fitzgerald, M. C., Chernushevich, I., Standing, K. G., Whitman, C. P. & Kent, S. B. Probing the oligomeric structure of an enzyme by electrospray ionization time-of-flight mass spectrometry. *Proc Natl Acad Sci U S A* 93, 6851-6 (1996).
90. Medzihradszky, K. F. et al. The characteristics of peptide collision-induced dissociation using a high-performance MALDI-TOF/TOF tandem mass spectrometer. *Anal Chem* 72, 552-558 (2000).
91. Qin, J. & Chait, B. T. Identification and characterization of posttranslational modifications of proteins by MALDI ion trap mass spectrometry. *Anal Chem* 69, 4002-9 (1997).
92. van Wijk, K. J. Challenges and prospects of plant proteomics. *Plant Physiol* 126, 501-8. (2001).
93. Harrison, P. & Black, C. Two-dimensional electrophoretic mapping of proteins of bundle sheath and mesophyll cells of the C4 grass *Digitaria sanguinalis* (L) Scop. (crabgrass). *Plant Physiol* 70, 1359-1366 (1982).
94. Wu, F. S. & Wang, M. Y. Extraction of proteins for sodium dodecyl sulfate-polyacrylamide gel electrophoresis from protease-rich plant tissues. *Anal Biochem* 139, 100-3. (1984).
95. Colas des Francs, C., Thiellement, H. & De Vienne, D. Analysis of leaf proteins by two-dimensional gel electrophoresis. *Plant Physiol* 78, 178-182 (1985).
96. Zivy, M., Thiellement, H., de Vienne, D. & Hofmann, J.-P. Study on nuclear and cytoplasmic genom expression in wheat by two-dimensional gel electrophoresis. 1. First results on 18 alloplasmic lines. *Theoretical and Applied Genetics* 66, 1-7 (1983).
97. Zivy, M., Thiellement, H., de Vienne, D. & Hofmann, J.-P. Study on nuclear and cytoplasmic genom expression in wheat by two-dimensional gel electrophoresis. 2. Genetic differences

- between two lines and two groups of cytoplasm at five developmental stages or organs. *Theoretical and Applied Genetics* 68, 335-345 (1984).
98. Bahrman, N., de Vienne, D., Thiellement, H. & Hofmann, J. P. Two-dimensional gel electrophoresis of proteins for genetic studies in Douglas fir (*Pseudotsuga menziesii*). *Biochemical Genetics* 23, 247-255 (1985).
 99. Leonardi, A., Damerval, C. & de Vienne, D. Organ-specific variability and inheritance of maize proteins revealed by two-dimensional electrophoresis. *Genetic Research Cambridge* 52, 97-103 (1988).
 100. David, J. L., Zivy, M., Cardin, M. L. & Brabant, P. Protein evolution in dynamically managed populations of wheat - adaptive responses to macro-environmental conditions. *Theoretical and Applied Genetics* 95, 932-941 (1997).
 101. Santoni, V., Bellini, C. & Caboche, M. Use of two-dimensional protein-pattern analysis for the characterization of *Arabidopsis thaliana* mutants. *Planta* 192, 557-566 (1994).
 102. Santoni, V., Delarue, M., Caboche, M. & Bellini, C. A comparison of two-dimensional electrophoresis data with phenotypical traits in *Arabidopsis* leads to the identification of a mutant (*cri1*) that accumulates cytokinins. *Planta* 202, 62-69 (1997).
 103. Herbik, A. et al. Iron and copper nutrition-dependent changes in protein expression in a tomato wild type and the nicotianamine-free mutant *chloronerva*. *Plant Physiol* 111, 533-540 (1996).
 104. Faure, J. D. et al. The *pasticcino* genes of *Arabidopsis thaliana* are involved in the control of cell division and differentiation. *Development* 125, 909-918 (1998).
 105. Tacchini, P., Fink, A., Thiellement, H. & Greppin, H. Analysis of leaf proteins in late flowering mutants of *Arabidopsis thaliana*. *Physiologia Plantarum* 93, 312-316 (1995).
 106. Damerval, C. & Le Guilloux, M. Characterization of novel proteins affected by the *o2* mutation and expressed during maize endosperm development. *Molecular & General Genetics* 257, 354-361 (1998).
 107. Kamo, M., Kawakami, T., Miyatake, N. & Tsugita, A. Separation and characterization of *Arabidopsis thaliana* proteins by two-dimensional gel electrophoresis. *Electrophoresis* 16, 423-300 (1995).
 108. Tsugita, A. et al. Separation and characterization of rice proteins. *Electrophoresis* 15, 708-720 (1994).
 109. Komatsu, S., Muhammad, A. & Rakwal, R. Separation and characterization of proteins from green and etiolated shoots of rice (*Oryza sativa* L.): towards a rice proteome. *Electrophoresis* 20, 630-636 (1999).
 110. Rakwal, R., Agrawal, G. K. & Yonekura, M. Separation of proteins from stressed rice (*Oryza sativa* L.) leaf tissues by two-dimensional polyacrylamide gel electrophoresis: induction of pathogenesis-related and cellular protectant proteins by jasmonic acid, UV irradiation and copper chloride. *Electrophoresis* 20, 3472-3478 (1999).
 111. Rakwal, R. & Komatsu, S. Role of jasmonate in the rice (*Oryza sativa* L.) self-defense mechanism using proteome analysis. *Electrophoresis* 21, 2492-2500 (2000).
 112. Ramani, S. & Apte, S. K. Transient expression of multiple genes in salinity-stressed young seedlings of rice (*Oryza sativa* L.) cv *Bura Rata*. *Biochemical & Biophysical Research Communications* 233, 663-667 (1997).
 113. Costa, P., Bahrman, N., Frigerio, J. M., Kremer, A. & Plomion, C. Water-deficit-responsive proteins in maritime pine. *Plant Molecular Biology* 38, 587-596 (1998).
 114. Riccardi, F., Gazeau, P., Devienne, D. & Zivy, M. Protein changes in response to progressive water deficit in maize - quantitative variation and polypeptide identification. *Plant Physiol* 117, 1253-1263 (1998).
 115. Hurkman, W., Tanaka, C. & Dupont, F. The effect of salt stress on polypeptides in membrane-fractions from barley roots. *Plant Physiol* 88, 1263-1273 (1988).
 116. Huber, L. A. Mapping cells and sub-cellular organelles on 2-D gels: 'new tricks for an old horse'. [Review]. *FEBS Letters* 369, 122-125 (1995).
 117. Pasquali, C., Fialka, I. & Huber, L. A. Subcellular fractionation, electromigration analysis and mapping of organelles. *Journal of Chromatography B* 722, 89-102 (1999).
 118. Cordwell, S. J., Nouwens, A. S., Verrills, N. M., Basseal, D. J. & Walsh, B. J. Subproteomics based upon protein cellular location and relative solubilities in conjunction with composite two-dimensional electrophoresis gels. *Electrophoresis* 21, 1094-1103. (2000).

119. Santoni, V. et al. Use of a proteome strategy for tagging proteins present at the plasma membrane. *Plant J* 16, 633-641 (1998).
120. Rouquie, D. et al. Construction of a directory of tobacco plasma membrane proteins by combined two-dimensional gel electrophoresis and protein sequencing. *Electrophoresis* 18, 654-60. (1997).
121. Seigneurin-Berny, D., Rolland, N., Garin, J. & Joyard, J. Technical Advance: Differential extraction of hydrophobic proteins from chloroplast envelope membranes: a subcellular-specific proteomic approach to identify rare intrinsic membrane proteins. *Plant J* 19, 217-228 (1999).
122. Santoni, V. et al. Towards the recovery of hydrophobic proteins on two-dimensional electrophoresis gels. *Electrophoresis* 20, 705-711 (1999).
123. Santoni, V., Molloy, M. & Rabilloud, T. Membrane proteins and proteomics: Un amour impossible? [Review]. *Electrophoresis* 21, 1054-1070 (2000).
124. Santoni, V., Kieffer, S., Desclaux, D., Masson, F. & Rabilloud, T. Membrane proteomics: Use of additive main effects with multiplicative interaction model to classify plasma membrane proteins according to their solubility and electrophoretic properties. *Electrophoresis* 21, 3329-3344 (2000).
125. Robertson, D. et al. Differential extraction and protein sequencing reveals major differences in patterns of primary cell wall proteins from plants. *J Biol Chem* 272, 15841-15848 (1997).
126. Peltier, J. B. et al. Central functions of the luminal and peripheral thylakoid proteome of Arabidopsis determined by experimentation and genome-wide prediction. *Plant Cell* 14, 211-236 (2002).
127. Peltier, J. B. et al. Proteomics of the chloroplast. Systematic identification and targeting analysis of luminal and peripheral thylakoid proteins. *Plant Cell* 12, 319-342 (2000).
128. Ferro, M. et al. Organic solvent extraction as a versatile procedure to identify hydrophobic chloroplast membrane proteins. *Electrophoresis* 21, 3517-26 (2000).
129. van Wijk, K. J. Proteomics of the chloroplast: experimentation and prediction. *Trends in Plant Science* 5, 420-425 (2000).
130. Schubert, M. et al. Proteome map of the chloroplast lumen of Arabidopsis thaliana. *J Biol Chem* 277, 8354-65 (2002).
131. Yamaguchi, K., von Knoblauch, K. & Subramanian, A. R. The plastid ribosomal proteins. Identification of all the proteins in the 30 S subunit of an organelle ribosome (chloroplast). *J Biol Chem* 275, 28455-65. (2000).
132. Yamaguchi, K. & Subramanian, A. R. The plastid ribosomal proteins. Identification of all the proteins in the 50 S subunit of an organelle ribosome (chloroplast). *J Biol Chem* 275, 28466-82. (2000).
133. Kruft, V., Eubel, H., Jansch, L., Werhahn, W. & Braun, H. P. Proteomic approach to identify novel mitochondrial proteins in Arabidopsis. *Plant Physiol* 127, 1694-710 (2001).
134. Millar, A. H., Sweetlove, L. J., Giege, P. & Leaver, C. J. Analysis of the Arabidopsis mitochondrial proteome. *Plant Physiol* 127, 1711-27 (2001).
135. Bardel, J. et al. A survey of the plant mitochondrial proteome in relation to development. *Proteomics* 2, 880-98 (2002).
136. Werhahn, W. & Braun, H. P. Biochemical dissection of the mitochondrial proteome from Arabidopsis thaliana by three-dimensional gel electrophoresis. *Electrophoresis* 23, 640-6 (2002).
137. Panter, S. et al. Identification with proteomics of novel proteins associated with the peribacteroid membrane of soybean root nodules. *Mol Plant Microbe Interact* 13, 325-33. (2000).
138. Schagger, H. & von Jagow, G. Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal Biochem* 199, 223-31 (1991).
139. Peltier, J. B., Ytterberg, J., Liberles, D. A., Roepstorff, P. & van Wijk, K. J. Identification of a 350-kDa ClpP protease complex with 10 different Clp isoforms in chloroplasts of Arabidopsis thaliana. *J Biol Chem* 276, 16318-16327 (2001).
140. Werhahn, W. et al. Purification and Characterization of the Preprotein Translocase of the Outer Mitochondrial Membrane from Arabidopsis. Identification of Multiple Forms of TOM20. *Plant Physiol* 125, 943-954 (2001).

141. Gavin, A. C. et al. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* 415, 141-7 (2002).
142. Ho, Y. et al. Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* 415, 180-3 (2002).
143. Cove, D. J., Knight, C. D. & Lamparter, T. Mosses as model systems. *Trends in Plant Science* 2, 99-105 (1997).
144. Schaefer, D. G. & Zryd, J. P. Efficient Gene Targeting in the Moss *Physcomitrella Patens*. *Plant J* 11, 1195-1206 (1997).
145. Reski, R. *Physcomitrella* and *Arabidopsis* - the David and Goliath of Reverse Genetics. *Trends in Plant Science* 3, 209-210 (1998).
146. Weschke, W. et al. Sucrose transport into barley seeds: molecular characterization of two transporters and implications for seed development and starch accumulation. *Plant J* 21, 455-67 (2000).
147. Kehr, J. et al. Analysis of phloem protein patterns from different organs of *Cucurbita maxima* Duch. by matrix-assisted laser desorption/ionization time of flight mass spectroscopy combined with sodium dodecyl sulfate polyacrylamide gel electrophoresis. *Planta* 207, 612-9 (1999).
148. Walz, C., Juenger, M., Schad, M. & Kehr, J. Evidence for the presence and activity of a complete antioxidant defence system in mature sieve tubes. *Plant J* 31, 189-97 (2002).
149. Kamp, R.M., Srinivasa, B. R., Vonknoblauch, K. & Subramanian, A. R. Occurrence of a Methylated Protein in Chloroplast Ribosomes. *Biochemistry* 26, 5866 (1987).
150. Giavalisco, P., Nordhoff, E., Lehrach, H., Gobom, J. & Klose, J. Extraction of proteins from plant tissue for two-dimensional electrophoresis analysis. *Electrophoresis* 24, 207-216 (2003).
151. Damerval, C., de Vienne, D., Zivy, M. & Thiellement, H. Technical improvements in two-dimensional electrophoresis increase the level of genetic variation detected in wheat-seedling proteins. *Electrophoresis* 7, 52-54 (1986).
152. Klose, J. in *2-D Proteom Analysis Protocols* (ed. Link, A. J.) 67-85 (Humana Press, Totowa, New Jersey, 1999).
153. Hippler, M., Klein, J., Fink, A., Allinger, T. & Hoerth, P. Towards functional proteomics of membrane protein complexes: analysis of thylakoid membranes from *Chlamydomonas reinhardtii*. *Plant J* 28, 595-606 (2001).
154. Klose, J. in *2-D Proteom Analysis Protocols* (ed. Link, A. J.) 147-172 (Humana Press, Totowa, New Jersey, 1999).
155. Bjellqvist, B. et al. Isoelectric focusing in immobilized pH gradients: principle, methodology and some applications. *Journal of Biochemical & Biophysical Methods* 6, 317-339 (1982).
156. Jungblut, P. R. & Seifert, R. Analysis by high-resolution two-dimensional electrophoresis of differentiation-dependent alterations in cytosolic protein pattern of HL-60 leukemic cells. *Journal of Biochemical and Biophysical Methods* 21, 47-58 (1990).
157. Shevchenko, A., Wilm, M., Vorm, O. & Mann, M. Mass spectrometric sequencing of proteins from silver stained polyacrylamide gels. *Anal Chem* 68, 850-858 (1996).
158. Fazekas de St Groth, S., Webster, R. G. & Datyner, A. Two new staining procedures for quantitative estimation of proteins on electrophoresis strips. *Biochim Biophys Acta* 71, 377 - 391 (1963).
159. Doherty, N. S. et al. Analysis of changes in acute-phase plasma proteins in an acute inflammatory response and in rheumatoid arthritis using two-dimensional gel electrophoresis. *Electrophoresis* 19, 355-363 (1998).
160. Castellanos-Serra, L., Proenza, W., Huerta, V., Moritz, R. L. & Simpson, R. J. Proteome analysis of polyacrylamide gel-separated proteins visualized by reversible negative staining using imidazole-zinc salts. *Electrophoresis* 20, 732-737 (1999).
161. Courchesne, P. L., Luethy, R. & Patterson, S. D. Comparison of in-gel and on-membrane digestion methods at low to sub-pmol level for subsequent peptide and fragment-ion mass analysis using matrix-assisted laser-desorption/ionization mass spectrometry. *Electrophoresis* 18, 369-381 (1997).
162. Steinberg, T. H., Haugland, R. P. & Singer, V. L. Applications of Sypro Orange and Sypro Red Protein Gel Stains. *Anal Biochem.* 239, 238-245 (1996).

163. Steinberg, T. H., Jones, L. J., Haugland, R. P. & Singer, V. L. Sypro Orange and Sypro Red Protein Gel Stains - One-Step Fluorescent Staining of Denaturing Gels For Detection of Nanogram Levels of Protein. *Anal Biochem.* 239, 223-237 (1996).
164. Berggren, K. N. et al. An improved formulation of SYPRO Ruby protein gel stain: Comparison with the original formulation and with a ruthenium II tris (bathophenanthroline disulfonate) formulation. *Proteomics* 2, 486-98 (2002).
165. Rabilloud, T., Strub, J. M., Luche, S., van Dorsselaer, A. & Lunardi, J. A comparison between Sypro Ruby and ruthenium II tris (bathophenanthroline disulfonate) as fluorescent stains for protein detection in gels. *Proteomics* 1, 699-704 (2001).
166. Malone, J. P., Radabaugh, M. R., Leimgruber, R. M. & Gerstenecker, G. S. Practical aspects of fluorescent staining for proteomic applications. *Electrophoresis* 22, 919-932 (2001).
167. Smilansky, Z. Automated registration for images of two-dimensional protein gels. *Electrophoresis* 22, 1616-1626 (2001).
168. Vorm, O., Roepstorff, P. & Mann, M. Improved Resolution and Very High Sensitivity in MALDI TOF of Matrix Surfaces Made by Fast Evaporation. *Anal Chem* 66, 3281 - 3287 (1994).
169. Schuerenberg, M. et al. Prestructured MALDI-MS sample supports. *Anal Chem* 72, 3436-42 (2000).
170. Gobom, J. et al. alpha-cyano-4-hydroxycinnamic acid affinity sample preparation. A protocol for MALDI-MS peptide analysis in proteomics. *Anal Chem* 73, 434-438 (2001).
171. Beavis, R., Chaudhary, T. & Chait, B. alpha-cyano-4-hydroxycinnamic acid as a matrix for Matrix-Assisted Laser Desorption Mass Spectrometry. *Organic Mass Spectrometry* 27, 156-158 (1992).
172. Kussmann, M. & Roepstorff, P. Sample preparation techniques for peptides and proteins analyzed by MALDI-MS. *Methods Mol Biol* 146, 405-24 (2000).
173. Strupat, K., Karas, M. & Hillenkamp, F. 2,5-Dihydroxybenzoic acid - A new Matrix for laser desorption ionization mass-spectrometry. *International Journal of Mass Spectrometry and Ion Processes* 111, 89-102 (1991).
174. Beavis, R. C. & Chait, B. T. Cinnamic acid derivatives as matrices for ultraviolet laser desorption mass spectrometry of proteins. *Rapid Commun Mass Spectrom* 3, 432-5. (1989).
175. Nordhoff, E. et al. Large-gel two-dimensional electrophoresis-matrix assisted laser desorption/ionization-time of flight-mass spectrometry: an analytical challenge for studying complex protein mixtures. *Electrophoresis* 22, 2844-55. (2001).
176. Gobom, J. et al. A calibration method that simplifies and improves accurate determination of peptide molecular masses by MALDI-TOF MS. *Anal Chem* 74, 3915 - 3923 (2002).
177. Perkins, D. N., Pappin, D. J., Creasy, D. M. & Cottrell, J. S. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* 20, 3551-67. (1999).
178. de Vienne, D., Leonardi, A. & Damerval, C. Genetic aspects of variation of protein amounts in maize and pea. *Electrophoresis* 9, 742-750 (1988).
179. Porubleva, L., Vander Velden, K., Kothari, S., Livier, D. J. & Chitnis, P. R. The proteome of maize: Use of gene sequence and expressed sequence tag data for identification of proteins with mass fingerprints. *Electrophoresis* 22, 1724-1738 (2001).
180. Herskovits, T. T., Jaillet, H. & Gadegebeku, B. On the structural stability and solvent denaturation of proteins. II. Denaturation by the ureas. *J Biol Chem* 245, 4544-50 (1970).
181. Rabilloud, T., Adessi, C., Giraudel, A. & Lunardi, J. Improvement of the solubilization of proteins in two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis* 18, 307-16. (1997).
182. Rabilloud, T. Use of thiourea to increase the solubility of membrane proteins in two-dimensional electrophoresis. *Electrophoresis* 19, 758-760 (1998).
183. Molloy, M. P. et al. Extraction of membrane proteins by differential solubilization for separation using two-dimensional gel electrophoresis. *Electrophoresis* 19, 837-844 (1998).
184. Lanne, B. et al. Thiourea enhances mapping of the proteome from murine white adipose tissue. *Proteomics* 1, 819-28. (2001).
185. Perdew, G. H., Schaup, H. W. & Selivonchick, D. P. The use of a zwitterionic detergent in two-dimensional gel electrophoresis of trout liver microsomes. *Anal Biochem* 135, 453-5. (1983).

186. Chevallet, M. et al. New zwitterionic detergents improve the analysis of membrane proteins by two-dimensional electrophoresis. *Electrophoresis* 19, 1901-1909 (1998).
187. Friso, G. & Wikstrom, L. Analysis of proteins from membrane-enriched cerebellar preparations by two-dimensional gel electrophoresis and mass spectrometry. *Electrophoresis* 20, 917-27. (1999).
188. Harry, J. L. et al. Proteomics: capacity versus utility. *Electrophoresis* 21, 1071-81 (2000).
189. Gegenheimer, P. Preparation of extracts from plants. *Methods Enzymol* 182, 174-193 (1990).
190. Rabilloud, T. Solubilization of proteins for electrophoretic analyses. [Review]. *Electrophoresis* 17, 813-829 (1996).
191. Radin, N. S. Extraction of tissue lipids with a solvent of low toxicity. *Methods Enzymol* 72, 5-7 (1981).
192. Gorg, A. et al. The current state of two-dimensional electrophoresis with immobilized pH gradients [Review]. *Electrophoresis* 21, 1037-1053 (2000).
193. Nawrocki, A. et al. Correlation of acidic and basic carrier ampholyte and immobilized pH gradient two-dimensional gel electrophoresis patterns based on mass spectrometric protein identification. *Electrophoresis* 19, 1024-1035 (1998).
194. O'Farrell, P. Z., Goodman, H. M. & O'Farrell, P. H. High resolution two-dimensional electrophoresis of basic as well as acidic proteins. *Cell* 12, 1133-41. (1977).
195. Rabilloud, T. Silver staining of 2-D electrophoresis gels. [Review]. *Methods in Molecular Biology* 112, 297-305 (1999).
196. Switzer, R. C., Merrill, C. R. & Shifrin, S. A highly sensitive silver stain for detecting proteins and peptides in polyacrylamide gels. *Anal Biochem* 98, 231-237 (1979).
197. Rabilloud, T. Detecting proteins separated by 2-D gel electrophoresis. *Anal Chem* 72, 48A-55A (2000).
198. Tonge, R. et al. Validation and development of fluorescence two-dimensional differential gel electrophoresis proteomics technology. *Proteomics* 1, 377-396 (2001).
199. Patton, W. F. & Beechem, J. M. Rainbow's end: the quest for multiplexed fluorescence quantitative analysis in proteomics. *Curr Opin Chem Biol* 6, 63-9 (2002).
200. Patton, W. F. Detection technologies in proteome analysis. *Journal of Chromatography B* 771, 3-31 (2002).
201. Rabilloud, T., Vuillard, L., Gilly, C. & Lawrence, J. J. Silver-staining of proteins in polyacrylamide gels: a general overview. [Review]. *Cellular & Molecular Biology* 40, 57-75 (1994).
202. Voss, T. & Haberl, P. Observations on the reproducibility and matching efficiency of two-dimensional electrophoresis gels: consequences for comprehensive data analysis. *Electrophoresis* 21, 3345-50 (2000).
203. Berggren, K. et al. Background-free, high sensitivity staining of proteins in one- and two-dimensional sodium dodecyl sulfate-polyacrylamide gels using luminescent ruthenium complex. *Electrophoresis* 21, 2509-2521 (2000).
204. Heukeshoven, J. & Dernick, R. Improved silver staining procedure for fast staining in PhastSystem Development Unit. I. Staining of sodium dodecyl sulfate gels. *Electrophoresis* 9, 28-32 (1988).
205. Cohen, S. L. & Chait, B. T. Influence of matrix solution conditions on the MALDI-MS analysis of peptides and proteins. *Anal Chem* 68, 31-7. (1996).
206. Perera, I. K., Kantartzoglou, S. & Dyer, P. E. Some Characteristics of Matrix-Assisted Uv Laser Desorption Ionization Mass Spectrometric Analysis of Large Proteins. *International Journal of Mass Spectrometry & Ion Processes*. 156, 151-172 (1996).
207. Kussmann, M. et al. Matrix-assisted laser desorption/ionization mass spectrometry sample preparation techniques designed for various peptide and protein analytes. *Journal of Mass Spectrometry* 32, 593-601 (1997).
208. Gobom, J., Nordhoff, E., Mirgorodskaya, E., Ekman, R. & Roepstorff, P. Sample purification and preparation technique based on nano-scale reversed-phase columns for the sensitive analysis of complex peptide mixtures by matrix-assisted laser desorption/ionization mass spectrometry. *J Mass Spectrom* 34, 105-16 (1999).
209. Karas, M. & Bahr, U. in *Selected topics in mass spectrometry in the biomolecular sciences*. (eds. Caprioli, R. M., Malorni, A. & Sindona, G.) 33-54 (Kluwer Academic Publishers, Dordrecht, Boston, London, 1996).

210. Keller, B. O. & Li, L. Discerning matrix-cluster peaks in matrix-assisted laser desorption/ionization time-of-flight mass spectra of dilute peptide mixtures. *Journal of the American Society for Mass Spectrometry* 11, 88-93 (2000).
211. Harris, W. A., Janecki, D. J. & Reilly, J. P. Use of matrix clusters and trypsin autolysis fragments as mass calibrants in matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Communications in Mass Spectrometry* 16, 1714-1722 (2002).
212. Karty, J. A., Ireland, M. M. E., Brun, Y. V. & Reilly, J. P. Artifacts and unassigned masses encountered in peptide mass mapping. *Journal of Chromatography B* 782, 363-383 (2002).
213. Fountoulakis, M. & Langen, H. Identification of proteins by matrix-assisted laser desorption ionization-mass spectrometry following in-gel digestion in low-salt, nonvolatile buffer and simplified peptide recovery. *Anal Biochem* 250, 153-6. (1997).
214. Staudenmann, W. et al. Sample handling for proteome analysis. *Electrophoresis* 19, 901-908 (1998).
215. Fountoulakis, M., Juranville, J.-F., Dierssen, M. & Lubec, G. Proteomic analysis of the fetal brain. *Proteomics* 2, 1547-1576 (2002).
216. Scheler, C. et al. Peptide mass fingerprint sequence coverage from differently stained proteins on two-dimensional electrophoresis patterns by matrix assisted laser desorption/ionization-mass spectrometry (MALDI-MS). *Electrophoresis* 19, 918-927 (1998).
217. Galvani, M., Hamdan, M., Herbert, B. & Righetti, P. G. Alkylation kinetics of proteins in preparation for two-dimensional maps: a matrix assisted laser desorption/ionization-mass spectrometry investigation. *Electrophoresis* 22, 2058-65 (2001).
218. Galvani, M., Rovatti, L., Hamdan, M., Herbert, B. & Righetti, P. G. Protein alkylation in the presence/absence of thiourea in proteome analysis: a matrix assisted laser desorption/ionization-time of flight-mass spectrometry investigation. *Electrophoresis* 22, 2066-74 (2001).
219. Herbert, B. et al. Reduction and alkylation of proteins in preparation for two-dimensional map analysis: Why, when, and how? *Electrophoresis* 22, 2046-2057 (2001).
220. Piubelli, C., Galvani, M., Hamdan, M., Domenici, E. & Righetti, P. G. Proteome analysis of rat polymorphonuclear leukocytes: A two-dimensional electrophoresis/ mass spectrometry approach. *Electrophoresis* 23, 298-310 (2002).
221. Herbert, B. R. et al. Improved protein solubility in two-dimensional electrophoresis using tributyl phosphine as reducing agent. *Electrophoresis* 19, 845-851 (1998).
222. Molloy, M. P. Two-dimensional electrophoresis of membrane proteins using immobilized pH gradients. *Anal Biochem* 280, 1-10 (2000).
223. Gianazza, E. Isoelectric-Focusing as a Tool for the Investigation of Posttranslational Processing and Chemical Modifications of Proteins. *Journal of Chromatography A* 705, 67-87 (1995).
224. Fountoulakis, M. & Suter, L. Proteomic analysis of the rat liver. *Journal of Chromatography B* 782, 197-218 (2002).
225. Spreitzer, R. J. & Salvucci, M. E. Rubisco: Structure, regulatory interactions, and possibilities for a better enzyme. *Annual Review Of Plant Biology* 53, 449-475 (2002).
226. Paulsen, I. T., Sliwinski, M. K., Nelissen, B., Goffeau, A. & Saier, M. H., Jr. Unified inventory of established and putative transporters encoded within the complete genome of *Saccharomyces cerevisiae*. *FEBS Letters* 430, 116-25 (1998).
227. Stevens, T. J. & Arkin, I. T. Do more complex organisms have a greater proportion of membrane proteins in their genomes? *Proteins* 39, 417-20. (2000).
228. Masson, F. & Rossignol, M. Basic plasticity of protein expression in tobacco leaf plasma membrane. *Plant J* 8, 77-85 (1995).
229. Rossignol, M. Analysis of the plant proteome. *Curr Opin Biotechnol* 12, 131-4 (2001).
230. Kehr, J., Hustiak, F., Walz, C., Willmitzer, L. & Fisahn, J. Transgenic plants changed in carbon allocation pattern display a shift in diurnal growth pattern. *Plant J* 16, 497-503 (1998).
231. Oparka, K. J. & Cruz, S. S. The great escape: Phloem transport and unloading of macromolecules. *Annual Review of Plant Physiol & Plant Molecular Biology* 51, 323-347 (2000).
232. Capel, M. S. & Bourque, D. P. Characterization of *Nicotiana tabacum* chloroplast and cytoplasmic ribosomal proteins. *J Biol Chem* 257, 7746-7755 (1982).

233. Rabilloud, T. & Chevallet, M. in *Proteome research: Two-dimensional gel electrophoresis and identification methods* (ed. Rabilloud, T.) 9-31 (Springer, Heidelberg, 2000).
234. Granzier, H. L. & Wang, K. Gel electrophoresis of giant proteins: solubilization and silver-staining of titin and nebulin from single muscle fiber segments. *Electrophoresis* 14, 56-64. (1993).
235. Dunn, M. (BIOS Scientific Publishers, Alden Press, Oxford, 1993).
236. Rabilloud, T., Valette, C. & Lawrence, J. J. Sample application by in-gel rehydration improves the resolution of two-dimensional electrophoresis with immobilized pH gradients in the first dimension. *Electrophoresis* 15, 1552-1558 (1994).
237. Anderson, N. L. & Anderson, N. G. Proteome and proteomics: new technologies, new concepts, and new words. [Review]. *Electrophoresis* 19, 1853-1861 (1998).
238. Blackstock, W. P. & Weir, M. P. Proteomics: quantitative and physical mapping of cellular proteins. [Review]. *Trends in Biotechnology* 17, 121-127 (1999).
239. Celis, J. E. & Gromov, P. 2D protein electrophoresis: can it be perfected? *Current Opinion in Biotechnology*. 10, 16-21 (1999).
240. Fey, S. J. & Larsen, P. M. 2D or not 2D. *Current Opinion in Chemical Biology* 5, 26-33 (2001).
241. Gygi, S. P., Corthals, G. L., Zhang, Y., Rochon, Y. & Aebersold, R. Evaluation of two-dimensional gel electrophoresis-based proteome analysis technology. *Proc Natl Acad Sci U S A* 97, 9390-9395 (2000).
242. Jungblut, P. et al. Resolution power of two-dimensional electrophoresis and identification of proteins from gels. *Electrophoresis* 17, 839-847 (1996).
243. Ong, S. E. & Pandey, A. An evaluation of the use of two-dimensional gel electrophoresis in proteomics. *Biomolecular Engineering* 18, 195-205 (2001).
244. Rabilloud, T. Two-dimensional gel electrophoresis in proteomics: Old, old fashioned, but still climbs up the mountain. *Proteomics* 2, 3-10 (2002).
245. Nawrocki, J. P. et al. Analysis of Combinatorial Libraries Using Electrospray Fourier Transform Ion Cyclotron Resonance Mass Spectrometry. *Rapid Communications in Mass Spectrometry*. 10, 1860-1864 (1996).
246. Lopez, M. F. Advantages of carrier ampholyte IEF. *Methods in Molecular Biology* 112, 109-110 (1999).
247. Gorg, A., Obermaier, C., Boguth, G. & Weiss, W. Recent developments in two-dimensional gel electrophoresis with immobilized pH gradients: wide pH gradients up to pH 12, longer separation distances and simplified procedures. *Electrophoresis* 20, 712-7. (1999).
248. Rabilloud, T., Gelfi, C., Bossi, M. & Righetti, P. Protein precipitation induced alkaline immobilines for isoelectric-focusing in immobilized pH gradients - caused and remedies. *Electrophoresis* 8, 305-312 (1987).
249. Adessi, C., Miege, C., Albrieux, C. & Rabilloud, T. Two-dimensional electrophoresis of membrane proteins: A current challenge for immobilized pH gradients. *Electrophoresis* 18, 127-135 (1997).
250. Rimpilainen, M. & Righetti, P. Membrane proteome analysis by isoelectrical-focusing in pH gradients. *Electrophoresis* 6, 419-422 (1985).
251. Duncan, R. & Hershey, J. W. Evaluation of isoelectric focusing running conditions during two-dimensional isoelectric focusing/sodium dodecyl sulfate-polyacrylamide gel electrophoresis: variation of gel patterns with changing conditions and optimized isoelectric focusing conditions. *Anal Biochem* 138, 144-55. (1984).
252. Righetti, P. G. & Bossi, A. Isoelectric focusing in immobilized pH gradients: an update. [Review]. *Journal of Chromatography* 699, 77-89 (1997).
253. Blomberg, A. et al. Interlaboratory reproducibility of yeast protein patterns analyzed by immobilized pH gradient two-dimensional gel electrophoresis. *Electrophoresis* 16, 1935 - 1945 (1995).
254. Lopez, M. F. & Patton, W. F. Reproducibility of polypeptide spot position in two-dimensional gels run using carrier ampholytes in the isoelectrical focusing dimension. *Electrophoresis* 18, 338-343 (1997).
255. Corbett, J. M., Dunn, M. J., Posch, A. & Gorg, A. Positional reproducibility of protein spots in two-dimensional polyacrylamide gel electrophoresis using immobilised pH gradient

- isoelectric focusing in the first dimension: an interlaboratory comparison. *Electrophoresis* 15, 1205-11. (1994).
256. Klose, J. et al. Genetic analysis of the mouse brain proteome. *Nat Genet* 30, 385-393 (2002).
 257. Hoving, S., Voshol, H. & van Oostrum, J. Towards high performance two-dimensional gel electrophoresis using ultrazoom gels. *Electrophoresis* 21, 2617-21 (2000).
 258. Gronborg, M. et al. A Mass Spectrometry-based Proteomic Approach for Identification of Serine/Threonine-phosphorylated Proteins by Enrichment with Phospho-specific Antibodies: Identification of a Novel Protein, Frigg, as a Protein Kinase A Substrate. *Mol Cell Proteomics* 1, 517-27 (2002).
 259. Zuo, X. & Speicher, D. W. A method for global analysis of complex proteomes using sample prefractionation by solution isoelectrofocusing prior to two-dimensional electrophoresis. *Anal Biochem* 284, 266-278 (2000).
 260. Righetti, P. G., Castagna, A. & Herbert, B. Prefractionation techniques in proteome analysis. *Anal Chem* 73, 320A-326A (2001).
 261. Link, A. J. et al. Direct analysis of protein complexes using mass spectrometry. *Nat Biotechnol* 17, 676-682 (1999).
 262. Yates, J. R., 3rd, Link, A. J. & Schieltz, D. Direct analysis of proteins in mixtures. Application to protein complexes. *Methods Mol Biol* 146, 17-26 (2000).
 263. Washburn, M. P., Wolters, D. & Yates, J. R., 3rd. Large-scale analysis of the yeast proteome by multidimensional protein identification technology. *Nat Biotechnol* 19, 242-7 (2001).
 264. Florens, L. et al. A proteomic view of the Plasmodium falciparum life cycle. *Nature* 419, 520-6 (2002).
 265. Koller, A. et al. Proteomic survey of metabolic pathways in rice. *Proc Natl Acad Sci U S A* 99, 11969-74 (2002).
 266. Hunter, T. C., Andon, N. L., Koller, A., Yates, J. R. & Haynes, P. A. The functional proteomics toolbox: methods and applications. *J. Chromatogr. B* 782, 165-181 (2002).
 267. Rabilloud, T. Mechanisms of protein silver staining in polyacrylamide gels: a 10-year synthesis. [Review]. *Electrophoresis* 11, 785-794 (1990).
 268. Rabilloud, T. et al. Two-dimensional electrophoresis of human placental mitochondria and protein identification by mass spectrometry: toward a human mitochondrial proteome. *Electrophoresis* 19, 1006-1014 (1998).
 269. Jungblut, P. & Thiede, B. Protein identification from 2-DE gels by MALDI mass spectrometry. *Mass Spectrometry Reviews* 16, 145-62 (1997).
 270. Hille, J., Free, A. & Wätzig, H. Possibilities to improve automation, speed and precision of proteome analysis: A comparison of two-dimensional electrophoresis and alternatives. *Electrophoresis* 22, 4035-4052 (2001).
 271. Mann, M., Hendrickson, R. C. & Pandey, A. Analysis of proteins and proteomes by mass spectrometry [Review]. *Annual Review of Biochemistry* 70, 437-473 (2001).
 272. Mamyurin, B., Karataev, V., Shmikk, D. & Zagulin, V. Mass-reflectron a new nonmagnetic Time-of-Flight high-resolution mass-spectrometer. *Zhurnal Eksperimentalnoi Teoreticheskoi Fiziki* 64, 82-89 (1973).
 273. Spengler, B., Kirsch, D. & Kaufmann, R. Fundamental-Aspects of Postsource Decay in Matrix-Assisted Laser Desorption Mass-Spectrometry.1. Residual-Gas Effects. *Journal of Physical Chemistry* 96, 9678-9684 (1992).
 274. Kaufmann, R., Chaurand, P., Kirsch, D. & Spengler, B. Post-Source Decay and Delayed Extraction in Matrix-Assisted Laser Desorption/Ionization/Reflectron Time-of-Flight Mass Spectrometry - Are There Trade-Offs. *Rapid Communications in Mass Spectrometry*. 10, 1199-1208 (1996).
 275. Spengler, B. in *Selected topics in mass spectrometry in the biomolecular sciences*. (eds. Caprioli, R. M., Malorni, A. & Sindona, G.) 377-398 (Kluwer Academic Publishers, Dordrecht, Boston, London, 1996).
 276. Keough, T., Youngquist, R. S. & Lacey, M. P. A method for high-sensitivity peptide sequencing using postsource decay matrix-assisted laser desorption ionization mass spectrometry. *Proc Natl Acad Sci U S A* 96, 7131-7136 (1999).
 277. Keough, T., Lacey, M. P. & Youngquist, R. S. Derivatization procedures to facilitate de novo sequencing of lysine-terminated tryptic peptides using postsource decay matrix-assisted laser desorption/ionization mass spectrometry. *Rapid Commun Mass Spectrom* 14, 2348-56 (2000).

278. Haebel, S. & Kehr, J. Matrix-assisted laser desorption/ionisation time of flight mass spectrometry peptide mass fingerprints and post source decay: a tool for the identification and analysis of phloem proteins from *Curcubita maxima* Duch. separated by two-dimensional polyacrylamide gel electrophoresis. *Planta* 213, 586-593 (2001).
279. Mewes, H. W. et al. MIPS: a database for genomes and protein sequences. *Nucleic Acids Res* 28, 37-40 (2000).
280. Mann, M. & Wilm, M. Error Tolerant Identification of Peptides in Sequence Databases By Peptide Sequence Tags. *Anal Chem*. 66, 4390-4399 (1994).
281. Eng, J., MacCormack, A. & Yates, J. An approach to correlate tandem mass-spectral data of peptides with amino-acid-sequences in a protein database. *Journal of the American Society for Mass Spectrometry* 5, 976-989 (1994).
282. Roepstorff, P. Mass spectrometry in protein studies from genome to function. [Review]. *Current Opinion in Biotechnology* 8, 6-13 (1997).
283. Loboda, A. V., Krutchinsky, A. N., Bromirski, M., Ens, W. & Standing, K. G. A tandem quadrupole/time-of-flight mass spectrometer with a matrix-assisted laser desorption/ionization source: design and performance. *Rapid Communications in Mass Spectrometry* 14, 1047-1057 (2000).
284. Krutchinsky, A. N., Kalkum, M. & Chait, B. T. Automatic identification of proteins with a MALDI-quadrupole ion trap mass spectrometer. *Anal Chem* 73, 5066-5077 (2001).
285. Vestal, M. L. Methods of ion generation. *Chemical Reviews* 101, 361-375 (2001).
286. Hunt, D. F., Yates, J. R. d., Shabanowitz, J., Winston, S. & Hauer, C. R. Protein sequencing by tandem mass spectrometry. *Proc Natl Acad Sci U S A* 83, 6233-7 (1986).
287. Walsh, B. J., Molloy, M. P. & Williams, K. L. The Australian Proteome Analysis Facility (APAF): assembling large scale proteomics through integration and automation. *Electrophoresis* 19, 1883-90. (1998).
288. Quadroni, M. & James, P. Proteomics and automation. *Electrophoresis* 20, 664-677 (1999).
289. Lopez, M. F. Better approaches to finding the needle in a haystack: optimizing proteome analysis through automation. *Electrophoresis* 21, 1082-93. (2000).
290. Mahon, P. & Dupree, P. Quantitative and reproducible two-dimensional gel analysis using Phoretix 2D. *Electrophoresis* 22, 2075-2085 (2001).
291. Jensen, O. N., Mortensen, P., Vorm, O. & Mann, M. Automation of matrix-assisted laser desorption/ionization mass spectrometry using fuzzy logic feedback control. *Anal Chem* 69, 1706-14 (1997).
292. The Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408, 796-815 (2000).
293. Corthals, G. L., Wasinger, V. C., Hochstrasser, D. F. & Sanchez, J. C. The dynamic range of protein expression: A challenge for proteomic research. *Electrophoresis* 21, 1104-1115 (2000).
294. Celis, J. E. et al. A comprehensive two-dimensional gel protein database of noncultured unfractionated normal human epidermal keratinocytes: towards an integrated approach to the study of cell proliferation, differentiation and skin diseases. *Electrophoresis* 12, 802-72. (1991).
295. Burggraf, D. & Lottspeich, F. The further construction of the two-dimensional database of common human proteins. *Electrophoresis* 16, 831-40. (1995).
296. Lefkovits, I., Kettman, J. R. & Frey, J. R. Global analysis of gene expression in cells of the immune system I. Analytical limitations in obtaining sequence information on polypeptides in two-dimensional gel spots. *Electrophoresis* 21, 2688-2693 (2000).
297. Suss, K. H., Arkona, C., Manteuffel, R. & Adler, K. Calvin cycle multienzyme complexes are bound to chloroplast thylakoid membranes of higher plants in situ. *Proc Natl Acad Sci USA* 90, 5514-8 (1993).
298. D'Aderem, A. & Ulevitch, R. J. Toll-like receptors in the induction of the innate immune response. *Nature* 406, 782-787 (2000).
299. Neves, S. R., Ram, P. T. & Iyengar, R. G protein pathways. *Science* 296, 1636-9 (2002).
300. Hunter, T. The Croonian Lecture 1997. The phosphorylation of proteins on tyrosine: its role in cell growth and disease. *Philos Trans R Soc Lond B Biol Sci* 353, 583-605. (1998).
301. Pabo, C. O. & Sauer, R. T. Transcription factors: structural families and principles of DNA recognition. *Annual Review of Biochemistry* 61, 1053-95 (1992).

302. Kadonaga, J. T. Eukaryotic transcription: an interlaced network of transcription factors and chromatin-modifying machines. *Cell* 92, 307-13 (1998).
303. Wallin, E. & von Heijne, G. Genome-wide analysis of integral membrane proteins from eubacterial, archaean, and eukaryotic organisms. *Protein Science* 7, 1029-38 (1998).
304. Santoni, V. et al. Large scale characterization of plant plasma membrane proteins. *Biochimie* 81, 655-661 (1999).
305. Ideker, T. et al. Integrated genomic and proteomic analyses of a systematically perturbed metabolic network. *Science* 292, 929-34 (2001).
306. Pandey, A. et al. Identification of a novel immunoreceptor tyrosine-based activation motif-containing molecule, STAM2, by mass spectrometry and its involvement in growth factor and cytokine receptor signaling pathways. *J Biol Chem* 275, 38633-9. (2000).
307. Castagna, A. et al. Comparative two-dimensional mapping of prion protein isoforms in human cerebrospinal fluid and central nervous system. *Electrophoresis* 23, 339-46 (2002).
308. Eulgem, T., Rushton, P. J., Robatzek, S. & Somssich, I. E. The WRKY superfamily of plant transcription factors. *Trends in Plant Sciences* 5, 199-206 (2000).
309. Wilson, N. L., Schulz, B. L., Karlsson, N. G. & Packer, N. H. Sequential analysis of N- and O-linked glycosylation of 2D-PAGE separated glycoproteins. *Journal of Proteome Research* 1, 521-529 (2002).
310. Hellmann, H. & Estelle, M. Plant development: regulation by protein degradation. *Science* 297, 793-7 (2002).
311. Klose, J. & Feller, M. Genetic variability of proteins from plasma membranes and cytosols of mouse organs. *Biochem Genet* 19, 859-70. (1981).
312. Bukau, B. & Horwich, A. L. The Hsp70 and Hsp60 chaperone machines. *Cell* 92, 351-66 (1998).
313. Baumeister, W., Walz, J., Zuhl, F. & Seemuller, E. The proteasome: paradigm of a self-compartmentalizing protease. *Cell* 92, 367-80 (1998).
314. Ross-Macdonald, P. et al. Large-scale analysis of the yeast genome by transposon tagging and gene disruption. *Nature* 402, 413-8. (1999).
315. Mann, M. Quantitative proteomics? *Nat Biotechnol* 17, 954-5. (1999).
316. Krishna, R. G. & Wold, F. Post-translational modifications of proteins. *Adv. Enzymol. Relat. Mol. Biol.* 67, 265-298 (1993).
317. Celis, J. E. et al. Gene expression profiling: monitoring transcription and translation products using DNA microarrays and proteomics. *FEBS Letters* 480, 2-16 (2000).
318. Unlu, M., Morgan, M. E. & Minden, J. S. Difference gel electrophoresis: A single gel method for detecting changes in protein extracts. *Electrophoresis* 18, 2071-2077 (1997).
319. Syrovy, I. & Hodny, Z. Staining and Quantification of Proteins Separated by Polyacrylamide-Gel Electrophoresis. *Journal of Chromatography* 569, 175-196 (1991).
320. Steen, H. & Pandey, A. Proteomics goes quantitative: measuring protein abundance. *Trends Biotechnol* 20, 361 - 364 (2002).
321. Gygi, S. P. et al. Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat Biotechnol* 17, 994-999 (1999).
322. MacBeath, G. Proteomics comes to the surface. *Nat Biotechnol* 19, 828-9. (2001).
323. MacBeath, G. Protein microarrays and proteomics. *Nat Genet* 32, 526-532 (2002).
324. de Wildt, R. M., Mundy, C. R., Gorick, B. D. & Tomlinson, I. M. Antibody arrays for high-throughput screening of antibody-antigen interactions. *Nat Biotechnol* 18, 989-994 (2000).
325. Zhu, H. et al. Analysis of yeast protein kinases using protein chips. *Nat Genet* 26, 283-9. (2000).
326. Uetz, P. & Hughes, R. E. Systematic and large-scale two hybrid screens [Review]. *Current Opinion in Microbiology* 3, 303-308 (2000).
327. Ito, T. et al. Toward a protein-protein interaction map of the budding yeast: A comprehensive system to examine two-hybrid interactions in all possible combinations between the yeast proteins. *Proc Natl Acad Sci U S A* 97, 1143-7. (2000).
328. Sanger, F., Nicklen, S. & Coulson, A. R. DNA Sequencing with Chain-Terminating Inhibitors. *Proc Natl Acad Sci U S A* 74, 5463-5467 (1977).