

8. Appendix

8.1 Abbreviations

2D	two dimensional
a	atto (10^{-18})
ABTS	2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt
AP	alkaline phosphatase
β -Gal	β -galactosidase
BSA	bovine serum albumin
CCD	charge-coupled device
cDNA	complementaty DNA
Cy3	indocarbocyanine
Cy5	indodicarbocyanine
DDAO	9H-(1,3-dichloro-9,9- dimethylacridin-2-one-7-yl)
ddH ₂ O	double distilled water
DNA	deoxyribonucleic acid
Dp	<i>Dermatophagoides pteronyssinus</i>
DTT	1,4-Dithio-DL-threitol
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
ESI	electrospray ionisation
ESI-MS	electrospray ionisation-mass spectrometry
f	femto (10^{-15})
FAST	fluorescent Array Surface Technology
FDG	fluorescein digalactoside
FRET	fluorescence resonance energy transfer
*g	times gravity (9.81 m/s^2)
g	gram
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
GFP	green fluorescent protein
h	hour(s)

HRP	horseradish peroxidase
HSA	human serum albumin
IC ₅₀	50% inhibition concentration
ICAT	isotope-coded affinity tag
IgG	immunoglobulin G
IPTG	isopropyl-β-D-thiogalactoside
Ki	inhibition constant
l	litre
LC	liquid chromatography
μ	micro (10^{-6})
m	milli (10^{-3}) or meter
M	molar (mole/litre)
MALDI	matrix-assisted laser desorption ionization
Min	minutes
MIST	multiple spotting technique
MPIMG	Max-Planck-Institute for Molecular Genetics
mRNA	messenger RNA
mol	mole
MS	mass spectrometry
MWCO	molecular weight cut-off
n	nano (10^{-9})
Ni-NTA	nickel-nitrilotriacetic acid
NHS	N-hydroxysuccinimide
p	pico (10^{-12})
PAGE	polyacrylamide gel electrophoresis
PBS	phosphate buffered saline
PBS-T	phosphate buffered saline with Tween-20
PCR	polymerase chain reaction
PDMS	poly-dimethylsiloxane
PEG	polyethyleneglycol
PETG	2-phenylethyl β-D-thiogalactoside
Phe	phenylalanine
Pro	proline
PVDF	polyvinylidenefluoride

RCA	rolling circle amplification
RNA	ribonucleic acid
rpm	revolutions per minute
RTS	rapid translation system
s	second(s)
scFv	single-chain variable fragment
SDS	sodium dodecyl sulfate
SELDI	surface enhanced laser/desorption ionization
SPR	surface plasmon resonance
TBS	tris-buffered saline
TBS-T	tris-buffered saline with Tween®-20
TEMED	N,N,N’,N’-tetramethyl-ethylenediamine
TOF	time of flight
Tris	tris-hydroxymethylaminomethane
Tween-20	polyoxyethylene(20) sorbitan monolaurate
U	unit
z	zepto (10^{-21})
ZPPD	Z-D-Phe-Pro-
(v/v)	methoxypropylboroglycinepinanediol ester
(w/v)	volume per volume
	weight per volume

8.2 List of Figures

Figure 1: Microarray with 512 spots	9
Figure 2: Interior of spotting robot	9
Figure 3: Spotting head with 16 solid steel pins	10
Figure 4: Spotting head with 16 piezoelectric nozzles (courtesy of Dr. L. Nyarsik, MPIIMG)	10
Figure 5: Confocal laser scanner	11
Figure 6: Principle of confocal scanner (35)	11
Figure 7: Left: Photograph of the nanowell plate comprising 12x8 wells with a well volume of 1.5 µl. Right: Enlarged photograph of the nanowells with a spacing of 2.25 mm between each well.....	13
Figure 8: Labelling strategies. A displays direct and indirect labelling for antibody microarrays. B shows indirect labelling applied with protein microarrays, while C display the detection of scFv’s by labelled protein L and the indirect detection of protein via labelled aptamers and fluorescence resonance energy transfer	21
Figure 9: Photograph of assembled slides with a 30-µm gap by metal spacers.....	39

Figure 10: Spatial arrangement of the dilution rows. The schematic assembly of the molecules is demonstrated next to the scan. The numbers below the scan indicate the spotted amount of capture molecules in amol/spot	51
Figure 11: Graph of mean signal intensity versus spotted concentration of protein chip surfaces	52
Figure 12: Graph of mean signal intensity versus spotted concentration of antibody chip surfaces	53
Figure 13: Detection limits of protein chip surfaces with standard deviation error bars	54
Figure 14: Detection limits of antibody chip surfaces with standard deviation error bars	54
Figure 15: Scan of one microarray used for determination of pin coefficients	56
Figure 16: Pin coefficients for 150 micron pins	56
Figure 17: Pin coefficients for 100 micron pins	57
Figure 18: Scan of an antibody chip used for the screening of different betaine concentrations in the MIST-buffer	58
Figure 19: Diagram of average signal intensity versus betaine concentration	58
Figure 20: Scan of an antibody chip used for the screening of different PEG concentrations in the MIST-buffer	59
Figure 21: Diagram of average signal intensity versus Cy5-labelled HSA concentration for different PEG 3350 concentrations.....	60
Figure 22: Scan of an antibody chip used for the screening of different glycerol concentrations in the MIST-buffer	60
Figure 23: Diagram of average signal intensity versus Cy5-labelled HSA concentration for different glycerol concentrations	61
Figure 24: Scan of two microarrays used for the optimisation of the washing procedure. Slide A displays a slide washed with TBS-TT, while slide B shows a slide washed with TBS-T	62
Figure 25: Scan of an epoxy-coated slide with different protein concentrations for determination of the optimal protein concentration for scFv screening	63
Figure 26: Scan of two slides for comparison of the TG-1 strain (A) and the HB2151 strain (B) for expression of scFv's.....	64
Figure 27: Mean signal intensity versus surface coating for an horseradish peroxidase assay	65
Figure 28: Immunosorbent assay on an antibody microarray using MIST. Left: Scan of an antibody chip. Right: Scheme of the second spotting with Cy5-labelled HSA and illustration of the molecular assembly on the chip. Numbers in squares indicate spotted amount of Cy5-labeled HSA in amol per spot.....	66
Figure 29: Average signal intensity versus spotted concentration for the microarray shown in Figure 28 with standard deviation error bars	67
Figure 30: Average signal intensity versus spotted concentration of a protein microarray detected with monoclonal anti-HSA antibodies.....	67
Figure 31: Average signal intensity versus spotted concentration of an protein microarray detected with polyclonal anti-fibrinogen antibodies.....	67
Figure 32: Scan and quantification of spotted monoclonal anti-HSA (A) and polyclonal anti-fibrinogen antibodies (B), which were spotted in equimolar ratio on a protein chip.....	68
Figure 33: Denaturing SDS-PAGE of cyclophilin A and poly-ubiquitin with three ubiquitin moieties	69
Figure 34: Denaturing SDS-PAGE of CALM2 and PDLIM1.....	69
Figure 35: Dilution row of anti-poly-ubiquitin clone from position B1 (A) and E1 (B) using chip data from scanning with low intensities and ELISA data from scanning	

after 15-minute incubation. Left: Scan of the dilution row as well as the negative control spotted in sets of 10 spots per concentration step. Numbers indicate the dilution. Right: Mean signal intensity and mean absorbance at 405 nm versus dilution factor with standard deviations for the chip data. Since ELISA data was obtained from duplicates, no standard deviations were calculated for the ELISA data	70
Figure 36: Dilution row of anti-poly-ubiquitin clone from position B1 (A) and E1 (B) using chip data from scanning with high intensities and ELISA data from scanning after overnight incubation. Left: Scan of the dilution row as well as the negative control spotted in sets of 10 spots per concentration step. Numbers indicate the dilution. Right: Mean signal intensity and mean absorbance at 405 nm versus dilution factor with standard deviations for the chip data. Since ELISA data was obtained from duplicates, no standard deviations were calculated for the ELISA data	71
Figure 37: Scatter plot of mean signal intensity of the microarray data versus mean absorbance at 405 nm data from the ELISA. Four selections comprising 96 monoclonal scFv's were screened in parallel. Two dashed lines indicating microarray signal intensities above 1000 and ELISA absorbance values above 0.5 were introduced to allow easy localisation of clear signals.....	72
Figure 38: Ordered graphs of relative intensity\absorbance of the best 15 binders for each protein. Ordering was done according to the signal intensity obtained from the microarray data.....	73
Figure 39: Scans of microarrays for the inhibition assay of HRP using sodium azide as inhibitor	74
Figure 40: Mean signal intensity versus time of incubation diagram with standard deviation error bars for inhibition assay of HRP. Scanning and analysis was performed without incubation and after 2 hours of incubation	74
Figure 41: Mean signal intensity versus molecules of HRP diagram with standard deviation error bars for the detection limit assay of HRP. Scanning and analysis was performed without incubation and after 2 hours of incubation	75
Figure 42: Mean signal intensity versus concentration of EDTA diagram with standard deviation error bars for the inhibition assay of AP. Scanning and analysis was performed without incubation	76
Figure 43: Mean signal intensity versus molecules of AP diagram with standard deviation error bars for the detection limit assay. Scanning and analysis was performed without incubation and after 2 hours of incubation	76
Figure 44: Mean signal intensity versus concentration of PETG diagram with standard deviation error bars for inhibition assay of β -galactosidase. Scanning and analysis was performed without incubation	77
Figure 45: Mean signal intensity versus moles of β -galactosidase diagram with standard deviation error bars for the detection limit assay. Scanning and analysis was performed without incubation and after 2 hours of incubation	77
Figure 46: Mean signal intensity versus moles of cathepsin D diagram with standard deviation error bars for the detection limit assay of cathepsin D. Scanning and analysis was performed without incubation and after 1 hour of incubation	78
Figure 47: Diagram of mean signal intensity versus anti-HSA antibodies for the enzymatic signal amplification	79
Figure 48: Scan of protein chip incubated with thrombin	80
Figure 49: Scan of protein chip incubated without thrombin.....	81
Figure 50: Diagram of relative signal intensity versus immobilised component of slides with and without thrombin incubation	82

Figure 51: Scan of protein chip with ZDPP applied using MIST	83
Figure 52: Diagram of signal intensity versus immobilised substance for thrombin inhibitor assay.....	84
Figure 53: Emission spectra of two dilutions of transcription and translation mix containing GFP	85
Figure 54: Emission spectra of transcription and translation mix containing β-galactosidase and 50 nM FDG.....	85
Figure 55: Scan of nanowells and diagram of signal intensity versus dispensed volume for two dilutions of an undiluted cell-free expression mix containing GFP.....	86
Figure 56: Scan of nanowells and diagram of signal intensity versus dispensed volume of two dilutions of a pre-diluted cell-free expression mix containing GFP	86
Figure 57: Scan of nanowells and diagram of signal intensity versus dispensed volume for a dilution of an undiluted cell-free expression mix containing β-galactosidase .	87
Figure 58: Scan of nanowells and diagram for the undiluted cell-free expression of GFP in the nanowell plate. Two batches were prepared, one with P_{T7}-GFP-DNA (A), one lacking DNA (B).....	88
Figure 59: Scan of nanowells and diagram for the undiluted cell-free expression of β-galactosidase in the nanowell plate. Two batches were prepared, one with P_{T7}-GFP-DNA (A), one lacking DNA (B)	88
Figure 60: Scan of nanowells and diagram for the enzymatic inhibition assay in nanowells. (A) Assay with duplicates of PBS, 66 μM, 6.6 μM and 0.66 μM PETG (from left to right). (B) Assay with duplicates of PBS, 6.6 μM, 0.66 μM and 0.066 μM PETG	89
Figure 61: Principle of the fluorescence spectrometer. A displays the case for the measurement of fluorescence and B displays the scattering of the laser beam by particles in the solution	104

8.3 List of Tables

Table 1: Properties of the dyes used (119)	41
Table 2: Coefficients of variation for protein and antibody microarrays.....	53

8.4 Curriculum Vitae

Name: Philipp Angenendt
Date and place of birth: 26th of April 1978 in Emmerich, Germany

EDUCATION

since March 2002	Max-Planck-Institute for Molecular Genetics - Berlin Ph.D. studies, Protein Group, Dept. Lehrach Thesis: Studies on the optimisation and application of protein arrays
------------------	---

August 2001 – February 2002	Max-Planck-Institute for Molecular Genetics - Berlin Bachelor student, Protein Group, Dept. Lehrach Thesis: Comparison of different antibodies and surfaces in a protein array system (Grade: 1.0)
September 1998 – February 2002	Mannheim University of Applied Sciences – Mannheim Bachelor of Science with Honours in Biotechnology (Grade: 1.4)
July 1997 – April 1998	Military Service – Flensburg
September 1989 – June 1997	Collegium Augustinianum Gaesdonck – Goch Abitur (Grade: 2.1)

EXPERIENCE ABROAD

February 2001 – July 2001	University of Teesside - Middlesbrough, England International semester within bachelor studies
September 1992 – December 1992	Glenstal Abbey School – Murroe, Ireland Exchange student

PRACTICAL EXPERIENCE

August 2000 – January 2001	German Cancer Research Center – Heidelberg Practical Semester, Functional Genome Analysis Group
October 1998 – October 1999	Mannheim University of Applied Sciences - Mannheim Assistant, Department of Chemistry
May 1998 – August 1998	Deutsche Lebensmittel Union GmbH – Kleve Assistant, Quality Control

May 1997

Deutsche Lebensmittel Union GmbH – Kleve
Internship in quality assurance (biological and chemical departments)

AWARDS

November 2003

Mannheim University of Applied Sciences - Mannheim
Best student of the B.Sc. (hons) in Biotechnology course

CONSULTING

September 2003 - present

Scientific Advisory Team on Microarray Technology
Genetix Ltd., New Milton, England

8.5 Publications

Parts of this work have been published:

Angenendt, P., Glökler, J., Sobek, J., Lehrach, H. & Cahill, D. J. (2003) The next generation of protein microarray support materials: an evaluation for protein and antibody microarray applications. *Journal of Chromatography A* **1009**, 97-104.

Angenendt, P., Glökler, J., Konthur, Z., Lehrach, H. & Cahill, D. J. (2003) 3D protein microarrays: performing multiplex immunoassays on a single chip. *Analytical Chemistry* **75**, 4368-4372.

Glökler, J. & Angenendt, P. (2003) Protein and Antibody Microarray Technology. *Journal of Chromatography B*, **797**, 229-40.

Angenendt, P. (2003) Multiplexed Protein and Antibody Microarrays: Harnessing the Multiple Spotting Technique. *SCREENING*, **6**, 39-41.

Angenendt, P. (2003) Generierung multiplexer Protein Mikroarrays mit Hilfe der Multiple Spotting Technique *BIOforum*, **1-2**, 23-25.

Angenendt, P. & Glökler, J. (2004) Evaluation of Antibodies and Microarray Coatings as a Prerequisite for the Generation of Optimised Antibody Microarrays. In *Methods in Molecular Biology*. Edited by E. Fung. Totowa: The Humana Press, *in press*.

Angenendt, P., Nyarsik, L., Szaflarski, W., Glökler, J., Nierhaus, K. N., Lehrach, H., Cahill, D. J., Lueking, A. (2003) Cell-free protein expression and functional assay in nanowell chip. *Analytical Chemistry, in press*.

Angenendt, P., Wilde, J., Kiljanka, G., Baars, S., Cahill, D. J., Kreutzberger, J., Lehrach, H., Konthur, Z., Glökler, J. (2003) Seeing better through MIST: Evaluation of monoclonal recombinant antibody fragments on microarrays. *submitted*.

Kersten, B., Feilner, T., Angenendt, P., Giavalisco, P., Brenner, W., Bürkle, L. (2004) Proteomic approaches in plant biology. *submitted*.

A patent has been filed describing techniques established in this work:

Glökler, J., Angenendt, P., Kreutzberger, J., Lehrach, H. (2004) Method of performing multiplex assays on microarrays. US Patent Application

8.6 Acknowledgements

I would like to thank the following persons for their friendliness and support. Without you this thesis would not appear in this present form and my time at the Max-Planck-Institute would not have been so joyous and memorable.

In particular:

Dr. Jörn Glökler, for his tutoring and advice. For his ideas and impetus towards interesting questions and for his huge scientific background and experience on which I could rely all the times. For his visions, his continuous motivation and appreciation. Finally, for making me not only concentrating on the specifics, but also catching a glance at the “big” picture.

Prof. Dr. Dolores Cahill, for her support and the possibility to work in her excellent group. For her preparedness to answer questions or to discuss matter no matter how late it was. For

her offer to conduct my PhD at the Max-Planck Institute and for the possibility to learn from her.

Prof. Dr. Hans Lehrach, for being a supervisor of my thesis. For all his support with regard to my PhD application at the FU Berlin and for the helpful discussions.

Prof. Dr. Volker Erdmann, for being an evaluator of my thesis.

Dr. Konrad Büssow, for being an advisor to my studies and for all the fruitful discussions.

Dr. Zoltán Konthur, for his help and advice with regard to my PhD applications and his support in our cooperations.

Dr. Jürgen Kreutzberger, for his support and assistance, especially in the time I worked for the University of Bochum.

Dr. Sabine Baars, for her friendliness, the pleasant cooperation and for her corrections of thesis thesis.

Dr. Claudia Falter and Johanna Belart, for finding the right expressions with regard to all formalities and for always bringing me the really good news.

Tanja Feilner, Claudia Gutjahr, Sabine Horn, Dr. Birgit Kersten, Andrea Koenig, Sigrid Steller, Gregor Kijanka, Armin Kramer, Alexandra Possling and Silke Wehrmeyer for their help and the fun we had in the lab. For their continuous willingness to celebrate publications with me.

Dr. James Adjaye, Dr. Knud Nierhaus, Witold Szaflarski, Dr .Lajos Nyarsik, Dr. Jens Sobek, Jeannine Wilde, Dr. Frauke Melchior, Tanja Büsgen, Dr. Uli Stelzl, Dr. Angelika Lüking, Dr. Sylvia Krobisch, Ute Nonhoff and Prof. Erich Wanker, for the interesting cooperations. Although, not all of them could be brought to an end, I enjoyed the planning and discussion very much.

To all of those, who are not mentioned here explicitly: Thanks.

My family, for their confidence in me and my decisions. For their continued love and support throughout the last 25 years.

Monika, for her love, understanding and belief in me. For making me enjoying every day with her.

8.7 References

- (1) Gygi, S. P.; Rochon, Y.; Franz, B. R.; Aebersold, R. (1999) Correlation between protein and mRNA abundance in yeast. *Mol Cell Biol* **19**, 1720-30.
- (2) Anderson, L.; Seilhamer, J. (1997) A comparison of selected mRNA and protein abundances in human liver. *Electrophoresis* **18**, 533-7.
- (3) Klose, J. (1975) Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. A novel approach to testing for induced point mutations in mammals. *Humangenetik* **26**, 231-43.
- (4) O'Farrell, P. H. (1975) High resolution two-dimensional electrophoresis of proteins. *J Biol Chem* **250**, 4007-21.
- (5) Harrington, M. G.; Lee, K. H.; Yun, M.; Zewert, T.; Bailey, J. E.; Hood, L. (1993) Mechanical precision in two-dimensional electrophoresis can improve protein spot positional reproducibility. *Appl Theor Electrophor* **3**, 347-53.
- (6) Langen, H.; Roder, D.; Juraville, J. F.; Fountoulakis, M. (1997) Effect of protein application mode and acrylamide concentration on the resolution of protein spots separated by two-dimensional gel electrophoresis. *Electrophoresis* **18**, 2085-90.
- (7) Gorg, A.; Obermaier, C.; Boguth, G.; Weiss, W. (1999) Recent developments in two-dimensional gel electrophoresis with immobilized pH gradients: wide pH gradients up to pH 12, longer separation distances and simplified procedures. *Electrophoresis* **20**, 712-7.
- (8) Wildgruber, R.; Harder, A.; Obermaier, C.; Boguth, G.; Weiss, W.; Fey, S. J.; Larsen, P. M.; Gorg, A. (2000) Towards higher resolution: two-dimensional electrophoresis of *Saccharomyces cerevisiae* proteins using overlapping narrow immobilized pH gradients. *Electrophoresis* **21**, 2610-6.
- (9) Fivaz, M.; Vilbois, F.; Pasquali, C.; van der Goot, F. G. (2000) Analysis of glycosyl phosphatidylinositol-anchored proteins by two-dimensional gel electrophoresis. *Electrophoresis* **21**, 3351-6.

- (10) Lehr, S.; Kotzka, J.; Knebel, B.; Schiller, M.; Krone, W.; Muller-Wieland, D. (2002) Primary skin fibroblasts as human model system for proteome analysis. *Proteomics* **2**, 280-7.
- (11) Wandschneider, S.; Fehring, V.; Jacobs-Emeis, S.; Thiesen, H. J.; Lohr, M. (2001) Autoimmune pancreatic disease: preparation of pancreatic juice for proteome analysis. *Electrophoresis* **22**, 4383-90.
- (12) Taniguchi, N.; Ekuni, A.; Ko, J. H.; Miyoshi, E.; Ikeda, Y.; Ihara, Y.; Nishikawa, A.; Honke, K.; Takahashi, M. (2001) A glycomic approach to the identification and characterization of glycoprotein function in cells transfected with glycosyltransferase genes. *Proteomics* **1**, 239-47.
- (13) Sarioglu, H.; Lottspeich, F.; Walk, T.; Jung, G.; Eckerskorn, C. (2000) Deamidation as a widespread phenomenon in two-dimensional polyacrylamide gel electrophoresis of human blood plasma proteins. *Electrophoresis* **21**, 2209-18.
- (14) Kaufmann, H.; Bailey, J. E.; Fussenegger, M. (2001) Use of antibodies for detection of phosphorylated proteins separated by two-dimensional gel electrophoresis. *Proteomics* **1**, 194-9.
- (15) Lopez, M. F. (2000) Better approaches to finding the needle in a haystack: optimizing proteome analysis through automation. *Electrophoresis* **21**, 1082-93.
- (16) Lilley, K. S.; Razzaq, A.; Dupree, P. (2002) Two-dimensional gel electrophoresis: recent advances in sample preparation, detection and quantitation. *Curr Opin Chem Biol* **6**, 46-50.
- (17) Gygi, S. P.; Corthals, G. L.; Zhang, Y.; Rochon, Y.; Aebersold, R. (2000) Evaluation of two-dimensional gel electrophoresis-based proteome analysis technology. *Proc Natl Acad Sci U S A* **97**, 9390-5.
- (18) Corthals, G. L.; Wasinger, V. C.; Hochstrasser, D. F.; Sanchez, J. C. (2000) The dynamic range of protein expression: a challenge for proteomic research. *Electrophoresis* **21**, 1104-15.
- (19) Schutt, F.; Ueberle, B.; Schnolzer, M.; Holz, F. G.; Kopitz, J. (2002) Proteome analysis of lipofuscin in human retinal pigment epithelial cells. *FEBS Lett* **528**, 217-21.
- (20) Hoving, S.; Voshol, H.; van Oostrum, J. (2000) Towards high performance two-dimensional gel electrophoresis using ultrazoom gels. *Electrophoresis* **21**, 2617-21.

- (21) Hoving, S.; Gerrits, B.; Voshol, H.; Muller, D.; Roberts, R. C.; van Oostrum, J. (2002) Preparative two-dimensional gel electrophoresis at alkaline pH using narrow range immobilized pH gradients. *Proteomics* **2**, 127-34.
- (22) Santoni, V.; Kieffer, S.; Desclaux, D.; Masson, F.; Rabilloud, T. (2000) Membrane proteomics: use of additive main effects with multiplicative interaction model to classify plasma membrane proteins according to their solubility and electrophoretic properties. *Electrophoresis* **21**, 3329-44.
- (23) Molloy, M. P.; Phadke, N. D.; Chen, H.; Tyldesley, R.; Garfin, D. E.; Maddock, J. R.; Andrews, P. C. (2002) Profiling the alkaline membrane proteome of *Caulobacter crescentus* with two-dimensional electrophoresis and mass spectrometry. *Proteomics* **2**, 899-910.
- (24) Patton, W. F. (1999) Proteome analysis. II. Protein subcellular redistribution: linking physiology to genomics via the proteome and separation technologies involved. *J Chromatogr B Biomed Sci Appl* **722**, 203-23.
- (25) Link, A. J.; Eng, J.; Schieltz, D. M.; Carmack, E.; Mize, G. J.; Morris, D. R.; Garvik, B. M.; Yates, J. R., 3rd (1999) Direct analysis of protein complexes using mass spectrometry. *Nat Biotechnol* **17**, 676-82.
- (26) Madoz-Gurpide, J.; Wang, H.; Misek, D. E.; Brichory, F.; Hanash, S. M. (2001) Protein based microarrays: a tool for probing the proteome of cancer cells and tissues. *Proteomics* **1**, 1279-87.
- (27) Gygi, S. P.; Rist, B.; Gerber, S. A.; Turecek, F.; Gelb, M. H.; Aebersold, R. (1999) Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat Biotechnol* **17**, 994-9.
- (28) Ochi, H.; Horiuchi, I.; Araki, N.; Toda, T.; Araki, T.; Sato, K.; Murai, H.; Osoegawa, M.; Yamada, T.; Okamura, K.; Ogino, T.; Mizumoto, K.; Yamashita, H.; Saya, H.; Kira, J. (2002) Proteomic analysis of human brain identifies alpha-enolase as a novel autoantigen in Hashimoto's encephalopathy. *FEBS Lett* **528**, 197-202.
- (29) Gehanne, S.; Cecconi, D.; Carboni, L.; Righetti, P. G.; Domenici, E.; Hamdan, M. (2002) Quantitative analysis of two-dimensional gel-separated proteins using isotopically marked alkylating agents and matrix-assisted laser desorption/ionization mass spectrometry. *Rapid Commun Mass Spectrom* **16**, 1692-8.
- (30) Chivasa, S.; Ndimba, B. K.; Simon, W. J.; Robertson, D.; Yu, X. L.; Knox, J. P.; Bolwell, P.; Slabas, A. R. (2002) Proteomic analysis of the *Arabidopsis thaliana* cell wall. *Electrophoresis* **23**, 1754-65.

- (31) Karas, M.; Hillenkamp, F. (1988) Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *Anal Chem* **60**, 2299-301.
- (32) Fenn, J. B.; Mann, M.; Meng, C. K.; Wong, S. F.; Whitehouse, C. M. (1989) Electrospray ionization for mass spectrometry of large biomolecules. *Science* **246**, 64-71.
- (33) Choudhary, J. S.; Blackstock, W. P.; Creasy, D. M.; Cottrell, J. S. (2001) Matching peptide mass spectra to EST and genomic DNA databases. *Trends Biotechnol* **19**, S17-22.
- (34) Soldatov, A. V.; Nabirochkina, E. N.; Georgieva, S. G.; Eickhoff, H. (2001) Adjustment of transfer tools for the production of micro- and macroarrays. *Biotechniques* **31**, 848, 850, 852, 854.
- (35) Schermer, M. J. In *DNA Microarrays*; Schena, M., Ed.; Oxford University Press: New York, 1999, pp 17 - 42.
- (36) Ge, H. (2000) UPA, a universal protein array system for quantitative detection of protein-protein, protein-DNA, protein-RNA and protein-ligand interactions. *Nucleic Acids Res* **28**, e3.
- (37) Lueking, A.; Horn, M.; Eickhoff, H.; Büssow, K.; Lehrach, H.; Walter, G. (1999) Protein microarrays for gene expression and antibody screening. *Anal Biochem* **270**, 103-11.
- (38) Mendoza, L. G.; McQuary, P.; Mongan, A.; Gangadharan, R.; Brignac, S.; Eggers, M. (1999) High-throughput microarray-based enzyme-linked immunosorbent assay (ELISA). *Biotechniques* **27**, 778-80, 782-6, 788.
- (39) de Wildt, R. M.; Mundy, C. R.; Gorick, B. D.; Tomlinson, I. M. (2000) Antibody arrays for high-throughput screening of antibody-antigen interactions. *Nat Biotechnol* **18**, 989-94.
- (40) MacBeath, G.; Schreiber, S. L. (2000) Printing proteins as microarrays for high-throughput function determination. *Science* **289**, 1760-3.
- (41) Haab, B. B.; Dunham, M. J.; Brown, P. O. (2001) Protein microarrays for highly parallel detection and quantitation of specific proteins and antibodies in complex solutions. *Genome Biol* **2**, RESEARCH0004.
- (42) Angenendt, P.; Glöckler, J.; Murphy, D.; Lehrach, H.; Cahill, D. J. (2002) Toward optimized antibody microarrays: a comparison of current microarray support materials. *Anal Biochem* **309**, 253-60.

- (43) Arenkov, P.; Kukhtin, A.; Gemmell, A.; Voloshchuk, S.; Chupeeva, V.; Mirzabekov, A. (2000) Protein microchips: use for immunoassay and enzymatic reactions. *Anal Biochem* **278**, 123-31.
- (44) Kiyonaka, S.; Sada, K.; Yoshimura, I.; Shinkai, S.; Kato, N.; Hamachi, I. (2004) Semi-wet peptide/protein array using supramolecular hydrogel. *Nat Mater* **3**, 58-64. Epub 2003 Dec 7.
- (45) Kusnezow, W.; Jacob, A.; Walijew, A.; Diehl, F.; Hoheisel, J. D. (2003) Antibody microarrays: An evaluation of production parameters. *Proteomics* **3**, 254-64.
- (46) Peluso, P.; Wilson, D. S.; Do, D.; Tran, H.; Venkatasubbaiah, M.; Quincy, D.; Heidecker, B.; Poindexter, K.; Tolani, N.; Phelan, M.; Witte, K.; Jung, L. S.; Wagner, P.; Nock, S. (2003) Optimizing antibody immobilization strategies for the construction of protein microarrays. *Anal Biochem* **312**, 113-24.
- (47) Zhu, H.; Klemic, J. F.; Chang, S.; Bertone, P.; Casamayor, A.; Klemic, K. G.; Smith, D.; Gerstein, M.; Reed, M. A.; Snyder, M. (2000) Analysis of yeast protein kinases using protein chips. *Nat Genet* **26**, 283-9.
- (48) Xia, Y.; Kim, E.; Zhao, X. M.; Rogers, J. A.; Prentiss, M.; Whitesides, G. M. (1996) Complex Optical Surfaces Formed by Replica Molding Against Elastomeric Masters. *Science* **273**, 347-9.
- (49) Biran, I.; Walt, D. R. (2002) Optical imaging fiber-based single live cell arrays: a high-density cell assay platform. *Anal Chem* **74**, 3046-54.
- (50) Bernhard, D. D.; Mall, S.; Pantano, P. (2001) Fabrication and characterization of microwell array chemical sensors. *Anal Chem* **73**, 2484-90.
- (51) Cohen, C. B.; Chin-Dixon, E.; Jeong, S.; Nikiforov, T. T. (1999) A microchip-based enzyme assay for protein kinase A. *Anal Biochem* **273**, 89-97.
- (52) Hadd, A. G.; Raymond, D. E.; Halliwell, J. W.; Jacobson, S. C.; Ramsey, J. M. (1997) Microchip device for performing enzyme assays. *Anal Chem* **69**, 3407-12.
- (53) Hutchens, T. W.; Yip, T.-T. (1993) New desorption strategies for the mass spectrometric analysis of macromolecules. *Rapid Comm. Mass Spectrom.* **7**, 576-580.
- (54) Espina, V.; Dettloff, K. A.; Cowherd, S.; Petricoin, E. R. E.; Liotta, L. A. (2004) Use of proteomic analysis to monitor responses to biological therapies. *Expert Opin Biol Ther* **4**, 83-93.
- (55) Zheng, Y.; Xu, Y.; Ye, B.; Lei, J.; Weinstein, M. H.; O'Leary, M. P.; Richie, J. P.; Mok, S. C.; Liu, B. C. (2003) Prostate carcinoma tissue proteomics for biomarker discovery. *Cancer* **98**, 2576-82.

- (56) Malmqvist, M.; Karlsson, R. (1997) Biomolecular interaction analysis: affinity biosensor technologies for functional analysis of proteins. *Curr Opin Chem Biol* **1**, 378-83.
- (57) Joos, T. O.; Schrenk, M.; Hopfl, P.; Kroger, K.; Chowdhury, U.; Stoll, D.; Schorner, D.; Durr, M.; Herick, K.; Rupp, S.; Sohn, K.; Hammerle, H. (2000) A microarray enzyme-linked immunosorbent assay for autoimmune diagnostics. *Electrophoresis* **21**, 2641-50.
- (58) Robinson, W. H.; DiGennaro, C.; Hueber, W.; Haab, B. B.; Kamachi, M.; Dean, E. J.; Fournel, S.; Fong, D.; Genovese, M. C.; de Vegvar, H. E.; Skriner, K.; Hirschberg, D. L.; Morris, R. I.; Muller, S.; Pruijn, G. J.; van Venrooij, W. J.; Smolen, J. S.; Brown, P. O.; Steinman, L.; Utz, P. J. (2002) Autoantigen microarrays for multiplex characterization of autoantibody responses. *Nat Med* **8**, 295-301.
- (59) Sreekumar, A.; Nyati, M. K.; Varambally, S.; Barrette, T. R.; Ghosh, D.; Lawrence, T. S.; Chinnaiyan, A. M. (2001) Profiling of cancer cells using protein microarrays: discovery of novel radiation-regulated proteins. *Cancer Res* **61**, 7585-93.
- (60) Hiller, R.; Laffer, S.; Harwanegg, C.; Huber, M.; Schmidt, W. M.; Twardosz, A.; Barletta, B.; Becker, W. M.; Blaser, K.; Breiteneder, H.; Chapman, M.; Crameri, R.; Duchene, M.; Ferreira, F.; Fiebig, H.; Hoffmann-Sommergruber, K.; King, T. P.; Kleber-Janke, T.; Kurup, V. P.; Lehrer, S. B.; Lidholm, J.; Muller, U.; Pini, C.; Reese, G.; Scheiner, O.; Scheynius, A.; Shen, H. D.; Spitzauer, S.; Suck, R.; Swoboda, I.; Thomas, W.; Tinghino, R.; Van Hage-Hamsten, M.; Virtanen, T.; Kraft, D.; Muller, M. W.; Valenta, R. (2002) Microarrayed allergen molecules: diagnostic gatekeepers for allergy treatment. *Faseb J* **16**, 414-6.
- (61) Kim, T. E.; Park, S. W.; Cho, N. Y.; Choi, S. Y.; Yong, T. S.; Nahm, B. H.; Lee, S.; Noh, G. (2002) Quantitative measurement of serum allergen-specific IgE on protein chip. *Exp Mol Med* **34**, 152-8.
- (62) Samsonova, J. V.; Baxter, G. A.; Crooks, S. R.; Small, A. E.; Elliott, C. T. (2002) Determination of ivermectin in bovine liver by optical immunobiosensor. *Biosens Bioelectron* **17**, 523-9.
- (63) Huang, R. P.; Huang, R.; Fan, Y.; Lin, Y. (2001) Simultaneous detection of multiple cytokines from conditioned media and patient's sera by an antibody-based protein array system. *Anal Biochem* **294**, 55-62.
- (64) Paweletz, C. P.; Charboneau, L.; Bichsel, V. E.; Simone, N. L.; Chen, T.; Gillespie, J. W.; Emmert-Buck, M. R.; Roth, M. J.; Petricoin, I. E.; Liotta, L. A. (2001) Reverse

- phase protein microarrays which capture disease progression show activation of pro-survival pathways at the cancer invasion front. *Oncogene* **20**, 1981-9.
- (65) Nishizuka, S.; Charboneau, L.; Young, L.; Major, S.; Reinhold, W. C.; Waltham, M.; Kourou-Mehr, H.; Bussey, K. J.; Lee, J. K.; Espina, V.; Munson, P. J.; Petricoin, E., 3rd; Liotta, L. A.; Weinstein, J. N. (2003) Proteomic profiling of the NCI-60 cancer cell lines using new high-density reverse-phase lysate microarrays. *Proc Natl Acad Sci U S A* **100**, 14229-34. Epub 2003 Nov 17.
- (66) Knezevic, V.; Leethanakul, C.; Bichsel, V. E.; Worth, J. M.; Prabhu, V. V.; Gutkind, J. S.; Liotta, L. A.; Munson, P. J.; Petricoin, E. F., 3rd; Krizman, D. B. (2001) Proteomic profiling of the cancer microenvironment by antibody arrays. *Proteomics* **1**, 1271-8.
- (67) Nielsen, U. B.; Cardone, M. H.; Sinskey, A. J.; MacBeath, G.; Sorger, P. K. (2003) Profiling receptor tyrosine kinase activation by using Ab microarrays. *Proc Natl Acad Sci U S A* **22**, 22.
- (68) Fang, Y.; Frutos, A. G.; Lahiri, J. (2002) Membrane protein microarrays. *J Am Chem Soc* **124**, 2394-5.
- (69) Houseman, B. T.; Huh, J. H.; Kron, S. J.; Mrksich, M. (2002) Peptide chips for the quantitative evaluation of protein kinase activity. *Nat Biotechnol* **20**, 270-4.
- (70) Zhu, H.; Bilgin, M.; Bangham, R.; Hall, D.; Casamayor, A.; Bertone, P.; Lan, N.; Jansen, R.; Bidlingmaier, S.; Houfek, T.; Mitchell, T.; Miller, P.; Dean, R. A.; Gerstein, M.; Snyder, M. (2001) Global analysis of protein activities using proteome chips. *Science* **293**, 2101-5.
- (71) Holt, L. J.; Bussow, K.; Walter, G.; Tomlinson, I. M. (2000) By-passing selection: direct screening for antibody-antigen interactions using protein arrays. *Nucleic Acids Res* **28**, E72.
- (72) Kersten, B.; Feilner, T.; Kramer, A.; Wehrmeyer, S.; Possling, A.; Witt, I.; Zanor, M. I.; Stracke, R.; Lueking, A.; Kreutzberger, J.; Lehrach, H.; Cahill, D. J. (2003) Generation of Arabidopsis protein chips for antibody and serum screening. *Plant Mol Biol*.
- (73) Pasqualini, R.; Arap, W. (2003) Hybridoma-free generation of monoclonal antibodies. *Proc Natl Acad Sci U S A* **19**, 19.
- (74) Nissim, A.; Hoogenboom, H. R.; Tomlinson, I. M.; Flynn, G.; Midgley, C.; Lane, D.; Winter, G. (1994) Antibody fragments from a 'single pot' phage display library as immunochemical reagents. *Embo J* **13**, 692-8.

- (75) Krebs, B.; Rauchenberger, R.; Reiffert, S.; Rothe, C.; Tesar, M.; Thomassen, E.; Cao, M.; Dreier, T.; Fischer, D.; Hoss, A.; Inge, L.; Knappik, A.; Marget, M.; Pack, P.; Meng, X. Q.; Schier, R.; Sohlemann, P.; Winter, J.; Wolle, J.; Kretzschmar, T. (2001) High-throughput generation and engineering of recombinant human antibodies. *Journal of Immunological Methods* **254**, 67-84.
- (76) Casey, J. L.; Keep, P. A.; Chester, K. A.; Robson, L.; Hawkins, R. E.; Begent, R. H. (1995) Purification of bacterially expressed single chain Fv antibodies for clinical applications using metal chelate chromatography. *J Immunol Methods* **179**, 105-16.
- (77) He, M.; Taussig, M. J. (1997) Antibody-ribosome-mRNA (ARM) complexes as efficient selection particles for in vitro display and evolution of antibody combining sites. *Nucleic Acids Res* **25**, 5132-4.
- (78) Hanes, J.; Schaffitzel, C.; Knappik, A.; Pluckthun, A. (2000) Picomolar affinity antibodies from a fully synthetic naive library selected and evolved by ribosome display. *Nat Biotechnol* **18**, 1287-92.
- (79) Barrick, J. E.; Takahashi, T. T.; Balakin, A.; Roberts, R. W. (2001) Selection of RNA-binding peptides using mRNA-peptide fusions. *Methods* **23**, 287-93.
- (80) Hochuli, E.; Dobeli, H.; Schacher, A. (1987) New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues. *J Chromatogr* **411**, 177-84.
- (81) Mukhija, R.; Rupa, P.; Pillai, D.; Garg, L. C. (1995) High-level production and one-step purification of biologically active human growth hormone in Escherichia coli. *Gene* **165**, 303-6.
- (82) Joshi, B.; Janda, L.; Stoytcheva, Z.; Tichy, P. (2000) PkWA, a WD-repeat protein, is expressed in spore-derived mycelium of Thermomonospora curvata and phosphorylation of its WD domain could act as a molecular switch. *Microbiology* **146**, 3259-67.
- (83) Negro, A.; Onisto, M.; Masiero, L.; Garbisa, S. (1995) Synthesis and refolding of human TIMP-2 from E. coli, with specific activity for MMP-2. *FEBS Lett* **360**, 52-6.
- (84) Zahn, R.; von Schroetter, C.; Wuthrich, K. (1997) Human prion proteins expressed in Escherichia coli and purified by high-affinity column refolding. *FEBS Lett* **417**, 400-4.
- (85) Büssow, K.; Nordhoff, E.; Lubbert, C.; Lehrach, H.; Walter, G. (2000) A human cDNA library for high-throughput protein expression screening. *Genomics* **65**, 1-8.

- (86) Voss, S.; Skerra, A. (1997) Mutagenesis of a flexible loop in streptavidin leads to higher affinity for the Strep-tag II peptide and improved performance in recombinant protein purification. *Protein Eng* **10**, 975-82.
- (87) Stofko-Hahn, R. E.; Carr, D. W.; Scott, J. D. (1992) A single step purification for recombinant proteins. Characterization of a microtubule associated protein (MAP 2) fragment which associates with the type II cAMP-dependent protein kinase. *FEBS Lett* **302**, 274-8.
- (88) Chong, S.; Mersha, F. B.; Comb, D. G.; Scott, M. E.; Landry, D.; Vence, L. M.; Perler, F. B.; Benner, J.; Kucera, R. B.; Hirvonen, C. A.; Pelletier, J. J.; Paulus, H.; Xu, M. Q. (1997) Single-column purification of free recombinant proteins using a self-cleavable affinity tag derived from a protein splicing element. *Gene* **192**, 271-81.
- (89) Smith, D. B.; Johnson, K. S. (1988) Single-step purification of polypeptides expressed in Escherichia coli as fusions with glutathione S-transferase. *Gene* **67**, 31-40.
- (90) di Guan, C.; Li, P.; Riggs, P. D.; Inouye, H. (1988) Vectors that facilitate the expression and purification of foreign peptides in Escherichia coli by fusion to maltose-binding protein. *Gene* **67**, 21-30.
- (91) Dickason, R. R.; Edwards, R. A.; Bryan, J.; Huston, D. P. (1995) Versatile E. coli thioredoxin specific monoclonal antibodies afford convenient analysis and purification of prokaryote expressed soluble fusion protein. *J Immunol Methods* **185**, 237-44.
- (92) Braun, P.; Hu, Y.; Shen, B.; Halleck, A.; Koundinya, M.; Harlow, E.; LaBaer, J. (2002) Proteome-scale purification of human proteins from bacteria. *Proc Natl Acad Sci U S A* **99**, 2654-9.
- (93) Lamla, T.; Erdmann, V. A. (2004) The Nano-tag, a streptavidin-binding peptide for the purification and detection of recombinant proteins. *Protein Expr Purif* **33**, 39-47.
- (94) Boettner, M.; Prinz, B.; Holz, C.; Stahl, U.; Lang, C. (2002) High-throughput screening for expression of heterologous proteins in the yeast *Pichia pastoris*. *J Biotechnol* **99**, 51-62.
- (95) Holz, C.; Hesse, O.; Bolotina, N.; Stahl, U.; Lang, C. (2002) A micro-scale process for high-throughput expression of cDNAs in the yeast *Saccharomyces cerevisiae*. *Protein Expr Purif* **25**, 372-8.
- (96) Allen, S. V.; Miller, E. S. (1999) RNA-binding properties of in vitro expressed histidine-tagged RB69 RegA translational repressor protein. *Anal Biochem* **269**, 32-7.

- (97) He, M.; Taussig, M. J. (2001) Single step generation of protein arrays from DNA by cell-free expression and in situ immobilisation (PISA method). *Nucleic Acids Res* **29**, E73-3.
- (98) Hansson, M.; Ringdahl, J.; Robert, A.; Power, U.; Goetsch, L.; Nguyen, T. N.; Uhlen, M.; Stahl, S.; Nygren, P. A. (1999) An in vitro selected binding protein (affibody) shows conformation-dependent recognition of the respiratory syncytial virus (RSV) G protein. *Immunotechnology* **4**, 237-52.
- (99) Koide, A.; Bailey, C. W.; Huang, X.; Koide, S. (1998) The fibronectin type III domain as a scaffold for novel binding proteins. *J Mol Biol* **284**, 1141-51.
- (100) Beste, G.; Schmidt, F. S.; Stibora, T.; Skerra, A. (1999) Small antibody-like proteins with prescribed ligand specificities derived from the lipocalin fold. *Proc Natl Acad Sci U S A* **96**, 1898-903.
- (101) Forrer, P.; Stumpp, M. T.; Binz, H. K.; Pluckthun, A. (2003) A novel strategy to design binding molecules harnessing the modular nature of repeat proteins. *FEBS Lett* **539**, 2-6.
- (102) Schneider, D.; Gold, L.; Platt, T. (1993) Selective enrichment of RNA species for tight binding to Escherichia coli rho factor. *Faseb J* **7**, 201-7.
- (103) Smith, D.; Collins, B. D.; Heil, J.; Koch, T. H. (2003) Sensitivity and specificity of photoaptamer probes. *Molecular & Cellular Proteomics* **2**, 11-8.
- (104) Golden, M. C.; Resing, K. A.; Collins, B. D.; Willis, M. C.; Koch, T. H. (1999) Mass spectral characterization of a protein-nucleic acid photocrosslink. *Protein Science* **8**, 2806-12.
- (105) Petach, H.; Gold, L. (2002) Dimensionality is the issue: use of photoaptamers in protein microarrays. *Current Opinion in Biotechnology* **13**, 309-14.
- (106) Schweitzer, B.; Wiltshire, S.; Lambert, J.; O'Malley, S.; Kukanskis, K.; Zhu, Z.; Kingsmore, S. F.; Lizardi, P. M.; Ward, D. C. (2000) Inaugural article: immunoassays with rolling circle DNA amplification: a versatile platform for ultrasensitive antigen detection. *Proc Natl Acad Sci U S A* **97**, 10113-9.
- (107) Shao, W.; Zhou, Z.; Laroche, I.; Lu, H.; Zong, Q.; Patel, D. D.; Kingsmore, S.; Piccoli, S. P. (2003) Optimization of Rolling-Circle Amplified Protein Microarrays for Multiplexed Protein Profiling. *J Biomed Biotechnol* **2003**, 299-307.
- (108) Wu, X.; Liu, H.; Liu, J.; Haley, K. N.; Treadway, J. A.; Larson, J. P.; Ge, N.; Peale, F.; Bruchez, M. P. (2003) Immunofluorescent labeling of cancer marker Her2 and other cellular targets with semiconductor quantum dots. *Nat Biotechnol* **21**, 41-6.

- (109) Brody, E. N.; Willis, M. C.; Smith, J. D.; Jayasena, S.; Zichi, D.; Gold, L. (1999) The use of aptamers in large arrays for molecular diagnostics. *Mol Diagn* **4**, 381-8.
- (110) Hamaguchi, N.; Ellington, A.; Stanton, M. (2001) Aptamer beacons for the direct detection of proteins. *Anal Biochem* **294**, 126-31.
- (111) Li, J. J.; Fang, X.; Tan, W. (2002) Molecular aptamer beacons for real-time protein recognition. *Biochem Biophys Res Commun* **292**, 31-40.
- (112) Nelson, B. P.; Liles, M. R.; Frederick, K. B.; Corn, R. M.; Goodman, R. M. (2002) Label-free detection of 16S ribosomal RNA hybridization on reusable DNA arrays using surface plasmon resonance imaging. *Environ Microbiol* **4**, 735-43.
- (113) Dare, T. O.; Davies, H. A.; Turton, J. A.; Lomas, L.; Williams, T. C.; York, M. J. (2002) Application of surface-enhanced laser desorption/ionization technology to the detection and identification of urinary parvalbumin-alpha: a biomarker of compound-induced skeletal muscle toxicity in the rat. *Electrophoresis* **23**, 3241-51.
- (114) Hallborn, J.; Carlsson, R. (2002) Automated screening procedure for high-throughput generation of antibody fragments. *Biotechniques Suppl*, 30-7.
- (115) Smith, D.; Collins, B. D.; Heil, J.; Koch, T. H. (2003) Sensitivity and specificity of photoaptamer probes. *Mol Cell Proteomics* **2**, 11-8.
- (116) Kukar, T.; Eckendorf, S.; Gu, Y.; Lian, W.; Megginson, M.; She, J. X.; Wu, D. (2002) Protein microarrays to detect protein-protein interactions using red and green fluorescent proteins. *Anal Biochem* **306**, 50-4.
- (117) Angenendt, P. , Mannheim University of Applied Sciences, Mannheim, Germany, 2003.
- (118) Jackson_ImmunoResearch_Laboratories (2001) Technical Information on Probes Conjugated to Our Antibodies and Other Proteins: Cyanine Dyes - Cy2, Cy3, and Cy5 [WWW Document]. URL <http://www.jacksonimmuno.com/2001site/home/catalog/f-cy3-5.htm> .
- (119) DPC_Biermann_GmbH (2001) New Secondary Antibodies Conjugated to Cyanine [WWW Document]. URL <http://www.acris-antibodies.de/deutsch/focus006.htm> .
- (120) Winter, G.; Griffiths, A. D.; Hawkins, R. E.; Hoogenboom, H. R. (1994) Making antibodies by phage display technology. *Annu Rev Immunol* **12**, 433-55.
- (121) Hoogenboom, H. R.; Griffiths, A. D.; Johnson, K. S.; Chiswell, D. J.; Hudson, P.; Winter, G. (1991) Multi-subunit proteins on the surface of filamentous phage: methodologies for displaying antibody (Fab) heavy and light chains. *Nucleic Acids Res* **19**, 4133-7.

- (122) Angenendt, P.; Glokler, J.; Sobek, J.; Lehrach, H.; Cahill, D. J. (2003) Next generation of protein microarray support materials: evaluation for protein and antibody microarray applications. *J Chromatogr A* **1009**, 97-104.
- (123) Ortiz de Montellano, P. R.; David, S. K.; Ator, M. A.; Tew, D. (1988) Mechanism-based inactivation of horseradish peroxidase by sodium azide. Formation of meso-azidoporphyrin IX. *Biochemistry* **27**, 5470-6.
- (124) Deininger, M. H.; Meyermann, R. (1998) Multiple epitope labeling by the exclusive use of alkaline phosphatase conjugates in immunohistochemistry. *Histochem Cell Biol* **110**, 425-30.
- (125) Fiering, S. N.; Roederer, M.; Nolan, G. P.; Micklem, D. R.; Parks, D. R.; Herzenberg, L. A. (1991) Improved FACS-Gal: flow cytometric analysis and sorting of viable eukaryotic cells expressing reporter gene constructs. *Cytometry* **12**, 291-301.
- (126) Burke, B. J.; Regnier, F. E. (2003) Stopped-flow enzyme assays on a chip using a microfabricated mixer. *Anal Chem* **75**, 1786-91.
- (127) Schulte, T.; Bohringer, S.; Schols, L.; Muller, T.; Fischer, C.; Riess, O.; Przuntek, H.; Berger, K.; Epplen, J. T.; Kruger, R. (2003) Modulation of disease risk according to a cathepsin D / apolipoprotein E genotype in Parkinson's disease. *J Neural Transm* **110**, 749-55.
- (128) Castilla, E. A.; Prayson, R. A.; Abramovich, C. M.; Cohen, M. L. (2003) Immunohistochemical expression of cathepsin D in meningiomas. *American Journal of Clinical Pathology* **119**, 123-8.
- (129) Kanber, Y.; Demirbag, N. R.; Sam, A. D.; Aydin, N. (2002) Cathepsin D expression in colorectal adenocarcinomas and adenomas. *International Journal of Biological Markers* **17**, 165-8.
- (130) Rochefort, H.; Garcia, M.; Glondu, M.; Laurent, V.; Liaudet, E.; Rey, J. M.; Roger, P. (2000) Cathepsin D in breast cancer: mechanisms and clinical applications, a 1999 overview. *Clinica Chimica Acta* **291**, 157-70.
- (131) Tapparelli, C.; Metternich, R.; Ehrhardt, C.; Zurini, M.; Claeson, G.; Scully, M. F.; Stone, S. R. (1993) In vitro and in vivo characterization of a neutral boron-containing thrombin inhibitor. *J Biol Chem* **268**, 4734-41.
- (132) Claeson, G.; Philipp, M.; Agner, E.; Scully, M. F.; Metternich, R.; Kakkar, V. V.; DeSoyza, T.; Niu, L. H. (1993) Benzyloxycarbonyl-D-Phe-Pro-methoxypropylboroglycine: a novel inhibitor of thrombin with high selectivity containing a neutral side chain at the P1 position. *Biochem J* **290**, 309-12.

- (133) Heim, R.; Prasher, D. C.; Tsien, R. Y. (1994) Wavelength mutations and posttranslational autoxidation of green fluorescent protein. *Proc Natl Acad Sci U S A* **91**, 12501-12504.
- (134) Russo-Marie, F.; Roederer, M.; Sager, B.; Herzenberg, L. A.; Kaiser, D. (1993) Beta-galactosidase activity in single differentiating bacterial cells. *Proc Natl Acad Sci U S A* **90**, 8194-8.
- (135) Schröder, D. H., 2003.
- (136) Axon Instruments, I. *GenePix Pro 4.1 User's Guide and Tutorial*, 2002.
- (137) Riechmann, L.; Holliger, P. (1997) The C-terminal domain of TolA is the coreceptor for filamentous phage infection of E. coli. *Cell* **90**, 351-60.
- (138) Click, E. M.; Webster, R. E. (1997) Filamentous phage infection: required interactions with the TolA protein. *J Bacteriol* **179**, 6464-71.
- (139) Zinder, N. D.; Model, P.; Boeke, J. D. In *US Patent 4,595,658*, 1986.
- (140) Molecular_Probes (2003) Produkt Information: Amplex Red Hydrogen Peroxide/Peroxidase Assay Kit (A-22188) [WWW Document]. <http://www.probes.com/media/pis/mp22188.pdf>.
- (141) Jones, L. J.; Upson, R. H.; Haugland, R. P.; Panchuk-Voloshina, N.; Zhou, M. (1997) Quenched BODIPY dye-labeled casein substrates for the assay of protease activity by direct fluorescence measurement. *Anal Biochem* **251**, 144-52.
- (142) Schloss, P. D.; Handelsman, J. (2003) Biotechnological prospects from metagenomics. *Curr Opin Biotechnol* **14**, 303-10.
- (143) Mungall, B. A.; Xu, X.; Klimov, A. (2003) Assaying susceptibility of avian and other influenza A viruses to zanamivir: comparison of fluorescent and chemiluminescent neuraminidase assays. *Avian Dis* **47**, 1141-4.
- (144) Huclova, J.; Satinsky, D.; Sklenarova, H.; Karliceck, R. (2003) Determination of salbutamol using on-line solid-phase extraction and sequential injection analysis. Comparison of chemiluminescence and fluorescence detection. *Anal Bioanal Chem* **376**, 448-54. Epub 2003 Feb 22.
- (145) Wu, J. H.; Diamond, S. L. (1995) A fluorescence quench and dequench assay of fibrinogen polymerization, fibrinogenolysis, or fibrinolysis. *Anal Biochem* **224**, 83-91.
- (146) Chalfie, M. (1995) Green fluorescent protein. *Photochem Photobiol* **62**, 651-6.