

8 Literaturverzeichnis

- Aggarwal BB, Eessalu TE (1987). Induction of receptors for tumor necrosis factor-alpha by interferons is not a major mechanism for their synergistic cytotoxic response. 262(21): 10000-10007
- Aggarwal BB, Eessalu TE, Hass PE (1985). Characterization of receptors for human tumour necrosis factor and their regulation by gamma-interferon. 318(6047): 665-667
- Ahdieh M, Vandenbos T, Youakim A (2001). Lung epithelial barrier function and wound healing are decreased by IL-4 and IL-13 and enhanced by IFN-gamma. Am J Physiol Cell Physiol 281(6): C2029-C2038
- Amasheh S, Meiri N, Gitter AH, Schoneberg T, Mankertz J, Schulzke JD, Fromm M (2002). Claudin-2 expression induces cation-selective channels in tight junctions of epithelial cells. J Cell Sci 115(Pt 24): 4969-4976
- Anderson JM, Van Itallie CM (1995). Tight junctions and the molecular basis for regulation of paracellular permeability. Am J Physiol 269(4 Pt 1): G467-G475
- Anderson JM, Van Itallie CM (1999). Tight junctions: closing in on the seal. Curr Biol 9(24): R922-R924
- Bamforth SD, Kniesel U, Wolburg H, Engelhardt B, Risau W (1999). A dominant mutant of occludin disrupts tight junction structure and function. J Cell Sci 112 (Pt 12): 1879-1888
- Barre-Sinoussi F, Chermann JC, Rey F, Nugeyre MT, Chamaret S, Gruest J, Dauguet C, Axler-Blin C, Vezinet-Brun F, Rouzioux C, Rozenbaum W, Montagnier L (1983). Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). 220(4599): 868-871
- Bartels M (2001). Untersuchungen zum Vorkommen von Antikörpern gegen das bovine Immunschwächevirus bei Rindern in Oberbayern. Dissertation, Tierärztliche Fakultät, München
- Batman PA, Fleming SC, Sedgwick PM, MacDonald TT, Griffin GE (1994). HIV infection of human fetal intestinal explant cultures induces epithelial cell proliferation. AIDS 8(2): 161-167
- Batman PA, Miller AR, Forster SM, Harris JR, Pinching AJ, Griffin GE (1989). Jejunal enteropathy associated with human immunodeficiency virus infection: quantitative histology. J Clin Pathol 42(3): 275-281
- Bement WM, Forscher P, Mooseker MS (1993). A novel cytoskeletal structure involved in purse string wound closure and cell polarity maintenance. J Cell Biol 121(3): 565-578
- Bendfeldt K (2000). Tumornekrosefaktor (TNF)- α -induzierte Barrierestörung an menschlichen Colonzellen (HT-29/B6): Lokalisierung mittels Conductance Scanning. Dissertation, Freie Universität Berlin

- Bandinelli M, Pistello M, Lombardi S, Poli A, Garzelli C, Matteucci D, Ceccherini-Nelli L, Malvaldi G, Tozzini F (1995). Feline immunodeficiency virus: an interesting model for AIDS studies and an important cat pathogen. 8(1): 87-112
- Booth BW, Adler KB, Bonner JC, Tournier F, Martin LD (2001). Interleukin-13 induces proliferation of human airway epithelial cells in vitro via a mechanism mediated by transforming growth factor-alpha. Am J Respir Cell Mol Biol 25(6): 739-743
- Brazille P, Dereuddre-Bosquet N, Leport C, Clayette P, Boyer O, Vilde JL, Dormont D, Benveniste O (2003). Decreases in plasma TNF -alpha level and IFN-gamma mRNA level in peripheral blood mononuclear cells (PBMC) and an increase in IL -2 mRNA level in PBMC are associated with effective highly active antiretroviral therapy in HIV -infected patients. Clin Exp Immunol 131(2): 304-311
- Bürk RR (1973). A factor from a transformed cell line that affects cell migration. Proc Nat Acad Sci USA 70(2): 369-372
- Callanan JJ, Hosie MJ, Jarrett O (1991). Transmission of feline immunodeficiency virus from mother to kitten. Vet Rec 128: 332-333
- Capobianchi MR, Fais S, Di Paolo MC, Agostini D, Paoluzi P, Pallone F, Dianzani F (1992). Absence of circulating interferon in patients with inflammatory bowel disease. Suggestion against an autoimmune etiology. Clin Exp Immunol 90(1): 85-87
- Cario E, Becker A, Sturm A, Goebell H, Dignass AU (1999). Peripheral blood mononuclear cells promote intestinal epithelial restitution in vitro through an interleukin-2/interferon-gamma-dependent pathway. Scand J Gastroenterol 34(11): 1132-1138
- Ceponis PJ, Botelho F, Richards CD, McKay DM (2000). Interleukins 4 and 13 increase intestinal epithelial permeability by a phosphatidylinositol 3-kinase pathway. Lack of evidence for STAT 6 involvement. J Biol Chem 275(37): 29132-29137
- Chen MC, Lee AT, Soll AH (1991). Mitogenic response of canine fundic epithelial cells in short-term culture to transforming growth factor alpha and insulinlike growth factor I. J Clin Invest 87(5): 1716-1723
- Chiassone DC, Simon PL, Smith PL (1990). Interleukin-1: effects on rabbit ileal mucosal ion transport in vitro. Eur J Pharmacol 180(2-3): 217-228
- Church DL, Sutherland LR, Gill MJ, Visser ND, Kelly JK (1992). Absence of an association between enteric parasites in the manifestations and pathogenesis of HIV enteropathy in gay men. The GI/HIV Study Group. Scand J Infect Dis 24(5): 567-575
- Ciacci C, Lind SE, Podolsky DK (1993). Transforming growth factor beta regulation of migration in wounded rat intestinal epithelial monolayers. Gastroenterol 105(1): 93-101

Clark RA (2003). Epithelial-mesenchymal networks in wounds: a hierarchical view. 120(6): IX -XI

Clavel F, Guetard D, Brun-Vezinet F, Chamaret S, Rey MA, Santos-Ferreira MO, Laurent AG, Dauguet C, Katlama C, Rouzioux C, et al. (1986). Isolation of a new human retrovirus from West African patients with AIDS. Science 233(4761): 343-346

Colebunders R, Lusakumuni K, Nelson AM, Gigase P, Lebughe I, van Marck E, Kapita B, Francis H, Salaun JJ, Quinn TC, et al. (1988). Persistent diarrhoea in Zairian AIDS patients: an endoscopic and histological study. Gut 29(12): 1687-1691

Corredor J, Yan F, Shen CC, Tong W, John SK, Wilson G, Whitehead R, Polk DB (2003). Tumor necrosis factor regulates intestinal epithelial cell migration by receptor-dependent mechanisms. Am J Physiol Cell Physiol 284(4): C953-C961

Cosentino G, Soprana E, Thienes CP, Siccardi AG, Viale G, Vercelli D (1995). IL-13 down-regulates CD14 expression and TNF-alpha secretion in normal human monocytes. J Immunol 155(6): 3145-3151

Courchamp F, Pontier D (1994). Feline immunodeficiency virus: an epidemiological review. C R Acad Sci III 317(12): 1123-1134

Cummins AG, LaBrooy JT, Stanley DP, Rowland R, Shearman DJ (1990). Quantitative histological study of enteropathy associated with HIV infection. Gut 31(3): 317-321

Dalgleish AG, Beverley PC, Clapham PR, Crawford DH, Greaves MF, Weiss RA (1984). The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. Nature 312(5996): 763-767

Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhardt M, Di Marzio P, Marmon S, Sutton RE, Hill CM, Davis CB, Peiper SC, Schall TJ, Littman DR, Landau NR (1996). Identification of a major co-receptor for primary isolates of HIV-1. Nature 381(6584): 661-666

Desrosiers RC (1990). The simian immunodeficiency viruses. Annu Rev Immunol 8: 557-578

Dieth V, Lutz H, Hauser B, Ossent P (1989). Pathologische Befunde bei mit Lentiviren infizierten Katzen. Schweiz Arch Tierheilk 131: 19-25

Dignass A, Stein J (2000). Chronisch entzündliche Darmerkrankungen. Gastroenterol Update 2000

Dignass AU (2001). Mechanisms and modulation of intestinal epithelial repair. Inflamm Bowel Dis 7(1): 68-77

Dignass AU, Lynch-Devaney K, Podolsky DK (1994). Hepatocyte growth factor/scatter factor modulates intestinal epithelial cell proliferation and migration. Biochem Biophys Res Commun 202(2): 701-709.

- Dignass AU, Podolsky DK (1993). Cytokine modulation of intestinal epithelial cell restitution: central role of transforming growth factor beta. *Gastroenterol* 105(5): 1323-1332
- Dignass AU, Stow JL, Babyatsky MW (1996). Acute epithelial injury in the rat small intestine *in vivo* is associated with expanded expression of transforming growth factor alpha and beta. *Gut* 38(5): 687-693
- Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, Parmentier M, Collman RG, Doms RW (1996). A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 85(7): 1149-1158
- Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, Cayanan C, Madden PJ, Koup RA, Moore JP, Paxton WA (1996). HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* 381(6584): 667-673
- Fanning AS, Jameson BJ, Jesaitis LA, Anderson JM (1998). The tight junction protein ZO-1 establishes a link between the transmembrane protein occludin and the actin cytoskeleton. *J Biol Chem* 273(45): 29745-29753.
- Feil W, Klimesch S, Karner P, Wenzl E, Starlinger M, Lancy ER, Schiessel R (1989). Importance of an alkaline microenvironment for rapid restitution of the rabbit duodenal mucosa *in vitro*. *Gastroenterol* 97(1): 112-122
- Feinberg MB (1996). Changing the natural history of HIV disease. *Lancet* 348(9022): 239-246
- Feng Y, Broder CC, Kennedy PE, Berger EA (1996). HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272(5263): 872-877
- Florian P (2002). Restitution von Einzelzell-Läsionen im Kolonepithel. Dissertation, Freie Universität Berlin
- Florian P, Gitter AH, Schulzke JD, Fromm M (2001). Recovery of colon epithelial barrier after single-cell lesion. *Pflügers Arch Suppl* 441(6): R 208
- Florian P, Schöneberg T, Schulzke JD, Fromm M, Gitter AH (2002). Single-cell epithelial defects close rapidly by an actinomyosin purse string mechanism with functional tight junctions. *J Physiol* 545(2): 485-499
- Furuse M, Fujita K, Hiiragi T, Fujimoto K, Tsukita S (1998). Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol* 141: 1539-1550
- Furuse M, Furuse K, Sasaki H, Tsukita S (2001). Conversion of Zonulae occludentes from tight to leaky strand type by introducing claudin-2 into Madin Darby Canine Kidney I cells. *J Cell Biol* 153(2): 263-272

- Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A, Sugitani Y, Noda T, Kubo A, Tsukita S (2002). Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J Cell Biol* 156(6): 1099-1111
- Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S (1993). Occludin: a novel integral membrane protein at tight junctions. *J Cell Biol* 123: 1777-1788
- Fuss IJ, Neurath M, Boirivant M, Klein JS, de la Motte C, Strong SA, Fiocchi C, Strober W (1996). Disparate CD4+ lamina propria (LP) lymphokine secretion profiles in inflammatory bowel disease. Crohn's disease LP cells manifest increased secretion of IFN-gamma, whereas ulcerative colitis LP cells manifest increased secretion of IL-5. *J Immunol* 157(3): 1261-1270
- Gallo RC, Sarin PS, Gelmann EP, Robert-Guroff M, Richardson E, Kalyanaraman VS, Mann D, Sidhu GD, Stahl RE, Zolla-Pazner S, Leibowitch J, Popovic M (1983). Isolation of human T-cell leukemia virus in acquired immune deficiency syndrome (AIDS). *Science* 220(4599): 865-867
- Gendelman HE, Baca LM, Turpin J, Kalter DC, Hansen B, Orenstein JM, Dieffenbach CW, Friedman RM, Meltzer MS (1990). Regulation of HIV replication in infected monocytes by IFN-alpha. Mechanisms for viral restriction. *J Immunol* 145(8): 2669-2676
- Gitter AH, Bendfeldt K, Schmitz H, Schulzke JD, Bentzel CJ, Fromm M (2000a). Epithelial Barrier Defects in HT-29/B6 Colonic Cell Monolayers Induced by Tumor Necrosis Factor- α . *Ann N Y Acad Sci* 915: 193-203
- Gitter AH, Bendfeldt K, Schulzke JD, Fromm M (2000b). Leaks in the epithelial barrier caused by spontaneous and TNF α -induced single-cell apoptosis. *FASEB J* 14: 1749-1753
- Gitter AH, Bertog M, Schulzke JD, Fromm M (1997). Measurement of paracellular epithelial conductivity by conductance scanning. *Pflügers Arch* 434: 830-840
- Gitter AH, Wullstein F, Fromm M, Schulzke JD (2001). Epithelial barrier defects in ulcerative colitis: characterization and quantification by electrophysiological imaging. *Gastroenterol* 121: 1320-1328
- Göke M, Podolsky DK (1996). Regulation of the mucosal epithelial barrier. *Baillieres Clin Gastroenterol* 10(3): 393-405.
- Gölz J, Mayr C, Heise W (1999). HIV und AIDS. Urban und Fischer Verlag, München
- Gonzalez-Mariscal L, Betanzos A, Nava P, Jaramillo BE (2003). Tight junction proteins. *Prog Biophys Mol Biol* 81(1): 1-44
- Greenson JK, Belitsos PC, Yardley JH, Bartlett JG (1991). AIDS enteropathy: occult enteric infections and duodenal mucosal alterations in chronic diarrhea. *Ann Intern Med* 114: 366-372

- Grindem CB, Corbett WT, Ammerman BE, Tomkins MT (1989). Seroepidemiologic survey of feline immunodeficiency virus infection in cats of Wake Country, North Carolina. *J Am Vet Med Assoc* 194: 226-228
- Grunfeld C, Kotler DP, Shigenaga JK, Doerrler W, Tierney A, Wang J, Pierson RN, Jr., Feingold KR (1991). Circulating interferon-alpha levels and hypertriglyceridemia in the acquired immunodeficiency syndrome. *Am J Med* 90(2): 154-162
- Gumbiner B, Stevenson B, Grimaldi A (1988). The role of the cell adhesion molecule uvomorulin in the formation and maintenance of the epithelial junctional complex. *J Cell Biol* 107(4): 1575-1587
- Haas DW, Lavelle J, Nadler JP, Greenberg SB, Frame P, Mustafa N, St Clair M, McKinnis R, Dix L, Elkins M, Rooney J (2000). A randomized trial of interferon alpha therapy for HIV type 1 infection. *AIDS Res Hum Retroviruses* 16(3): 183-190
- Hall PA, Coates PJ, Ansari B, Hopwood D (1994). Regulation of cell number in the mammalian gastrointestinal tract: the importance of apoptosis. *J Cell Sci* 107 (Pt 12): 3569-3577
- Hartmann K, Hinze K (1991). Epidemiologie und Klinik der FIV -Infektion in Bayern. *Tierärztl Praxis* 19: 545-551
- Hatzakis A, Gargalianos P, Kiosses V, Lazanas M, Sypsa V, Anastassopoulou C, Vigklis V, Sambatakou H, Botsi C, Paraskevis D, Stalgis C (2001). Low-dose IFN-alpha monotherapy in treatment-naive individuals with HIV-1 infection: evidence of potent suppression of viral replication. *J Interferon Cytokine Res* 21(10): 861-869
- Heath JP (1996). Epithelial cell migration in the intestine. *Cell Biol Int* 20(2): 139-146
- Heldin CH, Westermark B (1999). Mechanism of action and in vivo role of platelet-derived growth factor. *Physiol Rev* 79(4): 1283-1316
- Heller F, Fuss IJ, Nieuwenhuis EE, Blumberg RS, Strober W (2002). Oxazolone colitis, a Th2 colitis model resembling ulcerative colitis, is mediated by IL-13-producing NK-T cells. *Immunity* 17(5): 629-638
- Henrikson CK, Argenzio RA, Liacos JA, Khosla J (1989). Morphologic and functional effects of bile salt on the porcine colon during injury and repair. *Lab Invest* 60(1): 72-87
- Heyman M, Darmon N, Dupont C, Dugas B, Hirribaren A, Blaton MA, Desjeux JF (1994). Mononuclear cells from infants allergic to cow's milk secrete tumor necrosis factor- α , altering intestinal function. *Gastroenterol* 106: 1514-1523
- Hopper CD, Harbour D, Gruffydd-Jones TJ (1992). Evidence of vertical transmission of FIV in experimentally and naturally infected cats. European Commission: Concerted action on feline AIDS workshop, Lucca, Italy 18.-21.06.1992

- Hopper CD, Sparkes AH, Gruffydd-Jones TJ, Crispin SM, Muir P, Harbour D, Stokes CR (1989). Clinical and laboratory findings in cats infected with feline immunodeficiency virus. *Vet Rec* 125: 341-346
- Imanishi J, Kamiyama K, Iguchi I, Kita M, Sotozono C, Kinoshita S (2000). Growth factors: importance in wound healing and maintenance of transparency of the cornea. *Prog Retin Eye Res* 19(1): 113-129
- Jurriaans S, Van Gemen B, Weverling GJ, Van Strijp D, Nara P, Coutinho R, Koot M, Schuitemaker H, Goudsmit J (1994). The natural history of HIV-1 infection: virus load and virus phenotype independent determinants of clinical course? *Virology* 204(1): 223-233
- Kamiyama K, Iguchi I, Wang X, Imanishi J (1998). Effects of PDGF on the migration of rabbit corneal fibroblasts and epithelial cells. *Cornea* 17(3): 315-325
- Kapembwa MS, Fleming SC, Sewankambo N, Serwadda D, Lucas S, Moody A, Griffin GE (1991). Altered small-intestinal permeability associated with diarrhoea in human-immunodeficiency-virus-infected Caucasian and African subjects. *Clin Sci (Lond)* 81(3): 327-334
- Kato K, Chen MC, Nguyen M, Lehmann FS, Podolsky DK, Soll AH (1999). Effects of growth factors and trefoil peptides on migration and replication in primary oxyntic cultures. *Am J Physiol* 276(5 Pt 1): G1105-G1116
- Klatzmann D, Champagne E, Chamaret S, Gruet J, Guetard D, Hercend T, Gluckman JC, Montagnier L (1984). T-lymphocyte T4 molecule behaves as the receptor for human retrovirus LAV. *Nature* 312(5996): 767-768
- Kotler DP, Gaetz HP, Lange M, Klein EB, Holt PR (1984). Enteropathy associated with the acquired immunodeficiency syndrome. *Ann Intern Med* 101(4): 421-428
- Kotler DP, Orenstein JM (1994). Prevalence of intestinal microsporidiosis in HIV-infected individuals referred for gastroenterological evaluation. *Am J Gastroenterol* 89(11): 1998-2002
- Kotler DP, Reka S, Clayton F (1993). Intestinal mucosal inflammation associated with human immunodeficiency virus infection. *Dig Dis Sci* 38(6): 1119-1127
- Kraft W, Dürr UM (1996). Katzenkrankheiten, Klinik und Therapie.
- Kreusel KM, Fromm M, Schulzke JD, Hegel U (1991). Cl⁻ secretion in epithelial monolayers of mucus-forming human colon cell (HT-29/B6). *Am J Physiol* 261: C574-C582
- Kuwabara T, Perkins DG, Cogan DG (1976). Sliding of the epithelium in experimental corneal wounds. *Invest Ophthalmol* 15(1): 4-14
- Lacy ER (1988). Epithelial restitution in the gastrointestinal tract. *J Clin Gastroenterol* 10 Suppl 1: S72-S77.

- Lesur O, Brisebois M, Thibodeau A, Chagnon F, Lane D, Fullop T (2003). Role of Interferon-gamma and Interleukin-2 in rat lung epithelial cell migration and apoptosis after oxidant injury. *Am J Physiol Lung Cell Mol Physiol* 286(1): L4-L14
- Lotz MM, Rabinovitz I, Mercurio AM (2000). Intestinal restitution: progression of actin cytoskeleton rearrangements and integrin function in a model of epithelial wound healing. *Am J Pathol* 156(3): 985-996
- Lutz H, Egberink H, Arnold P, Winkler G, Wolfensberger C, Jarrett O, Parodi AL, Pedersen NC, Horzinek MC (1988). Felines T-lymphotopes Lentivirus (FTLV): Experimentelle Infektion und Vorkommen in einigen Ländern Europas. *Kleintierpraxis* 33: 445-492
- MacDonald TT, Hutchings P, Choy MY, Murch S, Cooke A (1990). Tumour necrosis factor-alpha and interferon-gamma production measured at the single cell level in normal and inflamed human intestine. *Clin Exp Immunol* 81(2): 301-305
- Madara JL, Stafford J (1989). Interferon-gamma directly affects barrier function of cultured intestinal epithelial monolayers. *J Clin Invest* 83: 724-727
- Mandracchia VJ, Sanders SM, Frerichs JA (2001). The use of becaplermin (rhPDGF-BB) gel for chronic nonhealing ulcers. A retrospective analysis. *Clin Podiatr Med Surg* 18(1): 189-209
- Martin P (1997). Wound healing--aiming for perfect skin regeneration. *Science* 276(5309): 75-81
- Martin-Padura I, Lostaglio S, Schneemann M, Williams L, Romano M, Fruscella P, Panzeri C, Stoppacciaro A, Ruco L, Villa A, Simmons D, Dejana E (1998). Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J Cell Biol* 142(1): 117-127
- Matsuda M, Ubels JL, Edelhauser HF (1985). A larger corneal epithelial wound closes at a faster rate. *Invest Ophthalmol Vis Sci* 26(6): 897-900
- McCarthy KM, Skare IB, Stankewich MC, Furuse M, Tsukita S, Rogers RA, Lynch RD, Schneeberger EE (1996). Occludin is a functional component of the tight junction. *J Cell Sci* 109 (Pt 9): 2287-2298
- McClain SA, Simon M, Jones E, Nandi A, Gailit JO, Tonnesen MG, Newman D, Clark RA (1996). Mesenchymal cell activation is the rate-limiting step of granulation tissue induction. *149(4): 1257-70*
- McCormack SA, Viar MJ, Johnson LR (1992). Migration of IEC-6 cells: a model for mucosal healing. *Am J Physiol* 263(3 Pt 1): G426-G435.
- McGowan I, Radford-Smith G, Jewell DP (1994). Cytokine gene expression in HIV-infected intestinal mucosa. *AIDS* 8(11): 1569-1575

- Medina R, Rahner C, Mitic LL, Anderson JM, Van Itallie CM (2000). Occludin localization at the tight junction requires the second extracellular loop. *J Membr Biol* 178(3): 235-247
- Med-World, zur A, Medizin Dv, im uG, Internet (2003). HIV - Infektion und AIDS. <http://www.medicine-worldwide.de/krankheiten/infektionskrankheiten/aids.html>
- Mellors JW, Kingsley LA, Rinaldo CR, Jr., Todd JA, Hoo BS, Kokka RP, Gupta P (1995). Quantitation of HIV-1 RNA in plasma predicts outcome after seroconversion. *Ann Intern Med* 122(8): 573-579
- Mellors JW, Rinaldo CR, Jr., Gupta P, White RM, Todd JA, Kingsley LA (1996). Prognosis in HIV-1 infection predicted by the quantity of virus in plasma. *Science* 272(5265): 1167-1170
- Meyer zum Büschenfelde D (2002). TFF3 dependent modulation of the E-Catherin/β-Catenin adhesion complex. Jahrestagung der Deutschen Gesellschaft für Laboratoriumsmedizin (DGML) und der Deutschen Gesellschaft für Klinische Chemie (DGKC), Düsseldorf
- Mildvan D, Bassiakos Y, Zucker ML, Hyslop N, Jr., Krown SE, Sacks HS, Zachary J, Paredes J, Fessel WJ, Rhame F, Kramer F, Fischl MA, Poiesz B, Wood K, Ruprecht RM, Kim J, Grossberg SE, Kasdan P, Berge P, Marshak A, Pettinelli C (1996). Synergy, activity and tolerability of zidovudine and interferon-alpha in patients with symptomatic HIV-1 infection: AIDS Clinical Trial Group 068. *Antivir Ther* 1(2): 77-88
- Miller CA, Debas HT (1995). Epidermal growth factor stimulates the restitution of rat gastric mucosa in vitro. *Exp Physiol* 80(6): 1009-1118
- Minty A, Chalon P, Derocq JM, Dumont X, Guillemot JC, Kaghad M, Labit C, Leplatois P, Liauzun P, Miloux B, et al. (1993). Interleukin-13 is a new human lymphokine regulating inflammatory and immune responses. *Nature* 362(6417): 248-250
- Mouser JF, Hyams JS (1999). Infliximab: a novel chimeric monoclonal antibody for the treatment of Crohn's disease. *Clin Ther* 21(6): 932-42; discussion 931
- Murch SH, Lamkin VA, Savage MO, Walker-Smith JA, MacDonald TT (1991). Serum concentrations of tumour necrosis factor alpha in childhood chronic inflammatory bowel disease. *Gut* 32(8): 913-917
- Nagai MK, Embil JM (2002). Bepacupermin: recombinant platelet derived growth factor, a new treatment for healing diabetic foot ulcers. *Expert Opin Biol Ther* 2(2): 211-218
- Nathan CF (1987). Secretory products of macrophages. *J Clin Invest* 79(2): 319-326
- Neurath MF, Schurmann G (2000). Immunopathogenesis of inflammatory bowel diseases. *Chirurg* 71(1): 30-40.

- Nusrat A, Delp C, Madara JL (1992). Intestinal epithelial restitution. Characterization of a cell culture model and mapping of cytoskeletal elements in migrating cells. *J Clin Invest* 89(5): 1501-1511
- Orenstein JM, Kotler DP (1995). Diarrheogenic bacterial enteritis in acquired immune deficiency syndrome: a light and electron microscopy study of 52 cases. *Hum Pathol* 26(5): 481-492
- Osborn L, Hession C, Tizard R, Vassallo C, Luhowskyj S, Chi-Rosso G, Lobb R (1989). Direct expression cloning of vascular cell adhesion molecule 1, a cytokine-induced endothelial protein that binds to lymphocytes. *Cell* 59(6): 1203-1211
- Ozaki H, Ishii K, Horiuchi H, Arai H, Kawamoto T, Okawa K, Iwamatsu A, Kita T (1999). Cutting edge: combined treatment of TNF-alpha and IFN-gamma causes redistribution of junctional adhesion molecule in human endothelial cells. *J Immunol* 163(2): 553-557
- Paimela H, Goddard PJ, Carter K, Khakee R, McNeil PL, Ito S, Silen W (1993). Restitution of frog gastric mucosa in vitro: effect of basic fibroblast growth factor. *Gastroenterol* 104(5): 1337-1345
- Pantaleo G, Graziosi C, Demarest JF, Butini L, Montroni M, Fox CH, Orenstein JM, Kotler DP, Fauci AS (1993). HIV infection is active and progressive in lymphoid tissue during the clinically latent stage of disease. *Nature* 362(6418): 355-358
- Parronchi P, Romagnani P, Annunziato F, Sampognaro S, Becchio A, Giannarini L, Maggi E, Pupilli C, Tonelli F, Romagnani S (1997). Type 1 T-helper cell predominance and interleukin-12 expression in the gut of patients with Crohn's disease. *Am J Pathol* 150(3): 823-832
- Pedersen G, Saermark T, Bendtzen K, Brynskov J (2000). Cultures of human colonic epithelial cells isolated from endoscopical biopsies from patients with inflammatory bowel disease. Effect of IFNgamma, TNFalpha and IL-1beta on viability, butyrate oxidation and IL-8 secretion. *Autoimmunity* 32(4): 255-263
- Pedersen NC, Ho EW, Brown ML, Yamamoto JK (1987). Isolation of a T-lymphotropic virus from domestic cats with an immunodeficiency-like syndrome. *Science* 235: 790-793
- Pedersen NC, Yamamoto JK, Ishida T, Hansen H (1989). Feline immunodeficiency virus infection. *Vet Immunol Immunopathol* 21: 111-129
- Pfister RR (1975). The healing of corneal epithelial abrasions in the rabbit: a scanning electron microscope study. *Invest Ophthalmol* 14(9): 648-661
- Piasecki E, Knysz B, Gasiorowski J, Gladysz A (1999). Decrease of enhanced interferon alpha levels in sera of HIV-infected and AIDS patients receiving combined antiretroviral therapy. *Arch Immunol Ther Exp (Warsz)* 47(1): 37-44

- Piazuelo E, Jimenez P, Lanas A, Garcia A, Esteva F, Sainz R (2000). Platelet-derived growth factor and epidermal growth factor play a major role in human colonic fibroblast repair activities. *Eur Surg Res* 32(3): 191-196
- Pitha J (1971). The fine structure of regenerating epithelium in the small intestine. *Virchows Arch B Cell Pathol* 7: 314-341
- Plevy SE, Landers CJ, Prehn J, Carramanzana NM, Deem RL, Shealy D, Targan SR (1997). A role for TNF-alpha and mucosal T helper-1 cytokines in the pathogenesis of Crohn's disease. *J Immunol* 159(12): 6276-6282
- Podolsky DK (2002). Inflammatory bowel disease. *N Engl J Med* 347(6): 417-429
- Poli G, Orenstein JM, Kinter A, Folks TM, Fauci AS (1989). Interferon-alpha but not AZT suppresses HIV expression in chronically infected cell lines. *Science* 244(4904): 575-577
- Radema SA, van Deventer SJ, Cerami A (1991). Interleukin 1 beta is expressed predominantly by enterocytes in experimental colitis. *Gastroenterol* 100(5 Pt 1): 1180-116
- Rajasekaran AK, Hojo M, Huima T, Rodriguez-Boulan E (1996). Catenins and zonula occludens-1 form a complex during early stages in the assembly of tight junctions. *J Cell Biol* 132(3): 451-463
- Rajasekaran AK, Rajasekaran SA (2003). Role of Na-K-ATPase in the assembly of tight junctions. *Am J Physiol Renal Physiol* 285(3): F388-F396
- Ramachandran A, Madesh M, Balasubramanian KA (2000). Apoptosis in the intestinal epithelium: its relevance in normal and pathophysiological conditions. *J Gastroenterol Hepatol* 15(2): 109-120
- Rance KS, Jordan HL, Kennedy-Stoskopf S, Tompkins MB (1994). Feline Immunodeficiency virus can be experimentally transmitted via milk during acute maternal infection. *J Virol* 68: 3380-3385
- Rene E, Marche C, Regnier B, Saimot AG, Vilde JL, Perrone C, Michon C, Wolf M, Chevalier T, Vallot T, et al. (1989). Intestinal infections in patients with acquired immunodeficiency syndrome. A prospective study in 132 patients. *Dig Dis Sci* 34(5): 773-780
- Riegler M, Feil W, Lindroth J, Wong M (1992). MIPSY: real-time morphometry to quantify the time course of rapid epithelial restitution. *Pathol Res Pract* 188(4-5): 443-8
- Riegler M, Sedivy R, Sogukoglu T, Cosentini E, Bischof G, Teleky B, Feil W, Schiessel R, Hamilton G, Wenzl E (1997). Effect of growth factors on epithelial restitution of human colonic mucosa in vitro. *Scand J Gastroenterol* 32(9): 925-932

- Rival Y, Del Maschio A, Rabiet MJ, Dejana E, Duperray A (1996). Inhibition of platelet endothelial cell adhesion molecule-1 synthesis and leukocyte transmigration in endothelial cells by the combined action of TNF-alpha and IFN-gamma. *J Immunol* 157(3): 1233-1241
- Ruggiero V, Tavernier J, Fiers W, Baglioni C (1986). Induction of the synthesis of tumor necrosis factor receptors by interferon-gamma. *J Immunol* 136(7): 2445-2450
- Saitou M, Furuse M, Sasaki H, Schulzke JD, Fromm M, Takano H, Noda T, Tsukita S (2000). Complex phenotype of mice lacking occludin, a component of tight junction strands. *11(12)*: 4131-4142
- Santos MF, McCormack SA, Guo Z, Okolicany J, Zheng Y, Johnson LR, Tigyi G (1997). Rho proteins play a critical role in cell migration during the early phase of mucosal restitution. *J Clin Invest* 100(1): 216-225.
- Sato Y, Rifkin DB (1988). Autocrine activities of basic fibroblast growth factor: regulation of endothelial cell movement, plasminogen activator synthesis, and DNA synthesis. *J Cell Biol* 107(3): 1199-1205
- Schmitz H, Fromm M, Bentzel CJ, Scholz P, Detjen K, Mankertz J, Bode H, Epple HJ, Riecken EO, Schulzke JD (1999). Tumor necrosis-factor-alpha (TNF- α) regulates the epithelial barrier in the human intestinal cell line HT-29/B6. *J Cell Sci* 112: 137-146
- Schmitz H, Fromm M, Bode H, Scholz P, Riecken EO, Schulzke JD (1996). Tumor necrosis factor-alpha induces Cl⁻ and K⁺ secretion in human distal colon driven by prostaglandin E₂. *Am J Physiol* 271: G669-G674
- Schmitz H, Rokos K, Florian P, Gitter AH, Fromm M, Scholz P, Ullrich R, Zeitz M, Pauli G, Schulzke JD (2002). Supernatants of HIV-infected immune cells affect the barrier function of human HT-29/B6 intestinal epithelial cells. *AIDS* 16(7): 983-991
- Schreiber S, Nikolaus S, Hampe J, Hamling J, Koop I, Groessner B, Lochs H, Raedler A (1999). Tumour necrosis factor alpha and interleukin 1beta in relapse of Crohn's disease. *Lancet* 353(9151): 459-461
- Schulzke JD, Fromm M, Hegel U (1986). Epithelial and subepithelial resistance of rat large intestine: segmental differences, effect of stripping, time course, and action of aldosterone. *Pflügers Arch* 407: 632-637
- Sheetz MP, Felsenfeld D, Galbraith CG, Choquet D (1999). Cell migration as a five-step cycle. *Biochem soc symp* 65: 233-243.
- Sher ME, D'Angelo AJ, Stein TA, Bailey B, Burns G, Wise L (1995). Cytokines in Crohn's colitis. *Am J Surg* 169(1): 133-136
- Simon DB, Lu Y, Choate KA, Velazquez H, Al-Sabban E, Praga M, Casari G, Bettinelli A, Colussi G, Rodriguez-Soriano J, McCredie D, Milford D, Sanjad S, Lifton RP (1999). Paracellin-1, a renal tight junction protein required for paracellular Mg²⁺ resorption. *Science* 285(5424): 103-106

- Snider TG, Hoyt PG, Coats KS, Graves KF, Cooper CR, Storts RWL, D.G., Jenny BF (2003). Natural bovine lentiviral type 1 infection in Holstein dairy cattle. I. Clinical, serological, and pathological observations. *Comp Immunol Microbiol Infect Dis* 26: 89-101
- Staehelin LA (1973). Further observations on the fine structure of freeze-cleaved tight junctions. *J Cell Sci* 13(3): 763-786
- Staehelin LA (1974). Structure and function of intercellular junctions. *Int Rev Cytol* 39: 191-283
- Stange EF (1999). Colitis ulcerosa - Morbus Crohn. UNI-MED Verlag AG 1. Auflage
- Stevenson BR, Anderspn JM, Bullivant S (1988). The epithelial tight junction: structure, function and preliminary biochemical characterization. *Mol Cell Biochem* 83: 129-145
- Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA (1986). Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biol* 103(3): 755-766
- Stockmann M, Schmitz H, Fromm M, Schmidt W, Pauli G, Scholz P, Riecken EO, Schulzke JD (2000). Mechanisms of epithelial barrier impairment in HIV infection. *Ann N Y Acad Sci* 915: 293-303
- Stout AJ, Gresser I, Thompson WD (1993). Inhibition of wound healing in mice by local interferon alpha/beta injection. *Int J Exp Pathol* 74(1): 79-85
- Sturm A, Sudermann T, Schulte KM, Goebell H, Dignass AU (1999). Modulation of intestinal epithelial wound healing in vitro and in vivo by lysophosphatidic acid. *Gastroenterol* 117(2): 368-377
- Stylianou E, Yndestad A, Sikkeland LI, Bjerkeli V, Damas JK, Haug T, Eiken HG, Aukrust P, Froland SS (2002). Effects of interferon-alpha on gene expression of chemokines and members of the tumour necrosis factor superfamily in HIV-infected patients. *Clin Exp Immunol* 130(2): 279-285
- Sumer N, Palabiyikoglu M (1995). Induction of remission by interferon-alpha in patients with chronic active ulcerative colitis. *Eur J Gastroenterol Hepatol* 7(7): 597-602
- Tepass U (2003). Claudin complexities at the apical junctional complex. *Nat Cell Biol* 5(7): 595-597
- Tsukita S, Furuse M (2000). The structure and function of claudins, cell adhesion molecules at tight junctions. *Ann N Y Acad Sci* 915: 129-135
- Ullrich R, Zeitz M, Heise W, L'Age M, Hoffken G, Riecken EO (1989). Small intestinal structure and function in patients infected with human immunodeficiency virus (HIV): evidence for HIV-induced enteropathy. *Ann Intern Med* 111(1): 15-21

- van Balkom BP, Schoon EJ, Stockbrugger RW, Wolters FL, van Hogezand RA, van Deventer SJ, Oldenburg B, van Dullemen HM, Russel MG (2002). Effects of anti-tumour necrosis factor-alpha therapy on the quality of life in Crohn's disease. *Aliment Pharmacol Ther* 16(6): 1101-1107
- Van Itallie C, Rahner C, Anderson J (2001). Regulated expression of claudin-4 decreases paracellular conductance through a selective decrease in sodium permeability. *J Clin Invest* 107(10): 1319-1327
- von Sydow M, Sonnerborg A, Gaines H, Strannegard O (1991). Interferon-alpha and tumor necrosis factor-alpha in serum of patients in various stages of HIV-1 infection. *AIDS Res Hum Retroviruses* 7: 375-380
- Voth R, Rossol S, Klein K, Hess G, Schutt KH, Schroder HC, Meyer zum Buschenfelde KH, Muller WE (1990). Differential gene expression of IFN-alpha and tumor necrosis factor-alpha in peripheral blood mononuclear cells from patients with AIDS related complex and AIDS. *J Immunol* 144(3): 970-975
- Vyakarnam A, Matear P, Meager A, Kelly G, Stanley B, Weller I, Beverley P (1991). Altered production of tumour necrosis factors alpha and beta and interferon gamma by HIV-infected individuals. *Clin Exp Immunol* 84(1): 109-115
- Wright NA, Irwin M (1982). The kinetics of villus cell populations in the mouse small intestine. I. Normal villi: the steady state requirement. *Cell Tissue Kinet* 15(6): 595-609
- Yanaka A, Muto H, Fukutomi H, Ito S, Silen W (1996). Role of transforming growth factor-beta in the restitution of injured guinea pig gastric mucosa in vitro. *Am J Physiol* 271(1 Pt 1): G75-G85
- Youakim A, Ahdieh M (1999). Interferon-gamma decreases barrier function in T84 cells by reducing ZO-1 levels and disrupting apical actin. *Am J Physiol* 276(5 Pt 1): G1279-G1288
- Zieske JD, Gipson IK (1986). Protein synthesis during corneal epithelial wound healing. *Invest Ophthalmol Vis Sci* 27(1): 1-7