

8. Literatur

- Agami, R. and R. Bernards (2000). "Distinct initiation and maintenance mechanisms cooperate to induce G1 cell cycle arrest in response to DNA damage." Cell **102**(1): 55-66.
- Anderson, C. W. (1993). "DNA damage and the DNA-activated protein kinase." Trends Biochem Sci **18**(11): 433-7.
- Anderson, C. W. and S. P. Lees-Miller (1992). "The nuclear serine/threonine protein kinase DNA-PK." Crit Rev Eukaryot Gene Expr **2**(4): 283-314.
- Arata, Y., M. Fujita, et al. (2000). "Cdk2-dependent and -independent pathways in E2F-mediated S phase induction." J Biol Chem **275**(9): 6337-45.
- Arrington, E. D., M. C. Caldwell, et al. (2000). "Enhanced sensitivity and long-term G2 arrest in hydrogen peroxide-treated Ku80-null cells are unrelated to DNA repair defects." Free Radic Biol Med **29**(11): 1166-76.
- Asano, M., J. R. Nevins, et al. (1996). "Ectopic E2F expression induces S phase and apoptosis in Drosophila imaginal discs." Genes Dev **10**(11): 1422-32.
- Blattner, C., A. Sparks, et al. (1999). "Transcription factor E2F-1 is upregulated in response to DNA damage in a manner analogous to that of p53." Mol Cell Biol **19**(5): 3704-13.
- Blattner, C., E. Tobiasch, et al. (1999). "DNA damage induced p53 stabilization: no indication for an involvement of p53 phosphorylation." Oncogene **18**(9): 1723-32.
- Blier, P. R., A. J. Griffith, et al. (1993). "Binding of Ku protein to DNA. Measurement of affinity for ends and demonstration of binding to nicks." J Biol Chem **268**(10): 7594-601.
- Cao, Q. P., S. Pitt, et al. (1994). "DNA-dependent ATPase from HeLa cells is related to human Ku autoantigen." Biochemistry **33**(28): 8548-57.
- Carter, T., I. Vancurova, et al. (1990). "A DNA-activated protein kinase from HeLa cell nuclei." Mol Cell Biol **10**(12): 6460-71.
- Cho, R. J., M. Huang, et al. (2001). "Transcriptional regulation and function during the human cell cycle." Nat Genet **27**(1): 48-54.

- Cloud, J. E., C. Rogers, et al. (2002). "Mutant mouse models reveal the relative roles of E2F1 and E2F3 in vivo." Mol Cell Biol **22**(8): 2663-72.
- Cress, W. D., D. G. Johnson, et al. (1993). "A genetic analysis of the E2F1 gene distinguishes regulation by Rb, p107, and adenovirus E4." Mol Cell Biol **13**(10): 6314-25.
- Dahm-Daphi, J., E. Dikomey, et al. (1998). "DNA-repair, cell killing and normal tissue damage." Strahlenther Onkol **174 Suppl 3**: 8-11.
- Darroudi, F. and A. T. Natarajan (1987). "Cytological characterization of Chinese hamster ovary X-ray-sensitive mutant cells xrs 5 and xrs 6. I. Induction of chromosomal aberrations by X-irradiation and its modulation with 3-aminobenzamide and caffeine." Mutat Res **177**(1): 133-48.
- de Vries, E., W. van Driel, et al. (1989). "HeLa nuclear protein recognizing DNA termini and translocating on DNA forming a regular DNA-multimeric protein complex." J Mol Biol **208**(1): 65-78.
- DeGregori, J., G. Leone, et al. (1997). "Distinct roles for E2F proteins in cell growth control and apoptosis." Proc Natl Acad Sci U S A **94**(14): 7245-50.
- DeSalle, L. M. and M. Pagano (2001). "Regulation of the G1 to S transition by the ubiquitin pathway." FEBS Lett **490**(3): 179-89.
- Dikomey, E., J. Dahm-Daphi, et al. (1998). "Correlation between cellular radiosensitivity and non-repaired double-strand breaks studied in nine mammalian cell lines." Int J Radiat Biol **73**(3): 269-78.
- Dyson, N. (1998). "The regulation of E2F by pRB-family proteins." Genes Dev **12**(15): 2245-62.
- Elledge, S. J. (1996). "Cell cycle checkpoints: preventing an identity crisis." Science **274**(5293): 1664-72.
- Falzon, M., J. W. Fewell, et al. (1993). "EBP-80, a transcription factor closely resembling the human autoantigen Ku, recognizes single- to double-strand transitions in DNA." J Biol Chem **268**(14): 10546-52.
- Fan, J. and J. R. Bertino (1997). "Functional roles of E2F in cell cycle regulation." Oncogene **14**(10): 1191-200.
- Field, S. J., F. Y. Tsai, et al. (1996). "E2F-1 functions in mice to promote apoptosis and suppress proliferation." Cell **85**(4): 549-61.
- Frolov, M. V., D. S. Huen, et al. (2001). "Functional antagonism between E2F family members." Genes Dev **15**(16): 2146-60.

- Germanier, M., M. Defais, et al. (2000). "Transcription-coupled repair is inducible in hamster cells." Nucleic Acids Res **28**(23): 4674-8.
- Giffin, W., H. Torrance, et al. (1996). "Sequence-specific DNA binding by Ku autoantigen and its effects on transcription." Nature **380**(6571): 265-8.
- Gu, Y., S. Jin, et al. (1997). "Ku70-deficient embryonic stem cells have increased ionizing radiosensitivity, defective DNA end-binding activity, and inability to support V(D)J recombination." Proc Natl Acad Sci U S A **94**(15): 8076-81.
- Gu, Y., K. J. Seidl, et al. (1997). "Growth retardation and leaky SCID phenotype of Ku70-deficient mice." Immunity **7**(5): 653-65.
- Harbour, J. W. and D. C. Dean (2000). "Rb function in cell-cycle regulation and apoptosis." Nat Cell Biol **2**(4): E65-7.
- Harper, J. W. and S. J. Elledge (1999). "Skipping into the E2F1-destruction pathway." Nat Cell Biol **1**(1): E5-7.
- Harrington, E. A., J. L. Bruce, et al. (1998). "pRB plays an essential role in cell cycle arrest induced by DNA damage." Proc Natl Acad Sci U S A **95**(20): 11945-50.
- Hateboer, G., R. M. Kerkhoven, et al. (1996). "Degradation of E2F by the ubiquitin-proteasome pathway: regulation by retinoblastoma family proteins and adenovirus transforming proteins." Genes Dev **10**(23): 2960-70.
- Hayes, S., P. Shivanov, et al. (1998). "DDB, a putative DNA repair protein, can function as a transcriptional partner of E2F1." Mol Cell Biol **18**(1): 240-9.
- He, S., B. L. Cook, et al. (2000). "E2F is required to prevent inappropriate S-phase entry of mammalian cells." Mol Cell Biol **20**(1): 363-71.
- Helin, K. (1998). "Regulation of cell proliferation by the E2F transcription factors." Curr Opin Genet Dev **8**(1): 28-35.
- Helin, K. and E. Harlow (1994). "Heterodimerization of the transcription factors E2F-1 and DP-1 is required for binding to the adenovirus E4 (ORF6/7) protein." J Virol **68**(8): 5027-35.
- Helin, K., C. L. Wu, et al. (1993). "Heterodimerization of the transcription factors E2F-1 and DP-1 leads to cooperative trans-activation." Genes Dev **7**(10): 1850-61.
- Hofferer, M., C. Wirbelauer, et al. (1999). "Increased levels of E2F-1-dependent DNA binding activity after UV- or gamma-irradiation." Nucleic Acids Res **27**(2): 491-5.

- Höfferer, M., C. Wirbelauer, et al. (1999). "Increased levels of E2F-1-dependent DNA binding activity after UV- or gamma-irradiation." Nucleic Acids Res **27**(2): 491-5.
- Hsieh, J. K., D. Yap, et al. (2002). "Novel function of the cyclin A binding site of E2F in regulating p53- induced apoptosis in response to DNA damage." Mol Cell Biol **22**(1): 78-93.
- Huang, H. and H. G. Claycamp (1993). "DNA excision repair as a component of adaptation to low doses of ionizing radiation in Escherichia coli." Int J Radiat Biol **64**(5): 613-9.
- Huang, Y., T. Ishiko, et al. (1997). "Role for E2F in DNA damage-induced entry of cells into S phase." Cancer Res **57**(17): 3640-3.
- Huber, H. E., G. Edwards, et al. (1993). "Transcription factor E2F binds DNA as a heterodimer." Proc Natl Acad Sci U S A **90**(8): 3525-9.
- Ikeda, M. A., L. Jakoi, et al. (1996). "A unique role for the Rb protein in controlling E2F accumulation during cell growth and differentiation." Proc Natl Acad Sci U S A **93**(8): 3215-20.
- Innocente, S. A., J. L. Abrahamson, et al. (1999). "p53 regulates a G2 checkpoint through cyclin B1." Proc Natl Acad Sci U S A **96**(5): 2147-52.
- Jackson, S. P. (1996). "DNA damage detection by DNA dependent protein kinase and related enzymes." Cancer Surv **28**: 261-79.
- Jeggo, P. A. (1997). "DNA-PK: at the cross-roads of biochemistry and genetics." Mutat Res **384**(1): 1-14.
- Jhappan, C., T. M. Yusufzai, et al. (2000). "The p53 response to DNA damage in vivo is independent of DNA-dependent protein kinase." Mol Cell Biol **20**(11): 4075-83.
- Jin, S., S. Inoue, et al. (1997). "Functions of the DNA dependent protein kinase." Cancer Surv **29**: 221-61.
- Jin, S. and D. T. Weaver (1997). "Double-strand break repair by Ku70 requires heterodimerization with Ku80 and DNA binding functions." Embo J **16**(22): 6874-85.
- Jost, C. A., D. Ginsberg, et al. (1996). "A conserved region of unknown function participates in the recognition of E2F family members by the adenovirus E4 ORF 6/7 protein." Virology **220**(1): 78-90.

- Kalma, Y., L. Marash, et al. (2001). "Expression analysis using DNA microarrays demonstrates that E2F-1 up-regulates expression of DNA replication genes including replication protein A2." Oncogene **20**(11): 1379-87.
- Knudsen, K. E., D. Booth, et al. (2000). "RB-dependent S-phase response to DNA damage." Mol Cell Biol **20**(20): 7751-63.
- Kohn, K. W. (1999). "Molecular interaction map of the mammalian cell cycle control and DNA repair systems." Mol Biol Cell **10**(8): 2703-34.
- Koike, M., T. Awaji, et al. (1999). "Differential subcellular localization of DNA-dependent protein kinase components Ku and DNA-PKcs during mitosis." J Cell Sci **112**(Pt 22): 4031-9.
- Kovesdi, I., R. Reichel, et al. (1986). "Identification of a cellular transcription factor involved in E1A trans-activation." Cell **45**(2): 219-28.
- Kowalik, T. F., J. DeGregori, et al. (1995). "E2F1 overexpression in quiescent fibroblasts leads to induction of cellular DNA synthesis and apoptosis." J Virol **69**(4): 2491-500.
- Krek, W., M. E. Ewen, et al. (1994). "Negative regulation of the growth-promoting transcription factor E2F-1 by a stably bound cyclin A-dependent protein kinase." Cell **78**(1): 161-72.
- Lavin, M. F. and Y. Shiloh (1997). "The genetic defect in ataxia-telangiectasia." Annu Rev Immunol **15**: 177-202.
- Lees, J. A., M. Saito, et al. (1993). "The retinoblastoma protein binds to a family of E2F transcription factors." Mol Cell Biol **13**(12): 7813-25.
- Lees-Miller, S. P. (1996). "The DNA-dependent protein kinase, DNA-PK: 10 years and no ends in sight." Biochem Cell Biol **74**(4): 503-12.
- Lees-Miller, S. P., Y. R. Chen, et al. (1990). "Human cells contain a DNA-activated protein kinase that phosphorylates simian virus 40 T antigen, mouse p53, and the human Ku autoantigen." Mol Cell Biol **10**(12): 6472-81.
- Lees-Miller, S. P., K. Sakaguchi, et al. (1992). "Human DNA-activated protein kinase phosphorylates serines 15 and 37 in the amino-terminal transactivation domain of human p53." Mol Cell Biol **12**(11): 5041-9.
- Leone, G., J. DeGregori, et al. (1998). "E2F3 activity is regulated during the cell cycle and is required for the induction of S phase." Genes Dev **12**(14): 2120-30.

- Leone, G., F. Nuckolls, et al. (2000). "Identification of a novel E2F3 product suggests a mechanism for determining specificity of repression by Rb proteins." Mol Cell Biol **20**(10): 3626-32.
- Leskov, K. S., T. Criswell, et al. (2001). "When X-ray-inducible proteins meet DNA double strand break repair." Semin Radiat Oncol **11**(4): 352-72.
- Li, L. J., G. S. Naeve, et al. (1993). "Temporal regulation of cyclin A-p107 and p33cdk2 complexes binding to a human thymidine kinase promoter element important for G1-S phase transcriptional regulation." Proc Natl Acad Sci U S A **90**(8): 3554-8.
- Lieber, M. R., U. Grawunder, et al. (1997). "Tying loose ends: roles of Ku and DNA-dependent protein kinase in the repair of double-strand breaks." Curr Opin Genet Dev **7**(1): 99-104.
- Lin, W. C., F. T. Lin, et al. (2001). "Selective induction of E2F1 in response to DNA damage, mediated by ATM- dependent phosphorylation." Genes Dev **15**(14): 1833-44.
- Marti, A., C. Wirbelauer, et al. (1999). "Interaction between ubiquitin-protein ligase SCFSKP2 and E2F-1 underlies the regulation of E2F-1 degradation." Nat Cell Biol **1**(1): 14-9.
- Meng, R. D., P. Phillips, et al. (1999). "p53-independent increase in E2F-1 expression enhances the cytotoxic effects of etoposide and of adriamycin." Int J Oncol **14**(1): 5-14.
- Messier, H., T. Fuller, et al. (1993). "p70 lupus autoantigen binds the enhancer of the T-cell receptor beta- chain gene." Proc Natl Acad Sci U S A **90**(7): 2685-9.
- Moberg, K., M. A. Starz, et al. (1996). "E2F-4 switches from p130 to p107 and pRB in response to cell cycle reentry." Mol Cell Biol **16**(4): 1436-49.
- Morkel, M., J. Wenkel, et al. (1997). "An E2F-like repressor of transcription." Nature **390**(6660): 567-8.
- Muller, H. and K. Helin (2000). "The E2F transcription factors: key regulators of cell proliferation." Biochim Biophys Acta **1470**(1): M1-12.
- Myung, K., C. Chen, et al. (2001). "Multiple pathways cooperate in the suppression of genome instability in *Saccharomyces cerevisiae*." Nature **411**(6841): 1073-6.
- Nevins, J. R. (1992). "E2F: a link between the Rb tumor suppressor protein and viral oncoproteins." Science **258**(5081): 424-9.

- Nevins, J. R. (2001). "The Rb/E2F pathway and cancer." Hum Mol Genet **10**(7): 699-703.
- Nevins, J. R., J. DeGregori, et al. (1997). "Functional analysis of E2F transcription factor." Methods Enzymol **283**: 205-19.
- Nevins, J. R., G. Leone, et al. (1997). "Role of the Rb/E2F pathway in cell growth control." J Cell Physiol **173**(2): 233-6.
- Nussenzweig, A., C. Chen, et al. (1996). "Requirement for Ku80 in growth and immunoglobulin V(D)J recombination." Nature **382**(6591): 551-5.
- O'Connor, D. J. and X. Lu (2000). "Stress signals induce transcriptionally inactive E2F-1 independently of p53 and Rb." Oncogene **19**(20): 2369-76.
- O'Connor, M. J., H. Zimmermann, et al. (1999). "Characterization of an E1A-CBP interaction defines a novel transcriptional adapter motif (TRAM) in CBP/p300." J Virol **73**(5): 3574-81.
- O'Connor, R. J. and P. Hearing (1994). "Mutually exclusive interaction of the adenovirus E4-6/7 protein and the retinoblastoma gene product with internal domains of E2F-1 and DP-1." J Virol **68**(11): 6848-62.
- Ormondroyd, E., S. de la Luna, et al. (1995). "A new member of the DP family, DP-3, with distinct protein products suggests a regulatory role for alternative splicing in the cell cycle transcription factor DRTF1/E2F." Oncogene **11**(8): 1437-46.
- Ouyang, H., A. Nussenzweig, et al. (1997). "Ku70 is required for DNA repair but not for T cell antigen receptor gene recombination In vivo." J Exp Med **186**(6): 921-9.
- Phillips, A. C., S. Bates, et al. (1997). "Induction of DNA synthesis and apoptosis are separable functions of E2F-1." Genes Dev **11**(14): 1853-63.
- Pierce, A. M., R. Schneider-Broussard, et al. (1999). "E2F1 has both oncogenic and tumor-suppressive properties in a transgenic model." Mol Cell Biol **19**(9): 6408-14.
- Polager, S., Y. Kalma, et al. (2002). "E2Fs up-regulate expression of genes involved in DNA replication, DNA repair and mitosis." Oncogene **21**(3): 437-46.
- Qin, X. Q., D. M. Livingston, et al. (1994). "Deregulated transcription factor E2F-1 expression leads to S-phase entry and p53-mediated apoptosis." Proc Natl Acad Sci U S A **91**(23): 10918-22.
- Reeves, W. H. (1992). "Antibodies to the p70/p80 (Ku) antigens in systemic lupus erythematosus." Rheum Dis Clin North Am **18**(2): 391-414.

- Ren, B., H. Cam, et al. (2002). "E2F integrates cell cycle progression with DNA repair, replication, and G(2)/M checkpoints." Genes Dev **16**(2): 245-56.
- Rogoff, H. A., M. T. Pickering, et al. (2002). "E2F1 induces phosphorylation of p53 that is coincident with p53 accumulation and apoptosis." Mol Cell Biol **22**(15): 5308-18.
- Ross, J. F., A. Naar, et al. (2001). "Active repression and E2F inhibition by pRB are biochemically distinguishable." Genes Dev **15**(4): 392-7.
- Ruscetti, T., B. E. Lehnert, et al. (1998). "Stimulation of the DNA-dependent protein kinase by poly(ADP-ribose) polymerase." J Biol Chem **273**(23): 14461-7.
- Shan, B., T. Durfee, et al. (1996). "Disruption of RB/E2F-1 interaction by single point mutations in E2F-1 enhances S-phase entry and apoptosis." Proc Natl Acad Sci U S A **93**(2): 679-84.
- Shin, E. K., S. G. Tevosian, et al. (1996). "The N-terminal region of E2F-1 is required for transcriptional activation of a new class of target promoter." J Biol Chem **271**(21): 12261-8.
- Slansky, J. E. and P. J. Farnham (1996). "Introduction to the E2F family: protein structure and gene regulation." Curr Top Microbiol Immunol **208**: 1-30.
- Smith, G. C. and S. P. Jackson (1999). "The DNA-dependent protein kinase." Genes Dev **13**(8): 916-34.
- Takata, M., M. S. Sasaki, et al. (1998). "Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells." Embo J **17**(18): 5497-508.
- Tuteja, N., R. Tuteja, et al. (1994). "Human DNA helicase II: a novel DNA unwinding enzyme identified as the Ku autoantigen." Embo J **13**(20): 4991-5001.
- Vandel, L. and T. Kouzarides (1999). "Residues phosphorylated by TFIIH are required for E2F-1 degradation during S-phase." Embo J **18**(15): 4280-91.
- Vassilev, A., J. Yamauchi, et al. (1998). "The 400 kDa subunit of the PCAF histone acetylase complex belongs to the ATM superfamily." Mol Cell **2**(6): 869-75.
- Wang, S., N. Nath, et al. (1999). "Prohibitin, a potential tumor suppressor, interacts with RB and regulates E2F function." Oncogene **18**(23): 3501-10.
- Wang, S., N. Nath, et al. (1999). "Rb and prohibitin target distinct regions of E2F1 for repression and respond to different upstream signals." Mol Cell Biol **19**(11): 7447-60.

-
- Weinberg, R. A. (1996). "E2F and cell proliferation: a world turned upside down." Cell **85**(4): 457-9.
- West, R. B., M. Yaneva, et al. (1998). "Productive and nonproductive complexes of Ku and DNA-dependent protein kinase at DNA termini." Mol Cell Biol **18**(10): 5908-20.
- Woo, R. A., K. G. McLure, et al. (1998). "DNA-dependent protein kinase acts upstream of p53 in response to DNA damage." Nature **394**(6694): 700-4.
- Wu, X. and A. J. Levine (1994). "p53 and E2F-1 cooperate to mediate apoptosis." Proc Natl Acad Sci U S A **91**(9): 3602-6.
- Wunderlich, M. and S. J. Berberich (2002). "Mdm2 inhibition of p53 induces E2F1 transactivation via p21." Oncogene **21**(28): 4414-21.
- Xu, M., K. A. Sheppard, et al. (1994). "Cyclin A/CDK2 binds directly to E2F-1 and inhibits the DNA-binding activity of E2F-1/DP-1 by phosphorylation." Mol Cell Biol **14**(12): 8420-31.
- Yaneva, M., T. Kowalewski, et al. (1997). "Interaction of DNA-dependent protein kinase with DNA and with Ku: biochemical and atomic-force microscopy studies." Embo J **16**(16): 5098-112.
- Zakian, V. A. (1995). "ATM-related genes: what do they tell us about functions of the human gene?" Cell **82**(5): 685-7.
- Zamanian, M. and N. B. La Thangue (1993). "Transcriptional repression by the Rb-related protein p107." Mol Biol Cell **4**(4): 389-96.
- Zhu, L., S. van den Heuvel, et al. (1993). "Inhibition of cell proliferation by p107, a relative of the retinoblastoma protein." Genes Dev **7**(7A): 1111-25.
- Ziebold, U., T. Reza, et al. (2001). "E2F3 contributes both to the inappropriate proliferation and to the apoptosis arising in Rb mutant embryos." Genes Dev **15**(4): 386-91.