

6 Literaturverzeichnis

- Albert, S., and D. Gallwitz. 1999. Two new members of a family of Ypt/Rab GTPase activating proteins. Promiscuity of substrate recognition. *J Biol Chem.* 274:33186-9.
- Albert, S., and D. Gallwitz. 2000. Msb4p, a protein involved in Cdc42p-dependent organization of the actin cytoskeleton, is a Ypt/Rab-specific GAP. *Biol Chem.* 381:453-6.
- Allan, B.B., B.D. Moyer, and W.E. Balch. 2000. Rab1 recruitment of p115 into a cis-SNARE complex: programming budding COPII vesicles for fusion. *Science.* 289:444-8.
- Alvarez, C., H. Fujita, A. Hubbard, and E. Sztul. 1999. ER to Golgi transport: Requirement for p115 at a pre-Golgi VTC stage. *J Cell Biol.* 147:1205-22.
- Andag, U., T. Neumann, and H.D. Schmitt. 2001. The coatomer-interacting protein Ds11p is required for Golgi-to-endoplasmic reticulum retrieval in yeast. *J Biol Chem.* 276:39150-60.
- Andag, U., and H.D. Schmitt. 2003. Ds11p, an essential component of the Golgi-endoplasmic reticulum retrieval system in yeast, uses the same sequence motif to interact with different subunits of the COPI vesicle coat. *J Biol Chem.* 278:51722-34.
- Antonin, W., I. Dulubova, D. Arac, S. Pabst, J. Plitzner, J. Rizo, and R. Jahn. 2002a. The N-terminal domains of syntaxin 7 and vti1b form three-helix bundles that differ in their ability to regulate SNARE complex assembly. *J Biol Chem.* 277:36449-56.
- Antonin, W., D. Fasshauer, S. Becker, R. Jahn, and T.R. Schneider. 2002b. Crystal structure of the endosomal SNARE complex reveals common structural principles of all SNAREs. *Nat Struct Biol.* 9:107-11.
- Antony, B., P. Gounon, R. Schekman, and L. Orci. 2003. Self-assembly of minimal COPII cages. *EMBO Rep.* 4:419-24.
- Antony, B., Madden D, Hamamoto S, Orci L, and S. R. 2001. Dynamics of the COPII coat with GTP and stable analogues. *Nat Cell Biol.* 3(6):531-7.
- Arac, D., I. Dulubova, J. Pei, I. Huryeva, N.V. Grishin, and J. Rizo. 2005. Three-dimensional structure of the rSly1 N-terminal domain reveals a conformational change induced by binding to Syntaxin 5. *J Mol Biol.* 346:589-601.
- Aridor, M., S.I. Bannykh, T. Rowe, and W.E. Balch. 1999. Cargo can modulate COPII vesicle formation from the endoplasmic reticulum. *J Biol Chem.* 274:4389-99.
- Aridor, M., J. Weissman, S. Bannykh, C. Nuoffer, and W.E. Balch. 1998. Cargo selection by the COPII budding machinery during export from the ER. *J Cell Biol.* 141:61-70.
- Ballou, L., P. Gopal, B. Krummel, M. Tammi, and C.E. Ballou. 1986. A mutation that prevents glucosylation of the lipid-linked oligosaccharide precursor leads to underglycosylation of secreted yeast invertase. *Proc Natl Acad Sci U S A.* 83:3081-5.
- Banfield, D.K., M.J. Lewis, and H.R. Pelham. 1995. A SNARE-like protein required for traffic through the Golgi complex. *Nature.* 375:806-9.
- Bannykh, S.I., T. Rowe, and W.E. Balch. 1996. The organization of endoplasmic reticulum export complexes. *J Cell Biol.* 135:19-35.
- Barlowe, C. 1997. Coupled ER to Golgi transport reconstituted with purified cytosolic proteins. *J. Cell Biol.* 139:1097-1108.
- Barlowe, C., L. Orci, T. Yeung, M. Hosobuchi, S. Hamamoto, N. Salama, M.F. Rexach, M. Ravazzola, M. Amherdt, and R. Schekman. 1994. COPII: a membrane coat formed by Sec proteins that drive vesicle budding from the endoplasmic reticulum. *Cell.* 77:895-907.
- Barlowe, C., and R. Schekman. 1993. SEC12 encodes a guanine-nucleotide-exchange factor essential for transport vesicle budding from the ER. *Nature.* 365:347-9.

- Barr, F.A., N. Nakamura, and G. Warren. 1998. Mapping the interaction between GRASP65 and GM130, components of a protein complex involved in the stacking of Golgi cisternae. *Embo J.* 17:3258-68.
- Barr, F.A., M. Puype, J. Vandekerckhove, and G. Warren. 1997. GRASP65, a protein involved in the stacking of Golgi cisternae. *Cell.* 91:253-62.
- Barrowman, J., M. Sacher, and S. Ferro-Novick. 2000. TRAPP stably associates with the Golgi and is required for vesicle docking. *Embo J.* 19:862-869.
- Becker, D.M., and L. Guarente. 1991. High-efficiency transformation of yeast by electroporation. *Methods Enzymol.* 194:182-7.
- Belden, W.J., and C. Barlowe. 2001. Role of Erv29p in collecting soluble secretory proteins into ER-derived transport vesicles. *Science.* 294:1528-31.
- Benli, M., F. Doring, D.G. Robinson, X. Yang, and D. Gallwitz. 1996. Two GTPase isoforms, Ypt31p and Ypt32p, are essential for Golgi function in yeast. *Embo J.* 15:6460-75.
- Bi, E., J.B. Chiavetta, H. Chen, G.C. Chen, C.S. Chan, and J.R. Pringle. 2000. Identification of novel, evolutionarily conserved Cdc42p-interacting proteins and of redundant pathways linking Cdc24p and Cdc42p to actin polarization in yeast. *Mol Biol Cell.* 11:773-93.
- Birnboim, H.C., and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7:1513-23.
- Bock, J.B., H.T. Matern, A.A. Peden, and R.H. Scheller. 2001. A genomic perspective on membrane compartment organization. *Nature.* 409:839-41.
- Boeddinghaus, C., A.J. Merz, R. Laage, and C. Ungermann. 2002. A cycle of Vam7p release from and PtdIns 3-P-dependent rebinding to the yeast vacuole is required for homotypic vacuole fusion. *J Cell Biol.* 157:79-89.
- Boehm, J., F. Letourneur, W. Ballensiefen, D. Ossipov, C. Demolliere, and H.D. Schmitt. 1997. Sec12p requires Rer1p for sorting to coatomer (COPI)-coated vesicles and retrieval to the ER. *J Cell Sci.* 110:991-1003.
- Boehm, J., H.D. Ulrich, R. Ossig, and H.D. Schmitt. 1994. Kex2-dependent invertase secretion as a tool to study the targeting of transmembrane proteins which are involved in ER-->Golgi transport in yeast. *Embo J.* 13:3696-710.
- Bonifacino, J.S., and B.S. Glick. 2004. The mechanisms of vesicle budding and fusion. *Cell.* 116:153-66.
- Bourne, H.R. 1988. Do GTPases direct membrane traffic in secretion? *Cell.* 53:669-71.
- Bracher, A., T. Dresbach, H. Betz, and W. Weissenhorn. 2000. Crystallization and preliminary X-ray analysis of squid neuronal Sec1. *Acta Crystallogr D Biol Crystallogr.* 56 (Pt 4):501-3.
- Bracher, A., and W. Weissenhorn. 2002. Structural basis for the Golgi membrane recruitment of Sly1p by Sed5p. *Embo J.* 21:6114-24.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 72:248-54.
- Brandhorst, D. 2005. Entwicklung eines *in vitro* Fluoreszenzassays zur Charakterisierung der Fusion von frühen Endosomen. Dissertation. Universität Oldenburg, Oldenburg, Germany.
- Brennwald, P., B. Kearns, K. Champion, S. Keranen, V. Bankaitis, and P. Novick. 1994. Sec9 is a SNAP-25-like component of a yeast SNARE complex that may be the effector of Sec4 function in exocytosis. *Cell.* 79:245-58.
- Brunger, A.T., and B. DeLaBarre. 2003. NSF and p97/VCP: similar at first, different at last. *FEBS Lett.* 555:126-33.
- Bryant, N.J., and D.E. James. 2001. Vps45p stabilizes the syntaxin homologue Tlg2p and positively regulates SNARE complex formation. *Embo J.* 20:3380-8.

- Burnette, W.N. 1981. "Western blotting": electrophoretic transfer of proteins from sodium dodecyl sulfate--polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein *A*. *Anal Biochem.* 112:195-203.
- Burri, L., and T. Lithgow. 2004. A complete set of SNAREs in yeast. *Traffic.* 5:45-52.
- Burri, L., O. Varlamov, C.A. Doege, K. Hofmann, T. Beilharz, J.E. Rothman, T.H. Sollner, and T. Lithgow. 2003. A SNARE required for retrograde transport to the endoplasmic reticulum. *Proc Natl Acad Sci U S A.* 100:9873-9877.
- Cao, X., N. Ballew, and C. Barlowe. 1998. Initial docking of ER-derived vesicles requires Uso1p and Ypt1p but is independent of SNARE proteins. *Embo J.* 17:2156-65.
- Cao, X., and C. Barlowe. 2000. Asymmetric requirements for a Rab GTPase and SNARE proteins in fusion of COPII vesicles with acceptor membranes. *J Cell Biol.* 149:55-66.
- Carroll, K.S., J. Hanna, I. Simon, J. Krise, P. Barbero, and S.R. Pfeffer. 2001. Role of Rab9 GTPase in facilitating receptor recruitment by TIP47. *Science.* 292:1373-6.
- Chantalat, S., R. Courbeyrette, F. Senic-Matuglia, C.L. Jackson, B. Goud, and A. Peyroche. 2003. A novel Golgi membrane protein is a partner of the ARF exchange factors Gea1p and Gea2p. *Mol Biol Cell.* 14:2357-71.
- Chavrier, P., and B. Goud. 1999. The role of ARF and Rab GTPases in membrane transport. *Curr Opin Cell Biol.* 11:466-75.
- Cheever, M.L., T.K. Sato, T. de Beer, T.G. Kutateladze, S.D. Emr, and M. Overduin. 2001. Phox domain interaction with PtdIns(3)P targets the Vam7 t-SNARE to vacuole membranes. *Nat Cell Biol.* 3:613-8.
- Chen, Y.A., S.J. Scales, S.M. Patel, Y.C. Doung, and R.H. Scheller. 1999. SNARE complex formation is triggered by Ca²⁺ and drives membrane fusion. *Cell.* 97:165-74.
- Chen, Y.A., and R.H. Scheller. 2001. SNARE-mediated membrane fusion. *Nat Rev Mol Cell Biol.* 2:98-106.
- Ciejek, E., and J. Thorner. 1979. Recovery of *S. cerevisiae* a cells from G1 arrest by alpha factor pheromone requires endopeptidase action. *Cell.* 18:623-35.
- Cohen, S.N., A.C. Chang, and L. Hsu. 1972. Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc Natl Acad Sci U S A.* 69:2110-4.
- Collins, K.M., N.L. Thorngren, R.A. Fratti, and W.T. Wickner. 2005. Sec17p and HOPS, in distinct SNARE complexes, mediate SNARE complex disruption or assembly for fusion. *Embo J.* 24:1775-86.
- Conibear, E., J.N. Cleck, and T.H. Stevens. 2003. Vps51p Mediates the Association of the GARP (Vps52/53/54) Complex with the Late Golgi t-SNARE Tlg1p. *Mol. Biol. Cell.* 14:1610-1623.
- Conibear, E., and T.H. Stevens. 2000. Vps52p, Vps53p, and Vps54p form a novel multisubunit complex required for protein sorting at the yeast late Golgi. *Mol Biol Cell.* 11:305-23.
- Cosson, P., C. Demolliere, S. Hennecke, R. Duden, and F. Letourneur. 1996. Delta- and zeta-COP, two coatomer subunits homologous to clathrin-associated proteins, are involved in ER retrieval. *Embo J.* 15:1792-8.
- Cosson, P., and F. Letourneur. 1994. Coatomer interaction with di-lysine endoplasmic reticulum retention motifs. *Science.* 263:1629-31.
- Cosson, P., S. Schröder-Kohne, D.S. Sweet, C. Demolliere, S. Hennecke, G. Frigerio, and F. Letourneur. 1997. The Sec20/Tip20p complex is involved in ER retrieval of di-lysine-tagged proteins. *Eur J Cell Biol.* 73:93-7.
- Cowles, C.R., S.D. Emr, and B.F. Horazdovsky. 1994. Mutations in the *VPS45* gene, a *SEC1* homologue, result in vacuolar protein sorting defects and accumulation of membrane vesicles. *J Cell Sci.* 107 (Pt 12):3449-59.

- Dacks, J.B., and W.F. Doolittle. 2002. Novel syntaxin gene sequences from *Giardia*, *Trypanosoma* and algae: implications for the ancient evolution of the eukaryotic endomembrane system. *J Cell Sci.* 115:1635-1642.
- Dascher, C., R. Ossig, D. Gallwitz, and H.D. Schmitt. 1991. Identification and structure of four yeast genes (*SLY*) that are able to suppress the functional loss of *YPT1*, a member of the RAS superfamily. *Mol Cell Biol.* 11:872-85.
- De Antoni, A., J. Schmitzova, H.H. Trepte, D. Gallwitz, and S. Albert. 2002. Significance of GTP hydrolysis in Ypt1p-regulated endoplasmic reticulum to Golgi transport revealed by the analysis of two novel Ypt1-GAPs. *J Biol Chem.* 277:41023-31.
- Dietrich, L.E., C. Boeddinghaus, T.J. LaGrassa, and C. Ungermann. 2003. Control of eukaryotic membrane fusion by N-terminal domains of SNARE proteins. *Biochim Biophys Acta.* 1641:111-9.
- Dietrich, L.E., R. Gurezka, M. Veit, and C. Ungermann. 2004. The SNARE Ykt6 mediates protein palmitoylation during an early stage of homotypic vacuole fusion. *Embo J.* 23:45-53.
- Dietrich, L.E., K. Peplowska, T.J. Lagrassa, H. Hou, J. Rohde, and C. Ungermann. 2005. The SNARE Ykt6 is released from yeast vacuoles during an early stage of fusion. *EMBO Rep.* 6:245-50.
- Dilcher, M., B. Kohler, and G.F. von Mollard. 2001. Genetic interactions with the yeast Q-SNARE *VT11* reveal novel functions for the R-SNARE *YKT6*. *J. Biol. Chem.* 276:34537-34544.
- Dilcher, M., B. Veith, S. Chidambaram, E. Hartmann, H.D. Schmitt, and G. Fischer von Mollard. 2003. Use1p is a yeast SNARE protein required for retrograde traffic to the ER. *Embo J.* 22:3664-74.
- Dogic, D., B. de Chassey, E. Pick, D. Cassel, Y. Lefkir, S. Hennecke, P. Cosson, and F. Letourneur. 1999. The ADP-ribosylation factor GTPase-activating protein Glo3p is involved in ER retrieval. *Eur J Cell Biol.* 78:305-10.
- Donaldson, J.G., D. Cassel, R.A. Kahn, and R.D. Klausner. 1992. ADP-ribosylation factor, a small GTP-binding protein, is required for binding of the coatomer protein beta-COP to Golgi membranes. *Proc Natl Acad Sci U S A.* 89:6408-12.
- Donaldson, J.G., A. Honda, and R. Weigert. 2005. Multiple activities for Arf1 at the Golgi complex. *Biochim Biophys Acta.* 1744:364-73.
- Dower, W.J., J.F. Miller, and C.W. Ragsdale. 1988. High efficiency transformation of *E. coli* by high voltage electroporation. *Nucleic Acids Res.* 16:6127-45.
- Du, L.L., R.N. Collins, and P.J. Novick. 1998. Identification of a Sec4p GTPase-activating protein (GAP) as a novel member of a Rab GAP family. *J Biol Chem.* 273:3253-6.
- Duden, R., M. Hosobuchi, S. Hamamoto, M. Winey, B. Byers, and R. Schekman. 1994. Yeast beta- and beta'-coat proteins (COP). Two coatomer subunits essential for endoplasmic reticulum-to-Golgi protein traffic. *J Biol Chem.* 269:24486-95.
- Duden, R., L. Kajikawa, L. Wuestehube, and R. Schekman. 1998. epsilon-COP is a structural component of coatomer that functions to stabilize alpha-COP. *Embo J.* 17:985-95.
- Dulubova, I., T. Yamaguchi, D. Arac, H. Li, I. Huryeva, S.W. Min, J. Rizo, and T.C. Sudhof. 2003. Convergence and divergence in the mechanism of SNARE binding by Sec1/Munc18-like proteins. *Proc Natl Acad Sci U S A.* 100:32-7.
- Dulubova, I., T. Yamaguchi, Y. Gao, S.W. Min, I. Huryeva, T.C. Sudhof, and J. Rizo. 2002. How Tlg2p/syntaxin 16 'snare' Vps45. *Embo J.* 21:3620-31.
- Fasshauer, D. 2003. Structural insights into the SNARE mechanism. *Biochim Biophys Acta.* 1641:87-97.
- Fasshauer, D., W. Antonin, M. Margittai, S. Pabst, and R. Jahn. 1999. Mixed and non-cognate SNARE complexes. Characterization of assembly and biophysical properties. *J Biol Chem.* 274:15440-6.

- Fasshauer, D., W. Antonin, V. Subramaniam, and R. Jahn. 2002. SNARE assembly and disassembly exhibit a pronounced hysteresis. *Nat Struct Biol.* 9:144-51.
- Fasshauer, D., W.K. Eliason, A.T. Brunger, and R. Jahn. 1998a. Identification of a minimal core of the synaptic SNARE complex sufficient for reversible assembly and disassembly. *Biochemistry.* 37:10354-62.
- Fasshauer, D., and M. Margittai. 2004. A transient N-terminal interaction of SNAP-25 and syntaxin nucleates SNARE assembly. *J Biol Chem.* 279:7613-21.
- Fasshauer, D., H. Otto, W.K. Eliason, R. Jahn, and A.T. Brunger. 1997. Structural changes are associated with soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptor complex formation. *J Biol Chem.* 272:28036-41.
- Fasshauer, D., R.B. Sutton, A.T. Brunger, and R. Jahn. 1998b. Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proc Natl Acad Sci U S A.* 95:15781-6.
- Ferro-Novick, S., and R. Jahn. 1994. Vesicle fusion from yeast to man. *Nature.* 370:191-193.
- Ferro-Novick, S., and P. Novick. 1993. The role of GTP-binding proteins in transport along the exocytic pathway. *Annu Rev Cell Biol.* 9:575-99.
- Fiebig, K.M., L.M. Rice, E. Pollock, and A.T. Brunger. 1999. Folding intermediates of SNARE complex assembly. *Nat Struct Biol.* 6:117-23.
- Filippini, F., V. Rossi, T. Galli, A. Budillon, M. D'Urso, and M. D'Esposito. 2001. Longins: a new evolutionary conserved VAMP family sharing a novel SNARE domain. *Trends Biochem Sci.* 26:407-9.
- Finger, F.P., and P. Novick. 2000. Synthetic interactions of the post-Golgi sec mutations of *Saccharomyces cerevisiae*. *Genetics.* 156:943-51.
- Fischer von Mollard, G., and T.H. Stevens. 1999. The *Saccharomyces cerevisiae* v-SNARE Vti1p is required for multiple membrane transport pathways to the vacuole. *Mol Biol Cell.* 10:1719-32.
- Franzusoff, A., K. Redding, J. Crosby, R.S. Fuller, and R. Schekman. 1991. Localization of components involved in protein transport and processing through the yeast Golgi apparatus. *J Cell Biol.* 112:27-37.
- Fukasawa, M., O. Varlamov, W.S. Eng, T.H. Sollner, and J.E. Rothman. 2004. Localization and activity of the SNARE Ykt6 determined by its regulatory domain and palmitoylation. *Proc Natl Acad Sci U S A.* 101:4815-4820.
- Fukuda, R., J.A. McNew, T. Weber, F. Parlati, T. Engel, W. Nickel, J.E. Rothman, and T.H. Sollner. 2000. Functional architecture of an intracellular membrane t-SNARE. *Nature.* 407:198-202.
- Furst, J., R.B. Sutton, J. Chen, A.T. Brunger, and N. Grigorieff. 2003. Electron cryomicroscopy structure of N-ethyl maleimide sensitive factor at 11 Å resolution. *Embo J.* 22:4365-74.
- Gallwitz, D., C. Donath, and C. Sander. 1983. A yeast gene encoding a protein homologous to the human c-has/bas proto-oncogene product. *Nature.* 306:704-7.
- Gallwitz, D., and R. Jahn. 2003. The riddle of the Sec1/Munc-18 proteins - new twists added to their interactions with SNAREs. *Trends Biochem Sci.* 28:113-6.
- Garcia, E., P. McPherson, T. Chilcote, K. Takei, and P. De Camilli. 1995. rbSec1A and B colocalize with syntaxin 1 and SNAP-25 throughout the axon, but are not in a stable complex with syntaxin. *J. Cell Biol.* 129:105-120.
- Gavin, A.C., M. Bosche, R. Krause, P. Grandi, M. Marzioch, A. Bauer, J. Schultz, J.M. Rick, A.M. Michon, C.M. Cruciat, M. Remor, C. Hofert, M. Schelder, M. Brajenovic, H. Ruffner, A. Merino, K. Klein, M. Hudak, D. Dickson, T. Rudi, V. Gnau, A. Bauch, S. Bastuck, B. Huhse, C. Leutwein, M.A. Heurtier, R.R. Copley, A. Edelmann, E. Querfurth, V. Rybin, G. Drewes, M. Raida, T. Bouwmeester, P. Bork, B. Seraphin, B. Kuster, G. Neubauer, and G. Superti-Furga. 2002. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature.* 415:141-7.

- Gaynor, E.C., T.R. Graham, and S.D. Emr. 1998. COPI in ER/Golgi and intra-Golgi transport: do yeast COPI mutants point the way? *Biochim Biophys Acta.* 1404:33-51.
- Gerrard, S.R., N.J. Bryant, and T.H. Stevens. 2000. *VPS21* controls entry of endocytosed and biosynthetic proteins into the yeast prevacuolar compartment. *Mol Biol Cell.* 11:613-26.
- Gietz, D., A. St Jean, R.A. Woods, and R.H. Schiestl. 1992. Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* 20:1425.
- Glick, B.S., T. Elston, and G. Oster. 1997. A cisternal maturation mechanism can explain the asymmetry of the Golgi stack. *FEBS Lett.* 414:177-81.
- Glick, B.S., and V. Malhotra. 1998. The curious status of the Golgi apparatus. *Cell.* 95:883-9.
- Gonzalez, L.C., Jr., W.I. Weis, and R.H. Scheller. 2001. A novel snare N-terminal domain revealed by the crystal structure of Sec22b. *J Biol Chem.* 276:24203-11.
- Gotte, M., and G.F. von Mollard. 1998. A new beat for the SNARE drum. *Trends Cell Biol.* 8:215-8.
- Gould, S.J., D. McCollum, A.P. Spong, J.A. Heyman, and S. Subramani. 1992. Development of the yeast *Pichia pastoris* as a model organism for a genetic and molecular analysis of peroxisome assembly. *Yeast.* 8:613-28.
- Graham, M.E., P. Washbourne, M.C. Wilson, and R.D. Burgoyne. 2001. SNAP-25 with mutations in the zero layer supports normal membrane fusion kinetics. *J Cell Sci.* 114:4397-4405.
- Gupta, G.D., and I.B. Heath. 2002. Predicting the distribution, conservation, and functions of SNAREs and related proteins in fungi. *Fungal Genet Biol.* 36:1-21.
- Hama, H., G.G. Tall, and B.F. Horazdovsky. 1999. Vps9p is a guanine nucleotide exchange factor involved in vesicle-mediated vacuolar protein transport. *J Biol Chem.* 274:15284-91.
- Hanson, P.I., J.E. Heuser, and R. Jahn. 1997. Neurotransmitter release - four years of SNARE complexes. *Curr Opin Neurobiol.* 7:310-5.
- Hara-Kuge, S., O. Kuge, L. Orci, M. Amherdt, M. Ravazzola, F.T. Wieland, and J.E. Rothman. 1994. En bloc incorporation of coatomer subunits during the assembly of COP-coated vesicles. *J Cell Biol.* 124:883-92.
- Hardwick, K., and H. Pelham. 1992. *SED5* encodes a 39-kD integral membrane protein required for vesicular transport between the ER and the Golgi complex. *J. Cell Biol.* 119:513-521.
- Hardwick, K.G., J.C. Boothroyd, A.D. Rudner, and H.R. Pelham. 1992. Genes that allow yeast cells to grow in the absence of the HDEL receptor. *Embo J.* 11:4187-95.
- Harlow, E., and D. Lane. 1988. Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Hay, J.C., and R.H. Scheller. 1997. SNAREs and NSF in targeted membrane fusion. *Curr Opin Cell Biol.* 9:505-12.
- Hemsley, A., N. Arnhheim, M.D. Toney, G. Cortopassi, and D.J. Galas. 1989. A simple method for site-directed mutagenesis using the polymerase chain reaction. *Nucleic Acids Res.* 17:6545-51.
- Hicke, L., and R. Schekman. 1989. Yeast Sec23p acts in the cytoplasm to promote protein transport from the endoplasmic reticulum to the Golgi complex *in vivo* and *in vitro*. *Embo J.* 8:1677-84.
- Hicke, L., T. Yoshihisa, and R. Schekman. 1992. Sec23p and a novel 105-kDa protein function as a multimeric complex to promote vesicle budding and protein transport from the endoplasmic reticulum. *Mol Biol Cell.* 3:667-76.
- Hinners, I., F. Wendler, H. Fei, L. Thomas, G. Thomas, and S.A. Tooze. 2003. AP-1 recruitment to VAMP4 is modulated by phosphorylation-dependent binding of PACS-1. *EMBO Rep.* 4:1182-9.
- Hirose, H., K. Arasaki, N. Dohmae, K. Takio, K. Hatsuzawa, M. Nagahama, K. Tani, A. Yamamoto, M. Tohyama, and M. Tagaya. 2004. Implication of ZW10 in membrane trafficking between the endoplasmic reticulum and Golgi. *Embo J.* 23:1267-78.

- Hoffman, C.S., and F. Winston. 1987. A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene*. 57:267-72.
- Hohl, T.M., F. Parlati, C. Wimmer, J.E. Rothman, T.H. Sollner, and H. Engelhardt. 1998. Arrangement of subunits in 20 S particles consisting of NSF, SNAPs, and SNARE complexes. *Mol Cell*. 2:539-48.
- Holthuis, J.C., B.J. Nichols, S. Dhruvakumar, and H.R. Pelham. 1998. Two syntaxin homologues in the TGN/endosomal system of yeast. *Embo J*. 17:113-26.
- Hong, W. 2005. SNAREs and traffic. *Biochim Biophys Acta*. 1744:493-517.
- Horazdovsky, B.F., G.R. Busch, and S.D. Emr. 1994. *VPS21* encodes a rab5-like GTP binding protein that is required for the sorting of yeast vacuolar proteins. *Embo J*. 13:1297-309.
- Hosobuchi, M., T. Kreis, and R. Schekman. 1992. *SEC21* is a gene required for ER to Golgi protein transport that encodes a subunit of a yeast coatomer. *Nature*. 360:603-5.
- Hughson, F.M. 1999. Membrane fusion: structure snared at last. *Curr Biol*. 9:R49-52.
- Ito, H., Y. Fukuda, K. Murata, and A. Kimura. 1983. Transformation of intact yeast cells treated with alkali cations. *J Bacteriol*. 153:163-8.
- Itoh, T., E.A. Toh, and Y. Matsui. 2004. Mmr1p is a mitochondrial factor for Myo2p-dependent inheritance of mitochondria in the budding yeast. *Embo J*. 23:2520-30.
- Itoh, T., A. Watabe, E.A. Toh, and Y. Matsui. 2002. Complex formation with Ypt11p, a rab-type small GTPase, is essential to facilitate the function of Myo2p, a class V myosin, in mitochondrial distribution in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 22:7744-57.
- Jahn, R. 2004. Principles of exocytosis and membrane fusion. *Ann NY Acad Sci*. 1014:170-178.
- Jahn, R., T. Lang, and T.C. Sudhof. 2003. Membrane fusion. *Cell*. 112:519-33.
- Jedd, G., C. Richardson, R. Litt, and N. Segev. 1995. The Ypt1 GTPase is essential for the first two steps of the yeast secretory pathway. *J Cell Biol*. 131:583-90.
- Jones, S., C. Newman, F. Liu, and N. Segev. 2000. The TRAPP Complex Is a Nucleotide Exchanger for Ypt1 and Ypt31/32. *Mol Biol Cell*. 11:4403-4411.
- Julius, D., R. Schekman, and J. Thorner. 1984. Glycosylation and processing of prepro-alpha-factor through the yeast secretory pathway. *Cell*. 36:309-18.
- Kahn, R.A., P. Randazzo, T. Serafini, O. Weiss, C. Rulka, J. Clark, M. Amherdt, P. Roller, L. Orci, and J.E. Rothman. 1992. The amino terminus of ADP-ribosylation factor (ARF) is a critical determinant of ARF activities and is a potent and specific inhibitor of protein transport. *J Biol Chem*. 267:13039-46.
- Kaiser, C.A., and R. Schekman. 1990. Distinct sets of *SEC* genes govern transport vesicle formation and fusion early in the secretory pathway. *Cell*. 61:723-33.
- Kassir, Y., and G. Simchen. 1991. Monitoring meiosis and sporulation in *Saccharomyces cerevisiae*. *Methods Enzymol*. Vol. 194. 94-110.
- Katz, L., and P. Brennwald. 2000. Testing the 3Q:1R "rule": mutational analysis of the ionic "zero" layer in the yeast exocytic SNARE complex reveals no requirement for arginine. *Mol Biol Cell*. 11:3849-58.
- Kee, Y., J.S. Yoo, C.D. Hazuka, K.E. Peterson, S.C. Hsu, and R.H. Scheller. 1997. Subunit structure of the mammalian exocyst complex. *Proc Natl Acad Sci U S A*. 94:14438-43.
- Khan, A.H., D.C. Thurmond, C. Yang, B.P. Ceresa, C.D. Sigmund, and J.E. Pessin. 2001. Munc18c regulates insulin-stimulated GLUT4 translocation to the transverse tubules in skeletal muscle. *J Biol Chem*. 276:4063-4069.
- Kim, D.-W., T. Massey, M. Sacher, M. Pypaert, and S. Ferro-Novick. 2001. Sgflp, a new component of the Sec34p/Sec35p complex. *Traffic*. 2:820-830.

- Kim, D.-W., M. Sacher, A. Scarpa, A.M. Quinn, and S. Ferro-Novick. 1999. High-copy suppressor analysis reveals a physical interaction between Sec34p and Sec35p, a protein implicated in vesicle docking. *Mol. Biol. Cell.* 10:3317-3329.
- Kim, M.-S., M.-J. Yi, K.-H. Lee, J. Wagner, C. Munger, Y.-G. Kim, M. Whiteway, M. Cygler, B.-H. Oh, and M. Sacher. 2005a. Biochemical and crystallographic studies reveal a specific interaction between TRAPP subunits Trs33p and Bet3p. *Traffic*. 6:1183-1195.
- Kim, Y.G., E.J. Sohn, J. Seo, K.J. Lee, H.S. Lee, I. Hwang, M. Whiteway, M. Sacher, and B.H. Oh. 2005b. Crystal structure of bet3 reveals a novel mechanism for Golgi localization of tethering factor TRAPP. *Nat Struct Mol Biol.* 12:38-45.
- Klenow, H., and I. Henningsen. 1970. Selective elimination of the exonuclease activity of the deoxyribonucleic acid polymerase from *Escherichia coli* B by limited proteolysis. *Proc Natl Acad Sci U S A.* 65:168-75.
- Kosodo, Y., Y. Noda, H. Adachi, and K. Yoda. 2002. Binding of Sly1 to Sed5 enhances formation of the yeast early Golgi SNARE complex. *J Cell Sci.* 115:3683-91.
- Kraynack, B.A., A. Chan, E. Rosenthal, M. Essid, B. Umansky, M.G. Waters, and H.D. Schmitt. 2005. Dsl1p, Tip20p, and the novel Dsl3(Sec39) protein are required for the stability of the Q/t-SNARE complex at the endoplasmic reticulum in yeast. *Mol. Biol. Cell.* 16:3963-3977.
- Kuehn, M.J., J.M. Herrmann, and R. Schekman. 1998. COPII-cargo interactions direct protein sorting into ER-derived transport vesicles. *Nature*. 391:187-90.
- Kuge, O., C. Dascher, L. Orci, T. Rowe, M. Amherdt, H. Plutner, M. Ravazzola, G. Tanigawa, J. Rothman, and W. Balch. 1994. Sar1 promotes vesicle budding from the endoplasmic reticulum but not Golgi compartments. *J. Cell Biol.* 125:51-65.
- Kummel, D., J.J. Muller, Y. Roske, R. Misselwitz, K. Bussow, and U. Heinemann. 2005. The structure of the TRAPP subunit TPC6 suggests a model for a TRAPP subcomplex. *EMBO Rep.* 6:787-93.
- Kweon, H.S., G.V. Beznoussenko, M. Micaroni, R.S. Polishchuk, A. Trucco, O. Martella, D. Di Giandomenico, P. Marra, A. Fusella, A. Di Pentima, E.G. Berger, W.J. Geerts, A.J. Koster, K.N. Burger, A. Luini, and A.A. Mironov. 2004. Golgi enzymes are enriched in perforated zones of golgi cisternae but are depleted in COPI vesicles. *Mol Biol Cell.* 15:4710-24.
- Kweon, Y., A. Rothe, E. Conibear, and T.H. Stevens. 2003. Ykt6p is a multifunctional yeast R-SNARE that is required for multiple membrane transport pathways to the vacuole. *Mol Biol Cell.* 14:1868-81.
- Ladinsky, M.S., D.N. Mastronarde, J.R. McIntosh, K.E. Howell, and L.A. Staehelin. 1999. Golgi structure in three dimensions: functional insights from the normal rat kidney cell. *J. Cell Biol.* 144:1135-1149.
- Ladinsky, M.S., C.C. Wu, S. McIntosh, J.R. McIntosh, and K.E. Howell. 2002. Structure of the Golgi and distribution of reporter molecules at 20 degrees C reveals the complexity of the exit compartments. *Mol Biol Cell.* 13:2810-25.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227:680-5.
- Lanoix, J., J. Ouwendijk, A. Stark, E. Szafer, D. Cassel, K. Dejgaard, M. Weiss, and T. Nilsson. 2001. Sorting of Golgi resident proteins into different subpopulations of COPI vesicles: a role for ArfGAP1. *J Cell Biol.* 155:1199-212.
- Lazar, T., M. Gotte, and D. Gallwitz. 1997. Vesicular transport: how many Ypt/Rab-GTPases make a eukaryotic cell? *Trends Biochem Sci.* 22:468-72.
- Lee, M.C., L. Orci, S. Hamamoto, E. Futai, M. Ravazzola, and R. Schekman. 2005. Sar1p N-terminal helix initiates membrane curvature and completes the fission of a COPII vesicle. *Cell.* 122:605-17.

- Lee, M.C.S., E.A. Miller, J. Goldberg, L. Orci, and R. Schekman. 2004. Bi-directional protein transport between the ER and Golgi. *Annu Rev Cell Dev Biol.* 20:87-123.
- LePecq, J.B., and C. Paoletti. 1967. A fluorescent complex between ethidium bromide and nucleic acids. Physical-chemical characterization. *J Mol Biol.* 27:87-106.
- Letourneur, F., E.C. Gaynor, S. Hennecke, C. Demolliere, R. Duden, S.D. Emr, H. Riezman, and P. Cosson. 1994. Coatomer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell.* 79:1199-207.
- Lewis, M.J., and H.R. Pelham. 1990. A human homologue of the yeast HDEL receptor. *Nature.* 348:162-3.
- Lewis, M.J., and H.R. Pelham. 1996. SNARE-mediated retrograde traffic from the Golgi complex to the endoplasmic reticulum. *Cell.* 85:205-15.
- Lewis, M.J., J.C. Rayner, and H.R. Pelham. 1997. A novel SNARE complex implicated in vesicle fusion with the endoplasmic reticulum. *Embo J.* 16:3017-24.
- Li, B., and J.R. Warner. 1996. Mutation of the Rab6 homologue of *Saccharomyces cerevisiae*, YPT6, inhibits both early Golgi function and ribosome biosynthesis. *J Biol Chem.* 271:16813-9.
- Linstedt, A.D., M. Foguet, M. Renz, H.P. Seelig, B.S. Glick, and H.P. Hauri. 1995. A C-terminally-anchored Golgi protein is inserted into the endoplasmic reticulum and then transported to the Golgi apparatus. *Proc Natl Acad Sci U S A.* 92:5102-5.
- Linstedt, A.D., and H.P. Hauri. 1993. Giantin, a novel conserved Golgi membrane protein containing a cytoplasmic domain of at least 350 kDa. *Mol Biol Cell.* 4:679-93.
- Linstedt, A.D., S.A. Jesch, A. Mehta, T.H. Lee, R. Garcia-Mata, D.S. Nelson, and E. Sztul. 2000. Binding relationships of membrane tethering components. The giantin N terminus and the GM130 N terminus compete for binding to the p115 C terminus. *J Biol Chem.* 275:10196-201.
- Littleton, J.T., E.R. Chapman, R. Kreber, M.B. Garment, S.D. Carlson, and B. Ganetzky. 1998. Temperature-sensitive paralytic mutations demonstrate that synaptic exocytosis requires SNARE complex assembly and disassembly. *Neuron.* 21:401-13.
- Liu, Y., and C. Barlowe. 2002. Analysis of Sec22p in endoplasmic reticulum/Golgi transport reveals cellular redundancy in SNARE protein function. *Mol. Biol. Cell.* 13:3314-3324.
- Lodish, H., A. Berk, S.L. Zipursky, P. Matsudaira, D. Baltimore, and J.E. Darnell. 1999. Molecular Cell Biology. W. H. Freeman & Co., New York.
- Lowe, M., and T.E. Kreis. 1995. *In vitro* assembly and disassembly of coatomer. *J Biol Chem.* 270:31364-71.
- Lowe, M., N. Nakamura, and G. Warren. 1998. Golgi division and membrane traffic. *Trends Cell Biol.* 8:40-4.
- Luo, Z., and D. Gallwitz. 2003. Biochemical and genetic evidence for the involvement of yeast Ypt6-GTPase in protein retrieval to different Golgi compartments. *J. Biol. Chem.* 278:791-799.
- Lupashin, V., and E. Sztul. 2005. Golgi tethering factors. *Biochim Biophys Acta.* 1744:325-39.
- Lupashin, V.V., I.D. Pokrovskaya, J.A. McNew, and M.G. Waters. 1997. Characterization of a novel yeast SNARE protein implicated in Golgi retrograde traffic. *Mol Biol Cell.* 8:2659-76.
- Lupashin, V.V., and M.G. Waters. 1997. t-SNARE activation through transient interaction with a rab-like guanosine triphosphatase. *Science.* 276:1255-8.
- Mammoto, A., T. Sasaki, Y. Kim, and Y. Takai. 2000. Physical and functional interaction of rabphilin-11 with mammalian Sec13 protein. Implication in vesicle trafficking. *J Biol Chem.* 275:13167-70.
- Martinez-Arca, S., S. Arold, R. Rudge, F. Laroche, and T. Galli. 2004. A mutant impaired in SNARE complex dissociation identifies the plasma membrane as first target of Synaptobrevin 2. *Traffic.* 5:371-382.

- Martinez-Arca, S., R. Rudge, M. Vacca, G. Raposo, J. Camonis, V. Proux-Gillardeaux, L. Daviet, E. Formstecher, A. Hamburger, F. Filippini, M. D'Esposito, and T. Galli. 2003. A dual mechanism controlling the localization and function of exocytic v-SNAREs. *Proc Natl Acad Sci U S A.* 100:9011-9016.
- Martinez-Menarguez, J.A., R. Prekeris, V.M. Oorschot, R. Scheller, J.W. Slot, H.J. Geuze, and J. Klumperman. 2001. Peri-Golgi vesicles contain retrograde but not anterograde proteins consistent with the cisternal progression model of intra-Golgi transport. *J Cell Biol.* 155:1213-24.
- Marz, K.E., J.M. Lauer, and P.I. Hanson. 2003. Defining the SNARE complex binding surface of (alpha)-SNAP: implications for SNARE complex disassembly. *J. Biol. Chem.* 278:27000-27008.
- Matsuoka, K., Y. Morimitsu, K. Uchida, and R. Schekman. 1998. Coat assembly directs v-SNARE concentration into synthetic COPII vesicles. *Mol Cell.* 2:703-8.
- McNew, J.A., F. Parlati, R. Fukuda, R.J. Johnston, K. Paz, F. Paumet, T.H. Sollner, and J.E. Rothman. 2000. Compartmental specificity of cellular membrane fusion encoded in SNARE proteins. *Nature.* 407:153-9.
- McNew, J.A., M. Sogaard, N.M. Lampen, S. Machida, R.R. Ye, L. Lacomis, P. Tempst, J.E. Rothman, and T.H. Sollner. 1997. Ykt6p, a prenylated SNARE essential for endoplasmic reticulum-Golgi transport. *J. Biol. Chem.* 272:17776-17783.
- Meyer, T.S., and B.L. Lamberts. 1965. Use of coomassie brilliant blue R250 for the electrophoresis of microgram quantities of parotid saliva proteins on acrylamide-gel strips. *Biochim Biophys Acta.* 107:144-5.
- Mezzacasa, A., and A. Helenius. 2002. The transitional ER defines a boundary for quality control in the secretion of tsO45 VSV glycoprotein. *Traffic.* 3:833-49.
- Miller, E., B. Antonny, S. Hamamoto, and R. Schekman. 2002. Cargo selection into COPII vesicles is driven by the Sec24p subunit. *Embo J.* 21:6105-13.
- Miller, E.A., T.H. Beilharz, P.N. Malkus, M.C. Lee, S. Hamamoto, L. Orci, and R. Schekman. 2003. Multiple cargo binding sites on the COPII subunit Sec24p ensure capture of diverse membrane proteins into transport vesicles. *Cell.* 114:497-509.
- Mironov, A.A., G.V. Beznoussenko, P. Nicoziani, O. Martella, A. Trucco, H.S. Kweon, D. Di Giandomenico, R.S. Polishchuk, A. Fusella, P. Lupetti, E.G. Berger, W.J. Geerts, A.J. Koster, K.N. Burger, and A. Luini. 2001. Small cargo proteins and large aggregates can traverse the Golgi by a common mechanism without leaving the lumen of cisternae. *J Cell Biol.* 155:1225-38.
- Mironov, A.A., G.V. Beznoussenko, R.S. Polishchuk, and A. Trucco. 2005. Intra-Golgi transport: a way to a new paradigm? *Biochim Biophys Acta.* 1744:340-50.
- Mironov, A.A., A.A. Mironov, Jr., G.V. Beznoussenko, A. Trucco, P. Lupetti, J.D. Smith, W.J. Geerts, A.J. Koster, K.N. Burger, M.E. Martone, T.J. Deerinck, M.H. Ellisman, and A. Luini. 2003. ER-to-Golgi carriers arise through direct en bloc protrusion and multistage maturation of specialized ER exit domains. *Dev Cell.* 5:583-94.
- Misura, K.M., J.B. Bock, L.C. Gonzalez, Jr., R.H. Scheller, and W.I. Weis. 2002. Three-dimensional structure of the amino-terminal domain of syntaxin 6, a SNAP-25 C homolog. *Proc Natl Acad Sci U S A.* 99:9184-9.
- Misura, K.M., R.H. Scheller, and W.I. Weis. 2000. Three-dimensional structure of the neuronal-Sec1-syntaxin 1a complex. *Nature.* 404:355-62.
- Morsomme, P., and H. Riezman. 2002. The Rab GTPase Ypt1p and tethering factors couple protein sorting at the ER to vesicle targeting to the Golgi apparatus. *Dev Cell.* 2:307-17.
- Mossessova, E., L.C. Bickford, and J. Goldberg. 2003. SNARE selectivity of the COPII coat. *Cell.* 114:483-95.

- Muniz, M., P. Morsomme, and H. Riezman. 2001. Protein sorting upon exit from the endoplasmic reticulum. *Cell.* 104:313-320.
- Munro, S., and H.R. Pelham. 1987. A C-terminal signal prevents secretion of luminal ER proteins. *Cell.* 48:899-907.
- Nakajima, H., A. Hirata, Y. Ogawa, T. Yonehara, K. Yoda, and M. Yamasaki. 1991. A cytoskeleton-related gene, uso1, is required for intracellular protein transport in *Saccharomyces cerevisiae*. *J Cell Biol.* 113:245-60.
- Nakajima, K.-i., H. Hirose, M. Taniguchi, H. Kurashina, K. Arasaki, M. Nagahama, K. Tani, A. Yamamoto, and M. Tagaya. 2004. Involvement of BNIP1 in apoptosis and endoplasmic reticulum membrane fusion. *Embo J.* 23:3216-3226.
- Nakamura, N., M. Lowe, T.P. Levine, C. Rabouille, and G. Warren. 1997. The vesicle docking protein p115 binds GM130, a cis-Golgi matrix protein, in a mitotically regulated manner. *Cell.* 89:445-55.
- Nakamura, N., C. Rabouille, R. Watson, T. Nilsson, N. Hui, P. Slusarewicz, T.E. Kreis, and G. Warren. 1995. Characterization of a cis-Golgi matrix protein, GM130. *J Cell Biol.* 131:1715-26.
- Nakano, A., D. Brada, and R. Schekman. 1988. A membrane glycoprotein, Sec12p, required for protein transport from the endoplasmic reticulum to the Golgi apparatus in yeast. *J. Cell Biol.* 107:851-863.
- Nakano, A., and M. Muramatsu. 1989. A novel GTP-binding protein, Sar1p, is involved in transport from the endoplasmic reticulum to the Golgi apparatus. *J. Cell Biol.* 109:2677-2691.
- Nicholson, K.L., M. Munson, R.B. Miller, T.J. Filip, R. Fairman, and F.M. Hughson. 1998. Regulation of SNARE complex assembly by an N-terminal domain of the t-SNARE Sso1p. *Nat Struct Biol.* 5:793-802.
- Nie, Z., D.S. Hirsch, and P.A. Randazzo. 2003. Arf and its many interactors. *Curr Opin Cell Biol.* 15:396-404.
- Nishikawa, S., and A. Nakano. 1993. Identification of a gene required for membrane protein retention in the early secretory pathway. *Proc Natl Acad Sci U S A.* 90:8179-83.
- Novick, P., C. Field, and R. Schekman. 1980. Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. *Cell.* 21:205-15.
- Nufer, O., F. Kappeler, S. Guldbrandsen, and H.P. Hauri. 2003. ER export of ERGIC-53 is controlled by cooperation of targeting determinants in all three of its domains. *J Cell Sci.* 116:4429-40.
- Oka, T., D. Ungar, F.M. Hughson, and M. Krieger. 2004. The COG and COPI complexes interact to control the abundance of GEARs, a subset of Golgi integral membrane proteins. *Mol. Biol. Cell.* 15:2423-2435.
- Oka, T., E. Vasile, M. Penman, C.D. Novina, D.M. Dykxhoorn, D. Ungar, F.M. Hughson, and M. Krieger. 2005. Genetic analysis of the subunit organization and function of the Conserved Oligomeric Golgi (COG) complex: studies of COG5- and COG7-deficient mammalian cells. *J. Biol. Chem.* 280:32736-32745.
- Orci, L., B.S. Glick, and J.E. Rothman. 1986. A new type of coated vesicular carrier that appears not to contain clathrin: its possible role in protein transport within the Golgi stack. *Cell.* 46:171-84.
- Orci, L., M. Ravazzola, P. Meda, C. Holcomb, H. Moore, L. Hicke, and R. Schekman. 1991. Mammalian Sec23p Homologue is restricted to the endoplasmic reticulum transitional cytoplasm. *Proc Natl Acad Sci U S A.* 88:8611-8615.
- Ossig, R., C. Dascher, H.H. Trepte, H.D. Schmitt, and D. Gallwitz. 1991. The yeast *SLY* gene products, suppressors of defects in the essential GTP-binding Ypt1 protein, may act in endoplasmic reticulum-to-Golgi transport. *Mol Cell Biol.* 11:2980-93.

- Ossig, R., W. Laufer, H.D. Schmitt, and D. Gallwitz. 1995. Functionality and specific membrane localization of transport GTPases carrying C-terminal membrane anchors of synaptobrevin-like proteins. *Embo J.* 14:3645-53.
- Ossig, R., H.D. Schmitt, B. de Groot, D. Riedel, S. Keranen, H. Ronne, H. Grubmuller, and R. Jahn. 2000. Exocytosis requires asymmetry in the central layer of the SNARE complex. *Embo J.* 19:6000-10.
- Otte, S., and C. Barlowe. 2004. Sorting signals can direct receptor-mediated export of soluble proteins into COPII vesicles. *Nat Cell Biol.* 6:1189-94.
- Palade, G. 1975. Intracellular aspects of the process of protein synthesis. *Science.* 189:347-58.
- Palmer, D.J., J.B. Helms, C.J. Beckers, L. Orci, and J.E. Rothman. 1993. Binding of coatomer to Golgi membranes requires ADP-ribosylation factor. *J Biol Chem.* 268:12083-9.
- Paravicini, G., B.F. Horazdovsky, and S.D. Emr. 1992. Alternative pathways for the sorting of soluble vacuolar proteins in yeast: a vps35 null mutant missorts and secretes only a subset of vacuolar hydrolases. *Mol Biol Cell.* 3:415-27.
- Parlati, F., J.A. McNew, R. Fukuda, R. Miller, T.H. Sollner, and J.E. Rothman. 2000. Topological restriction of SNARE-dependent membrane fusion. *Nature.* 407:194-8.
- Parlati, F., O. Varlamov, K. Paz, J.A. McNew, D. Hurtado, T.H. Sollner, and J.E. Rothman. 2002. Distinct SNARE complexes mediating membrane fusion in Golgi transport based on combinatorial specificity. *Proc Natl Acad Sci U S A.* 99:5424-5429.
- Paumet, F., B. Brugger, F. Parlati, J.A. McNew, T.H. Sollner, and J.E. Rothman. 2001. A t-SNARE of the endocytic pathway must be activated for fusion. *J Cell Biol.* 155:961-8.
- Paumet, F., V. Rahimian, and J.E. Rothman. 2004. The specificity of SNARE-dependent fusion is encoded in the SNARE motif. *Proc Natl Acad Sci U S A.* 101:3376-3380.
- Peden, A.A., G.Y. Park, and R.H. Scheller. 2001. The Di-leucine motif of vesicle-associated membrane protein 4 is required for its localization and AP-1 binding. *J Biol Chem.* 276:49183-7.
- Pelham, H.R. 1996. The dynamic organisation of the secretory pathway. *Cell Struct Funct.* 21:413-9.
- Pelham, H.R., K.G. Hardwick, and M.J. Lewis. 1988. Sorting of soluble ER proteins in yeast. *Embo J.* 7:1757-62.
- Peng, R., and D. Gallwitz. 2002. Sly1 protein bound to Golgi syntaxin Sed5p allows assembly and contributes to specificity of SNARE fusion complexes. *J Cell Biol.* 157:645-55.
- Pereira-Leal, J.B., and M.C. Seabra. 2000. The mammalian Rab family of small GTPases: definition of family and subfamily sequence motifs suggests a mechanism for functional specificity in the Ras superfamily. *J Mol Biol.* 301:1077-87.
- Peterson, M.R., C.G. Burd, and S.D. Emr. 1999. Vac1p coordinates Rab and phosphatidylinositol 3-kinase signaling in Vps45p-dependent vesicle docking/fusion at the endosome. *Curr Biol.* 9:159-62.
- Pevsner, J., S.C. Hsu, J.E. Braun, N. Calakos, A.E. Ting, M.K. Bennett, and R.H. Scheller. 1994. Specificity and regulation of a synaptic vesicle docking complex. *Neuron.* 13:353-61.
- Peyroche, A., S. Paris, and C.L. Jackson. 1996. Nucleotide exchange on ARF mediated by yeast Gea1 protein. *Nature.* 384:479-81.
- Pfeffer, S.R. 2001. Rab GTPases: specifying and deciphering organelle identity and function. *Trends Cell Biol.* 11:487-91.
- Poon, P.P., D. Cassel, A. Spang, M. Rotman, E. Pick, R.A. Singer, and G.C. Johnston. 1999. Retrograde transport from the yeast Golgi is mediated by two ARF GAP proteins with overlapping function. *Embo J.* 18:555-64.
- Prekeris, R., J. Klumperman, and R.H. Scheller. 2000. Syntaxin 11 is an atypical SNARE abundant in the immune system. *Eur J Cell Biol.* 79:771-80.

- Preuss, D., J. Mulholland, A. Franzusoff, N. Segev, and D. Botstein. 1992. Characterization of the *Saccharomyces* Golgi complex through the cell cycle by immunoelectron microscopy. *Mol Biol Cell.* 3:789-803.
- Pullikuth, A.K., and P.J. Weidman. 2002. *In vitro* transport on *cis* and *trans* sides of the Golgi involves two distinct types of coatomer and ADP-ribosylation factor-independent transport intermediates. *J Biol Chem.* 277:50355-64.
- Rabouille, C., T.P. Levine, J.M. Peters, and G. Warren. 1995. An NSF-like ATPase, p97, and NSF mediate cisternal regrowth from mitotic Golgi fragments. *Cell.* 82:905-14.
- Ram, R.J., B. Li, and C.A. Kaiser. 2002. Identification of Sec36p, Sec37p, and Sec38p: components of yeast complex that contains Sec34p and Sec35p. *Mol. Biol. Cell.* 13:1484-1500.
- Randazzo, P.A., and R.A. Kahn. 1994. GTP hydrolysis by ADP-ribosylation factor is dependent on both an ADP-ribosylation factor GTPase-activating protein and acid phospholipids. *J Biol Chem.* 269:10758-63.
- Redding, K., C. Holcomb, and R.S. Fuller. 1991. Immunolocalization of Kex2 protease identifies a putative late Golgi compartment in the yeast *Saccharomyces cerevisiae*. *J Cell Biol.* 113:527-38.
- Rein, U., U. Andag, R. Duden, H.D. Schmitt, and A. Spang. 2002. ARF-GAP-mediated interaction between the ER-Golgi v-SNAREs and the COPI coat. *J Cell Biol.* 157:395-404.
- Rice, L.M., and A.T. Brunger. 1999. Crystal structure of the vesicular transport protein Sec17: implications for SNAP function in SNARE complex disassembly. *Mol Cell.* 4:85-95.
- Richardson, C.J., S. Jones, R.J. Litt, and N. Segev. 1998. GTP hydrolysis is not important for Ypt1 GTPase function in vesicular transport. *Mol Cell Biol.* 18:827-38.
- Rizo, J., and T.C. Sudhof. 2002. Snakes and Munc18 in synaptic vesicle fusion. *Nat Rev Neurosci.* 3:641-53.
- Rose, M.D., L.M. Misra, and J.P. Vogel. 1989. *KAR2*, a karyogamy gene, is the yeast homolog of the mammalian BiP/GRP78 gene. *Cell.* 57:1211-21.
- Rossanese, O.W., J. Soderholm, B.J. Bevis, I.B. Sears, J. O'Connor, E.K. Williamson, and B.S. Glick. 1999. Golgi structure correlates with transitional endoplasmic reticulum organization in *Pichia pastoris* and *Saccharomyces cerevisiae*. *J. Cell Biol.* 145:69-81.
- Rossi, G., A. Salminen, L.M. Rice, A.T. Brunger, P. Brennwald, and J.E. Gerst. 1997. Analysis of a yeast SNARE complex reveals remarkable similarity to the neuronal SNARE complex and a novel function for the C terminus of the SNAP-25 homolog, Sec9: Conserved alpha-helical segments on yeast homologs of the synaptobrevin/VAMP family of v-SNAREs mediate exocytic function. *J Biol Chem.* 272:16610-7.
- Rossi, V., D.K. Banfield, M. Vacca, L.E. Dietrich, C. Ungermann, M. D'Esposito, T. Galli, and F. Filippini. 2004a. Longins and their longin domains: regulated SNAREs and multifunctional SNARE regulators. *Trends Biochem Sci.* 29:682-8.
- Rossi, V., R. Picco, M. Vacca, M. D'Esposito, M. D'Urso, T. Galli, and F. Filippini. 2004b. VAMP subfamilies identified by specific R-SNARE motifs. *Biol Cell.* 96:251-6.
- Rothman, J.E. 1994. Mechanisms of intracellular protein transport. *Nature.* 372:55-63.
- Rothstein, R.J. 1983. One-step gene disruption in yeast. *Methods Enzymol.* 101:202-11.
- Rowe, T., C. Dascher, S. Bannykh, H. Plutner, and W.E. Balch. 1998. Role of vesicle-associated syntaxin 5 in the assembly of pre-Golgi intermediates. *Science.* 279:696-700.
- Sacher, M., J. Barrowman, D. Schieltz, J.R. Yates, 3rd, and S. Ferro-Novick. 2000. Identification and characterization of five new subunits of TRAPP. *Eur J Cell Biol.* 79:71-80.
- Sacher, M., J. Barrowman, W. Wang, J. Horecka, Y. Zhang, M. Pypaert, and S. Ferro-Novick. 2001. TRAPP I implicated in the specificity of tethering in ER-to-Golgi transport. *Mol Cell.* 7:433-42.

- Sacher, M., Y. Jiang, J. Barrowman, A. Scarpa, J. Burston, L. Zhang, D. Schieltz, J.R. Yates, 3rd, H. Abelsonich, and S. Ferro-Novick. 1998. TRAPP, a highly conserved novel complex on the *cis*-Golgi that mediates vesicle docking and fusion. *Embo J.* 17:2494-503.
- Sacher, M., S. Stone, and S. Ferro-Novick. 1997. The synaptobrevin-related domains of Bos1p and Sec22p bind to the syntaxin-like region of Sed5p. *J Biol Chem.* 272:17134-8.
- Saifee, O., L. Wei, and M.L. Nonet. 1998. The *Caenorhabditis elegans* unc-64 locus encodes a syntaxin that interacts genetically with synaptobrevin. *Mol Biol Cell.* 9:1235-52.
- Salama, N.R., J.S. Chuang, and R.W. Schekman. 1997. Sec31 encodes an essential component of the COPII coat required for transport vesicle budding from the endoplasmic reticulum. *Mol Biol Cell.* 8:205-17.
- Salama, N.R., T. Yeung, and R.W. Schekman. 1993. The Sec13p complex and reconstitution of vesicle budding from the ER with purified cytosolic proteins. *Embo J.* 12:4073-82.
- Salinovich, O., and R.C. Montelaro. 1986. Reversible staining and peptide mapping of proteins transferred to nitrocellulose after separation by sodium dodecylsulfate-polyacrylamide gel electrophoresis. *Anal Biochem.* 156:341-7.
- Salminen, A., and P.J. Novick. 1987. A ras-like protein is required for a post-Golgi event in yeast secretion. *Cell.* 49:527-38.
- Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sapperstein, S.K., D.M. Walter, A.R. Grosvenor, J.E. Heuser, and M.G. Waters. 1995. p115 is a general vesicular transport factor related to the yeast endoplasmic reticulum to Golgi transport factor Uso1p. *Proc Natl Acad Sci U S A.* 92:522-6.
- Sato, K., and A. Nakano. 2005. Dissection of COPII subunit-cargo assembly and disassembly kinetics during Sar1p-GTP hydrolysis. *Nat Struct Mol Biol.* 12:167-174.
- Sato, M., K. Sato, and A. Nakano. 1996. Endoplasmic reticulum localization of Sec12p is achieved by two mechanisms: Rer1p-dependent retrieval that requires the transmembrane domain and Rer1p-independent retention that involves the cytoplasmic domain. *J. Cell Biol.* 134:279-293.
- Sato, T.K., T. Darsow, and S.D. Emr. 1998. Vam7p, a SNAP-25-like molecule, and Vam3p, a syntaxin homolog, function together in yeast vacuolar protein trafficking. *Mol Cell Biol.* 18:5308-19.
- Sato, T.K., P. Rehling, M.R. Peterson, and S.D. Emr. 2000. Class C Vps protein complex regulates vacuolar SNARE pairing and is required for vesicle docking/fusion. *Mol Cell.* 6:661-71.
- Scales, S.J., Y.A. Chen, B.Y. Yoo, S.M. Patel, Y.C. Doung, and R.H. Scheller. 2000. SNAREs contribute to the specificity of membrane fusion. *Neuron.* 26:457-64.
- Scales, S.J., B.Y. Yoo, and R.H. Scheller. 2001. The ionic layer is required for efficient dissociation of the SNARE complex by alpha-SNAP and NSF. *Proc Natl Acad Sci U S A.* 98:14262-14267.
- Schimmoller, F., I. Simon, and S.R. Pfeffer. 1998. Rab GTPases, directors of vesicle docking. *J Biol Chem.* 273:22161-4.
- Schröder, S., F. Schimmoller, B. Singer-Kruger, and H. Riezman. 1995. The Golgi-localization of yeast Emp47p depends on its di-lysine motif but is not affected by the ret1-1 mutation in alpha-COP. *J Cell Biol.* 131:895-912.
- Schuyler, S.C., and D. Pellman. 2002. Analysis of the size and shape of protein complexes from yeast. *Methods Enzymol.* 351:150-68.
- Seals, D.F., G. Eitzen, N. Margolis, W.T. Wickner, and A. Price. 2000. A Ypt/Rab effector complex containing the Sec1 homolog Vps33p is required for homotypic vacuole fusion. *Proc Natl Acad Sci U S A.* 97:9402-7.

- Seeburg, P.H., J. Shine, J.A. Martial, A. Ullrich, J.D. Baxter, and H.M. Goodman. 1977. Nucleotide sequence of part of the gene for human chorionic somatomammotropin: purification of DNA complementary to predominant mRNA species. *Cell.* 12:157-65.
- Seemann, J., E.J. Jokitalo, and G. Warren. 2000. The role of the tethering proteins p115 and GM130 in transport through the Golgi apparatus in vivo. *Mol Biol Cell.* 11:635-45.
- Segev, N. 2001. Ypt and Rab GTPases: insight into functions through novel interactions. *Curr Opin Cell Biol.* 13:500-11.
- Semenza, J.C., K.G. Hardwick, N. Dean, and H.R. Pelham. 1990. *ERD2*, a yeast gene required for the receptor-mediated retrieval of luminal ER proteins from the secretory pathway. *Cell.* 61:1349-57.
- Serafini, T., L. Orci, M. Amherdt, M. Brunner, R.A. Kahn, and J.E. Rothman. 1991. ADP-ribosylation factor is a subunit of the coat of Golgi-derived COP-coated vesicles: a novel role for a GTP-binding protein. *Cell.* 67:239-53.
- Sevrioukov, E.A., J.P. He, N. Moghrabi, A. Sunio, and H. Kramer. 1999. A role for the deep orange and carnation eye color genes in lysosomal delivery in *Drosophila*. *Mol Cell.* 4:479-86.
- Sharp, P.A., B. Sugden, and J. Sambrook. 1973. Detection of two restriction endonuclease activities in *Haemophilus parainfluenzae* using analytical agarose-ethidium bromide electrophoresis. *Biochemistry.* 12:3055-63.
- Sherman, F. 1991. Getting started with yeast. *Methods Enzymol.* 194:3-20.
- Sherman, F., and J. Hicks. 1991. Micromanipulation and dissection of asci. *Methods Enzymol.* 194:21-37.
- Shorter, J., M.B. Beard, J. Seemann, A.B. Dirac-Svejstrup, and G. Warren. 2002. Sequential tethering of Golgins and catalysis of SNAREpin assembly by the vesicle-tethering protein p115. *J Cell Biol.* 157:45-62.
- Sikorski, R.S., and J.D. Boeke. 1991. *In vitro* mutagenesis and plasmid shuffling: from clone gene to mutant yeast. *Methods Enzymol.* 194:302-318.
- Sikorski, R.S., and P. Hieter. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics.* 122:19-27.
- Simons, K., and M. Zerial. 1993. Rab proteins and the road maps for intracellular transport. *Neuron.* 11:789-99.
- Singer-Kruger, B., H. Stenmark, A. Dusterhoff, P. Philippse, J.S. Yoo, D. Gallwitz, and M. Zerial. 1994. Role of three rab5-like GTPases, Ypt51p, Ypt52p, and Ypt53p, in the endocytic and vacuolar protein sorting pathways of yeast. *J Cell Biol.* 125:283-98.
- Siniossoglou, S., S.Y. Peak-Chew, and H.R. Pelham. 2000. Ric1p and Rgp1p form a complex that catalyses nucleotide exchange on Ypt6p. *Embo J.* 19:4885-94.
- Siniossoglou, S., and H.R.B. Pelham. 2002. Vps51p Links the VFT Complex to the SNARE Tlg1p. *J Biol. Chem.* 277:48318-48324.
- Skehel, J.J., and D.C. Wiley. 1998. Coiled coils in both intracellular vesicle and viral membrane fusion. *Cell.* 95:871-4.
- Søgaard, M., K. Tani, R.R. Ye, S. Geromanos, P. Tempst, T. Kirchhausen, J.E. Rothman, and T. Söllner. 1994. A rab protein is required for the assembly of SNARE complexes in the docking of transport vesicles. *Cell.* 78:937-48.
- Söllner, T., M.K. Bennett, S.W. Whiteheart, R.H. Scheller, and J.E. Rothman. 1993a. A protein assembly-disassembly pathway in vitro that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell.* 75:409-18.
- Söllner, T., S.W. Whiteheart, M. Brunner, H. Erdjument-Bromage, S. Geromanos, P. Tempst, and J.E. Rothman. 1993b. SNAP receptors implicated in vesicle targeting and fusion. *Nature.* 362:318-324.

- Sönnichsen, B., M. Lowe, T. Levine, E. Jamsa, B. Dirac-Svejstrup, and G. Warren. 1998. A role for giantin in docking COPI vesicles to Golgi membranes. *J Cell Biol.* 140:1013-21.
- Spang, A., and R. Schekman. 1998. Reconstitution of retrograde transport from the Golgi to the ER in vitro. *J Cell Biol.* 143:589-99.
- Spelbrink, R.G., and S.F. Nothwehr. 1999. The yeast GRD20 gene is required for protein sorting in the trans-Golgi network/endosomal system and for polarization of the actin cytoskeleton. *Mol. Biol. Cell.* 10:4263-4281.
- Springer, S., and R. Schekman. 1998. Nucleation of COPII vesicular coat complex by endoplasmic reticulum to Golgi vesicle SNAREs. *Science.* 281:698-700.
- Stearns, T., R.A. Kahn, D. Botstein, and M.A. Hoyt. 1990. ADP ribosylation factor is an essential protein in *Saccharomyces cerevisiae* and is encoded by two genes. *Mol Cell Biol.* 10:6690-9.
- Stegmann, T. 2000. Membrane fusion mechanisms: the influenza hemagglutinin paradigm and its implications for intracellular fusion. *Traffic.* 1:598-604.
- Stenbeck, G., R. Schreiner, D. Herrmann, S. Auerbach, F. Lottspeich, J.E. Rothman, and F.T. Wieland. 1992. Gamma-COP, a coat subunit of non-clathrin-coated vesicles with homology to Sec21p. *FEBS Lett.* 314:195-8.
- Stevens, T., B. Esmon, and R. Schekman. 1982. Early stages in the yeast secretory pathway are required for transport of carboxypeptidase Y to the vacuole. *Cell.* 30:439-48.
- Stone, S., M. Sacher, Y. Mao, C. Carr, P. Lyons, A.M. Quinn, and S. Ferro-Novick. 1997. Bet1p activates the v-SNARE Bos1p. *Mol Biol Cell.* 8:1175-81.
- Strom, M., P. Vollmer, T.J. Tan, and D. Gallwitz. 1993. A yeast GTPase-activating protein that interacts specifically with a member of the Ypt/Rab family. *Nature.* 361:736-9.
- Sutton, R.B., D. Fasshauer, R. Jahn, and A.T. Brunger. 1998. Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature.* 395:347-53.
- Suvorova, E.S., R. Duden, and V.V. Lupashin. 2002. The Sec34/Sec35p complex, a Ypt1p effector required for retrograde intra-Golgi trafficking, interacts with Golgi SNAREs and COPI vesicle coat proteins. *J. Cell Biol.* 157:631-643.
- Sweet, D.J., and H.R. Pelham. 1992. The *Saccharomyces cerevisiae SEC20* gene encodes a membrane glycoprotein which is sorted by the HDEL retrieval system. *Embo J.* 11:423-32.
- Sweet, D.J., and H.R. Pelham. 1993. The *TIP1* gene of *Saccharomyces cerevisiae* encodes an 80 kDa cytoplasmic protein that interacts with the cytoplasmic domain of Sec20p. *Embo J.* 12:2831-40.
- Tai, G., L. Lu, T.L. Wang, B.L. Tang, B. Goud, L. Johannes, and W. Hong. 2004. Participation of the Syntaxin 5/Ykt6/GS28/GS15 SNARE complex in transport from the early/recycling endosome to the trans-Golgi network. *Mol. Biol. Cell.* 15:4011-4022.
- Tai, W.C., and D.K. Banfield. 2001. AtBS14a and AtBS14b, two Bet1/Sft1-like SNAREs from *Arabidopsis thaliana* that complement mutations in the yeast *SFT1* gene. *FEBS Lett.* 500:177-82.
- Tall, G.G., H. Hama, D.B. DeWald, and B.F. Horazdovsky. 1999. The phosphatidylinositol 3-phosphate binding protein Vac1p interacts with a Rab GTPase and a Sec1p homologue to facilitate vesicle-mediated vacuolar protein sorting. *Mol Biol Cell.* 10:1873-89.
- Teasdale, R.D., and M.R. Jackson. 1996. Signal-mediated sorting of membrane proteins between the endoplasmic reticulum and the Golgi apparatus. *Annu Rev Cell Dev Biol.* 12:27-54.
- TerBush, D.R., T. Maurice, D. Roth, and P. Novick. 1996. The Exocyst is a multiprotein complex required for exocytosis in *Saccharomyces cerevisiae*. *Embo J.* 15:6483-94.
- Tochio, H., M.M. Tsui, D.K. Banfield, and M. Zhang. 2001. An autoinhibitory mechanism for nonsyntaxin SNARE proteins revealed by the structure of Ykt6p. *Science.* 293:698-702.

- Toonen, R.F., and M. Verhage. 2003. Vesicle trafficking: pleasure and pain from SM genes. *Trends Cell Biol.* 13:177-86.
- Trucco, A., R.S. Polishchuk, O. Martella, A. Di Pentima, A. Fusella, D. Di Giandomenico, E. San Pietro, G.V. Beznoussenko, E.V. Polishchuk, M. Baldassarre, R. Buccione, W.J. Geerts, A.J. Koster, K.N. Burger, A.A. Mironov, and A. Luini. 2004. Secretory traffic triggers the formation of tubular continuities across Golgi sub-compartments. *Nat Cell Biol.* 6:1071-81.
- Tsui, M.M., and D.K. Banfield. 2000. Yeast Golgi SNARE interactions are promiscuous. *J Cell Sci.* 113 (Pt 1):145-52.
- Tsui, M.M., W.C. Tai, and D.K. Banfield. 2001. Selective formation of Sed5p-containing SNARE complexes is mediated by combinatorial binding interactions. *Mol Biol Cell.* 12:521-38.
- Turnbull, A.P., D. Kummel, B. Prinz, C. Holz, J. Schultchen, C. Lang, F.H. Niesen, K.P. Hofmann, H. Delbruck, J. Behlke, E.C. Muller, E. Jarosch, T. Sommer, and U. Heinemann. 2005. Structure of palmitoylated BET3: insights into TRAPP complex assembly and membrane localization. *Embo J.* 24:875-84.
- Uemura, T., T. Ueda, R.L. Ohniwa, A. Nakano, K. Takeyasu, and M.H. Sato. 2004. Systematic analysis of SNARE molecules in Arabidopsis: dissection of the post-Golgi network in plant cells. *Cell Struct Funct.* 29:49-65.
- Ullrich, A., J. Shine, J. Chirgwin, R. Pictet, E. Tischer, W.J. Rutter, and H.M. Goodman. 1977. Rat insulin genes: construction of plasmids containing the coding sequences. *Science.* 196:1313-9.
- Ungar, D., and F.M. Hughson. 2003. SNARE protein structure and function. *Annu Rev Cell Dev Biol.* 19:493-517.
- Ungar, D., T. Oka, E.E. Brittle, E. Vasile, V.V. Lupashin, J.E. Chatterton, J.E. Heuser, M. Krieger, and M.G. Waters. 2002. Characterization of a mammalian Golgi-localized protein complex, COG, that is required for normal Golgi morphology and function. *J. Cell Biol.* 157:405-415.
- Ungar, D., T. Oka, E. Vasile, M. Krieger, and F.M. Hughson. 2005. Subunit architecture of the conserved oligomeric Golgi complex. *J. Biol. Chem.* 280:32736-32745.
- Ungermann, C., B.J. Nichols, H.R. Pelham, and W. Wickner. 1998a. A vacuolar v-t-SNARE complex, the predominant form *in vivo* and on isolated vacuoles, is disassembled and activated for docking and fusion. *J Cell Biol.* 140:61-9.
- Ungermann, C., K. Sato, and W. Wickner. 1998b. Defining the functions of *trans*-SNARE pairs. *Nature.* 396:543-8.
- VanRheenen, S.M., X. Cao, V.V. Lupashin, C. Barlowe, and M. Gerard Waters. 1998. Sec35p, a novel peripheral membrane protein, is required for ER to Golgi vesicle docking. *J. Cell Biol.* 141:1107-1119.
- VanRheenen, S.M., X. Cao, S.K. Sapperstein, E.C. Chiang, V.V. Lupashin, C. Barlowe, and M.G. Waters. 1999. Sec34p, a protein required for vesicle tethering to the yeast Golgi apparatus, is in a complex with Sec35p. *J. Cell Biol.* 147:729-742.
- Vanrheenen, S.M., B.A. Reilly, S.J. Chamberlain, and M.G. Waters. 2001. Dsl1p, an essential protein required for membrane traffic at the endoplasmic reticulum/Golgi interface in yeast. *Traffic.* 2:212-31.
- Veit, M. 2004. The human SNARE protein Ykt6 mediates its own palmitoylation at C-terminal cysteine residues. *Biochem J.* 384:233-7.
- Veit, M., T.H. Sollner, and J.E. Rothman. 1996. Multiple palmitoylation of synaptotagmin and the t-SNARE SNAP-25. *FEBS Lett.* 385:119-23.
- Verhage, M., A.S. Maia, J.J. Plomp, A.B. Brussaard, J.H. Heeroma, H. Vermeer, R.F. Toonen, R.E. Hammer, T.K. van den Berg, M. Missler, H.J. Geuze, and T.C. Sudhof. 2000. Synaptic assembly of the brain in the absence of neurotransmitter secretion. *Science.* 287:864-9.
- Voets, T., R.F. Toonen, E.C. Brian, H. de Wit, T. Moser, J. Rettig, T.C. Sudhof, E. Neher, and M. Verhage. 2001. Munc18-1 promotes large dense-core vesicle docking. *Neuron.* 31:581-91.

- Vollmer, P., E. Will, D. Scheglmann, M. Strom, and D. Gallwitz. 1999. Primary structure and biochemical characterization of yeast GTPase-activating proteins with substrate preference for the transport GTPase Ypt7p. *Eur J Biochem.* 260:284-90.
- von Mollard, G.F., S.F. Nothwehr, and T.H. Stevens. 1997. The yeast v-SNARE Vti1p mediates two vesicle transport pathways through interactions with the t-SNAREs Sed5p and Pep12p. *J Cell Biol.* 137:1511-24.
- Vovis, G.F., and S. Lacks. 1977. Complementary action of restriction enzymes endo R-DpnI and Endo R-DpnII on bacteriophage f1 DNA. *J Mol Biol.* 115:525-38.
- Wach, A., A. Brachat, R. Poehlmann, and P. Philippsen. 1994. New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast.* 10:1793-1808.
- Walch-Solimena, C., R.N. Collins, and P.J. Novick. 1997. Sec2p mediates nucleotide exchange on Sec4p and is involved in polarized delivery of post-Golgi vesicles. *J Cell Biol.* 137:1495-509.
- Wang, W., M. Sacher, and S. Ferro-Novick. 2000. TRAPP stimulates guanine nucleotide exchange on Ypt1p. *J. Cell Biol.* 151:289-296.
- Wang, Y., I. Dulubova, J. Rizo, and T.C. Sudhof. 2001. Functional analysis of conserved structural elements in yeast syntaxin Vam3p. *J Biol Chem.* 276:28598-605.
- Waters, M.G., D.O. Clary, and J.E. Rothman. 1992. A novel 115-kD peripheral membrane protein is required for intercisternal transport in the Golgi stack. *J Cell Biol.* 118:1015-26.
- Waters, M.G., and F.M. Hughson. 2000. Membrane tethering and fusion in the secretory and endocytic pathways. *Traffic.* 1:588-97.
- Waters, M.G., and S.R. Pfeffer. 1999. Membrane tethering in intracellular transport. *Curr Opin Cell Biol.* 11:453-9.
- Watson, P., and D.J. Stephens. 2005. ER-to-Golgi transport: form and formation of vesicular and tubular carriers. *Biochim Biophys Acta.* 1744:304-15.
- Weber, T., B.V. Zemelman, J.A. McNew, B. Westermann, M. Gmachl, F. Parlati, T.H. Sollner, and J.E. Rothman. 1998. SNAREpins: minimal machinery for membrane fusion. *Cell.* 92:759-72.
- Weimbs, T., S.H. Low, S.J. Chapin, K.E. Mostov, P. Bucher, and K. Hofmann. 1997. A conserved domain is present in different families of vesicular fusion proteins: a new superfamily. *Proc Natl Acad Sci U S A.* 94:3046-51.
- Weimbs, T., K. Mostov, S.H. Low, and K. Hofmann. 1998. A model for structural similarity between different SNARE complexes based on sequence relationships. *Trends Cell Biol.* 8:260-2.
- Weiss, M., and T. Nilsson. 2000. Protein sorting in the Golgi apparatus: a consequence of maturation and triggered sorting. *FEBS Lett.* 486:2-9.
- Whyte, J.R., and S. Munro. 2001. The Sec34/35 Golgi transport complex is related to the exocyst, defining a family of complexes involved in multiple steps of membrane traffic. *Dev Cell.* 1:527-37.
- Whyte, J.R., and S. Munro. 2002. Vesicle tethering complexes in membrane traffic. *J Cell Sci.* 115:2627-37.
- Wichmann, H., L. Hengst, and D. Gallwitz. 1992. Endocytosis in yeast: evidence for the involvement of a small GTP-binding protein (Ypt7p). *Cell.* 71:1131-42.
- Wickner, W., and A. Haas. 2000. Yeast homotypic vacuole fusion: a window on organelle trafficking mechanisms. *Annu Rev Biochem.* 69:247-75.
- Wimmer, C., T.M. Hohl, C.A. Hughes, S.A. Muller, T.H. Sollner, A. Engel, and J.E. Rothman. 2001. Molecular mass, stoichiometry, and assembly of 20 S particles. *J. Biol. Chem.* 276:29091-29097.
- Wu, S.K., K. Zeng, I.A. Wilson, and W.E. Balch. 1996. Structural insights into the function of the Rab GDI superfamily. *Trends Biochem Sci.* 21:472-6.

- Wurmser, A.E., T.K. Sato, and S.D. Emr. 2000. New component of the vacuolar class C-Vps complex couples nucleotide exchange on the Ypt7 GTPase to SNARE-dependent docking and fusion. *J Cell Biol.* 151:551-62.
- Xu, Y., S. Martin, D.E. James, and W. Hong. 2002. GS15 forms a SNARE complex with syntaxin 5, GS28, and Ykt6 and is implicated in traffic in the early cisternae of the Golgi apparatus. *Mol Biol Cell.* 13:3493-507.
- Yaffe, M.P., and G. Schatz. 1984. Two nuclear mutations that block mitochondrial protein import in yeast. *Proc Natl Acad Sci U S A.* 81:4819-23.
- Yamaguchi, T., I. Dulubova, S.W. Min, X. Chen, J. Rizo, and T.C. Sudhof. 2002. Sly1 binds to Golgi and ER syntaxins via a conserved N-terminal peptide motif. *Dev Cell.* 2:295-305.
- Yamakawa, H., D.H. Seog, K. Yoda, M. Yamasaki, and T. Wakabayashi. 1996. Uso1 protein is a dimer with two globular heads and a long coiled-coil tail. *J Struct Biol.* 116:356-65.
- Yang, B., L. Gonzalez, Jr., R. Prekeris, M. Steegmaier, R.J. Advani, and R.H. Scheller. 1999. SNARE interactions are not selective. Implications for membrane fusion specificity. *J Biol Chem.* 274:5649-53.
- Yang, X., H.T. Matern, and D. Gallwitz. 1998. Specific binding to a novel and essential Golgi membrane protein (Yip1p) functionally links the transport GTPases Ypt1p and Ypt31p. *Embo J.* 17:4954-63.
- Yoshihisa, T., C. Barlowe, and R. Schekman. 1993. Requirement for a GTPase-activating protein in vesicle budding from the endoplasmic reticulum. *Science.* 259:1466-8.
- Zeng, Q., T.T.H. Tran, H.-X. Tan, and W. Hong. 2003. The cytoplasmic domain of Vamp4 and Vamp5 is responsible for their correct subcellular targeting: the N-terminal extension of Vamp4 contains a dominant autonomous targeting signal for the *trans*-Golgi network. *J. Biol. Chem.* 278:23046-23054.
- Zerial, M., and H. McBride. 2001. Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol.* 2:107-17.
- Zhang, T., and W. Hong. 2001. Ykt6 forms a SNARE complex with syntaxin 5, GS28, and Bet1 and participates in a late stage in endoplasmic reticulum-Golgi transport. *J Biol Chem.* 276:27480-7.
- Zhang, T., S.H. Wong, B.L. Tang, Y. Xu, and W. Hong. 1999. Morphological and functional association of Sec22b/ERS-24 with the pre-Golgi intermediate compartment. *Mol Biol Cell.* 10:435-53.
- Zhang, T., S.H. Wong, B.L. Tang, Y. Xu, F. Peter, V.N. Subramaniam, and W. Hong. 1997. The mammalian protein (rbet1) homologous to yeast Bet1p is primarily associated with the pre-Golgi intermediate compartment and is involved in vesicular transport from the endoplasmic reticulum to the Golgi apparatus. *J Cell Biol.* 139:1157-68.
- Zwilling, D. 2005. Characterization of the early endosomal SNARE complex. Doctoral Thesis in the Molecular Biology & Neurosciences Programme. Georg-August-Universität, Göttingen.