

7 Literaturverzeichnis

1. Antognini JF, Schwartz K: Exaggerated anesthetic requirements in the preferentially anesthetized brain. *Anesthesiology* 1993; 79: 1244-9
2. Antognini JF. The relationship among brain, spinal cord and anesthetic requirements. In: *Med Hypotheses* 1997; 48 (1): 83-87
3. Antognini JF, Carstens E, Buzin V. Isofluran Depresses Motoneuron Excitability by a Direct Spinal Action: An F-Wave Study. In: *Anesth Analg* 1999; 88: 681-685
4. Billard V, Gambus PL, Chamoun N, Stanski DR, Shafer SL. A comparison of spectral edge, delta power and Bispectralindex as EES measures of Alfentanil, Propofol and Midazolam drug effect. *Clin Pharmacol Ther*, 1997; 45-58
5. Bischoff P, Kochs E, Droese D, Meyer-Moldenhauer WH, Schulte am Esch J. Topographic – quantitative EEG- analysis of the paradoxical arousal reaction. EEG changes during urologic surgical producere using isoflurane/N₂O anesthesia. *Anästhesist*, 1993; 42 (3): 142-148
6. Bischoff P, Kochs E, Haferkorn D, Schulte am Esch J. Intraoperative EEG changes in relation to the surgical procedure during isofluran- nitrous oxide anesthesia: Hysterectomy versus mastectomy. *J Clin Anesth*, 1996; 8 (1): 36 43
7. Bosniak ZJ, Seagard JI, Wu A, Kampine JP: The effect of halothane on sympathetic ganglionic transmission. *Anesthesiology* 1982; 57: 473-79
8. Borges M, Antognini JF. Does the brain influence somatic responses to noxious stimuli during isofluran anesthesia. In: *Anesthesiology*, 1994; 81:1511-1515
9. Brigham, E.O. 1974, *The Fast Fourier Transform* (Englewood Cliffs, NJ: Prentice-Hall Brigham, E.O. 1974. Chapter 13
10. Clark DL, Rosner BS. Neurophysiologic effects of general anaesthetics: The electroencephalogram and sensory evoked responses in man. *Anesthesiology*, 1973. 38: 564-582
11. Cooley JW et al., 1965; Cooley, J. W., R. L. Tukey, "An Algorithm for the Machine Calculation of Complex Fourier Series", *Mathematics of Computation* (1965), Vol. 19, No. 90, pp. 297-301

12. Cooper R, Osselton JW, Shaw J. Elektroenzephalographie. Stuttgart- New York: Fischer, 1978
13. Coulthard P, Rood JP. Midazolam and somatosensory evoked potentials. Br J Oral Maxillofac Surg, 1993;31 (1): 28-31
14. De Jong Rh, Robles R, Corbin RW, Nace RA. Effect of inhalation anesthetics on monosynaptic and polynaptic transmission in the spinal cord. In: J Pharmacol Exp Ther, 1968; 162; 326-330
15. De Jong RH, Hershey WN, Wagman IH: Measurement of a spinal reflex response (H-reflex) during general anesthesia in man. Association between reflex depression and muscular relaxation. Anesthesiology 1967; 28: 382-9
16. Doi M, Gajraj RJ, Mantzaridis H, Kenny GN. Relationship between calculated blood concentration of Propofol and electrophysiological variables during emergence from anaesthesia : comparison of bispectralindex , spectral edge frequency, median frequency and auditory evoked potential index. In: Br J Anaest 1997; 78: 180-184
17. Doi M, Gajraj RJ, Mantzaridis H, Kenny GN: Prediction of movement at laryngeal mask airway insertion: comparison of auditory evoked potential index, bispectralindex, spectral edge frequency and median frequency. Br.J.Anaesth. 1999; 82: 203-7
18. Dutton RC, Smith WD, Smith NT: EEG Predicts movement response to surgical stimuli during general anesthesia with combinations of isoflurane, 70% N2O, and fentanyl. J.Clin.Monit. 1996; 12: 127-39
19. Dwyer RC, Rampil IJ, Eger EI II, Benett HL. The electroencephalogram does not predict depth of isofluran anesthesia. In: Anesthesiology 1994; 81: 403-409
20. Eger EI II, Saidman LJ, Brandstater B. Minimum alveolar anesthetic concentration: A Standard of anesthetic potency. In: Anesthesiology,1965; 26:756-763
21. Entholzer E, Schneck HJ, Hargasser S, Hipp R, Tempel G. Electroencephalographic demonstration of central nervous system effects of different premedication regimes. Anästhesist, 1994; 43 (7): 431-440
22. Fischer MA. AAEM Minimonography 13. H-reflexes and F-waves: Physiology and clinical indications. In: Muscle and Nerv 1992; 15: 1223-1233
23. Frank NB, Lieb WR. Molecular and cellular mechanisms of general anesthesia. Nature 1994;367 (6464): 607- 614

24. Freund FG, Martin WG, Hornbein TF. The H-Reflex as a measure of anesthetic potency in man. In : Anesthesiology 1969; 30: 642-647
25. Glass PS, Bloom M, Kearse L, Rosow C, Sebel P, Manberg P. Bispectral analysis measures sedation and memory effects of Propofol, Midazolam, isofluran and Alfentanil in healthy volunteers. Anesthesiology, 1997; 86: 836-847
26. Gravenstein MA, Sasse F, Hogan K. Effects of hypocapnia on canine spinal, subcortical, and cortical somatosensory- evoked potentials during isoflurane anesthesia. J clin Monit, 1992; 8 (2): 126- 130
27. Hochhaus G, Barth J, al-Fayoumi S, et al. Pharmacokinetics and pharmacodynamics of dexamethasone sodium-m-sulfobenzoate (DS) after intravenous and intramuscular administration: a comparison with dexamethasone phosphate (DP). J Clin Pharmacol (United States), Apr 2001, 41(4) p425-34
28. Hoffmann P. Über die Beziehung der Sehnenreflexe zur willkürlichen Bewegung und zum Tonus. Z. Biol ,1918,68: 351-370
29. Hoffmann P. Untersuchung über die Eigenreflexe (Sehnenreflexe) menschlicher Muskeln. 1922 Springer: Berlin .
30. Hoffmann P , Keller CJ. Über gleichzeitige willkürliche und künstliche Reizung von Nerven. Z. Biol., 1928, 87: 527- 536
31. Hoffmann P. Die physiologischen Eigenschaften der Eigenreflexe. Ergeb.Physiol., 1934, 36,15 108
32. Hoffmann P. Die Aufklärung der Wirkung des Jendrassikschen Handgriffes durch die Arbeiten von Sommer und Kuffler. Dtsch Z. Nervenheilk, 1951, 166: 60-64
33. Hwang IS Assessment of soleus motoneuronal excitability using the joint angle dependent H reflex in humans.J Electromyogr Kinesiol (England), Oct 2002, 12(5) p361-6
34. Jasper HH. The Twenty Electrode System of the international Federation.
35. Johansen JW, Sebel PS. Development and clinical application of electroencephalographic bispectrum monitoring. In Anesthesiology 2000; 93: 1336-1344

36. Johnsen CB, Taylor PM. Comparison of the effects of halothane, isofluran and methoxyfluran on the electroencephalogram of the horse. *Br J Anaesth* 1998; 81 (5): 748-753
37. Kano T Shimoji K. The effects of ketamine and neuroleptanalgesia on the evoked electrospinogram and electromyogram in man. *Anesthesiology* 40(3): 234-240, 1974
38. Kato T, Suzuki A, Ikeda K: Electroencephalographic derivatives as a tool for predicting the depth of sedation and anesthesia induced by sevoflurane. *Anesthesiology* 1998; 88: 642-50
39. Kennealy JA, Penovich PE, Moore- Nease SE. EEG and spectral analysis in acute hyperventilation. *Electroencephalogr Clin Neurophysiol.*, 1986; 63 (2): 98- 106
40. Kearse, LA, Manberg, P, Chamoun, N, de Bros, F, Zaslavsky A Bispectral Analysis of the Electroencephalogram Correlates with Patient Movement to Skin Incision During Propofol/Nitrous Oxide. *Anesthesiology*, 1994, 81: 1365-1370
41. Kendig JJ, Savola MK, Woodley SJ, et al. Alpha 2-adrenoceptors inhibit a nociceptive response in neonatal rat spinal cord. *Eur J Pharmacol (Netherlands)*, Jan 10 1991, 192(2) p293-300
42. Kerz T , Hennes HJ, Decq P, Filipetti P, Duvaldestin P. Effects of propofol on h-Reflex in humans. *Anesthesiology*, 2001; 94: 32-37
43. Kimura J. The H-Reflex and other late responses, *Electrodiagnosis in Diseases of Nerve and Muscle. Principle and Practice*; Philadelphia, FA Davis 1983: 379- 398
44. King BS, Rampil IJ: Anesthetic depression of spinal Motoneurons may contribute to lack of movement in response to noxious stimuli. In: *Anesthesiology* 1994; 81: 1484-1492
45. Klass DW, Daly DD. *Klinische Enzephalographie*. Stuttgart- New York: Fischer, 1984
46. Kochs E, Kalkman CJ, Thornton C, Newton D, Bischoff P, Kuppe H, Abke J, Konecny E, Nahm W, Stockmanns G: Middle latency auditory evoked responses and electroencephalographic derived variables do not predict movement to noxious stimulation during 1 minimum alveolar anesthetic

- concentration isoflurane/nitrous oxide anesthesia. *Anesth. Analg.* 1999; 88: 1412-7
47. Koitabashi T, Ochiai R, Takeda J, Fukushima K. Quantitative analysis of electroencephalographic (EEG) activity during Sevofluran anesthesia. *Masui*, 1992 1946-1950
 48. Kröll W. Narkosetiefe. In: List WF, Metzler H, Pasch T, Hrsg.. *Monitoring in Anästhesie und Intensivmedizin*. Berlin: Springer; 1995: 508-521
 49. Leis AA, Zhou HH, Mehta M, Harkey HL III, Paske WC. Behaviour of the H-Reflex in humans following mechanical perturbation or injury to rostral spinal cord. In: *Muscle and Nerve*, 1996; 19: 1373-1382
 50. Levy WJ, Shapiro HM, Maruchak G, Meathe E. Automated EEG processing for intraoperativ monitoring. *Anesthesiology* 1980; 53 (3): 223-236
 51. Liu J , Singh H, White PF: Electroencephalogram bispectral predicts the depth of midazolam-induced sedation. *Anesthesiology* 1996; 84:64-69
 52. Maccaebec PJ, Levine DB, Pinkhasov EI, Cracco RQ, T Sairis P. Evoked potentials recorded from scalp and spinous process during spinal column surgery. *Electroencephalograph Clin Neurophysiol*, 1983 a; 56 (6): 569-582
 53. Maccaebec PJ, Pinkhasov EI, Cracco RQ. Short latency evoked potential to median nerve stimulation: Effect of low frequency filters. *Electroencephalogr Clin Neurophysiol*, 1983 b; 55 (1): 34-44
 54. Madej TH, Paasuke RT. Anesthetic premedikation: aims, assessment and methods. *Can J Anaesth*, 1987; 34 (3): 259-273
 55. Magladery JW, Porter WE, Park AM, Teasdall RD. Electrophysiological studies of nerve and reflex activity in normal man. Identification of certain reflexes in the electromyogram and the conduction velocity of peripheral nerve fibres. In: *Bull Johns Hopkins Hosp* 1950; 86: 265-29
 56. Manberg PJ, Zratek D, Kovitch L, Christman L. Awareness during anesthesia with BIS monitoring. *Anesthesiology*, ASA Annual Meeting, October 14-18, 2000 San Francisco A 1371, 2000
 57. Mavroudakis N , Vandesteene A, Brunko E, Defevrimont M, Zegers de Beyl D: Spinal and brain-stem SEPs and H-Reflex during enflurane anestheasia. *Electroencephalogr Clin Neurophysiol* 1994; 2:82-5

58. Mayer RF and Mosser RS. Maturation of Human Reflexes. Edited by Desmedt JE. New Developments in Elektromyography and clinical Neurophysiology 3, 294-307. 1973 Basel Karger
59. Meibohm B, Derendorf H Basic concepts of pharmacokinetic/ pharmacodynamic (PK/PD) modelling. Int J Clin Pharmacol Ther (Germany), Oct 1997, 35(10) p401-13
60. Nagashima R, Levy G Effect of perfusion rate and distribution factors on drug elimination kinetics in a perfused organ system. J Pharm Sci (United States), Nov 1968, 57(11) p1991-3
61. Neigh JL, Garman JK, Harp JR. The Electroencephalographic pattern during anesthesia with ethrane: Effects of depth of anesthesia, PaCO₂, and nitrous oxide. Anesthesiology, 1971: 35 (5). 482-487
62. O'Connor MF, Daves SM, Tung A, Cook RI, Thisted R, Apfelbaum J. BIS monitoring to prevent awareness during general anesthesia. In: Anesthesiology 2001; 94: 520-522
63. Osawa M, Shingun K, Murakawa M, Adachi T, Kurata J, Seo N, Murayama T, Nakao S, Mori K. Effects of sevofluran on central nervous system electrical activity in cats. Anesth. Analg, 1994;79 (1): 52-57
64. Pereon Y, Bernard JM, Nguyen The Tich S, et al. The effects of desflurane on the nervous system: from spinal cord to muscles. Anesth Analg (United States), Aug 1999, 89(2) p490-5
65. Perine DL, Fontana JL. Use of BIS monitor does not decrease wake up recovery room times. In: Anesthesiology ASA Annual Meeting, October 14-18, 2000 San Francisco A309, 2000
66. Pestelak-Price C. Das internationale 10-20-System zur Elektrodenplatzierung: Begründung, praktische Anwendung zu den Meßschritten und Hinweise zum Setzen der Elektroden. EEG-Labor 1983; 5: 49-72
67. Pichelmayr I, Jeck-Thole S. EEG-Leitfaden für Anästhesisten. Stuttgart: Thieme, 1983
68. Pocock G, Richards CD: Cellular mechanisms in general anesthesia. Br J Anaesth 1991; 66: 116-28
69. Proakis JG et al., 1992, Rader CM, Ling F, Nikias CL: Signal analysis with higher order spectra, Advanced Digital Signal Processing. New York. Macmillan, 1992, pp 550-89

70. Quasha AL, Eger EI II, Tinker JH. Determination and applications of MAC. In: *Anesthesiology* 1980; 53: 315-334
71. Rampil IJ, Sasse F, Smith NT. Spectral edge frequency – a new correlate of anaesthetic depth. *Anesthesiology*, 1980; 53:12
72. Rampil IJ, Holzer JA, Quest DO. Prognostic value of computerized EEG analysis during carotid endarterectomy. *Anesth Anal*, 1983; 62 (2): 186-192
73. Rampil IJ, Matteo RS. Changes in EEG spectral edge frequency correlate with the haemodynamic response to laryngoscopy and intubation. *Anesthesiology*, 1987; 67 (1): 139-142
74. Rampil IJ, Laster MJ. No correlation between quantitative electroencephalographic measurements and movement response to noxious stimuli during isofluran anesthesia in rats. In *Anesthesiology*, 1992; 77: 920-925
75. Rampil IJ, King BS. Volatile anesthetics depress spinal motor neurons. In: *Anesthesiology* 1996; 85: 129-134
76. Rampil IJ. A primer for EEG signal processing in anesthesia. In: *Anesthesiology* 1998; 980-1002
77. Rampil IJ, Mason P, Sing H. Anesthetic potency (MAC) is independent forebrain structures in the rat. In: *Anesthesiology* 1993; 78:707-712
78. Rampil IJ. Anesthetic potency is not altered after hypothermic spinal cord transaction in rats. In: *Anesthesiology*; 1994, 80: 606-610
79. Rehberg B, Bouillon T, Zinserling J, Hoeft A: Comparative pharmacodynamic modelling of the EEG-slowing effect of isoflurane, sevoflurane and desflurane. *Anesthesiology* 1999a; 91:397-405
80. Rehberg B, Bouillon T, Gruenewald M: Comparison of the concentration-dependent effect of sevoflurane on the spinal H-reflex and the EEG in humans. *Acta Anaesthesiol Scand*. 2004 May;48(5):569-76.
81. Röpke H, Schwilden H. Interaction of isofluran and nitrous oxide combinations similar for the median electroencephalographic frequency and clinical anesthesia. In: *Anesthesiology* 1996; 84 (4): 782-788
82. Russ W, Krumholz W, Kling D, Hempelmann G. Erfahrungen mit einem neuen EEG-Spektralanalysator in der Karotischirurgie. *Anästhesist* 1985; 34 (2): 85-90

83. Rundshagen I, Kochs E, Schulte am Esch J. Surgical stimulation increases median nerve somatosensory evoked responses during isoflurane-nitrous oxide anaesthesia. *Br J Anaesth* 1995; 75: 598–602
84. Schwilden H, Stoeckel H. Untersuchungen über verschiedene EEG-Parameter als Indikatoren des Narkosezustandes. *Anästhn Intensiv Notfallmed* 1980; 15 (4): 279-286
85. Schiepatti M. The Hoffmann reflex: A means assessing spinal reflex excitability and its descending control in man. In: *Progr Neurobiol* 1987; 28: 345-376
86. Schwender D, Dauderer M, Klasing S, Finsterer U, Peter K. Power spectral analysis of the electroencephalogram during increasing end- expiratory concentrations of isoflurane, desflurane and sevoflurane. *Anesthesia*, 1998; 53 (4):335- 342
87. Sebel PS, Lang E, Rampil IJ, White PF, Cork R, Jopling M, Smith NT, Glass PSA, Manberg P. A multicenter study of bispectral electroencephalogram analysis for monitoring anesthetic effect. In: *Anesth Analog* 1997; 84: 891-899
88. Sharpe RM, Nathwani D, Pal SK, Brunner MD, Thornton C, Dore CJ, Newton DEF. Auditory evoked response, median frequency and 95% spectral edge during anaesthesia with desflurane and nitrous oxide. *Br J Anaesth* 1997; 78: 282-285
89. Sheiner LB, Stanski DR, Vozeh S, Miller RD, Ham J: Simultaneous modelling of pharmacokinetics and pharmacodynamics: application to d-tubocurarine. *Clin. Pharmacol. Ther.* 1979; 25: 358-71
90. Sleight JW, Donovan J. Comparison of Bispectralindex, 95% spectral edge frequency and approximate entropy of the EEG, with changes in heart rate variability during induction of general anaesthesia. In: *Br J Anaesth* 1999; 82: 659-662
91. Smith NT, Dec-Silver H, Sanford TJ Jr. EEG during high-dose fentanyl-, sufentanil-, or morphine-oxygen in anesthesia. *Anesth Analg* 1984; 63 (4): 386-393
92. Smith NT, Westover CJ Jr., Quinn M. An electroencephalographic comparison of Alfentanil with other narcotics and with thiopental. *J Clin Monit*, 1985; 1: 236-244

93. Soriano SG, Logigian EL, Scott RM, Prael PA, Madsen JR: Nitrous oxide depresses the H-reflex in children with cerebral palsy. *Anesth Analg* 1995; 80: 239-41
94. Stern RC, Towler SC, White PF, Evers AS. Elimination kinetics of sevoflurane and halothane from blood, brain, and adipose tissue in the rat. *Anesth Analg*, 1990; 71 (6): 658- 664
95. Sokoll MD; Hoyt JL, Gergis SD. Studies in muscle rigidity, nitrous oxide, and narcotic analgesic agents. In: *Anesth Analg*, 1972; 51: 16-20
96. Soriano SG, Logigian EL, Scott RM, Prael PA, Madsen JR: Nitrous oxide depresses the H-reflex in children with cerebral palsy. *Anesth Analg* 1995; 80: 239-41
97. Struys M, Versichelen L, Byttebier G, Mortier E, Moerman A, Rolly G. Clinical usefulness of the bispectral index for titrating propofol target effect-site concentration. *Anaesthesia* 1998; 53: 4-12
98. Sundt TM Jr, Sharbrough FW, Piepgras Dg. Correlation of cerebral blood flow and electroencephalographic changes during carotid endarterectomy with results of surgery and hemodynamics of cerebral ischemia. *Mayo Clin Proc* 1981; 56 (9): 533-543
99. Tarnow J. Prämedikation Anästhesiologie und Intensivmedizin, 1985; 26: 174-181
100. Tatsumi K, Hirai K, Furuya H, Okuda T. Effects of sevoflurane on the middle latency auditory evoked response and the electroencephalographic power spectrum. *Anesth Analg*, 1995; 80 (5): 940-943
101. Thornton C, Konieczko K, Jones JG, Jordan C, Dore CJ, Heneghan CP: Effect of surgical stimulation on the auditory evoked response. *Br. J. Anaesth.* 1988; 60: 372-8
102. Van de Worp HB, Kraaier V, Wieneke GH, van Huffelen AC. Quantitative EEG during progressive hypocarbia and hypoxia. Hyperventilation induced EEG changes reconsidered. *Electroencephalogr Clin Neurophysiology*, 1991; 79 (5): 335- 341
103. Vernon JM, Lang E, Sebel PS, Manberg, PJ. Prediction of Movement Using Bispectral EEG Analysis during Propofol/Alfentanil or Isoflurane/ Alfentanil Anesthesia and Analgesia, 1995, 80:780-785

104. Veselis RA, Reinsel R, Alagesan R, Heino R, Bedford RF. The EEG as monitor of midazolam amnesia: changes in power and topography as a function of amnesic state. *Anaesthesiology* , 1991; 74 (5): 866- 874
105. Zhou HH, Mehta M, Leis AA. Spinal cord motoneuron excitability during isofluran and nitrous oxide anesthesia. In: *Anesthesiology* 1997; 86: 302-307
106. Zhou HH, Turndorf H. Hyper- and hypoventilation affects spinal motor neuron excitability during isofluran anesthesia. In: *Anesthesiology* 1998; 87(2): 407-410
107. Zhou HH, Jin TT, Qin B, Turndorf H. Suppression of the spinal cord motoneuron excitability correlates with surgical immobility during isofluran anesthesia. In: *Anesthesiology* 1998; 88(4): 955-961