

## 6. Literaturverzeichnis

- Aldred MA, Huang Y, Liyanarachchi S, Pellegata NS, Gimm O, Jhiang S, Davuluri RV, de la Chapelle A, Eng C. (2004). Papillary and follicular thyroid carcinomas show distinctly different microarray expression profiles and can be distinguished by a minimum of five genes. *J Clin Oncol.* **22**, 3531-9.
- Anderson JM, Stevenson BR, Jesaitis LA, Goodenough DA, Mooseker MS. (1988). Characterization of ZO-1, a protein component of the tight junction from mouse liver and Madin-Darby canine kidney cells. *J Cell Biol.* **106**, 1141-9.
- Anderson JM. (1996). Cell signalling: MAGUK magic. *Curr Biol.* **6**, 382-4.
- Anderson JM. (2001). Molecular structure of tight junctions and their role in epithelial transport. *News Physiol Sci.* **16**, 126-30.
- Andersson K, Hamalainen M, Malmqvist M. (1999). Identification and optimization of regeneration conditions for affinity-based biosensor assays. A multivariate cocktail approach. *Anal Chem.* **71**, 2475-81.
- Ando-Akatsuka Y, Saitou M, Hirase T, Kishi M, Sakakibara A, Itoh M, Yonemura S, Furuse M, Tsukita S. (1996). Interspecies diversity of the occludin sequence: cDNA cloning of human, mouse, dog, and rat-kangaroo homologues. *J Cell Biol.* **133**, 43-7.
- Ando-Akatsuka Y, Yonemura S, Itoh M, Furuse M, Tsukita S. (1999). Differential behavior of E-cadherin and occludin in their colocalization with ZO-1 during the establishment of epithelial cell polarity. *J Cell Physiol.* **179**, 115-25.
- Andreeva AY, Krause E, Muller EC, Blasig IE, Utepbergenov DI. (2001). Protein kinase C regulates the phosphorylation and cellular localization of occludin. *J Biol Chem.* **276**, 38480-6.
- Antonetti DA, Barber AJ, Hollinger LA, Wolpert EB, Gardner TW. (1999). Vascular endothelial growth factor induces rapid phosphorylation of tight junction proteins occludin and zonula occludens 1. A potential mechanism for vascular permeability in diabetic retinopathy and tumors. *J Biol Chem.* **274**, 23463-7.
- Avila-Flores A, Rendon-Huerta E, Moreno J, Islas S, Betanzos A, Robles-Flores M, Gonzalez-Mariscal L. (2001). Tight-junction protein zonula occludens 2 is a target of phosphorylation by protein kinase C. *Biochem J.* **360**, 295-304.
- Balda MS, Anderson JM. (1993). Two classes of tight junctions are revealed by ZO-1 isoforms. *Am J Physiol.* **264**, C918-24.
- Balda MS, Gonzalez-Mariscal L, Matter K, Cereijido M, Anderson JM. (1993). Assembly of the tight junction: the role of diacylglycerol. *J Cell Biol.* **123**, 293-302.
- Balda MS, Anderson JM, Matter K. (1996a). The SH3 domain of the tight junction protein ZO-1 binds to a serine protein kinase that phosphorylates a region C-terminal to this domain. *FEBS Lett.* **399**, 326-32.
- Balda MS, Whitney JA, Flores C, Gonzalez S, Cereijido M, Matter K. (1996b). Functional dissociation of paracellular permeability and transepithelial electrical resistance and disruption of the apical-basolateral intramembrane diffusion barrier by expression of a mutant tight junction membrane protein. *J Cell Biol.* **134**, 1031-49.
- Balda MS, Garrett MD, Matter K. (2003). The ZO-1-associated Y-box factor ZONAB regulates epithelial cell proliferation and cell density. *J Cell Biol.* **160**, 423-32.
- Bamforth SD, Kniesel U, Wolburg H, Engelhardt B, Risau W. (1999). A dominant mutant of occludin disrupts tight junction structure and function. *J Cell Sci.* **112**, 1879-88.
- Baumann G, Frommel C, Sander C. (1989). Polarity as a criterion in protein design. *Protein Eng.* **2**, 329-34.
- Bazzoni G, Martinez-Estrada OM, Orsenigo F, Cordenonsi M, Citi S, Dejana E. (2000). Interaction of junctional adhesion molecule with the tight junction components ZO-1, cingulin, and occludin. *J Biol Chem.* **275**, 20520-6.

- Behlke J, Ristau O. (1997). Molecular mass determination by sedimentation velocity experiments and direct fitting of the concentration profiles. *Biophys J.* **72**, 428-34.
- Ben-Yosef T, Belyantseva IA, Saunders TL, Hughes ED, Kawamoto K, Van Itallie CM, Beyer LA, Halsey K, Gardner DJ, Wilcox ER, Rasmussen J, Anderson JM, Dolan DF, Forge A, Raphael Y, Camper SA, Friedman TB. (2003). Claudin 14 knockout mice, a model for autosomal recessive deafness DFNB29, are deaf due to cochlear hair cell degeneration. *Hum Mol Genet.* **12**, 2049-61.
- Berkes J, Viswanathan VK, Savkovic SD, Hecht G. (2003). Intestinal epithelial responses to enteric pathogens: effects on the tight junction barrier, ion transport, and inflammation. *Gut.* **52**, 439-51.
- Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. (2000). The Protein Data Bank. *Nucleic Acids Res.* **28**, 235-42.
- Blaszczyk J, Li Y, Yan H, Ji X. (2001). Crystal structure of unligated guanylate kinase from yeast reveals GMP-induced conformational changes. *J Mol Biol.* **307**, 247-57.
- Boado RJ, Li JY, Nagaya M, Zhang C, Pardridge WM. (1999). Selective expression of the large neutral amino acid transporter at the blood-brain barrier. *Proc Natl Acad Sci U S A.* **96**, 12079-84.
- Bourne PE, Weissig H. (Eds.) (2003). Structural Bioinformatics. Wiley Liss, Inc. Hoboken, New Jersey, USA.
- Britt DE, Yang DF, Yang DQ, Flanagan D, Callanan H, Lim YP, Lin SH, Hixson DC. (2004). Identification of a novel protein, LYRIC, localized to tight junctions of polarized epithelial cells. *Exp Cell Res.* **300**, 134-48.
- Brown JH, Cohen C, Parry DA. (1996). Heptad breaks in alpha-helical coiled coils: stutters and stammers. *Proteins.* **26**, 134-45.
- Budisa N, Steipe B, Demange P, Eckerskorn C, Kellermann J, Huber R. (1995). High-level biosynthetic substitution of methionine in proteins by its analogs 2-aminohexanoic acid, selenomethionine, telluromethionine and ethionine in Escherichia coli. *Eur J Biochem.* **230**, 788-96.
- Bullions LC, Notterman DA, Chung LS, Levine AJ. (1997). Expression of wild-type alpha-catenin protein in cells with a mutant alpha-catenin gene restores both growth regulation and tumor suppressor activities. *Mol Cell Biol.* **17**, 4501-8.
- Burkhard P, Stetefeld J, Strelkov SV. (2001). Coiled coils: a highly versatile protein folding motif. *Trends Cell Biol.* **11**, 82-8.
- Cereijido M, Shoshani L, Contreras RG. (2000). Molecular physiology and pathophysiology of tight junctions. I. Biogenesis of tight junctions and epithelial polarity. *Am J Physiol Gastrointest Liver Physiol.* **279**, G477-82.
- Chen YH, Yang JT, Martinez HM. (1972). Determination of the secondary structures of proteins by circular dichroism and optical rotatory dispersion. *Biochemistry.* **11**, 4120-31.
- Chen Y, Merzdorf C, Paul DL, Goodenough DA. (1997). COOH terminus of occludin is required for tight junction barrier function in early Xenopus embryos. *J Cell Biol.* **138**, 891-9.
- Chen YH, Lu Q, Goodenough DA, Jeansson B. (2002). Nonreceptor tyrosine kinase c-Yes interacts with occludin during tight junction formation in canine kidney epithelial cells. *Mol Biol Cell.* **13**, 1227-37.
- Claude P. (1978). Morphological factors influencing transepithelial permeability: a model for the resistance of the zonula occludens. *J Membr Biol.* **39**, 219-32.
- Colegio OR, Van Itallie C, Rahner C, Anderson JM. (2003). Claudin extracellular domains determine paracellular charge selectivity and resistance but not tight junction fibril architecture. *Am J Physiol Cell Physiol.* **284**, C1346-54.

- Cordenonsi M, Mazzon E, De Rigo L, Baraldo S, Meggio F, Citi S. (1997). Occludin dephosphorylation in early development of *Xenopus laevis*. *J Cell Sci.* **110**, 3131-9.
- Cordenonsi M, Turco F, D'atri F, Hammar E, Martinucci G, Meggio F, Citi S. (1999a). *Xenopus laevis* occludin. Identification of in vitro phosphorylation sites by protein kinase CK2 and association with cingulin. *Eur J Biochem.* **264**, 374-84.
- Cordenonsi M, D'Atri F, Hammar E, Parry DA, Kendrick-Jones J, Shore D, Citi S. (1999b). Cingulin contains globular and coiled-coil domains and interacts with ZO-1, ZO-2, ZO-3, and myosin. *J Cell Biol.* **147**, 1569-82.
- Coyne CB, Vanhook MK, Gambling TM, Carson JL, Boucher RC, Johnson LG. (2002). Regulation of airway tight junctions by proinflammatory cytokines. *Mol Biol Cell.* **13**, 3218-34.
- Crick FHC. (1953). The packing of  $\alpha$ -helices: simple coiled-coils. *Acta Cryst.* **6**, 689-97.
- D'Atri F, Nadalutti F, Citi S. (2002). Evidence for a functional interaction between cingulin and ZO-1 in cultured cells. *J Biol Chem.* **277**, 27757-64.
- Davies DC. (2002). Blood-brain barrier breakdown in septic encephalopathy and brain tumours. *J Anat.* **200**, 639-46.
- DeMaio L, Chang YS, Gardner TW, Tarbell JM, Antonetti DA. (2001). Shear stress regulates occludin content and phosphorylation. *Am J Physiol Heart Circ Physiol.* **281**, H105-13.
- De Vries HE, Kuiper J, de Boer AG, Van Berkel TJ, Breimer DD. (1997). The blood-brain barrier in neuroinflammatory diseases. *Pharmacol Rev.* **49**, 143-55.
- Diamond JM. (1977). Twenty-first Bowditch lecture. The epithelial junction: bridge, gate, and fence. *Physiologist.* **20**, 10-8.
- Ebnet K, Schulz CU, Meyer Zu Brickwedde MK, Pendl GG, Vestweber D. (2000). Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J Biol Chem.* **275**, 27979-88.
- Epstein CJ, Goldberger RF, Anfinsen CB. (1963). The genetic control of tertiary protein structure: studies with model systems. *Cold Spring Harb Symp Quant Biol* **28**, 439.
- Fairbanks G, Steck TL, Wallach DF. (1971). Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry.* **10**, 2606-17.
- Fanning AS, Jameson BJ, Jesaitis LA, Anderson JM. (1998). The tight junction protein ZO-1 establishes a link between the transmembrane protein occludin and the actin cytoskeleton. *J Biol Chem.* **273**, 29745-53.
- Fanning AS, Ma TY, Anderson JM. (2002). Isolation and functional characterization of the actin-binding region in the tight junction protein ZO-1. *FASEB J.* **16**, 1835-7.
- Fasano A. (2002). Toxins and the gut: role in human disease. *Gut.* **50** Suppl 3:III9-14.
- Fischer D, Barret C, Bryson K, Elofsson A, Godzik A, Jones D, Karplus KJ, Kelley LA, MacCallum RM, Pawowski K, Rost B, Rychlewski L, Sternberg M. (1999). CAFASP-1: critical assessment of fully automated structure prediction methods. *Proteins. Suppl.* **3**, 209-17.
- Fischetti VA, Landau GM, Schmidt JP, Sellers P. (1993). Identifying periodic occurrences of a template with applications to protein structure. *Information Processing Letters.* **45**, 11-18.
- Florian P, Schoneberg T, Schulzke JD, Fromm M, Gitter AH. (2002). Single-cell epithelial defects close rapidly by an actinomyosin purse string mechanism with functional tight junctions. *J Physiol.* **545**, 485-99.
- Frank R. (2002). The SPOT-synthesis technique. Synthetic peptide arrays on membrane supports--principles and applications. *J Immunol Methods.* **267**, 13-26.
- Fujimoto K. (1995). Freeze-fracture replica electron microscopy combined with SDS digestion for cytochemical labeling of integral membrane proteins. Application to the immunogold labeling of intercellular junctional complexes. *J Cell Sci.* **108**, 3443-9.

- Fujita K, Katahira J, Horiguchi Y, Sonoda N, Furuse M, Tsukita S. (2000). Clostridium perfringens enterotoxin binds to the second extracellular loop of claudin-3, a tight junction integral membrane protein. *FEBS Lett.* **476**, 258-61.
- Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S, Tsukita S. (1993). Occludin: a novel integral membrane protein localizing at tight junctions. *J Cell Biol.* **123**, 1777-88.
- Furuse M, Itoh M, Hirase T, Nagafuchi A, Yonemura S, Tsukita S, Tsukita S. (1994). Direct association of occludin with ZO-1 and its possible involvement in the localization of occludin at tight junctions. *J Cell Biol.* **127**, 1617-26.
- Furuse M, Fujita K, Hiiragi T, Fujimoto K, Tsukita S. (1998a). Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol.* **141**, 1539-50.
- Furuse M, Sasaki H, Fujimoto K, Tsukita S. (1998b). A single gene product, claudin-1 or -2, reconstitutes tight junction strands and recruits occludin in fibroblasts. *J Cell Biol.* **143**, 391-401.
- Furuse M, Sasaki H, Tsukita S. (1999). Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J Cell Biol.* **147**, 891-903.
- Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A, Sugitani Y, Noda T, Kubo A, Tsukita S. (2002). Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J Cell Biol.* **156**, 1099-111.
- Garrod DR, Merritt AJ, Nie Z. (2002). Desmosomal cadherins. *Curr Opin Cell Biol.* **14**, 537-45.
- Ghassemifar MR, Sheth B, Papenbrock T, Leese HJ, Houghton FD, Fleming TP. (2002). Occludin TM4(-): an isoform of the tight junction protein present in primates lacking the fourth transmembrane domain. *J Cell Sci.* **115**, 3171-80.
- Giepmans BN. Gap junctions and connexin-interacting proteins. (2004). *Cardiovasc Res.* **62**, 233-45.
- Gilula NB. (1974). Junction between cells. In *Cell Communication* (R.P. Cox, ed.) Wiley, New York.
- Gonzalez L Jr, Woolfson DN, Alber T. (1996). Buried polar residues and structural specificity in the GCN4 leucine zipper. *Nat Struct Biol.* **3**, 1011-8.
- Gonzalez-Mariscal L, Islas S, Contreras RG, Garcia-Villegas MR, Betanzos A, Vega J, Diaz-Quinonez A, Martin-Orozco N, Ortiz-Navarrete V, Cereijido M, Valdes J. (1999). Molecular characterization of the tight junction protein ZO-1 in MDCK cells. *Exp Cell Res.* **248**, 97-109.
- Gonzalez-Mariscal L, Namorado MC, Martin D, Luna J, Alarcon L, Islas S, Valencia L, Muriel P, Ponce L, Reyes JL. (2000a). Tight junction proteins ZO-1, ZO-2, and occludin along isolated renal tubules. *Kidney Int.* **57**, 2386-402.
- Gonzalez-Mariscal L, Betanzos A, Avila-Flores A. (2000b). MAGUK proteins: structure and role in the tight junction. *Semin Cell Dev Biol.* **11**, 315-24.
- Gonzalez-Mariscal L, Avila-Flores A, Betanzos A. (2001). The relationship between structure and function of tight junctions. In Cereijido M, Anderson JM. (Eds.), *Tight junctions*. Boca Raton.
- González-Mariscal L, Betanzos A, Nava P, Jaramillo BE. (2003). Tight junction proteins. *Prog Biophys Mol Biol.* **81**, 1-44.
- Gottardi CJ, Arpin M, Fanning AS, Louvard D. (1996). The junction-associated protein, zonula occludens-1, localizes to the nucleus before the maturation and during the remodeling of cell-cell contacts. *Proc Natl Acad Sci U S A.* **93**, 10779-84.
- Gow A, Southwood CM, Li JS, Pariali M, Riordan GP, Brodie SE, Danias J, Bronstein JM, Kachar B, Lazzarini RA. (1999). CNS myelin and sertoli cell tight junction strands are absent in Osp/claudin-11 null mice. *Cell.* **99**, 649-59.

- Gruber M, Lupas AN. (2003). Historical review: another 50th anniversary--new periodicities in coiled coils. *Trends Biochem Sci.* **28**, 679-85.
- Gumbiner B, Lowenkopf T, Apatira D. (1991). Identification of a 160-kDa polypeptide that binds to the tight junction protein ZO-1. *Proc Natl Acad Sci U S A.* **88**, 3460-4.
- Guo P, Weinstein AM, Weinbaum S. (2003). A dual-pathway ultrastructural model for the tight junction of rat proximal tubule epithelium. *Am J Physiol Renal Physiol.* **285**, F241-57.
- Hamazaki Y, Itoh M, Sasaki H, Furuse M, Tsukita S. (2002). Multi-PDZ domain protein 1 (MUPP1) is concentrated at tight junctions through its possible interaction with claudin-1 and junctional adhesion molecule. *J Biol Chem.* **277**, 455-61.
- Harbury PB, Zhang T, Kim PS, Alber T. (1993). A switch between two-, three-, and four-stranded coiled coils in GCN4 leucine zipper mutants. *Science.* **262**, 1401-7.
- Harhaj NS, Antonetti DA. (2004). Regulation of tight junctions and loss of barrier function in pathophysiology. *Int J Biochem Cell Biol.* **36**, 1206-37.
- Haskins J, Gu L, Wittchen ES, Hibbard J, Stevenson BR. (1998). ZO-3, a novel member of the MAGUK protein family found at the tight junction, interacts with ZO-1 and occludin. *J Cell Biol.* **141**, 199-208.
- Herrenknecht K, Ozawa M, Eckerskorn C, Lottspeich F, Lenter M, Kemler R. (1991). The uvomorulin-anchorage protein alpha-catenin is a vinculin homologue. *Proc Natl Acad Sci U S A.* **88**, 9156-60.
- Hibbs K, Skubitz KM, Pambuccian SE, Casey RC, Burleson KM, Oegema TR Jr, Thiele JJ, Grindle SM, Bliss RL, Skubitz AP. (2004). Differential gene expression in ovarian carcinoma: identification of potential biomarkers. *Am J Pathol.* **165**, 397-414.
- Hicks MR, Holberton DV, Kowalczyk C, Woolfson DN. (1997). Coiled-coil assembly by peptides with non-heptad sequence motifs. *Fold Des.* **2**, 149-58.
- Hirase T, Kawashima S, Wong EY, Ueyama T, Rikitake Y, Tsukita S, Yokoyama M, Staddon JM. (2001). Regulation of tight junction permeability and occludin phosphorylation by Rhoa-p160ROCK-dependent and -independent mechanisms. *J Biol Chem.* **276**, 10423-31.
- Hollande F, Lee DJ, Choquet A, Roche S, Baldwin GS. (2003). Adherens junctions and tight junctions are regulated via different pathways by progastrin in epithelial cells. *J Cell Sci.* **116**, 1187-97.
- Holtzhauer M. (Hrsg.). (1996) Methoden in der Proteinanalytik. Springer Verlag Berlin
- Hough CD, Woods DF, Park S, Bryant PJ. (1997). Organizing a functional junctional complex requires specific domains of the Drosophila MAGUK Discs large. *Genes Dev.* **11**, 3242-53.
- Huber D, Balda MS, Matter K. (2000). Occludin modulates transepithelial migration of neutrophils. *J Biol Chem.* **275**, 5773-8.
- Huber O, Krohn M, Kemler R. (1997). A specific domain in alpha-catenin mediates binding to beta-catenin or plakoglobin. *J Cell Sci.* **110**, 1759-65.
- Hynes RO. (1999). Cell adhesion: old and new questions. *Trends Cell Biol.* **9**, M33-7.
- Ikari A, Hirai N, Shiroma M, Harada H, Sakai H, Hayashi H, Suzuki Y, Degawa M, Takagi K. (2004). Association of paracellin-1 with ZO-1 augments the reabsorption of divalent cations in renal epithelial cells. *J Biol Chem.* **279**, 54826-32.
- Ikenouchi J, Matsuda M, Furuse M, Tsukita S. (2003). Regulation of tight junctions during the epithelium-mesenchyme transition: direct repression of the gene expression of claudins/occludin by Snail. *J Cell Sci.* **116**, 1959-67.
- Imamura Y, Itoh M, Maeno Y, Tsukita S, Nagafuchi A. (1999). Functional domains of alpha-catenin required for the strong state of cadherin based cell adhesion. *J Cell Biol.* **144**, 1311-22.

- Itoh M, Nagafuchi A, Yonemura S, Kitani-Yasuda T, Tsukita S, Tsukita S. (1993). The 220-kD protein colocalizing with cadherins in non-epithelial cells is identical to ZO-1, a tight junction-associated protein in epithelial cells: cDNA cloning and immunoelectron microscopy. *J Cell Biol.* **121**, 491-502.
- Itoh M, Nagafuchi A, Moroi S, Tsukita S. (1997). Involvement of ZO-1 in cadherin-based cell adhesion through its direct binding to alpha-catenin and actin filaments. *J Cell Biol.* **138**, 181-92.
- Itoh M, Morita K, Tsukita S. (1999a). Characterization of ZO-2 as a MAGUK family member associated with tight as well as adherens junctions with a binding affinity to occludin and alpha-catenin. *J Biol Chem.* **274**, 5981-6.
- Itoh M, Furuse M, Morita K, Kubota K, Saitou M, Tsukita S. (1999b). Direct binding of three tight junction-associated MAGUKs, ZO-1, ZO-2, and ZO-3, with the COOH termini of claudins. *J Cell Biol.* **147**, 1351-63.
- Itoh M, Sasaki H, Furuse M, Ozaki H, Kita T, Tsukita S. (2001). Junctional adhesion molecule (JAM) binds to PAR-3: a possible mechanism for the recruitment of PAR-3 to tight junctions. *J Cell Biol.* **154**, 491-7.
- Ivanov AI, Nusrat A, Parkos CA. (2004). Endocytosis of epithelial apical junctional proteins by a clathrin-mediated pathway into a unique storage compartment. *Mol Biol Cell.* **15**, 176-88.
- Izard T, Evans G, Borgon RA, Rush CL, Bricogne G, Bois PR. (2004). Vinculin activation by talin through helical bundle conversion. *Nature.* **427**, 171-5.
- Janek K, Rothmund S, Gast K, Beyermann M, Zipper J, Fabian H, Bienert M, Krause E. (2001). Study of the conformational transition of A beta(1-42) using D-amino acid replacement analogues. *Biochemistry.* **40**, 5457-63.
- Jeanssonne B, Lu Q, Goodenough DA, Chen YH. (2003). Claudin-8 interacts with multi-PDZ domain protein 1 (MUPP1) and reduces paracellular conductance in epithelial cells. *Cell Mol Biol (Noisy-le-grand).* **49**, 13-21.
- Johnson WC Jr. (1990). Protein secondary structure and circular dichroism: a practical guide. *Proteins.* **7**, 205-14.
- Jou TS, Nelson WJ. (1998). Effects of regulated expression of mutant RhoA and Rac1 small GTPases on the development of epithelial (MDCK) cell polarity. *J Cell Biol.* **142**, 85-100.
- Jou TS, Schneeberger EE, Nelson WJ. (1998). Structural and functional regulation of tight junctions by RhoA and Rac1 small GTPases. *J Cell Biol.* **142**, 101-15.
- Karlsson R, Michaelsson A, Mattsson L. (1991). Kinetic analysis of monoklonal antibody-antigen interactions with a new biosensor based analytic system. *J Immunol Methods.* **145**, 229-40.
- Katsube T, Takahisa M, Ueda R, Hashimoto N, Kobayashi M, Togashi S. (1998). Cortactin associates with the cell-cell junction protein ZO-1 in both Drosophila and mouse. *J Biol Chem.* **273**, 29672-7.
- Kevil CG, Oshima T, Alexander B, Coe LL, Alexander JS. (2000). H<sub>2</sub>O<sub>2</sub>-mediated permeability: role of MAPK and occludin. *Am J Physiol Cell Physiol.* **279**, C21-30.
- Kim J, Gye MC, Kim MK. (2004). Role of occludin, a tight junction protein, in blastocoel formation, and in the paracellular permeability and differentiation of trophectoderm in preimplantation mouse embryos. *Mol Cells.* **17**, 248-54.
- Kiuchi-Saishin Y, Gotoh S, Furuse M, Takasuga A, Tano Y, Tsukita S. (2002). Differential expression patterns of claudins, tight junction membrane proteins, in mouse nephron segments. *J Am Soc Nephrol.* **13**, 875-86.
- Kobayashi J, Inai T, Shibata Y. (2002). Formation of tight junction strands by expression of claudin-1 mutants in their ZO-1 binding site in MDCK cells. *Histochem Cell Biol.* **117**, 29-39.

- Kobielak A, Fuchs E. (2004). Alpha-catenin: at the junction of intercellular adhesion and actin dynamics. *Nat Rev Mol Cell Biol.* **5**, 614-25.
- Kojima T, Sawada N, Chiba H, Kokai Y, Yamamoto M, Urban M, Lee GH, Hertzberg EL, Mochizuki Y, Spray DC. (1999). Induction of tight junctions in human connexin 32 (hCx32)-transfected mouse hepatocytes: connexin 32 interacts with occludin. *Biochem Biophys Res Commun.* **266**, 222-9.
- Kojima T, Kokai Y, Chiba H, Osanai M, Kuwahara K, Mori M, Mochizuki Y, Sawada N. (2001). Occludin and claudin-1 concentrate in the midbody of immortalized mouse hepatocytes during cell division. *J Histochem Cytochem.* **49**, 333-40.
- Kojima T, Yamamoto T, Murata M, Chiba H, Kokai Y, Sawada N. (2003). Regulation of the blood-biliary barrier: interaction between gap and tight junctions in hepatocytes. *Med Electron Microsc.* **36**, 157-64.
- Kominsky SL, Vali M, Korz D, Gabig TG, Weitzman SA, Argani P, Sukumar S. (2004). Clostridium perfringens enterotoxin elicits rapid and specific cytolysis of breast carcinoma cells mediated through tight junction proteins claudin 3 and 4. *Am J Pathol.* **164**, 1627-33.
- Koslov ER, Maupin P, Pradhan D, Morrow JS, Rimm DL. (1997). Alpha-catenin can form asymmetric homodimeric complexes and/or heterodimeric complexes with beta-catenin. *J Biol Chem.* **272**, 27301-6.
- Kramer A, Schneider-Mergener J. (1998). Synthesis and screening of peptide libraries on continuous cellulose membrane supports. *Methods Mol Biol.* **87**, 25-39.
- Kuhlendahl S, Spangenberg O, Konrad M, Kim E, Garner CC. (1998). Functional analysis of the guanylate kinase-like domain in the synapse-associated protein SAP97. *Eur J Biochem.* **252**, 305-13.
- Kuwabara H, Kokai Y, Kojima T, Takakuwa R, Mori M, Sawada N. (2001). Occludin regulates actin cytoskeleton in endothelial cells. *Cell Struct Funct.* **26**, 109-16.
- Lacaz-Vieira F, Jaeger MM, Farshori P, Kachar B. (1999). Small synthetic peptides homologous to segments of the first external loop of occludin impair tight junction resealing. *J Membr Biol.* **168**, 289-97.
- Lapierre LA, Tuma PL, Navarre J, Goldenring JR, Anderson JM. (1999). VAP-33 localizes to both an intracellular vesicle population and with occludin at the tight junction. *J Cell Sci.* **112**, 3723-32.
- Laskowski RA, MacArthur MW, Moss DS, Thornton JM. (1993). PROCHECK: a program to check the stereochemical quality of protein structures. *J Appl Cryst.* **26**, 283-291.
- Lottspeich F, Zorbas H. (1998). Bioanalytik. Spektrum Akademischer Verlag Heidelberg Berlin.
- Lui WY, Mruk D, Lee WM, Cheng CY. (2003). Sertoli cell tight junction dynamics: their regulation during spermatogenesis. *Biol Reprod.* **68**, 1087-97.
- Lupas A, Van Dyke M, Stock J. (1991). Predicting coiled coils from protein sequences. *Science.* **252**, 1162-4.
- Lupas A. (1996a). Coiled coils: new structures and new functions. *Trends Biochem Sci.* **21**, 375-82.
- Lupas A. (1996b). Prediction and analysis of coiled-coil structures. *Methods Enzymol.* **266**, 513-25.
- Lupas A. (1997). Predicting coiled-coil regions in proteins. *Curr Opin Struct Biol.* **7**, 388-93.
- Mandel LJ, Bacallao R, Zampighi G. (1993). Uncoupling of the molecular 'fence' and paracellular 'gate' functions in epithelial tight junctions. *Nature.* **361**, 552-5.
- Mankertz J, Tavalali S, Schmitz H, Mankertz A, Riecken EO, Fromm M, Schulzke JD. (2000). Expression from the human occludin promoter is affected by tumor necrosis factor alpha and interferon gamma. *J Cell Sci.* **113**, 2085-90.

- Mankertz J, Waller JS, Hillenbrand B, Tavalali S, Florian P, Schoneberg T, Fromm M, Schulzke JD. (2002). Gene expression of the tight junction protein occludin includes differential splicing and alternative promoter usage. *Biochem Biophys Res Commun.* **298**, 657-66.
- Marfatia SM, Leu RA, Branton D, Chishti AH. (1995). Identification of the protein 4.1 binding interface on glycophorin C and p55, a homologue of the Drosophila discs-large tumor suppressor protein. *J Biol Chem.* **270**, 715-9.
- Masuko N, Makino K, Kuwahara H, Fukunaga K, Sudo T, Araki N, Yamamoto H, Yamada Y, Miyamoto E, Saya H. (1999). Interaction of NE-dlg/SAP102, a neuronal and endocrine tissue-specific membrane-associated guanylate kinase protein, with calmodulin and PSD-95/SAP90. A possible regulatory role in molecular clustering at synaptic sites. *J Biol Chem.* **274**, 5782-90.
- Mattagajasingh SN, Huang SC, Hartenstein JS, Benz EJ Jr. (2000). Characterization of the interaction between protein 4.1R and ZO-2. A possible link between the tight junction and the actin cytoskeleton. *J Biol Chem.* **275**, 30573-85.
- Matter K, Balda MS. (1998). Biogenesis of tight junctions: the C-terminal domain of occludin mediates basolateral targeting. *J Cell Sci.* **111**, 511-9.
- Mayhan WG. (1999). VEGF increases permeability of the blood-brain barrier via a nitric oxide synthase/cGMP-dependent pathway. *Am J Physiol.* **276**, C1148-53.
- McCarthy KM, Skare IB, Stankewich MC, Furuse M, Tsukita S, Rogers RA, Lynch RD, Schneeberger EE. (1996). Occludin is a functional component of the tight junction. *J Cell Sci.* **109**, 2287-98.
- McGee AW, Dakoji SR, Olsen O, Bredt DS, Lim WA, Prehoda KE. (2001). Structure of the SH3-guanylate kinase module from PSD-95 suggests a mechanism for regulated assembly of MAGUK scaffolding proteins. *Mol Cell.* **8**, 1291-301.
- McLaughlin M, Hale R, Ellston D, Gaudet S, Lue RA, Viel A. (2002). The distribution and function of alternatively spliced insertions in hDlg. *J Biol Chem.* **277**, 6406-12.
- Medina R, Rahner C, Mitic LL, Anderson JM, Van Itallie CM. (2000). Occludin localization at the tight junction requires the second extracellular loop. *J Membr Biol.* **178**, 235-47.
- Meyer TN, Schwesinger C, Denker BM. (2002). Zonula occludens-1 is a scaffolding protein for signaling molecules. Galpha(12) directly binds to the Src homology 3 domain and regulates paracellular permeability in epithelial cells. *J Biol Chem.* **277**, 24855-8.
- Misura KM, Gonzalez LC Jr, May AP, Scheller RH, Weis WI. (2001). Crystal structure and biophysical properties of a complex between the N-terminal SNARE region of SNAP25 and syntaxin 1a. *J Biol Chem.* **276**, 41301-9.
- Mitic LL, Schneeberger EE, Fanning AS, Anderson JM. (1999). Connexin-occludin chimeras containing the ZO-binding domain of occludin localize at MDCK tight junctions and NRK cell contacts. *J Cell Biol.* **146**, 683-93.
- Mitic LL, Van Itallie CM. (2001). Occludin and claudins: transmembrane proteins of the tight junction. In Cereijido M, Anderson JM. (Eds.), Tight junctions. Boca Raton.
- Mitic LL, Unger VM, Anderson JM. (2003). Expression, solubilization, and biochemical characterization of the tight junction transmembrane protein claudin-4. *Protein Sci.* **12**, 218-27.
- Miwa N, Furuse M, Tsukita S, Niikawa N, Nakamura Y, Furukawa Y. (2000). Involvement of claudin-1 in the beta-catenin/Tcf signaling pathway and its frequent upregulation in human colorectal cancers. *Oncol Res.* **12**, 469-76.
- Morita K, Sasaki H, Fujimoto K, Furuse M, Tsukita S. (1999). Claudin-11/OSP-based tight junctions of myelin sheaths in brain and Sertoli cells in testis. *J Cell Biol.* **145**, 579-88.
- Morita K, Tsukita S, Miyachi Y. (2004). Tight junction-associated proteins (occludin, ZO-1, claudin-1, claudin-4) in squamous cell carcinoma and Bowen's disease. *Br J Dermatol.* **151**, 328-34.

- Morton TA, Myszka DG, Chaiken IM. (1995). Interpreting complex binding kinetics from optical biosensors: a comparison of analysis by linearization, the integrated rate equation, and numerical integration. *Anal Biochem.* **227**, 176-85.
- Müller D, Kausalya PJ, Claverie-Martin F, Meij IC, Eggert P, Garcia-Nieto V, Hunziker W. (2003). A novel claudin 16 mutation associated with childhood hypercalciuria abolishes binding to ZO-1 and results in lysosomal mistargeting. *Am J Hum Genet.* **73**, 1293-301.
- Muresan Z, Paul DL, Goodenough DA. (2000). Occludin 1B, a variant of the tight junction protein occludin. *Mol Biol Cell.* **11**, 627-34.
- Myszka DG, Morton TA, Doyle ML, Chaiken IM. (1997). Kinetic analysis of a protein antigen-antibody interaction limited by mass transport on an optical biosensor. *Biophys Chem.* **64**, 127-37.
- Myszka DG, He X, Dembo M, Morton TA, Goldstein B. (1998). Extending the range of rate constants available from BIACORE: interpreting mass transport-influenced binding data. *Biophys J.* **75**, 583-94.
- Nagata K, Handa H. (Eds.) (2000). Real-Time Analysis of Biomolecular Interaction – Applications of BIACORE, Springer-Verlag Tokyo
- Nieba L, Krebber A, Pluckthun A. (1996). Competition BIACore for measuring true affinities: large differences from values determined from binding kinetics. *Anal Biochem.* **234**, 155-65.
- Nieset JE, Redfield AR, Jin F, Knudsen KA, Johnson KR, Wheelock MJ. (1997). Characterization of the interactions of alpha-catenin with alpha-actinin and beta-catenin/plakoglobin. *J Cell Sci.* **110**, 1013-22.
- Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, Furuse M, Tsukita S. (2003). Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol.* **161**, 653-60.
- Nusrat A, Parkos CA, Verkade P, Foley CS, Liang TW, Innis-Whitehouse W, Eastburn KK, Madara JL. (2000a). Tight junctions are membrane microdomains. *J Cell Sci.* **113**, 1771-81.
- Nusrat A, Chen JA, Foley CS, Liang TW, Tom J, Cromwell M, Quan C, Mrsny RJ. (2000b). The coiled-coil domain of occludin can act to organize structural and functional elements of the epithelial tight junction. *J Biol Chem.* **275**, 29816-22.
- Nix SL, Chishti AH, Anderson JM, Walther Z. (2000). hCASK and hDlg associate in epithelia, and their src homology 3 and guanylate kinase domains participate in both intramolecular and intermolecular interactions. *J Biol Chem.* **275**, 41192-200.
- Obama H, Ozawa M. (1997). Identification of the domain of alpha-catenin involved in its association with beta-catenin and plakoglobin (gamma-catenin). *J Biol Chem.* **272**, 11017-20.
- Ohnishi H, Nakahara T, Furuse K, Sasaki H, Tsukita S, Furuse M. (2004). JACOP, a novel plaque protein localizing at the apical junctional complex with sequence similarity to cingulin. *J Biol Chem.* **279**, 46014-22.
- Oshima T, Blaschuk O, Gour B, Symonds M, Elrod JW, Sasaki M, Jackson TH, Alexander JS. (2003). Tight junction peptide antagonists enhance neutrophil trans-endothelial chemotaxis. *Life Sci.* **73**, 1729-40.
- Paarmann I, Spangenberg O, Lavie A, Konrad M. (2002). Formation of complexes between Ca<sup>2+</sup>-calmodulin and the synapse-associated protein SAP97 requires the SH3 domain-guanylate kinase domain-connecting HOOK region. *J Biol Chem.* **277**, 40832-8.
- Patel SD, Chen CP, Bahna F, Honig B, Shapiro L. (2003). Cadherin-mediated cell-cell adhesion: sticking together as a family. *Curr Opin Struct Biol.* **13**, 690-8.
- Pauling L, Corey RB. (1953). Compound helical configurations of polypeptide chains: structure of proteins of the alpha-keratin type. *Nature.* **171**, 59-61.

- Peng BH, Lee JC, Campbell GA. (2003). In vitro protein complex formation with cytoskeleton-anchoring domain of occludin identified by limited proteolysis. *J Biol Chem.* **278**, 49644-51.
- Pfefferkorn G. (1968). Elektronenmikroskopie. Institut für medizinische Physik
- Plattner H, Hentschel J. (1997). Taschenlehrbuch Zellbiologie, Thieme Verlag, Stuttgart
- Pokutta S, Weis WI. (2000). Structure of the Dimerization and  $\beta$ -Catenin Binding Region of  $\alpha$ -Catenin. *Mol Cell.* **5**, 533-43.
- Pokutta S, Weis WI. (2002). The cytoplasmic face of cell contact sites. *Curr Opin Struct Biol.* **12**, 255-62.
- Pollard TD, Earnshaw WC. (2002). Cell Biology Saunders (an Imprint from Elsevier Science), Philadelphia
- Pradhan D, Lombardo CR, Roe S, Rimm DL, Morrow JS. (2001). Alpha-Catenin binds directly to spectrin and facilitates spectrin-membrane assembly in vivo. *J Biol Chem.* **276**, 4175-81.
- Rajasekaran AK, Hojo M, Huima T, Rodriguez-Boulan E. (1996). Catenins and zonula occludens-1 form a complex during early stages in the assembly of tight junctions. *J Cell Biol.* **132**, 451-63.
- Rao RK, Basuroy S, Rao VU, Karnaky Jr KJ, Gupta A. (2002). Tyrosine phosphorylation and dissociation of occludin-ZO-1 and E-cadherin-beta-catenin complexes from the cytoskeleton by oxidative stress. *Biochem J.* **368**, 471-81.
- Raschperger E, Engstrom U, Pettersson RF, Fuxe J. (2004). CLMP, a novel member of the CTX family and a new component of epithelial tight junctions. *J Biol Chem.* **279**, 796-804.
- Reichert M, Muller T, Hunziker W. (2000). The PDZ domains of zonula occludens-1 induce an epithelial to mesenchymal transition of Madin-Darby canine kidney I cells. Evidence for a role of beta-catenin/Tcf/Lef signaling. *J Biol Chem.* **275**, 9492-500.
- Rimm DL, Koslov ER, Kebriaei P, Cianci CD, Morrow JS. (1995). Alpha-1(E)-catenin is an actin-binding and -bundling protein mediating the attachment of F-actin to the membrane adhesion complex. *Proc Natl Acad Sci U S A.* **92**, 8813-7.
- Roh MH, Liu CJ, Laurinec S, Margolis B. (2002). The carboxyl terminus of zona occludens-3 binds and recruits a mammalian homologue of discs lost to tight junctions. *J Biol Chem.* **277**, 27501-9.
- Rousseau F, Schymkowitz JW, Itzhaki LS. (2003). The unfolding story of three-dimensional domain swapping. *Structure (Camb).* **11**, 243-51.
- Rubin LL, Staddon JM. (1999). The cell biology of the blood-brain barrier. *Annu Rev Neurosci.* **22**, 11-28.
- Rüffer C, Gerke V. (2004). The C-terminal cytoplasmic tail of claudins 1 and 5 but not its PDZ-binding motif is required for apical localization at epithelial and endothelial tight junctions. *Eur J Cell Biol.* **83**, 135-44.
- Ryeom SW, Paul D, Goodenough DA. (2000). Truncation mutants of the tight junction protein ZO-1 disrupt corneal epithelial cell morphology. *Mol Biol Cell.* **11**, 1687-96.
- Saitou M, Fujimoto K, Doi Y, Itoh M, Fujimoto T, Furuse M, Takano H, Noda T, Tsukita S. (1998). Occludin-deficient embryonic stem cells can differentiate into polarized epithelial cells bearing tight junctions. *J Cell Biol.* **141**, 397-408.
- Saitou M, Furuse M, Sasaki H, Schulzke JD, Fromm M, Takano H, Noda T, Tsukita S. (2000). Complex phenotype of mice lacking occludin, a component of tight junction strands. *Mol Biol Cell.* **11**, 4131-42.
- Sakakibara A, Furuse M, Saitou M, Ando-Akatsuka Y, Tsukita S. (1997). Possible involvement of phosphorylation of occludin in tight junction formation. *J Cell Biol.* **137**, 1393-401.

- Sambrook J, Fritsch EF, Maniatis T. (1989). Molecular cloning: A laboratory manual. Cold Spring Harbour Laboratory Press, USA.
- Sanchez-Pulido L, Martin-Belmonte F, Valencia A, Alonso MA. (2002). MARVEL: a conserved domain involved in membrane apposition events. *Trends Biochem Sci.* **27**, 599-601.
- Sasaki H, Matsui C, Furuse K, Mimori-Kiyosue Y, Furuse M, Tsukita S. (2003). Dynamic behavior of paired claudin strands within apposing plasma membranes. *Proc Natl Acad Sci U S A.* **100**, 3971-6.
- Schmidt A, Utepbergenov DI, Krause G, Blasig IE. (2001). Use of surface plasmon resonance for real-time analysis of the interaction of ZO-1 and occludin. *Biochem Biophys Res Commun.* **288**, 1194-9.
- Schmidt A. Dissertationsschrift im Fach Biochemie an der FU Berlin 2002.
- Schmidt A, Utepbergenov DI, Mueller SL, Beyermann M, Schneider-Mergener J, Krause G, Blasig IE. (2004). Occludin binds to the SH3-hinge-GuK unit of zonula occludens protein 1: potential mechanism of tight junction regulation. *Cell Mol Life Sci.* **61**, 1354-65.
- Schmidt RF, Thews G. (Hrsg.) (1995). Physiologie des Menschen Springer Verlag Berlin
- Schultz J, Milpetz F, Bork P, Ponting CP. (1998). SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A.* **95**, 5857-64.
- Simon DB, Lu Y, Choate KA, Velazquez H, Al-Sabban E, Praga M, Casari G, Bettinelli A, Colussi G, Rodriguez-Soriano J, McCredie D, Milford D, Sanjad S, Lifton RP. (1999). Paracellin-1, a renal tight junction protein required for paracellular Mg<sup>2+</sup> resorption. *Science.* **285**, 103-6.
- Simons KT, Bonneau R, Ruczinski I, Baker D. (1999). Ab initio protein structure prediction of CASP III targets using ROSETTA. *Proteins. Suppl.* **3**, 171-6.
- Sonoda N, Furuse M, Sasaki H, Yonemura S, Katahira J, Horiguchi Y, Tsukita S. (1999). Clostridium perfringens enterotoxin fragment removes specific claudins from tight junction strands: Evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol.* **147**, 195-204.
- Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA. (1986). Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biol.* **103**, 755-66.
- Tang VW, Goodenough DA. (2003). Paracellular ion channel at the tight junction. *Biophys J.* **84**, 1660-73.
- Tavares GA, Panepucci EH, Brunger AT. (2001). Structural characterization of the intramolecular interaction between the SH3 and guanylate kinase domains of PSD-95. *Mol Cell.* **8**, 1313-25.
- Tavelin S, Hashimoto K, Malkinson J, Lazorova L, Toth I, Artursson P. (2003). A new principle for tight junction modulation based on occludin peptides. *Mol Pharmacol.* **64**, 1530-40.
- Tewes BJ, Galla HJ. (2001). Lipid polarity in brain capillary endothelial cells. *Endothelium.* **8**, 207-20.
- Thompson JD, Higgins DG, Gibson TJ. (1994). ClustalW: improving the sensitivity of progressive multiple sequence alignments through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673-80
- Toepert F, Knaute T, Guffler S, Pires JR, Matzdorf T, Oschkinat H, Schneider-Mergener J. (2003). Combining SPOT synthesis and native peptide ligation to create large arrays of WW protein domains. *Angew Chem Int Ed Engl.* **42**, 1136-40.

- Traweger A, Fang D, Liu YC, Stelzhammer W, Krizbai IA, Fresser F, Bauer HC, Bauer H. (2002). The tight junction-specific protein occludin is a functional target of the E3 ubiquitin-protein ligase itch. *J Biol Chem.* **277**, 10201-8.
- Traweger A, Fuchs R, Krizbai IA, Weiger TM, Bauer HC, Bauer H. (2003). The tight junction protein ZO-2 localizes to the nucleus and interacts with the heterogeneous nuclear ribonucleoprotein scaffold attachment factor-B. *J Biol Chem.* **278**, 2692-700.
- Tsukamoto T, Nigam SK. (1997). Tight junction proteins form large complexes and associate with the cytoskeleton in an ATP depletion model for reversible junction assembly. *J Biol Chem.* **272**, 16133-9.
- Tsukamoto T, Nigam SK. (1999). Role of tyrosine phosphorylation in the reassembly of occludin and other tight junction proteins. *Am J Physiol.* **276**, F737-50.
- Tsukita S, Furuse M. (1999). Occludin and claudins in tight-junction strands: leading or supporting players? *Trends Cell Biol.* **9**, 268-73.
- Tsukita S, Furuse M. (2000). Pores in the wall: claudins constitute tight junction strands containing aqueous pores. *J Cell Biol.* **149**, 13-6.
- Tsukita S, Furuse M, Itoh M. (2001). Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol.* **2**, 285-93.
- Turner JR. (2000). 'Putting the squeeze' on the tight junction: understanding cytoskeletal regulation. *Semin Cell Dev Biol.* **11**, 301-8.
- Umeda K, Matsui T, Nakayama M, Furuse K, Sasaki H, Furuse M, Tsukita S. (2004). Establishment and characterization of cultured epithelial cells lacking expression of ZO-1. *J Biol Chem.* **279**, 44785-94.
- Van Itallie CM, Anderson JM. (1997). Occludin confers adhesiveness when expressed in fibroblasts. *J Cell Sci.* **110**, 1113-21.
- Van Itallie CM, Colegio OR, Anderson JM. (2004). The cytoplasmic tails of claudins can influence tight junction barrier properties through effects on protein stability. *J Membr Biol.* **199**, 29-38.
- Vasioukhin V, Fuchs E. (2001). Actin dynamics and cell-cell adhesion in epithelia. *Curr Opin Cell Biol.* **13**, 76-84.
- Viator I, Bader T, Paiha K, Huber LA. (2001). Perturbation of the tight junction permeability barrier by occludin loop peptides activates beta-catenin/TCF/LEF-mediated transcription. *EMBO Rep.* **2**, 306-12.
- Vogelmann R, Amieva MR, Falkow S, Nelson WJ. (2004). Breaking into the epithelial apical-junctional complex--news from pathogen hackers. *Curr Opin Cell Biol.* **16**, 86-93.
- Vriend G. (2000). Professional Gambling (über das Erstellen von Homologiemodellen) aus <http://www.cmbi.kun.nl/gv/articles/text/gambling.html>
- Wachtel M, Frei K, Ehler E, Fontana A, Winterhalter K, Gloor SM. (1999). Occludin proteolysis and increased permeability in endothelial cells through tyrosine phosphatase inhibition. *J Cell Sci.* **112**, 4347-56.
- Wan H, Winton HL, Soeller C, Tovey ER, Gruenert DC, Thompson PJ, Stewart GA, Taylor GW, Garrod DR, Cannell MB, Robinson C. (1999). Derp1 facilitates transepithelial allergen delivery by disruption of tight junctions. *J Clin Invest.* **104**, 123-33.
- Watabe-Uchida M, Uchida N, Imamura Y, Nagafuchi A, Fujimoto K, Uemura T, Vermeulen S, van Roy F, Adamson ED, Takeichi M. (1998). Alpha-Catenin-vinculin interaction functions to organize the apical junctional complex in epithelial cells. *J Cell Biol.* **142**, 847-57.
- Wilcox ER, Burton QL, Naz S, Riazuddin S, Smith TN, Ploplis B, Belyantseva I, Ben-Yosef T, Liburd NA, Morell RJ, Kachar B, Wu DK, Griffith AJ, Riazuddin S, Friedman TB. (2001). Mutations in the gene encoding tight junction claudin-14 cause autosomal recessive deafness DFNB29. *Cell.* **104**, 165-72.

- Witt KA, Mark KS, Hom S, Davis TP. (2003). Effects of hypoxia-reoxygenation on rat blood-brain barrier permeability and tight junctional protein expression. *Am J Physiol Heart Circ Physiol.* **285**, H2820-31.
- Wittchen ES, Haskins J, Stevenson BR. (1999). Protein interactions at the tight junction. Actin has multiple binding partners, and ZO-1 forms independent complexes with ZO-2 and ZO-3. *J Biol Chem.* **274**, 35179-85.
- Wong V. (1997). Phosphorylation of occludin correlates with occludin localization and function at the tight junction. *Am J Physiol.* **273**, C1859-67.
- Wong V, Gumbiner BM. (1997). A synthetic peptide corresponding to the extracellular domain of occludin perturbs the tight junction permeability barrier. *J Cell Biol.* **136**, 399-409.
- Wu H, Reissner C, Kuhlendahl S, Coblenz B, Reuver S, Kindler S, Gundelfinger ED, Garner CC. (2000). Intramolecular interactions regulate SAP97 binding to GKAP. *EMBO J.* **19**, 5740-51.
- Wu Z, Nybom P, Magnusson KE. (2000). Distinct effects of Vibrio cholerae haemagglutinin/protease on the structure and localization of the tight junction-associated proteins occludin and ZO-1. *Cell Microbiol.* **2**, 11-7.
- Yamamoto T, Harada N, Kano K, Taya S, Canaani E, Matsuura Y, Mizoguchi A, Ide C, Kaibuchi K. (1997). The Ras target AF-6 interacts with ZO-1 and serves as a peripheral component of tight junctions in epithelial cells. *J Cell Biol.* **139**, 785-95.
- Yamanaka T, Horikoshi Y, Suzuki A, Sugiyama Y, Kitamura K, Maniwa R, Nagai Y, Yamashita A, Hirose T, Ishikawa H, Ohno S. (2001). PAR-6 regulates aPKC activity in a novel way and mediates cell-cell contact-induced formation of the epithelial junctional complex. *Genes Cells.* **6**, 721-31.
- Yang J, Dokurno P, Tonks NK, Barford D. (2001). Crystal structure of the M-fragment of alpha-catenin: implications for modulation of cell adhesion. *EMBO J.* **20**, 3645-56.
- Yokoyama S, Tachibana K, Nakanishi H, Yamamoto Y, Irie K, Mandai K, Nagafuchi A, Monden M, Takai Y. (2001). Alpha-catenin-independent recruitment of ZO-1 to nectin-based cell-cell adhesion sites through afadin. *Mol Biol Cell.* **12**, 1595-609.
- Zegers I, Deswarte J, Wyns L. (1999). Trimeric domain-swapped barnase. *Proc Natl Acad Sci U S A.* **96**, 818-22.