

G. LITERATURE

- Aboitiz,F., Montiel,J., and Lopez,J. (2002). Critical steps in the early evolution of the isocortex: insights from developmental biology. *Braz.J Med.Biol.Res.* 35, 1455-1472.
- Arber,S., Burden,S.J., and Harris,A.J. (2002). Patterning of skeletal muscle. *Curr.Opin.Neurobiol.* 12, 100-103.
- Arribas,J. and Borroto,A. (2002). Protein ectodomain shedding. *Chem.Rev* 102, 4627-4638.
- Arribas,J., Coodly,L., Vollmer,P., Kishimoto,T.K., Rose-John,S., and Massague,J. (1996). Diverse cell surface protein ectodomains are shed by a system sensitive to metalloprotease inhibitors. *J Biol.Chem.* 271, 11376-11382.
- Arribas,J., Lopez-Casillas,F., and Massague,J. (1997). Role of the juxtamembrane domains of the transforming growth factor-alpha precursor and the beta-amyloid precursor protein in regulated ectodomain shedding. *J Biol.Chem.* 272, 17160-17165.
- Ausubel,F.M., Brent,R., Kingston,R.E., Moore,D.D., Seidman,J.G., Smith,J.A., and Struhl,K. (1994a). Enzymatic manipulation of DNA and RNA. In *Current Protocols of Molecular Biology*, F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl, eds. John Wiley & Sons.Inc).
- Ausubel,F.M., Brent,R., Kingston,R.E., Moore,D.D., Seidman,J.G., Smith,J.A., and Struhl,K. (1994b). Preparation and analysis of DNA. In *Current Protocols of Molecular Biology*, F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl, eds. John Wiley & Sons.Inc).
- Bailey,C.H., Bartsch,D., and Kandel,E.R. (1996). Toward a molecular definition of long-term memory storage. *Proc.Natl.Acad.Sci.U.S.A* 93, 13445-13452.
- Bailey,C.H., Giustetto,M., Huang,Y.Y., Hawkins,R.D., and Kandel,E.R. (2000). Is heterosynaptic modulation essential for stabilizing Hebbian plasticity and memory? *Nat.Rev.Neurosci.* 1, 11-20.
- Bailey,C.H., Kandel,E.R., and Si,K. (2004). The persistence of long-term memory; a molecular approach to self-sustaining changes in learning-induced synaptic growth. *Neuron* 44, 49-57.
- Biederer,T., Sara,Y., Mozhayeva,M., Atasoy,D., Liu,X., Kavalali,E.T., and Sudhof,T.C. (2002). SynCAM, a synaptic adhesion molecule that drives synapse assembly. *Science* 297, 1525-1531.
- Bizzozero,O.A. (1997). The mechanism and functional roles of protein palmitoylation in the nervous system. *Neuropediatrics* 28, 23-26.
- Bloom,F.E. and Swanson,L.W. (2003). Neuroscience. In *Fundamental neuroscience*, L. R. Squire, F. E. Bloom, S. K. McConnell, J. L. Roberts, N. C. Spitzer, and M. J. Zigmond, eds. (San Diego, USA: Academic press), pp. 1-45.

- Borrell-Pages,M., Rojo,F., Albanell,J., Baselga,J., and Arribas,J. (2003). TACE is required for the activation of the EGFR by TGF-alpha in tumors. *EMBO J* 22, 1114-1124.
- Bottjer,S.W., Brady,J.D., and Cribbs,B. (2000). Connections of a motor cortical region in zebra finches: relation to pathways for vocal learning. *J Comp Neurol.* 420, 244-260.
- Boulanger,L.M., Huh,G.S., and Shatz,C.J. (2001). Neuronal plasticity and cellular immunity: shared molecular mechanisms. *Curr.Opin.Neurobiol.* 11, 568-578.
- Boulanger,L.M. and Shatz,C.J. (2004). Immune signalling in neural development, synaptic plasticity and disease. *Nature Reviews Neuroscience* 5, 521-531.
- Boutin,J.A. (1997). Myristoylation. *Cell Signal.* 9, 15-35.
- Broadie,K. and Bate,M. (1993). Innervation directs receptor synthesis and localization in *Drosophila* embryo synaptogenesis. *Nature* 361, 350-353.
- Broadie,K., Prokop,A., Bellen,H.J., O'Kane,C.J., Schulze,K.L., and Sweeney,S.T. (1995). Syntaxin and synaptobrevin function downstream of vesicle docking in *Drosophila*. *Neuron* 15, 663-673.
- Brorson,J.R., Manzolillo,P.A., and Miller,R.J. (1994). Ca²⁺ entry via AMPA/KA receptors and excitotoxicity in cultured cerebellar Purkinje cells. *J Neurosci* 14, 187-197.
- Brown,C.L., Meise,K.S., Plowman,G.D., Coffey,R.J., and Dempsey,P.J. (1998). Cell surface ectodomain cleavage of human amphiregulin precursor is sensitive to a metalloprotease inhibitor. Release of a predominant N-glycosylated 43-kDa soluble form. *J Biol.Chem.* 273, 17258-17268.
- Burrone,J., O'Byrne,M., and Murthy,V.N. (2002). Multiple forms of synaptic plasticity triggered by selective suppression of activity in individual neurons. *Nature* 420, 414-418.
- Cadas,H., Gaillet,S., Beltramo,M., Venance,L., and Piomelli,D. (1996). Biosynthesis of an endogenous cannabinoid precursor in neurons and its control by calcium and cAMP. *J Neurosci.* 16, 3934-3942.
- Cantaloops,I., Haas,K., and Cline,H.T. (2000). Postsynaptic CPG15 promotes synaptic maturation and presynaptic axon arbor elaboration in vivo. *Nat.Neurosci.* 3, 1004-1011.
- Carlin,R.K., Grab,D.J., Cohen,R.S., and Siekevitz,P. (1980). Isolation and characterization of postsynaptic densities from various brain regions: enrichment of different types of postsynaptic densities. *J.Cell Biol* 86, 831-845.
- Catalano,S.M. and Shatz,C.J. (1998). Activity-dependent cortical target selection by thalamic axons. *Science* 281, 559-562.
- Chen,L., Yang,C., and Mower,G.D. (2001). Developmental changes in the expression of GABA(A) receptor subunits (alpha(1), alpha(2), alpha(3)) in the cat visual cortex and the effects of dark rearing. *Mol.Brain Res.* 88, 135-143.
- Cline,H.T. (2001). Dendritic arbor development and synaptogenesis. *Curr.Opin.Neurobiol.* 11, 118-126.

- Codony-Servat,J., Albanell,J., Lopez-Talavera,J.C., Arribas,J., and Baselga,J. (1999). Cleavage of the HER2 ectodomain is a pervanadate-activatable process that is inhibited by the tissue inhibitor of metalloproteases-1 in breast cancer cells. *Cancer Res.* 59, 1196-1201.
- Cohen-Cory,S. (2002). The developing synapse: construction and modulation of synaptic structures and circuits. *Science* 298, 770-776.
- Coligan,J.E., Kruisbeek,A.M., Margulies,D.H., Shevach,E.M., and Strober,W. (1994). Biotinylation and analysis of membrane bound and soluble proteins. In *Current Protocols in Immunology*, J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, and W. Strober, eds. John Wiley & Sons. Inc).
- Colledge,M., Snyder,E.M., Crozier,R.A., Soderling,J.A., Jin,Y., Langeberg,L.K., Lu,H., Bear,M.F., and Scott,J.D. (2003). Ubiquitination regulates PSD-95 degradation and AMPA receptor surface expression. *Neuron* 40, 595-607.
- Contestabile,A. (2000). Roles of NMDA receptor activity and nitric oxide production in brain development. *Brain Res.Brain Res.Rev* 32, 476-509.
- Corriveau,R.A. (1999). Electrical activity and gene expression in the development of vertebrate neural circuits. *J Neurobiol.* 41, 148-157.
- Corriveau,R.A., Shatz,C.J., and Nedivi,E. (1999). Dynamic regulation of cpg15 during activity-dependent synaptic development in the mammalian visual system. *J Neurosci* 19, 7999-8008.
- Costell,M., Mann,K., Yamada,Y., and Timpl,R. (1997). Characterization of recombinant perlecan domain I and its substitution by glycosaminoglycans and oligosaccharides. *Eur.J Biochem.* 243, 115-121.
- Crowley,J.C. and Katz,L.C. (2002). Ocular dominance development revisited. *Curr.Opin.Neurobiol.* 12, 104-109.
- Davis,G.W. and Bezprozvanny,I. (2001). Maintaining the stability of neural function: a homeostatic hypothesis. *Annu.Rev Physiol* 63, 847-869.
- Debski,E.A. and Cline,H.T. (2002). Activity-dependent mapping in the retinotectal projection. *Curr.Opin.Neurobiol.* 12, 93-99.
- DeChiara,T.M., Bowen,D.C., Valenzuela,D.M., Simmons,M.V., Poueymirou,W.T., Thomas,S., Kinetz,E., Compton,D.L., Rojas,E., Park,J.S., Smith,C., DiStefano,P.S., Glass,D.J., Burden,S.J., and Yancopoulos,G.D. (1996). The receptor tyrosine kinase MuSK is required for neuromuscular junction formation in vivo. *Cell* 85, 501-512.
- Dempsey,P.J., Meise,K.S., Yoshitake,Y., Nishikawa,K., and Coffey,R.J. (1997). Apical enrichment of human EGF precursor in Madin-Darby canine kidney cells involves preferential basolateral ectodomain cleavage sensitive to a metalloprotease inhibitor. *J Cell Biol.* 138, 747-758.
- Desai,N.S., Cudmore,R.H., Nelson,S.B., and Turrigiano,G.G. (2002). Critical periods for experience-dependent synaptic scaling in visual cortex. *Nat.Neurosci.* 5, 783-789.

- DiAntonio,A. and Hicke,L. (2004). Ubiquitin-dependent regulation of the synapse. Annual Review of Neuroscience 27, 223-246.
- Diaz-Rodriguez,E., Esparis-Ogando,A., Montero,J.C., Yuste,L., and Pandiella,A. (2000). Stimulation of cleavage of membrane proteins by calmodulin inhibitors. Biochem.J 346 Pt 2, 359-367.
- Diaz-Rodriguez,E., Montero,J.C., Esparis-Ogando,A., Yuste,L., and Pandiella,A. (2002). Extracellular signal-regulated kinase phosphorylates tumor necrosis factor alpha-converting enzyme at threonine 735: a potential role in regulated shedding. Mol.Biol.Cell 13, 2031-2044.
- Dityatev,A., Dityateva,G., and Schachner,M. (2000). Synaptic strength as a function of post- versus presynaptic expression of the neural cell adhesion molecule NCAM. Neuron 26, 207-217.
- Dolmetsch,R.E., Pajvani,U., Fife,K., Spotts,J.M., and Greenberg,M.E. (2001). Signaling to the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase pathway. Science 294, 333-339.
- Dong,J. and Wiley,H.S. (2000). Trafficking and proteolytic release of epidermal growth factor receptor ligands are modulated by their membrane-anchoring domains. J Biol.Chem. 275, 557-564.
- Dunning,D.D., Hoover,C.L., Soltesz,I., Smith,M.A., and O'Dowd,D.K. (1999). GABA(A) receptor-mediated miniature postsynaptic currents and alpha-subunit expression in developing cortical neurons. J.Neurophysiol. 82, 3286-3297.
- Ehlers,M.D. (2000). Reinsertion or degradation of AMPA receptors determined by activity-dependent endocytic sorting. Neuron 28, 511-525.
- Ehlers,M.D. (2003). Activity level controls postsynaptic composition and signaling via the ubiquitin-proteasome system. Nat.Neurosci. 6, 231-242.
- Ehrman,M. and Clausen,T. (2004). Proteolysis as a Regulatory Mechanism. Annu.Rev Genet.
- El-Husseini,A.E.-D. and Bredt,D.S. (2002). Protein palmitoylation: A regulator of neuronal development and function. Nature Reviews Neuroscience 3, 791-802.
- Facchinetto,F., Ciani,E., Dall'Olio,R., Virgili,M., Contestabile,A., and Fonnum,F. (1993). Structural, neurochemical and behavioural consequences of neonatal blockade of NMDA receptor through chronic treatment with CGP 39551 or MK-801. Brain Res.Dev.Brain Res. 74, 219-224.
- Fan,H. and Deryck,R. (1999). Ectodomain shedding of TGF-alpha and other transmembrane proteins is induced by receptor tyrosine kinase activation and MAP kinase signaling cascades. EMBO J 18, 6962-6972.
- Farazi,T.A., Waksman,G., and Gordon,J.I. (2001). The biology and enzymology of protein N-myristoylation. J Biol.Chem. 276, 39501-39504.

- Feller,M.B. (1999). Spontaneous correlated activity in developing neural circuits. *Neuron* 22, 653-656.
- Fernandez,M.T., Zitko,V., Gascon,S., and Novelli,A. (1991). The marine toxin okadaic acid is a potent neurotoxin for cultured cerebellar neurons. *Life Sci.* 49, L157-L162.
- Fischbach,G.D. and Rosen,K.M. (1997). ARIA: a neuromuscular junction neuregulin. *Annu.Rev.Neurosci.* 20, 429-458.
- Fruman,D.A., Klee,C.B., Bierer,B.E., and Burakoff,S.J. (1992). Calcineurin phosphatase activity in T lymphocytes is inhibited by FK 506 and cyclosporin A. *Proc.Natl.Acad.Sci.U.S.A* 89, 3686-3690.
- Fujisawa,K., Onoue,H., Abe,K., and Ito,Y. (1999). Multiple calcium channels regulate neurotransmitter release from vagus nerve terminals in the cat bronchiole. *British Journal of Pharmacology* 128, 262-268.
- Gallea-Robache,S., Morand,V., Millet,S., Bruneau,J.M., Bhatnagar,N., Chouaib,S., and Roman-Roman,S. (1997). A metalloproteinase inhibitor blocks the shedding of soluble cytokine receptors and processing of transmembrane cytokine precursors in human monocytic cells. *Cytokine* 9, 340-346.
- Gechman,Z., Alonso,J.L., Raab,G., Ingber,D.E., and Klagsbrun,M. (1999). The shedding of membrane-anchored heparin-binding epidermal-like growth factor is regulated by the Raf/mitogen-activated protein kinase cascade and by cell adhesion and spreading. *J.Biol.Chem.* 274, 28828-28835.
- Ghosh,A. and Greenberg,M.E. (1995). Calcium signaling in neurons: molecular mechanisms and cellular consequences. *Science* 268, 239-247.
- Goda,Y. and Davis,G.W. (2003). Mechanisms of synapse assembly and disassembly. *Neuron* 40, 243-264.
- Goggi,J., Pullar,I.A., Carney,S.L., and Bradford,H.F. (2003). Signalling pathways involved in the short-term potentiation of dopamine release by BDNF. *Brain Research* 968, 156-161.
- Graef,I.A., Mermelstein,P.G., Stankunas,K., Neilson,J.R., Deisseroth,K., Tsien,R.W., and Crabtree,G.R. (1999). L-type calcium channels and GSK-3 regulate the activity of NF-ATc4 in hippocampal neurons. *Nature* 401, 703-708.
- Gremo,F., Porru,S., and Vernadakis,A. (1984). Effects of corticosterone on chick embryonic retinal cells in culture. *Brain Res.* 317, 45-52.
- Hamburger,F. and Hamilton,H.L. (1951). A series of normal stages in the development of the chick embryo. *Journal of Morphology* 88, 49-92.
- Hammond,C. (2001). Cellular and molecular biology. Academic press), pp. 67-141.
- Han,B. and Fischbach,G.D. (1999). Processing of ARIA and release from isolated nerve terminals. *Philos.Trans.R.Soc.Lond B Biol.Sci.* 354, 411-416.

- Hanahan,D. (1985). Techniques for transformation of E.coli. In DNA cloning, a practical approach, D. M. Glover, ed. IRL press, Oxford, Washington D.C), pp. 109-135.
- Harris,B.T., Costa,E., and Grayson,D.R. (1995). Exposure of neuronal cultures to K⁺ depolarization or to N-methyl-D-aspartate increases the transcription of genes encoding the α 1 and α 5 GABA_A receptor subunits. Mol.Brain Res. 28, 338-342.
- Hasclowicz,T., Murai,N., Matsufuji,S., and Murakami,Y. (2002). Regulation of ornithine decarboxylase by antizymes and antizyme inhibitor in zebrafish (*Danio rerio*). Biochim.Biophys.Acta 1578, 21-28.
- Hering,H. and Kroger,S. (1996). Formation of synaptic specializations in the inner plexiform layer of the developing chick retina. J Comp Neurol. 375, 393-405.
- Hirata,M., Umata,T., Takahashi,T., Ohnuma,M., Miura,Y., Iwamoto,R., and Mekada,E. (2001). Identification of serum factor inducing ectodomain shedding of proHB-EGF and sStudies of noncleavable mutants of proHB-EGF. Biochem.Biophys.Res.Commun. 283, 915-922.
- Hooper,N.M. (1994). Families of zinc metalloproteases. FEBS Lett. 354, 1-6.
- Horton,J.C. and Hocking,D.R. (1996). An adult-like pattern of ocular dominance columns in striate cortex of newborn monkeys prior to visual experience. J Neurosci 16, 1791-1807.
- HUBEL,D.H. and Wiesel,T.N. (1963). Receptive fields of cells in striate cortex of very young, visually inexperienced kittens. J Neurophysiol. 26, 994-1002.
- Huh,G.S., Boulanger,L.M., Du,H., Riquelme,P.A., Brotz,T.M., and Shatz,C.J. (2000). Functional requirement for class I MHC in CNS development and plasticity. Science 290, 2155-2159.
- Hundhausen,C., Misztela,D., Berkhout,T.A., Broadway,N., Saftig,P., Reiss,K., Hartmann,D., Fahrenholz,F., Postina,R., Matthews,V., Kallen,K.J., Rose-John,S., and Ludwig,A. (2003). The disintegrin-like metalloproteinase ADAM10 is involved in constitutive cleavage of CX3CL1 (fractalkine) and regulates CX3CL1-mediated cell-cell adhesion. Blood 102, 1186-1195.
- Ientile,R., Macaione,V., Teletta,M., Pedale,S., Torre,V., and Macaione,S. (2001). Apoptosis and necrosis occurring in excitotoxic cell death in isolated chick embryo retina. Journal of Neurochemistry 79, 71-78.
- Inatani,M., Tanihara,H., Oohira,A., Otori,Y., Nishida,A., Honjo,M., Kido,N., and Honda,Y. (2000). Neuroglycan C, a neural tissue-specific transmembrane chondroitin sulfate proteoglycan, in retinal neural network formation. Invest Ophthalmol.Vis.Sci. 41, 4338-4346.
- Ishida,I. and Deguchi,T. (1983). Effect of depolarizing agents on choline acetyltransferase and acetylcholinesterase activities in primary cell cultures of spinal cord. J Neurosci 3, 1818-1823.
- Itai,T., Tanaka,M., and Nagata,S. (2001). Processing of tumor necrosis factor by the membrane-bound TNF-alpha-converting enzyme, but not its truncated soluble form. Eur.J.Biochem. 268, 2074-2082.

- Ito,H., Yoshimotoa,M., Alberta,J.S., Yamamotoa,N., and Sawaia,N. (1999a). Retinal projections and retinal ganglion cell distribution patterns in a sturgeon (*Acipenser transmontanus*), a non-teleost actinopterygian fish. *Brain Behav.Evol.* 53, 127-141.
- Ito,K., Okamoto,I., Araki,N., Kawano,Y., Nakao,M., Fujiyama,S., Tomita,K., Mimori,T., and Saya,H. (1999b). Calcium influx triggers the sequential proteolysis of extracellular and cytoplasmic domains of E-cadherin, leading to loss of beta-catenin from cell-cell contacts. *Oncogene* 18, 7080-7090.
- Itoh,K., Ozaki,M., Stevens,B., and Fields,R.D. (1997). Activity-dependent regulation of N-cadherin in DRG neurons: differential regulation of N-cadherin, NCAM, and L1 by distinct patterns of action potentials. *J.Neurobiol.* 33, 735-748.
- Jung,K.M., Tan,S., Landman,N., Petrova,K., Murray,S., Lewis,R., Kim,P.K., Kim,D.S., Ryu,S.H., Chao,M.V., and Kim,T.W. (2003). Regulated Intramembrane Proteolysis of the p75 Neurotrophin Receptor Modulates Its Association with the TrkA Receptor. *Journal of Biological Chemistry* 278, 42161-42169.
- Kandel,E.R. (1997). Genes, synapses, and long-term memory. *J Cell Physiol* 173, 124-125.
- Kandel,E.R. (2000). The brain and behaviour. In *Principles of Neural Science*, E. R. Kandel, J. H. Schwartz, and T. M. Jessell, eds. McGraw-Hill Companies, Inc), pp. 5-19.
- Kang,T., Park,H.I., Suh,Y., Zhao,Y.G., Tschesche,H., and Sang,Q.X. (2002). Autolytic processing at Glu586-Ser587 within the cysteine-rich domain of human adamalysin 19/disintegrin-metalloproteinase 19 is necessary for its proteolytic activity. *J Biol.Chem.* 277, 48514-48522.
- Karkkainen,I., Rybnikova,E., Pelto-Huikko,M., and Huovila,A.P. (2000). Metalloprotease-disintegrin (ADAM) genes are widely and differentially expressed in the adult CNS. *Mol.Cell Neurosci* 15, 547-560.
- Katsuki,H., Tomita,M., Takenaka,C., Shirakawa,H., Shimazu,S., Ibi,M., Kume,T., Kaneko,S., and Akaike,A. (2001). Superoxide dismutase activity in organotypic midbrain-striatum co-cultures is associated with resistance of dopaminergic neurons to excitotoxicity. *Journal of Neurochemistry* 76, 1336-1345.
- Katz,L.C. and Crowley,J.C. (2002). Development of cortical circuits: lessons from ocular dominance columns. *Nature Reviews Neuroscience* 3, 34-42.
- Katz,L.C. and Shatz,C.J. (1996). Synaptic activity and the construction of cortical circuits. *Science* 274, 1133-1138.
- Kavalali,E.T., Klingauf,J., and Tsien,R.W. (1999). Activity-dependent regulation of synaptic clustering in a hippocampal culture system. *Proc.Natl.Acad.Sci.U.S.A* 96, 12893-12900.
- Klee,C.B., Ren,H., and Wang,X. (1998). Regulation of the calmodulin-stimulated protein phosphatase, calcineurin. *J Biol.Chem.* 273, 13367-13370.
- Laemmli,U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.

- Le Gall,S.M., Auger,R., Dreux,C., and Mauduit,P. (2003). Regulated cell surface pro-EGF ectodomain shedding is a zinc metalloprotease-dependent process. *J.Biol Chem.* 278, 45255-45268.
- Lemjabbar,H., Li,D., Gallup,M., Sidhu,S., Drori,E., and Basbaum,C. (2003). Tobacco smoke-induced lung cell proliferation mediated by tumor necrosis factor alpha-converting enzyme and amphiregulin. *J Biol.Chem.* 278, 26202-26207.
- LeVay,S., Stryker,M.P., and Shatz,C.J. (1978). Ocular dominance columns and their development in layer IV of the cat's visual cortex: a quantitative study. *J Comp Neurol.* 179, 223-244.
- Levitan,I.B. (1999). It is calmodulin after all! Mediator of the calcium modulation of multiple ion channels. *Neuron* 22, 645-648.
- Li,H., Gu,X., Dawson,V.L., and Dawson,T.M. (2004). Identification of calcium- and nitric oxide-regulated genes by differential analysis of library expression (DAzLE). *Proc.Natl.Acad.Sci.U.S.A* 101, 647-652.
- Li,M., Jia,M., Fields,R.D., and Nelson,P.G. (1996). Modulation of calcium currents by electrical activity. *J Neurophysiol.* 76, 2595-2607.
- Lin,H., Wada,K., Yonezawa,M., Shinoki,K., Akamatsu,T., Tsukui,T., and Sakamoto,C. (2003). Tomoregulin ectodomain shedding by proinflammatory cytokines. *Life Sci.* 73, 1617-1627.
- Lin,J.W., Ju,W., Foster,K., Lee,S.H., Ahmadian,G., Wyszynski,M., Wang,Y.T., and Sheng,M. (2000). Distinct molecular mechanisms and divergent endocytotic pathways of AMPA receptor internalization. *Nat.Neurosci.* 3, 1282-1290.
- Lin,W., Burgess,R.W., Dominguez,B., Pfaff,S.L., Sanes,J.R., and Lee,K.F. (2001). Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse. *Nature* 410, 1057-1064.
- Lissin,D.V., Gomperts,S.N., Carroll,R.C., Christine,C.W., Kalman,D., Kitamura,M., Hardy,S., Nicoll,R.A., Malenka,R.C., and Von Zastrow,M. (1998). Activity differentially regulates the surface expression of synaptic AMPA and NMDA glutamate receptors. *Proc.Natl.Acad.Sci.U.S.A* 95, 7097-7102.
- Liu,J., Farmer,J.D., Jr., Lane,W.S., Friedman,J., Weissman,I., and Schreiber,S.L. (1991). Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* 66, 807-815.
- Loeb,J.A. and Fischbach,G.D. (1995). ARIA can be released from extracellular matrix through cleavage of a heparin-binding domain. *J.Cell Biol* 130, 127-135.
- Lowry,O.H., ROSEBROUGH,N.J., FARR,A.L., and RANDALL,R.J. (1951). Protein measurement with the Folin phenol reagent. *J Biol.Chem.* 193, 265-275.
- Man,H.Y., Lin,J.W., Ju,W.H., Ahmadian,G., Liu,L., Becker,L.E., Sheng,M., and Wang,Y.T. (2000). Regulation of AMPA receptor-mediated synaptic transmission by clathrin-dependent receptor internalization. *Neuron* 25, 649-662.

- Mathews,G.C. and Diamond,J.S. (2003). Neuronal Glutamate Uptake Contributes to GABA Synthesis and Inhibitory Synaptic Strength. *Journal of Neuroscience* 23, 2040-2048.
- Matthews,G.G. (2001). Mechanism of nerve action potential. In *Neurobiology*, (Malden, Massachusetts: Blackwell Science, Inc), pp. 66-91.
- McLaughlin,T., Torborg,C.L., Feller,M.B., and O'leary,D.D. (2003). Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development. *Neuron* 40, 1147-1160.
- McMahan,U.J. (1990). The agrin hypothesis. *Cold Spring Harb.Symp.Quant.Biol.* 55, 407-418.
- Mechtersheimer,S., Gutwein,P., Agmon-Levin,N., Stoeck,A., Oleszewski,M., Riedle,S., Postina,R., Fahrenholz,F., Fogel,M., Lemmon,V., and Altevogt,P. (2001). Ectodomain shedding of L1 adhesion molecule promotes cell migration by autocrine binding to integrins. *J.Cell Biol* 155, 661-673.
- Meffert,M.K., Chang,J.M., Wiltgen,B.J., Fanselow,M.S., and Baltimore,D. (2003). NF-kappa B functions in synaptic signaling and behavior. *Nat.Neurosci.* 6, 1072-1078.
- Meyer-Franke,A., Wilkinson,G.A., Kruttgen,A., Hu,M., Munro,E., Hanson,M.G., Jr., Reichardt,L.F., and Barres,B.A. (1998). Depolarization and cAMP elevation rapidly recruit TrkB to the plasma membrane of CNS neurons. *Neuron* 21, 681-693.
- Ming,G., Henley,J., Tessier-Lavigne,M., Song,H., and Poo,M. (2001). Electrical activity modulates growth cone guidance by diffusible factors. *Neuron* 29, 441-452.
- Misgeld,T., Burgess,R.W., Lewis,R.M., Cunningham,J.M., Lichtman,J.W., and Sanes,J.R. (2002). Roles of neurotransmitter in synapse formation: development of neuromuscular junctions lacking choline acetyltransferase. *Neuron* 36, 635-648.
- Miyata,S., Matsumoto,N., Taguchi,K., Akagi,A., Iino,T., Funatsu,N., and Maekawa,S. (2003). Biochemical and ultrastructural analyses of IgLON cell adhesion molecules, Kilon and OBCAM in the rat brain. *Neuroscience* 117, 645-658.
- Mizoguchi,A., Nakanishi,H., Kimura,K., Matsubara,K., Ozaki-Kuroda,K., Katata,T., Honda,T., Kiyohara,Y., Heo,K., Higashi,M., Tsutsumi,T., Sonoda,S., Ide,C., and Takai,Y. (2002). Nectin: an adhesion molecule involved in formation of synapses. *J Cell Biol.* 156, 555-565.
- Molkentin,J.D. (2004). Calcineurin-NFAT signaling regulates the cardiac hypertrophic response in coordination with the MAPKs. *Cardiovasc.Res.* 63, 467-475.
- Montero,J.C., Yuste,L., Diaz-Rodriguez,E., Espanis-Ogando,A., and Pandiella,A. (2002). Mitogen-activated protein kinase-dependent and -independent routes control shedding of transmembrane growth factors through multiple secretases. *Biochem.J* 363, 211-221.
- Moody,W.J. (1998). Control of spontaneous activity during development. *J.Neurobiol.* 37, 97-109.

- Moro,M.A., Hurtado,O., Cardenas,A., Romera,C., Madrigal,J.L., Fernandez-Tome,P., Leza,J.C., Lorenzo,P., and Lizasoain,I. (2003). Expression and function of tumour necrosis factor-alpha-converting enzyme in the central nervous system. *Neurosignals*. 12, 53-58.
- Moss,M.L. and Lambert,M.H. (2002). Shedding of membrane proteins by ADAM family proteases. *Essays Biochem.* 38, 141-153.
- Moss,M.L., White,J.M., Lambert,M.H., and Andrews,R.C. (2001). TACE and other ADAM proteases as targets for drug discovery. *Drug Discov.Today* 6, 417-426.
- Mullberg,J., Oberthur,W., Lottspeich,F., Mehl,E., Dittrich,E., Graeve,L., Heinrich,P.C., and Rose-John,S. (1994). The soluble human IL-6 receptor. Mutational characterization of the proteolytic cleavage site. *J Immunol.* 152, 4958-4968.
- Muller,S., Hoege,C., Pyrowolakis,G., and Jentsch,S. (2001). SUMO, ubiquitin's mysterious cousin. *Nat Rev Mol.Cell Biol.* 2, 202-210.
- Munno,D.W. and Syed,N.I. (2003). Synaptogenesis in the CNS: an odyssey from wiring together to firing together. *J Physiol* 552, 1-11.
- Myhre,G.M., Toruner,M., Abraham,S., and Egan,L.J. (2004). Metalloprotease disintegrin-mediated ectodomain shedding of EGFR ligands promotes intestinal epithelial restitution. *Am J Physiol Gastrointest.Liver Physiol.*
- Naeve,G.S., Ramakrishnan,M., Kramer,R., Hevroni,D., Citri,Y., and Theill,L.E. (1997). Neuritin: a gene induced by neural activity and neurotrophins that promotes neuritogenesis. *Proc.Natl.Acad.Sci.U.S.A* 94, 2648-2653.
- Nastrom,J., Schneider,S.P., and Perl,E.R. (1994). Differential L-glutamate responsiveness among superficial dorsal horn neurons. *J Neurophysiol.* 72, 2956-2965.
- Nedivi,E. (1999). Molecular analysis of developmental plasticity in neocortex. *J Neurobiol.* 41, 135-147.
- Nedivi,E., Wu,G.Y., and Cline,H.T. (1998). Promotion of dendritic growth by CPG15, an activity-induced signaling molecule. *Science* 281, 1863-1866.
- Nguyen,P.V. and Kandel,E.R. (1996). A Macromolecular Synthesis-Dependent Late Phase of Long-Term Potentiation Requiring cAMP in the Medial Perforant Pathway of Rat Hippocampal Slices. *Journal of Neuroscience* 16, 3189-3198.
- Norris,C.M., Blalock,E.M., Chen,K.C., Porter,N.M., and Landfield,P.W. (2002). Calcineurin enhances L-type Ca(2+) channel activity in hippocampal neurons: increased effect with age in culture. *Neuroscience* 110, 213-225.
- O'Brien,R., Xu,D., Mi,R., Tang,X., Hopf,C., and Worley,P. (2002). Synaptically targeted Narp plays an essential role in the aggregation of AMPA receptors at excitatory synapses in cultured spinal neurons. *J Neurosci.* 22, 4487-4498.
- O'Brien,R.J., Xu,D., Petralia,R.S., Steward,O., Huganir,R.L., and Worley,P. (1999). Synaptic clustering of AMPA receptors by the extracellular immediate-early gene product Narp. *Neuron* 23, 309-323.

- O'Donovan,M.J., Chub,N., and Wenner,P. (1998). Mechanisms of spontaneous activity in developing spinal networks. *J Neurobiol.* 37, 131-145.
- Okada,M., Onodera,K., Van Renterghem,C., Sieghart,W., and Takahashi,T. (2000). Functional correlation of GABA(A) receptor alpha subunits expression with the properties of IPSCs in the developing thalamus. *J Neurosci* 20, 2202-2208.
- Oohira,A., Shuo,T., Tokita,Y., Nakanishi,K., and Aono,S. (2004). Neuroglycan C, a brain-specific part-time proteoglycan, with a particular multidomain structure. *Glycoconj.J* 21, 53-57.
- Ozaki,M., Itoh,K., Miyakawa,Y., Kishida,H., and Hashikawa,T. (2004). Protein processing and releases of neuregulin-1 are regulated in an activity-dependent manner. *J Neurochem* 91, 176-188.
- Pak,D.T. and Sheng,M. (2003). Targeted protein degradation and synapse remodeling by an inducible protein kinase. *Science* 302, 1368-1373.
- Pandiella,A., Bosenberg,M.W., Huang,E.J., Besmer,P., and Massague,J. (1992). Cleavage of membrane-anchored growth factors involves distinct protease activities regulated through common mechanisms. *J.Biol Chem.* 267, 24028-24033.
- Pandiella,A. and Massague,J. (1991). Multiple signals activate cleavage of the membrane transforming growth factor-alpha precursor. *J Biol.Chem.* 266, 5769-5773.
- Paul,S., Nairn,A.C., Wang,P., and Lombroso,P.J. (2003). NMDA-mediated activation of the tyrosine phosphatase STEP regulates the duration of ERK signaling. *Nat Neurosci* 6, 34-42.
- Peiretti,F., Bernot,D., Lopez,S., Bonardo,B., Deprez-Beauclair,P., Juhan-Vague,I., and Nalbone,G. (2003). Modulation of PAI-1 and proMMP-9 syntheses by soluble TNFalpha and its receptors during differentiation of the human monocytic HL-60 cell line. *J Cell Physiol* 196, 346-353.
- Peled,A. and Geva,A.B. (1999). Brain Organization and Psychodynamics. *Journal of Psychotherapy Practice and Research* 8, 24-39.
- Penn,A.A., Riquelme,P.A., Feller,M.B., and Shatz,C.J. (1998). Competition in retinogeniculate patterning driven by spontaneous activity. *Science* 279, 2108-2112.
- Phong,M.C., Gutwein,P., Kadel,S., Hexel,K., Altevogt,P., Linderkamp,O., and Brenner,B. (2003). Molecular mechanisms of L-selectin-induced co-localization in rafts and shedding [corrected]. *Biochem.Biophys.Res.Commun.* 300, 563-569.
- Pittenger,C. and Kandel,E.R. (2003). In search of general mechanisms for long-lasting plasticity: Aplysia and the hippocampus. *Philos.Trans.R.Soc.Lond B Biol.Sci.* 358, 757-763.
- Poo,M.M. (2001). Neurotrophins as synaptic modulators. *Nat.Rev.Neurosci.* 2, 24-32.
- Purves,D., Augustine,G.J., Fitzpatrick,D., Katz,L.C., LaMantia,A.-S., and McNamara,J.O. (1997). Electrical signals of nerve cells. In *Neuroscience*, (Sunderland, Massachusetts: Sinauer Associates, Inc), pp. 37-50.

- Quinlan,E.M., Olstein,D.H., and Bear,M.F. (1999). Bidirectional, experience-dependent regulation of N-methyl-D-aspartate receptor subunit composition in the rat visual cortex during postnatal development. *Proc.Natl.Acad.Sci.U.S.A* 96, 12876-12880.
- Ranscht,B. (2000). Cadherins: molecular codes for axon guidance and synapse formation. *Int.J Dev.Neurosci* 18, 643-651.
- Raper,J.A. and Tessier-Lavigne,M. (1999). Growth cone and Axon pathfinding. In Fundamental Neuroscience, M. J. Zigmond, F. E. Bloom, S. C. Landis, J. L. Roberts, and L. R. Squire, eds. Academic Press), pp. 519-546.
- Rathjen,F.G., Wolff,J.M., Frank,R., Bonhoeffer,F., and Rutishauser,U. (1987). Membrane glycoproteins involved in neurite fasciculation. *J Cell Biol.* 104, 343-353.
- Reiss,Y., Kroger,S., Grassi,J., Tsim,K.W., Willbold,E., and Layer,P.G. (1996). Extracellular and asymmetric forms of acetylcholinesterase are expressed on cholinergic and noncholinergic terminal neuropil of the developing chick retina. *Cell Tissue Res.* 286, 13-22.
- Riederer,B.M., Monnet-Tschudi,F., and Honegger,P. (1992). Development and maintenance of the neuronal cytoskeleton in aggregated cell cultures of fetal rat telencephalon and influence of elevated K⁺ concentrations. *J Neurochem* 58, 649-658.
- Robache-Gallea,S., Bruneau,J.M., Robbe,H., Morand,V., Capdevila,C., Bhatnagar,N., Chouaib,S., and Roman-Roman,S. (1997). Partial purification and characterization of a tumor necrosis factor-alpha converting activity. *Eur.J Immunol.* 27, 1275-1282.
- Roerig,B. and Feller,M.B. (2000). Neurotransmitters and gap junctions in developing neural circuits. *Brain Res.Brain Res.Rev* 32, 86-114.
- Ron,D. (2004). Signaling cascades regulating NMDA receptor sensitivity to ethanol. *Neuroscientist*. 10, 325-336.
- Rougon,G. and Hobert,O. (2003). New insights into the diversity and function of neuronal immunoglobulin superfamily molecules. *Annu.Rev Neurosci* 26, 207-238.
- Sahin,U., Weskamp,G., Kelly,K., Zhou,H.M., Higashiyama,S., Peschon,J., Hartmann,D., Saftig,P., and Blobel,C.P. (2004). Distinct roles for ADAM10 and ADAM17 in ectodomain shedding of six EGFR ligands. *The Journal of Cell Biology* 164, 769-779.
- Saitoe,M., Schwarz,T.L., Umbach,J.A., Gunderson,C.B., and Kidokoro,Y. (2001). Absence of junctional glutamate receptor clusters in Drosophila mutants lacking spontaneous transmitter release. *Science* 293, 514-517.
- Saitoe,M., Tanaka,S., Takata,K., and Kidokoro,Y. (1997). Neural activity affects distribution of glutamate receptors during neuromuscular junction formation in Drosophila embryos. *Dev.Biol.* 184, 48-60.
- Salinas,P.C. (2003). Neurobiology: backchat at the synapse. *Nature* 425, 464-467.
- Sandrock,A.W., Jr., Dryer,S.E., Rosen,K.M., Gozani,S.N., Kramer,R., Theill,L.E., and Fischbach,G.D. (1997). Maintenance of acetylcholine receptor number by neuregulins at the neuromuscular junction in vivo. *Science* 276, 599-603.

- Sanes,D.H., Reh,T.A., and Harris,W.A. (2000). Synapse formation and electric function. In Development of the nervous system, D. H. Sanes, T. A. Reh, and W. A. Harris, eds. Academic press), pp. 289-348.
- Sanes,J.R. and Lichtman,J.W. (1999). Development of the vertebrate neuromuscular junction. *Annu.Rev Neurosci* 22, 389-442.
- Sanes,J.R. and Lichtman,J.W. (2001). Induction, assembly, maturation and maintenance of a postsynaptic apparatus. *Nature Reviews Neuroscience* 2, 791-805.
- Sargiacomo,M., Lisanti,M., Graeve,L., Le Bivic,A., and Rodriguez-Boulan,E. (1989). Integral and peripheral protein composition of the apical and basolateral membrane domains in MDCK cells. *J Membr.Biol.* 107, 277-286.
- Schlondorff,J. and Blobel,C.P. (1999). Metalloprotease-disintegrins: modular proteins capable of promoting cell-cell interactions and triggering signals by protein-ectodomain shedding. *J.Cell Sci.* 112 (Pt 21), 3603-3617.
- Schramm,M., Eimerl,S., and Costa,E. (1990). Serum and depolarizing agents cause acute neurotoxicity in cultured cerebellar granule cells: role of the glutamate receptor responsive to N-methyl-D-aspartate. *Proc.Natl.Acad.Sci.U.S.A* 87, 1193-1197.
- Schumacher,S., Volkmer,H., Buck,F., Otto,A., Tarnok,A., Roth,S., and Rathjen,F.G. (1997). Chicken acidic leucine-rich EGF-like domain containing brain protein (CALEB), a neural member of the EGF family of differentiation factors, is implicated in neurite formation. *J.Cell Biol* 136, 895-906.
- Schwartz,D.C. and Hochstrasser,M. (2003). A superfamily of protein tags: ubiquitin, SUMO and related modifiers. *Trends Biochem.Sci.* 28, 321-328.
- Serafini,T. (1999). Finding a partner in a crowd: neuronal diversity and synaptogenesis. *Cell* 98, 133-136.
- Shah,B.H., Farshori,M.P., and Catt,K.J. (2004). Neuropeptide-induced transactivation of a neuronal epidermal growth factor receptor is mediated by metalloprotease-dependent formation of heparin-binding epidermal growth factor. *J.Biol Chem.* 279, 414-420.
- Sharma,S.K., Bagnall,M.W., Sutton,M.A., and Carew,T.J. (2003). Inhibition of calcineurin facilitates the induction of memory for sensitization in Aplysia: requirement of mitogen-activated protein kinase. *Proc.Natl.Acad.Sci.U.S.A* 100, 4861-4866.
- Sheng,M., McFadden,G., and Greenberg,M.E. (1990). Membrane depolarization and calcium induce c-fos transcription via phosphorylation of transcription factor CREB. *Neuron* 4, 571-582.
- Sheng,M., Thompson,M.A., and Greenberg,M.E. (1991). CREB: a Ca(2+)-regulated transcription factor phosphorylated by calmodulin-dependent kinases. *Science* 252, 1427-1430.
- Shi,J., Townsend,M., and Constantine-Paton,M. (2000). Activity-dependent induction of tonic calcineurin activity mediates a rapid developmental downregulation of NMDA receptor currents. *Neuron* 28, 103-114.

- Shirakabe,K., Wakatsuki,S., Kurisaki,T., and Fujisawa-Sehara,A. (2001). Roles of Meltrin beta /ADAM19 in the processing of neuregulin. *J Biol.Chem.* 276, 9352-9358.
- Srour,N., Lebel,A., McMahon,S., Fournier,I., Fugere,M., Day,R., and Dubois,C.M. (2003). TACE/ADAM-17 maturation and activation of sheddase activity require proprotein convertase activity. *FEBS Letters* 554, 275-283.
- Stevens,C.F. (1994). CREB and memory consolidation. *Neuron* 13, 769-770.
- Sunnarborg,S.W., Hinkle,C.L., Stevenson,M., Russell,W.E., Raska,C.S., Peschon,J.J., Castner,B.J., Gerhart,M.J., Paxton,R.J., Black,R.A., and Lee,D.C. (2002). Tumor necrosis factor-alpha converting enzyme (TACE) regulates epidermal growth factor receptor ligand availability. *J.Biol.Chem.* 277, 12838-12845.
- Swanson,L.W., Lufkin,T., and Colman,D.R. (1999). Organization of nervous systems. In Fundamental neuroscience, M. J. Zigmond, F. E. Bloom, S. C. Landis, J. L. Roberts, and L. R. Squire, eds. Academic press), pp. 9-37.
- Tanokura,M. and Yamada,K. (1986). Effects of trifluoperazine on calcium binding by calmodulin. Heat capacity and entropy changes. *J Biol.Chem.* 261, 10749-10752.
- Tao,X., West,A.E., Chen,W.G., Corfas,G., and Greenberg,M.E. (2002). A calcium-responsive transcription factor, CaRF, that regulates neuronal activity-dependent expression of BDNF. *Neuron* 33, 383-395.
- Tartaglia,N., Du,J., Tyler,W.J., Neale,E., Pozzo-Miller,L., and Lu,B. (2001). Protein synthesis-dependent and -independent regulation of hippocampal synapses by brain-derived neurotrophic factor. *J Biol.Chem.* 276, 37585-37593.
- Thabard,W., Collette,M., Bataille,R., and Amiot,M. (2001). Protein kinase C delta and eta isoenzymes control the shedding of the interleukin 6 receptor alpha in myeloma cells. *Biochem.J.* 358, 193-200.
- Tian,N. and Copenhagen,D.R. (2003). Visual stimulation is required for refinement of ON and OFF pathways in postnatal retina. *Neuron* 39, 85-96.
- Towbin,H., Staehelin,T., and Gordon,J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc.Natl.Acad.Sci.U.S.A* 76, 4350-4354.
- Tsakadze,N.L., Sen,U., Zhao,Z., Sithu,S.D., English,W.R., and D'Souza,S.E. (2004). Signals mediating cleavage of intercellular adhesion molecule-1. *AJP - Cell Physiology* 287, C55-C63.
- Turrigiano,G., Abbott,L.F., and Marder,E. (1994). Activity-dependent changes in the intrinsic properties of cultured neurons. *Science* 264, 974-977.
- Umata,T., Hirata,M., Takahashi,T., Ryu,F., Shida,S., Takahashi,Y., Tsuneoka,M., Miura,Y., Masuda,M., Horiguchi,Y., and Mekada,E. (2001). A dual signaling cascade that regulates the ectodomain shedding of heparin-binding epidermal growth factor-like growth factor. *J Biol.Chem.* 276, 30475-30482.

- Vallano,M.L., Lambolez,B., Audinat,E., and Rossier,J. (1996). Neuronal activity differentially regulates NMDA receptor subunit expression in cerebellar granule cells. *Journal of Neuroscience* 16, 631-639.
- Verstreken,P. and Bellen,H.J. (2002). Meaningless minis? Mechanisms of neurotransmitter-receptor clustering. *Trends Neurosci* 25, 383-385.
- von Bohlen,O., Halbach, and Dermietzel,R. (1999). Methoden der Neurohistologie. Spektrum Akademisher Verlag GmbH), pp. 43-46.
- Waltereit,R., Dammermann,B., Wulff,P., Scafidi,J., Staubli,U., Kauselmann,G., Bundman,M., and Kuhl,D. (2001). Arg3.1/Arc mRNA Induction by Ca²⁺ and cAMP Requires Protein Kinase A and Mitogen-Activated Protein Kinase/Extracellular Regulated Kinase Activation. *Journal of Neuroscience* 21, 5484-5493.
- Watanabe,E., Maeda,N., Matsui,F., Kushima,Y., Noda,M., and Oohira,A. (1995). Neuroglycan C, a novel membrane-spanning chondroitin sulfate proteoglycan that is restricted to the brain. *J Biol.Chem.* 270, 26876-26882.
- Weiss,S. (1988). Forskolin attenuates the evoked release of [³H]GABA from striatal neurons in primary culture. *Brain Res.* 463, 182-186.
- Weskamp,G., Schlondorff,J., Lum,L., Becherer,J.D., Kim,T.W., Saftig,P., Hartmann,D., Murphy,G., and Blobel,C.P. (2004). Evidence for a critical role of the tumor necrosis factor alpha convertase (TACE) in ectodomain shedding of the p75 neurotrophin receptor (p75NTR). *J Biol.Chem.* 279, 4241-4249.
- West,A.E., Chen,W.G., Dalva,M.B., Dolmetsch,R.E., Kornhauser,J.M., Shaywitz,A.J., Takasu,M.A., Tao,X., and Greenberg,M.E. (2001). Calcium regulation of neuronal gene expression. *Proc.Natl.Acad.Sci.U.S.A* 98, 11024-11031.
- White,J.M. (2003). ADAMs: modulators of cell-cell and cell-matrix interactions. *Curr.Opin.Cell Biol.* 15, 598-606.
- Wong,R.O., Chernjavsky,A., Smith,S.J., and Shatz,C.J. (1995). Early functional neural networks in the developing retina. *Nature* 374, 716-718.
- Xu,D., Hopf,C., Reddy,R., Cho,R.W., Guo,L., Lanahan,A., Petralia,R.S., Wenthold,R.J., O'Brien,R.J., and Worley,P. (2003). Narp and NP1 form heterocomplexes that function in developmental and activity-dependent synaptic plasticity. *Neuron* 39, 513-528.
- Yamagata,M., Sanes,J.R., and Weiner,J.A. (2003). Synaptic adhesion molecules. *Curr.Opin.Cell Biol.* 15, 621-632.
- Yang,X., Arber,S., William,C., Li,L., Tanabe,Y., Jessell,T.M., Birchmeier,C., and Burden,S.J. (2001). Patterning of muscle acetylcholine receptor gene expression in the absence of motor innervation. *Neuron* 30, 399-410.
- Yardin,C., Terro,F., Lesort,M., Esclaire,F., and Hugon,J. (1998). FK506 antagonizes apoptosis and c-jun protein expression in neuronal cultures. *Neuroreport* 9, 2077-2080.

- Zaidi,N.F., Berezovska,O., Choi,E.K., Miller,J.S., Chan,H., Lilliehook,C., Hyman,B.T., Buxbaum,J.D., and Wasco,W. (2002). Biochemical and immunocytochemical characterization of calsenilin in mouse brain. *Neuroscience* 114, 247-263.
- Zhang,J., Kornecki,E., Jackman,J., and Ehrlich,Y.H. (1988). ATP secretion and extracellular protein phosphorylation by CNS neurons in primary culture. *Brain Res.Bull.* 21, 459-464.
- Zhang,L.I. and Poo,M.M. (2001). Electrical activity and development of neural circuits. *Nat.Neurosci.* 4 Suppl, 1207-1214.
- Zheng,J.Q. (2000). Turning of nerve growth cones induced by localized increases in intracellular calcium ions. *Nature* 403, 89-93.
- Zhu,W.J., Vicini,S., Harris,B.T., and Grayson,D.R. (1995). NMDA-mediated modulation of gamma-aminobutyric acid type A receptor function in cerebellar granule neurons. *J Neurosci* 15, 7692-7701.