

9 Literaturverzeichnis

Akers R. M., Beal W. E., McFadden T. B., Capuco A. V. (1990)

Morphometric analysis of involuting bovine mammary tissue after 21 or 42 days on non-suckling.

J Anim Sci. 11: 3604-3613.

Annen E. L., McGuire M. A., Vicini J. L., Collier R. J. (2003)

Effect of Posilac⁷ (bST) and dry period management strategy on milk yield.

J. Dairy Sci. 86 (Suppl. 1): 154(Abstr.).

Bachman K. C. (2004)

Dry period length: How short is long enough?

National Mastitis Council Annual Meeting Proceedings 2004.

Bachman K. C. (2002)

Milk production of dairy cows treated with estrogen at the onset of a short dry period.

J. Dairy Sci. 85: 797-803.

Barkema H. W., SchukkenY. H., Lam T. J., Beiboer M. L., Benedictus G., Brand A.

(1999)

Management practices associated with the incidence rate of clinical mastitis.

J Dairy Sci. 82: 1643-1654.

Bascom S. S., Young A. J. (1998)

A summary of the reasons why farmers cull cows.

J Dairy Sci. 81: 2299-2305.

Berry E. A., Hillerton J. E. (2002)

The effect of an intramammary teat seal on new intramammary infections.

J Dairy Sci. 85: 2512-2520.

Berry E. A., Johnston W. T., Hillerton J. E. (2003)

Prophylactic effects of two selective dry cow strategies accounting for interdependence of quarter.

J Dairy Sci. 86: 3912-3919.

Binde M., Bakke H. (1984)

Relationships between teat characteristics and udder health. A field survey.
Nord Vet Med. 36 (3-4): 111-116.

Bitman J., Wood D. L., Bright S. A., Miller R. H. (1988)

Lipid composition of bovine teat canal keratin.
J Dairy Sci. 71 (5): 1389-1395.

Boyum A., Wilk P., Gustavsson E., Veiby O. P., Reseland J., Haugen A. H.,

Optad P. K. (1996)

The effect of strenuous exercise and calorie deficiency and sleep deprivation on
white blood cells, plasma immunoglobulins and cytokines.

Scand. J. Immunol. 43: 228-235.

Bradley A. J., Green M. J. (2000)

A study of the incidence and significance of intramammary enterobacterial
infections acquired during dry period.

J Dairy Sci. 83: 1957-1965.

Bradley A. J., Green M. J. (2002)

The pint plan - Managing Dry Cows to Maximise Milk Quality.
Proceedings of the British Mastitis Conference 2002 Brockworth: 89-90.

Bramley A. J. (1985)

The sources of mastitis pathogens for dairy herd and their control.
Kieler Milchwirtschaftliche Forschungsberichte 37: 375-385.

Bright S. A., Bitman J., Capuco A. V., Wood D. L., Miller R. H. (1990)

Methods of collection and lipid composition of teat canal keratin in dry and
lactating cows.

J Dairy Sci. 73 (1): 98-106.

Browning J. W., Mein G. A., Barton M., Nicholls T. J., Brightling P. O. (1990)

Effects of antibiotic therapy at drying off on mastitis in the dry period and early
lactation.

Aust Vet J 67: 440.

Bundesgesetzblatt (2000)

Verordnung über Hygiene und Qualitätsanforderungen an Milch und Erzeugnisse auf Milchbasis.

Bundesgesetzblatt Teil I Nr. 36 vom 31.Juli 2000.

Canena J., Reis J., Pinto A. S., Santos A. M., Leitao J., Pinheiro T., Quina M. G. (1998)

Distribution of bismuth in the rat after oral dosing with ranitidine bismuth citrate and bismuth subcitrate.

J Pharm Pharmacol. 50: 279-283.

Capuco A. V., Akers R. M., Smith J. J. (1997)

Mammary growth in Holstein cows during the dry period: quantification of nucleic acids and histology.

J Dairy Sci. 80 (3): 477-487.

Capuco A. V., Bright S. A., Pankey J. W., Wood D. L., Miller R. H., Bitman J. (1992)

Increased susceptibility to intramammary infection following removal of teat canal keratin.

J Dairy Sci. 75 (8): 2126-2130.

Capuco A. V., Wood D. L., Bright S. A., Miller R. H., Bitman J. (1990)

Regeneration of teat canal keratin in lactating dairy cows.

J Dairy Sci. 73 (7): 1745-1750.

Capuco A. V., Wood D. L., Quast J. W. (2000)

Effects of teatcup liner tension on teat canal keratin and teat condition in Cows.

J Dairy Res. 67: 319-327.

Comalli M. P., Eberhart R. J., Griet L. C. Jr., Rothenbacher H. (1984)

Changes in the microscopic anatomy of the bovine teat canal during mammary Involution.

Am J Vet Res 45: 2236-2242.

Concha C. (1986)

Cell types and their immunological functions in bovine mammary tissues and secretions - a review of literature.

Nordisk Veterinaermedicin. 38: 257-272.

Cook N. B., Bennett T. B., Emery K. M., Nordlund K. V. (2002)

Monitoring nonlactating cow intramammary infection dynamics using DHI somatic cell count data.

J Dairy Sci. 85: 1119-1126.

Cook N. B., Wilkinson A., Gajewski K., Weigel D., Sharp P., Pionek D. (2004)

The Prevention of new intramammary infections during the dry period when using an internal teat sealant in conjunction with a dry cow antibiotic.

National Mastitis Council Annual Meeting Proceedings 2004: 292-293.

Coppock C. E., Everett R. W., Natzke R. P., Ainslie H. R. (1974)

Effect of dry period length on Holstein milk production and selected disorders at parturition.

J. Dairy Sci. 57: 712-718.

Cousins C. L., Higgs T. M., Jackson E. R., Neave F. K., Dodd F. H. (1980)

Susceptibility of the bovine udder to bacterial infection in the dry period.

J Dairy Res. 47: 11-18.

Craven N. (1991)

Is treatment necessary?

British Mast. Conf., Stoneleigh: 42-50.

Deutsche Veterinärmedizinische Gesellschaft (1994)

Leitlinien zur Bekämpfung der Mastitis des Rindes als Herdenproblem 3. ed.

DVG Verlag, Gießen.

Dingwell R. T., Duffield T. F., Leslie K. E., Keefe G. P., DesCoteaux L., Kelton D. F., Lissemore K. D., Schukken Y. H., Dick P., Bagg R. (2002)

The efficacy of intramammary tilmicosin at drying-off, and other risk factors for the prevention of new intramammary infections during the dry period.

J Dairy Sci. 85: 3250-3259.

- Dingwell R. T., Kelton D. F., Leslie K. E., Edge V. L. (2001)
Deciding to dry-off: does level of production matter?
National Mastitis Council Annual Meeting Proceedings 2001: 69-79.
- Dingwell R. T., Leslie K. E., Duffield T. F., Schukken Y. H., DesCoteaux L., Keefe G. P., Kelton D. F., Lissemore K. D., Shewfelt W., Dick P., Bagg R. (2003)
Efficacy if intramammary Tilmicosin and risk factors for cure of *Staphylococcus aureus* infection in the dry period.
J Dairy Sci. 86: 159-168.
- Dingwell R. T., Leslie K. E., Sargeant J. M., Schukken Y. H., Timms L. L. (2001)
Impact of milk production and important management factors on the process of dry-off in lactating dairy cows.
Dairy day 2001, Hutchinson, Report of Progress: 26-30.
- Dingwell R. T., Leslie K. E., Schukken Y. H., Sargeant J. M., Timms L. L., Duffield T. F., Keefe G. P., Kelton D. F., Lissemore K. D., Conklin J. (2004)
Association of cow and quarter-level factors at drying-off with new intramammary infections during the dry period.
Prev Vet Med. 63: 75-89.
- Dingwell R. T., Leslie K. E., Schukken Y. H., Sargeant J. M., Timms L. L. (2003)
Evaluation of the California mastitis test to detect an intramammary infection with a major pathogen in early lactation dairy cows.
Can Vet J. 44: 413-415.
- Dingwell R. T., Leslie K. E., Schukken Y. H., Sargeant J. M., Timms L. L., Duffield T. F., Keefe G. P., Kelton D. F., Lissemore K. D., Conklin J. (2004)
Association of cow and quarter-level factors at drying-off with new intramammary infections during the dry period.
Prev Vet Med. 63: 75-89.
- Dohoo I. R., Meek A. H., Martin S. W., Barnum D. A. (1981)
Use of total and differential somatic cell counts from composite milk samples to detect mastitis in individual cows.
Can J Comp Med. 45: 8-14.

Döpfer D., Schukken Y. H., Metzner R. M., Petersen D. (1993)

Betreuungsstrategien zur Sanierung von Milchviehbetrieben mit hohem Zellgehalt
in der Tankmilch.

Prakt. Tierarzt 74, Coll. Vet. XXIV: 70-72.

Eberhart R. J. (1986)

Management of dry cows to reduce mastitis.

J. Dairy Sci. 69: 1721-1732.

Enevoldsen C., Sørensen J. T. (1992)

Effects of dry period length on clinical mastitis and other major clinical health
disorders.

J Dairy Sci. 75: 1007-1014.

Erskine R. J., Bartlett P. C., Crawshaw P. C., Gombas D. M. (1994)

Efficacy of intramuscular oxytetracycline as adry cow treatment for
Staphylococcus aureus mastitis.

J Dairy Sci. 77: 3347.

Esslemont R. J., Kossaibati M. A. (1997)

Culling in 50 dairy herds in England.

Vet Rec. 140: 36-39.

Falkenberg U., Tenhagen B. A., Baumgartner B., Heuwieser W. (2002)

Prevention of intramammary infections in dairy cows by the use of a premilking
teat dip method with a foaming iodophor dip agent.

Dtsch. Tierarztl. Wochenschr 109: 423-427.

Fehlings K., Deneke J. (2000)

Hygienemanagement im Melkbereich aus tierärztlicher Sicht.

Tierärztl. Umschau. 55: 451-456.

Ferens W. A., Davis W. C., Hamilton M. J., Park Y. H., Deobald C. F., Fox L.,
Bohach G. (1998)

Activation of bovine lymphocyte subpopulations by staphylococcal enterotoxin C.
Infect Immun. 66: 573-580.

Fetrow J., Stewart S., Eicker S., Farnsworth R., Bey R. (2000)

Mastitis an economic consideration.

National Mastitis Council Annual Meeting Proceedings 2000: 3-28.

Filar J., Kandefer-Szerszen M., Szuster-Ciesielska A., Rzeski W. (1992)

Effects of cold treatment and ketosis induced by starvation on interferon production in leukocytes of lactating cows.

Dtsch Tierarztl Wochenschr. 99: 210-213.

Fox L. K., Hancock D. D., Weems C. W., Toma W., Chang E. (1997)

The effect of intramammary antibiotic therapy at calving on udder health traits.

J Dairy Sci. 70 (8): 1696-1700.

Funk D. A., Freeman A. E., Berger P. J. (1987)

Effects of previous days open, previous days dry, and present days open on lactation yield.

J Dairy Sci. 70: 2366-2373.

Gane E., Sutton M. M., Pybus J., Hamilton I. (1996)

Hepatic and cerebrospinal fluid accumulation of aluminium and bismuth in volunteers taking short course anti-ulcer therapy.

J Gastroenterol Hepatol. 111: 911-915.

Godden S., Rapnicki P., Stewart S., Fetrow J., Johnson A., Bey R., Farnsworth R. (2003)

Effectiveness of an internal teatseal in the prevention of new intramammary infections during the dry and early-lactation periods in dairy cows when used with a dry cow intramammary antibiotic.

J Dairy Sci 86: 3899-3911.

Godden S., Reneau J., Farnsworth R., Bey R. (2005)

Giving Dry Cow Mastitis the Boot.

http://www.extension.umn.edu/dairy/05dairydays/dry_cow_mastitis-DairyDays05.pdf.

- Green M. J., Green L. E., Medley G. F., Schukken Y. H., Bradley A. J. (2002)
Influence of dry period bacterial intramammary infection on clinical mastitis in
dairy cows.
J Dairy Sci 85: 2589–2599.
- Gulay M. S., Hayen M. J., Bachman K. C., Beloso T., Liboni M., Head H. H. (2003)
Milk production and feed intake of Holstein cows given short (30-d) or normal
(60-d) dry periods.
J Dairy Sci. 86: 2030-2038.
- Gulay M. S., Hayen M. J., Bachman K. C. (2003)
Milk production from Holstein halfudders after concurrent 30- and 70-d dry
periods.
J. Dairy Sci. 86 (Suppl. 1): 154(Abstr.).
- Gumen A., Rastani R. R., Grummer R. R., Wiltbank M. C. (2003)
Effects of varying dry period length and prepartm diet on reproduction in dairy
cattle.
J. Dairy Sci. 86 (Suppl.1): 239(Abstr.).
- Hamann J. (1987)
Effect of machine milking on teat condition.
IDF Bulletin 215: 33-53.
- Hamann J. (1987)
The role of machine milking factors in the aetiology & pathogenesis of mastitis.
Hohenheimer Arbeiten, Research on Milk Production: 22-56.
- Hamann J. (1996)
Strategie der Mastitisprävention – eine Übersicht – .
Deutsche Veterinärmedizinische Gesellschaft e.V., Tagung der Fachgruppe
„Milchhygiene, Arbeitskreis „Eutergesundheit“: 1- 10.
- Hamann J. (2001)
Relationships between somatic cell count and milk composition.
IDF World Summit 2001, Conference Proceedings.

Hamann J., Østeras O. (1994)

Special aspects. Teat tissue reactions to machine milking and new infection risk.
Bull. Int. Dairy Fed. 297: 35–41.

Hamann J., Claessens I., Krömker V., Nogai K. (1998)

Kompendium der Milchhygiene.
Zentrumsabteilung Hygiene und Technologie der Milch, Zentrum für
Lebensmittelwissenschaften der Tierärztlichen Hochschule Hannover.

Hamann J., Reichmuth J. (1990)

Compensatory milk production within the bovine udder: Effects of short-term non-milking of single quarters.
Journal of Dairy Research. 57 (1): 17-22.

Harmon R. J. (1994)

Physiology of mastitis and factors affecting somatic cell counts.
J Dairy Sci. 77: 2103-2112.

Harmon R. J., Reneau J. K. (1993)

Factors affecting somatic cell counts in milk.
National Mastitis Council Annual Meeting Proceedings 1993: 48-57.

Hassan Z., Daniel R. C., O'Boyle D., Frost A. J. (1999)

Effects of dry cow intramammary therapy on quarter infections in the dry period.
Vet Rec. 145: 635-639.

Heinz M., Michel G. (1991)

The structure of the bovine mammary gland during the dry period with special reference to the process of involution. 2. Electron microscopic studies.
Anat Histol Embryol. 20: 193-204.

Hillerton J. E., West J. G., Shearn M. F. (1992)

The cost of summer mastitis.
Vet Rec. 131: 315-317.

Hoblet K. H., Schnitkey G. D., Arbaugh D., Hogan J. S., Smith K. L., Schoenberger P. S., Todhunter D. A., Hueston W. D., Pritchard D. E., Bowman G. L. (1991)

Costs associated with selected preventive practices and with episodes of clinical mastitis in nine herds with low somatic cell count.

J Am Vet Med Assoc 199: 190-196.

Hogan J. S., Duthie A. H., Pankey J. W. (1986)

Fatty acid composition of bovine teat canal keratin.

J Dairy Sci. 69 (9): 2424-2427.

Hogan J. S., Smith K. L. (1999)

Environmental streptococcal mastitis.

Research and Reviews: Dairy Special Circular: 169-175. Bulletin Extension Research, Ohio State University.

Hogan J. S., Smith K. L., Todhunter D. A., Schoenberger P. S. (1988)

Growth responses of environmental mastitis pathogens to long-chain fatty acids.

J Dairy Sci. 71 (1): 245-249.

Hogan J. S., Smith K. L., Todhunter D. A., Schoenberger P. S. (1995)

Efficacy of a barrier teat dip containing 0,55 % chlorhexidine for prevention of bovine mastitis.

J Dairy Sci. 78: 2502-2506.

Hurley W. L. (1995)

Mammary Gland Involution and the Dry Period.

<http://classes.aces.uiuc.edu/AnSci308/involution.html>.

Hurley W. L. (1989)

Mammary function during involution.

J. Dairy Sci. 72: 1637-1646.

Hurley W. L., Aslam M., Hegarty H. M., Morkoc A. (1994)

Synthesis of lactoferrin and casein by explants of bovine mammary tissue.

Cell. Biol. Int. 18: 629-637.

Huxley J. N., Green M. J., Bradley A. J. (2002)

C. bovis – friend or foe?

Proc British mast Conf., Brockworth: 94-95.

Huxley J. N., Green M.. J., Green L. E., Bradley A. J. (2002)

Evaluation of the efficacy of an internal teat sealer during the dry period.

J Dairy Sci. 85: 551-561.

Kandefer-Szerszen M., Filar J., Szuster-Ciesielska A., Rzeski W. (1992)

Suppression of interferon response of bovine leukocytes during clinical and subclinical ketosis in lactating cows.

Dtsch. Tieraerztl. Wochenschr. 99: 440-443.

Keown J. F., Everett. R. W. (1986)

Effect of days carried calf, days dry and weight of first calf heifers on yield.

J. Dairy Sci. 69: 1891-1896.

Klucinski W., Degorski W., Miernik-Degorska E., Targowski S., Winnicka A. (1988)

Effect of ketone bodies on the phagocytic activity of bovine milk macrophages and polymorphonuclear leukocytes.

J. Vet. Med. A 35: 632-639.

Koch K. M., Kerr B. M., Gooding A. E., Davis I. M. (1996)

Pharmacokinetics of bismuth and ranitidine following multiple doses of ranitidine bismuth citrate.

Br J Clin Pharmacol. 42: 207-211.

Krömker V., Hamann J. (2001)

Mastitisfrüherkennung.

Milchpraxis 2/2001: 78-80.

Krömker V., Pfannenschmidt F. (2003)

Interner Zitzenversiegler.

Milchpraxis 3/2003: 124-126.

Literaturverzeichnis

Kutila T., Pyorala S., Kaartinen L., Isomaki R., Vahtola K., Myllykoski L., Saloniemi H. (2003)

Lactoferrin and citrate concentrations at drying-off and during early mammary involution of dairy cows.

J Vet Med A Physiol Pathol Clin Med. 50: 350-353.

Lacy-Hulbert S. J., Hillerton J. E. (1995)

Physical characteristics of the bovine teat canal and their influence on susceptibility to streptococcal infection.

J Dairy Res. 62: 395-404.

Lacy-Hulbert S. J., Woolford M. W. (2000)

Keratin plug in the teat canal after drying off. Intern. Symposium on Immunology of ruminant mammary gland.

IDF, Stresa.: 193-194.

Leslie K. E., Duffield T. F., Schukken T. H., LeBlanc S. J. (2000)

The influence of negative energy balance on udder health.

Proc. Regional Meeting of the National Mastitis Council: 25-33.

Loiseau P., Henry P., Jallon P., Legroux M. (1976)

Iatrogenic myoclonic encephalopathies caused by bismuth salts.

J Neurol Sci. 27: 133-143.

Kuhn M. T., Hutchison J. L., Norman H. D. (2005)

Characterization of Days Dry for United States Holsteins.

J. Dairy Sci. 88: 1147-1155.

Makuza S. M., McDaniel B. T. (1996)

Effects of days dry, previous days open, and current days open on milk yields of cows in Zimbabwe and North Carolina.

J. Dairy Sci. 79: 702-709.

Marshall, Edmondson (1993)

G3653, Using the California Mastitis Test.

<http://muextension.missouri.edu/explore/agguides/dairy/g03653.htm>.

Maunsell F. P., Morin D. E., Constable P. D., Hurley W. L., McCoy G. C., Kakoma I., Isaacson R. E. (1998)

Effects of mastitis on the volume and composition of colostrum produced by Holstein cows.

J Dairy Sci. 81: 1291-1299.

McAuliffe O., Ryan M. P., Ross R. P., Hill C., Breeuwer P., Abee T. (1998)

Lacticin 3147, a Broad-Spectrum Bacteriocin Which Selectively Dissipates the Membrane Potential.

Appl Environ Microbiol 64: 439-445.

Meaney W. J., Twomey D. P., Flynn J., Hill C., Ross R. P. (2001)

The use of a bismuth-based teat seal and the bacteriocin lacticin 3147 to prevent dry period mastitis in dairy cows.

Proc of the British Mastitis Conference, Garstang: 24-32.

Mein G.A., Williams D.M., Reinemann D. J. (2003)

Effects of milking on teat-end hyperkeratosis: 1. Mechanical forces applied by the teatcup liners and responses of the teat.

National Mastitis Council Annual Meeting Proceedings 2003: 114-123.

Mein G. A., Thompson P. D. (1993)

Milking the 30,000 pound herd.

J Dairy Sci. 76: 3294-3300.

Michel G., Heinz M.

The structure of the mammary gland of cattle during the dry period with special reference to the involutional process. 1. Light microscopic studies.

Anat Histol Embryol. 20 (2): 101-110.

Mielke H., Koblenz, C., Beuche W. (1991)

Verhalten der somatischen Zellen im Eutersekret während der Trockenperiode bei eutergesunden und euterkranken Kühen und Systematisierung der Phasen der Trockenperiode.

Der prakt. Tierarzt 72: 785-798.

Literaturverzeichnis

- Miller R. H., Bitman J., Bright S. A., Wood D. L., Capuco A. V. (1992)
Effect of clinical and subclinical mastitis on lipid composition of teat canal keratin.
J Dairy Sci. 75: 1436-1442.
- Monti G., Tenhagen B. A., Heuwieser W. (1999)
Culling policies in dairy herds.
Zentralbl Veterinarmed A. 46: 1-11.
- Mullarky I. K., Su C., Frieze N., Park Y. H., Sordillo L. M. (2001)
Staphylococcus aureus agr Genotypes with Enterotoxin Production Capabilities
Can Resist Neutrophil Bactericidal Activity.
Infect Immun. 69: 45-51.
- National Mastitis Council (1990)
Procedures for collecting milk samples
<http://www.nmconline.org/sampling.htm>
- Neave F. K., Dodd F. H., Kingwill R. G. (1966)
A method for controlling udder disease.
Vet Rec. 78: 521-523.
- Neijenhuis F. (1998)
Teat end callosity classification system.
Proc. 4th Intl. Dairy Housing Conference, ASAE, St Louis: 117 –123.
- Neijenhuis F., Barkema H. W., Hogeveen H., Noordhuizen J. P. (2000)
Classification and longitudinal examination of callused teat ends in dairy cows.
J Dairy Sci. 83: 2795-2804.
- Neijenhuis F., Barkema H. W., Hogeveen H., Noordhuizen J. P. (2001)
Relationship between teat-end callosity and occurrence of clinical mastitis.
J Dairy Sci. 84: 2664-2672.
- Nickerson S. C. (1989)
Immunological aspects of mammary involution.
J Dairy Sci. 72: 1665-1678.
- Nickerson S. C., Watts J. L., Boddie R. L., Pankey J. W. (1986)
Evaluation of 0.5 % and 1 % iodophor teat dips on commercial dairies.

J Dairy Sci. 69: 1693-1698.

Ohge H., Furne J. K., Springfield J., Sueda T., Madoff R. D., Levitt M. D. (2003)

The effect of antibiotics and bismuth on fecal hydrogen sulfide and sulfate-reducing bacteria in the rat.

FEMS Microbiol Lett. 228: 137-1342.

Ohnstad I. C., Mein G. A., Neijenhuis F., Hillerton J. E., Baines J. R (2003)

Assessing The Scale of Teat-End Problems and Their Likely Causes.

National Mastitis Council Annual Meeting Proceedings 2003: 128-135.

Oliver J., Dodd F. H., Neave F. K. (1956)

Udder Infections in the Dry Period IV

The relationship between the new infection rate in the early dry period and the daily milk yield at drying off when lactation was ended by either intermittent or abrupt cessation of milking.

J. Dairy Research 23: 204-210.

Östensson K., Hageltorn M., Aström G. (1988)

Differential cell counting in fraction-collected milk from dairy cows.

Acta Vet Scand. 29: 493-500.

Østeras O., Edge V. L. (2000)

Factors prior to dry period associated with high and low levels of cow milk somatic cell counts in next lactation.

Acta Vet Scand. 41: 63-77.

Østeras O., Edge V. L., Martin S. W. (1999)

Determinants of success or failure in the elimination of major mastitis pathogens in selective dry cow therapy.

J Dairy Sci. 82: 1221-1231.

Østeras O., Edge V. L., Martin S. W. (1999)

Determinants of success or failure in the elimination of major mastitis pathogens in selective dry cow therapy.

J Dairy Sci. 82: 1221-1231.

Paulrud C. O. (2005)

Basic concepts of the bovine teat canal.

Vet Res Commun. 29: 215-245.

Paulrud C. O., Morten Rasmussen D. (2001)

Dependence of bovine teat canal (Ductus papillaris mammae) length and diameter on keratin retrieval.

National Mastitis Council Annual Meeting Proceedings 2001: 255-257.

Pearson J. K. (1950)

The use of penicillin in the prevention of *C. pyogenes* infection in the non-lactating udder.

Vet.Rec. 62: 166-168.

Pearson J. K. (1951)

Further experiments in the use of penicillin in the prevention of *C. pyogenes* infection in the non-lactating udder.

Vet.Rec. 63: 215-220.

Pedersen L. H., Stoltenberg M., Ernst E., West M. J. (2003)

Leydig cell death in rats exposed to bismuth subnitrate.

J Appl Toxicol. 23: 235-238.

Postle D. S., Natzke R. P. (1974)

Efficacy of antibiotic treatment in the bovine udder as determined from field Studies.

Vet Med Small Anim Clin. 69: 1535-1539.

Pyörälä S., Taponen S., Jantunen A., Pyörälä E. (2000)

Efficacy of targeted 5-day parenteral and intramammary treatment of clinical *staphylococcus aureus* mastitis caused by penicillin-susceptible or penicillin-resistant bacterial strain.

Proc. World Buiatrics Congress, Punta del Esta: 6501-6504.

Rajala-Schultz P. J., Hogan J. S., Smith K. L. (2005)

Short communication: association between milk yield at dry-off and probability of intramammary infections at calving.

J Dairy Sci. 88 (2): 577-579.

Rao N., Brown P. W., Yerino P., Chang J., Hwang K. K. (1997)

Comparative absorption of bismuth in Sprague-Dawley rats following oral administration of preparations containing bismuth sucrose octasulfate, bismuth subsalicylate, and bismuth subcitrate.

Biopharm Drug Dispos. 18: 1-8.

Rastani R. R., Grummer R. R. (2003)

Shorter dry periods look good.

Hoard's Dairyman: 599.

Rastani R. R., Grummer R. R., Bertics S. J., Gumen A., Wiltbank M. C., Mashek D.

G., Schwab M. C. (2005)

Reducing dry period length to simplify feeding transition cows: milk production, energy balance, and metabolic profiles.

J Dairy Sci. 88: 1004-1014.

Redetzky R., Hamann J. (2003)

Möglichkeiten und Grenzen für die Anwendung von Teat Sealern zur Mastitisbekämpfung.

Praktischer Tierarzt 84: 530-538.

Rejman J. J., Hurley W. L., Bahr J. M. (1989)

Enzyme-linked immunosorbent assays of bovine lactoferrin and a 39-kilodalton protein found in mammary secretions during involution.

J Dairy Sci. 72: 555-560.

Rindsig R. B., Rodewald R. G., Smith A. R., Spahr S. L. (1978)

Complete versus selective dry cow therapy for mastitis control.

J Dairy Sci. 61: 1483-1497.

Ross J. F., Switzer R. C., Poston M. R., Lawhorn G. T. (2002)

Distribution of bismuth in the brain after intraperitoneal dosing of bismuth subnitrate in mice: implications for routes of entry of xenobiotic metals into the brain.

Brain Res. 725: 137-154.

Ruegg P. L., Sekito L. (2004)

Test characteristics and comparison of methods used to detect subclinical mastitis.

National Mastitis Council Annual Meeting Proceedings 2004: 361-362.

Ryan M. P., Flynn J., Hill C., Ross R. P., Meaney W. J. (1999)

The natural food grade inhibitor, lacticin 3147, reduced the incidence of mastitis after experimental challenge with *Streptococcus dysgalactiae* in nonlactating dairy cows.

J Dairy Sci. 82: 2625-2631.

Ryan M. P., Meaney W. J., Ross R. P., Hill C. (1998)

Evaluation of Lacticin 3147 and a Teat Seal Containing This Bacteriocin for Inhibition of Mastitis Pathogens.

Appl Environ Microbiol. 64: 2287-2290.

Ryan M. P., Rea M. C., Hill C., Ross R. P. (1996)

An application in cheddar cheese manufacture for a strain of *Lactococcus lactis* producing a novel broad-spectrum bacteriocin, lacticin 3147.

Appl. Environ. Microbiol. 62: 612-619.

Schairer M. L. (2001)

Estrogen treatments for the initiation of dryoff in dairy cows.

Master's thesis. Univ. Florida, Gainesville.

Schukken Y. H. (2002)

Future Mastitis control strategies for the dry cow.

World Buiatrics Congress 2002.

Schukken Y. H., Tikovsky I., Wilson D., Østeras O. (2001)

Factors affecting the success of antibiotic treatment at dry off.

National Mastitis Council Annual Meeting Proceedings 2001: 80-87.

Schukken Y. N., Vanvliet J. Vandegeer D., Grommers F. J. (1993)

A randomized blind trial on dry cow antibiotic infusion in a low somatic cell count herd.

J Dairy Sci 76: 2925-2930.

- Schukken Y., Tikovsky L., Wilson D., Østerås O. (2001)
Factors affecting the success of antibiotic treatment at dry off.
National Mastitis Council Annual Meeting Proceedings 2001: 80-87.
- Senft B., Neudecker J. (1991)
Defense mechanisms of the bovine mammary gland.
Tierarztl Prax. 19: 357-363.
- Shearn M. F., Hillerton J. E. (1996)
Hyperkeratosis of the teat duct orifice in the dairy cow.
J Dairy Res. 63 (4): 525-532.
- Sinkevitch M. G., Barto P. B., Bush L. J., Wells M. E., Adams G. D. (1974)
Effectiveness of antibiotic infusion at drying off in preventing new mastitis infections in cows.
Bovine Pract 9: 43-46.
- Slikkerveer A., de Wolff F. A. (1989)
Pharmacokinetics and toxicity of bismuth compounds.
Med Toxicol Adverse Drug Exp 4: 303-323.
- Slikkerveer A., Helmich R. B., van Der Voet G. B., de Wolff F. A. (1995)
Absorption of bismuth from several bismuth compounds during in vivo perfusion of rat small intestine.
J Pharm Sci. 84: 512-515.
- Smith A., Westgarth D. R., Jones M. R., Neave F. K., Dodd F. H., Brander G. C. (1967)
Methods of reducing the incidence of udder infection in dry cows.
Vet Rec. 81: 504-510.
- Smith K. L., Hogan J. S. (1999)
Managing dry cows to control mastitis.
Proc British Mastits Conf: 79-86.

Literaturverzeichnis

- Smith K. L., Todhunter D. A., Schoenberger P. S. (1985)
Environmental pathogens and intramammary infection during the dry period.
J Dairy Sci. 68: 402-417.
- Sol J., Sampimon O. C., Snoep J. J., Schukken Y. H. (1997)
Factors associated with bacteriological cure during lactation after therapy for
subclinical mastitis caused by *Staphylococcus aureus*.
J Dairy Sci. 80: 2803-2808.
- Sol J., Sampimon O. C., Snoep J. J., Schukken Y. H. (1994)
Factors associated with bacteriological cure after dry cow treatment of subclinical
staphylococcal mastitis with antibiotics.
J Dairy Sci. 77: 75-79.
- Sordillo L. M., Nickerson S. C. (1988)
Morphologic changes in the bovine mammary gland during involution and
lactogenesis.
Am J Vet Res. 49: 1112-1120.
- Sorensen J. T., Enevoldsen C. (1991)
Effect of dry period length on milk production in subsequent lactation.
J Dairy Sci. 74: 1277-1283.
- Sorenson J. T., Enevoldsen C. (1991)
Effect of dry period length on milk production in subsequent lactation.
J. Dairy Sci. 74: 1277-1283.
- Stoltenberg M., Flyvbjerg A., Sondergaard L. G., Rungby J. (2002)
Decreased serum testosterone levels in rats exposed intraperitoneally to bismuth
subnitrate.
J Appl Toxicol. 22: 111-115.
- Stoltenberg M., Hutson J. C. (2004)
Bismuth uptake in rat testicular macrophages: a follow-up observation suggesting
that bismuth alters interactions between testicular macrophages and Leydig cells.
J Histochem Cytochem. 52: 1241-1243.
- Suriyasathaporn W., Daemen A. J., Noordhuizen-Stassen E. N., Dieleman S. J.,

Nielen M., Schukken Y. H. (1999)

Beta-hydroxybutyrate levels in peripheral blood and ketone bodies supplemented in culture media affect the in vitro chemotaxis of bovine leukocytes.

Vet. Immunol. Immunopathol. 68: 177-186.

Suriyasathaporn W., Daemen A. J., Schukken Y. H. (1998)

Effect of serum β -hydroxybutyrate levels on in vitro chemotaxis of bovine leukocytes.

10th international conference of production diseases in farm animals: 266.

Suriyasathaporn W., Heuer C., Noordhuizen-Stassen E. N., Schukken Y. H. (2000)

Hyperketonemia and the impairment of udder defense: a review.

Vet Res. 31: 397-412.

Swanson E. W. (1965)

Comparing continuous milking with sixty-day dry periods in successive lactations.

J Dairy Sci 48: 1205-1209.

Timms L. L. (1997)

Efficacy of barrier teat dips in preventing dry period mastitis.

Proc. of the National Mastitis Council, Regional Meeting, Syracruse 1997: 10-17.

Timms L. L., Ackermann M., Kehrli M. (1997)

Characterization of teat end lesions observed on dairy cows during winter.

National Mastitis Council Annual Meeting Proceedings 1997: 204-209.

Timms L. L., Van der Maaten M. J., Kehrli M. E. Jr., Ackermann M. R. (1998)

Histologic features and results of virus isolation tests of tissues obtained from teat lesions that developed in dairy cattle during winter.

J Am Vet Med Assoc. 213: 862-865.

Todhunter D., Smith K. L., Hogan J. S. (1990)

Growth of gram-negative bacteria in dry cow secretion.

J Dairy Sci. 73: 363-372.

Todhunter D. A., Smith K. L., Hogan J. S. (1995)

Environmental streptococcal intramammary infections of the bovine mammary gland.

J Dairy Sci 78: 2366-2374.

Todhunter D. A., Smith K. L., Hogan J. S., Schoenberger P. S. (1991)

Gram-negative bacterial infections of the mammary gland in cows.

Am J Vet Res. 52: 184-188.

Tollersrud T., Kenny K., Caugant D. A., Lund A. (2000)

Characterisation of isolates of *Staphylococcus aureus* from acute, chronic and subclinical mastitis in cows in Norway.

Acta Pathologica, Microbiologica et Immunologica Scandinavica 108: 565-572.

Twomey D. P., Wheelock A. I., Flynn J., Meaney W. J., Hill C., Ross R. P. (2000)

Protection against *Staphylococcus aureus* mastitis in dairy cows using a bismuth-based teat seal containing the bacteriocin, lacticin 3147.

J Dairy Sci. 83 (9): 1981-1988.

Valde J. P., Østeras O., Simensen E. (2005)

Description of herd level criteria for good and poor udder health in norwegian dairy cows.

J Dairy Sci. 88: 86-92.

van der Merwe N. J. (1985)

Some observations on the morphology of the bovine teat canal (Ductus papillaris mammae).

J S Afr Vet Assoc. 56: 13-16.

Wendt K., Bostedt H., Mielke H., Fuchs H. W. (1994)

Euter- und Gesäugekrankheiten.

Gustav Fischer Verlag Jena, Stuttgart.

Wilkinson A. (2003)

To seal or not to seal: Teat Sealant Strategies.

National Mastitis Council Regional Meeting Proceedings 2003: 16-20.

Williamson J. (2002)

Dry period and heifer mastitis, role of internal and external sealants.

DeLaval Symposium, Kansas City, Proceedings: 91-99.

Wolter W., Kloppert B., Zschöck M., Castaneda V. H. (2003)

Stellenwert der zytobakteriologischen Untersuchung von Milchproben im rahmen
der Eutergesundheitsüberwachung.

Deutsche Veterinärmedizinische Gesellschaft (DVG), Tagung des Arbeitskreises
Eutergesundheit, Kiel, 2003: 53-64.

Woolford M. W., Williamson J. H., Day A. M., Copeman P. J. (1998)

The prophylactic effect of a teat sealer on bovine mastitis during the dry period and
the following lactation.

New Zealand Veterinary Journal 46: 12-19.

World Health Organization (1994)

WHO scientific working group on monitoring and management of bacterial
resistance an antimicrobial agents.

World Health Organization 1994 Geneva WHO/CDS/BVI/95.7.

Zecconi A., Hamann J., Bronzo V., Ruffo G. (1992)

Machine-induced teat tissue reactions and infection risk in a dairy herd free
from contagious mastitis pathogens.

J Dairy Res. 59 (3): 265-271.