

8 Literaturverzeichnis

Ader, P., Wessmann, A. und Wolffram, S. (2000)

Bioavailability and metabolism of the flavonol quercetin in the pig.

Free Rad. Biol. Med. 7: 1056-1067

Agullo, G., Gamet-Payraastre, L., Fernandez, Y., Anciaux, N., Demigné, C. und Rémésy, C. (1996)

Comparative effects of flavonoids on the growth, viability and metabolism of colonic adenocarcinoma cell line (HT29 cells).

Cancer lett. 105: 61-70

Afanas'ev, I.B., Dorozhko, A.I., Brodskii, A.V., Kostyuk, V.A. und Potapovitch, A.I. (1989)

Chelating and free radical scavenging mechanism of inhibitory action of rutin and quercetin in lipid peroxidation.

Biochem. Pharmacol. 38: 1763-1769

Anderson, J.J.B. und Garner, S.C. (1997)

The effect of phytoestrogens on bone.

Nutr. Res. 17: 1617-1632

Andlauer, W. und Fürst P. (1998)

Antioxidative power of phytochemicals with special reference to cereals.

Cereal foods World 43: 356-360

Aoshima, A., Yokoyama, T., Tanizaki, J., Izu, H., Yamada, M. (1997)

The sugar specificity of Na⁺/glucose cotransporter from rat jejunum.

Biosci. Biotech. Biochem 61: 979-983

Aviram, M. und Fuhrmann, B. (1998).

Polyphenolic flavonoids inhibit macrophage-mediated oxidation of LDL and attenuate atherogenesis.

Atherosclerosis. 137: 45-50

Aziz, A.A., Edwards, C.A., Lean, M.E.J., Crozier, A. (1998).

Absorption and excretion of conjugated flavonols, including Quercetin-4'-O- β -glucoside and Isorhamnetin-4'-O- β -glucoside by human volunteers after the consumption of onions.

Free Rad. Res. 29: 257-269

Barnes, S. (1998)

Phytoestrogens and breast cancer.

Baillière's Clin. Endocrinol. Met. 12: 559-579

Behrens, H. (1987)

Östrogenvergiftung.

In: *Lehrbuch der Schafkrankheiten*. pp. 260-261, 3. Auflage, Verlag Paul Parey, Berlin und Hamburg

Berhow, M.A. und Vaughn, S.F. (1999)

Higher plant flavonoids: biosynthesis and chemical ecology.

In: *Principles and Practices of Plant Ecology*. (W. V. Dashk, ed.), pp. 425-438; CRC Press Inc., Boca Raton, Florida

Block, G., Patterson, B. und Subar, A. (1992)

Fruit, vegetables and cancer prevention: a review of the epidemiological evidence.

Nutr. Cancer 18: 1-29

Bors, W., Heller, W., Michel, C. und Saran, M. (1992)

Struktural principles of flavonoid antioxidants.

In: *Free Radicals and the Liver*. (G. Csomòs & J. Fehèr, eds.), pp. 77-95, Springer-Verlag, Berlin

Boulton, D.W., Walle K., Walle T. (1998)

Extensive binding of the bioflavonoid quercetin to human plasma proteins.

J. Pharm. Pharmacol. 50: 243-249

Boulton, D.W., Walle, U.K. und Walle, T. (1999)

Fate of the flavonoid quercetin in human cell lines: Chemical instability and metabolism.

J. Pharm. Pharmacol. 51: 353-359

Bravo, L. (1998)

Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance.

Chem. Toxicol. 26: 99-103

Caldwell, J. (1985)

Glucuronic acid conjugation in the context of the metabolic conjugation of xenobiotics.

In: *Advances in Glucuronide Conjugation: Proceedings of the 40th Falk Symposium* held at Titisee, Germany, 1984.

(Matern, S., Bock, K.W., Gerock, W., eds.), pp. 7-20, Lancaster, England: MTP Press Limited, 1985

Cermak, R., Föllmer, U. und Wolfram, S. (1998)

The dietary flavonol quercetin induces chlorid secretion in rat colon.

Am J. Physiol. 275: G1166-G1172

Cermak, R., Vujcic, Z. und Wolfram, S. (2000)

The secretory response of the rat colon to the flavonol quercetin depends on the Ca^{2+} /calmodulin.

Exp. Physiol.: in press

Crespy, V., Morant, C., Manach, C., Besson, C., Demingè, C., und Rèmèsy, C. (1999)

Part of quercetin absorbed in the small intestine is conjugated and further secreted in the intestinal lumen.

Am. J. Physiol. 277: G120-G126

Criqui, M.H., Ringel, B.L. (1994)

Does diet or alcohol explain the french paradoxon.

The Lancet 344: 1719-1723

Choudhury, R., Srai, S.K., Debnam, E. und Rice-Evans, C.A. (1999)

Urinary excretion of hydroxycinnamates and flavonoids after oral and intravenous administration.

Free. Rad. Biol. Med. 27: 278-286

Das, D.K. (1994)

Naturally occurring flavonoids: structure, chemistry, and high-performance liquid chromatography methods for separation and characterisation.

In: *Methods in Enzymology*. Vol.234, pp. 410-420, Academic Press Inc., New York.

Day, A.J., Canada, F.J., Diaz, J.C., Kroon, P.A., Melauchlan, R., Faulds, C.B., Plumb, G.W., Morgan, M.R.A., Williamson, G. (2000)

Dietary flavonoid and isoflavone glycosides are hydrolysed by the lactase side of lactase phlorizin hydrolase.

FEBS Letters 468: 166-170

Day, A.J., Dupont, M.S., Ridley, S., Rhodes, M., Rhodes, M.J.C., Morgan, M.R.A. und Williamson, G. (1998)

Deglycosylation of flavonoid and isoflavonoid glycosides by human small intestine and liver beta-glucosidase activity.

FEBS Lett. 436: 71-75

de Walley, C.V., Rankin, S.M., Houtt, J.R.S., Jessup, W., Leake, D.S. (1990).

Flavonoids inhibit the oxidative modification of low density lipoproteins by macrophages.

Biochem. Pharmacol. 39: 1743-1740

Diener, M., Mestres, P., Bridges, R.J. und Rummel, W. (1989)

Funktional and morphological changes during electric field stimulation of rat colon descendens.

In: *Nerves and Gastrointestinal Tract*. (Singer, M.V. and Goebell, H., eds.), pp. 705-712, MTP Press, Lancaster, Boston, USA

Duthie, G.G., Duthie, S.J., Kyle, J.A.M. (2000)

Plant polyphenols in cancer and heart disease: implications as nutritional antioxidants.

Nutrition Research Reviews (2000) 13: 79-106

Fahey, G.H. und Jung, H.J.G. (1989)

Phenolic compounds in forages and fibrous foodstuffs.

In: *Toxicants of plant origin*. (Cheeke, P.R., ed.), pp 123-190, CRC Press, Inc., Boca Raton Florida, USA

Ferrali, M., Signorini, C., Caciotti, B., Sugherini, L., Ciccoli, L., Giachetti, D und Comporti, M. (1997)

Protection against oxidative damage of erythrocyte membrane by the flavonoid quercetin and its relation to iron chelating activity.

FEBS Lett. 416: 123-129

Formica, J.V. und Regelson, W. (1995)

Review of the biology of Quercetin and related bioflavonoids.

Fd. Chem. Toxic. 33: 1061-1080

Fotsis, T., Pepper, M.S., Montesano, R., Aktas, E., Breit, S., Schweigerer, L., Rasku, S., Wähälä, K. und Adlercreutz, H. (1998).

Phytoestrogens and inhibition of angiogenesis.

Baillière's Clin. Endocrinol. Metab. 12: 649-666

Fragner, J. (1964)

Bioflavonoide.

In: *Vitamine: Biochemie und Chemie*. (J. Fragner, ed.), pp. 487-549, Gustav Fischer Verlag, Jena

Frankel, E.N., Kanner, J., German, J.B., Parks, E. und Kinsella, J.E. (1993)

Inhibition of oxidation of human low density lipoprotein by phenolic substances in red wine.

Lancet 341: 454-457

Fuhr, U. (1996)

Metabolismus von Fremdstoffen.

In: *Phytopharmaka: Forschung und klinische Anwendung*. Hrsg.: Dieter Loew, Norbert Rietbrock HRG: Darmstadt Steinkopf 1-9

Gee, J.M., Du Pont, M.S., Rhodes, M.J.C. und Johnson, I.T. (1998)

Quercetin glucosides interact with the intestinal glucose transport pathway.

Free Radic. Biol. Med. 25: 19-25

Gee, J.M., Du Pont, M.S., Day, A.J., Plump, G.W., Williamson, G. und Johnson, I.T. (2000).

Intestinal transport of quercetin glycosides in rat involves both deglycosylation and interaction with the hexose transport pathway.

J. Nutr. 130: 2765-2771

Graefe, E.U., Derendorf, H., Veeit, M. (1999)

Pharmacokinetics and bioavailability of the flavonol quercetin in humans.

Int. J. Clin. Pharmacol. Ther. 37: 219-233

Griffith, L.A. (1982)

Mammalian metabolism of flavonoids.

In: *The flavonoids: Advances in research, Harborne*. (J.B., Marbry, T.J., eds.), pp. 681-718, Chapman and Hall, London, New-York

Gugler, R. Leschik, M., Dengler, H.J. (1975)

Disposition of quercetin in man after single oral and intravenous doses.

Europ. J. clin. Pharmacol. 9: 229-234

Guyton, J.R. (1995)

The role of lipoproteins in atherogenesis.

In: *Nutrition and Biotechnology in Heart Disease and Cancer*. (Longenecker, J.B., ed.), pp. 29-38, Plenum Press, New York

Halliwel, B. (1999)

Establishing the significance and optimal intake of dietary antioxidants: the biomarker concept.

Nut. Rev. 57: 104-113

Harborne, J.B. (1976)

In: *Chemistry and Biochemistry of Plant Pigments*. (T. W. Goodwin, ed.), pp. 736-779, Academic Press, London

Heilmann, J. und Merfort, I. (1998)

Aktueller Kenntnisstand zum Metabolismus von Flavonoiden. I.,
Resorption und Metabolismus von Flavonolen.

Pharmazie in unserer Zeit 27: 58-65

Heilmann, J. und Merfort, I. (1998a)

Aktueller Kenntnisstand zum Metabolismus von Flavonoiden. II.
Resorption und Metabolismus von Flavonen, Flavanonen, Flavanen,
Proanthocyanidinen Isoflavonoiden.

Pharmazie in unserer Zeit 27: 173-183

Herrmann, K. (1988)

On the occurrence of flavonole and flavone glycoside in vegetables.

Z. Lebensm. Unters. Forsch. 186: 1-5

Herrmann, K. (1990)

Vorkommen und Gehalte der Flavonoide in Obst.

Erwerbstobstanbau 32: 31-37

Hertog, M.G.L. (1996)

Epidemiological evidence on potential health properties of flavonoids.

Proceedings of the Nutrition Society 55: 385-397

Hertog, M.G.L., Feskens, E.J.M., Hollman, P.C.H., Katan, M.B., Kromhaut, D. (1993)

Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly study.

The Lancet 342: 1007-1011

Hertog, M.G.L., Hollman, P.C.H., van de Putte, B. (1993a)

Content of potentially anticarcinogenic flavonoids of tea infusions, wines, and fruit juices.

J. Agric. Fd. Food Chem. 41: 1242-1246

Hollman P.C.H. und Arts I.C.W. (2000)

Review: Flavonols, flavones and flavanols-nature, occurrence and dietary burden.

J. Sci Food. Agric. 80: 1081-1093

Hollman P.C.H., Buysman, M.P., van Gameren, Y., Cnossen, E., de Vries, J. und Katan, M.B. (1999)

The sugar moiety is a major determinant of the absorption of dietary flavonoid glycosides in man.

Free. Rad. Res. 31: 569-573

Hollman, P.C.H., de Vries, J.H.M., van Leeuwen, S.D., Mengelers, M.J.B., Katan, M.B. (1995).

Absorption of dietary quercetin glycosides and quercetin in healthy ileostomy volunteers.

Am. J. Clin. Nutr. 62: 1276-1282

Hollmann, P.C.H., Feskens, E.J.M. und Katan, M.B. (1999)

Tea flavonols in cardiovascular disease and cancer epidemiology.

P.S.E.B.M. 220: 198-202

Hollmann, P.C.H., Hertog, M.G.L., Katan, M.B. (1996)

Analysis and health effects of flavonoids.

Food Chemistry 57: 43-46

Hollman, P.C.H. und Katan, M.B. (1997)

Absorption, metabolism and health effects of dietary flavonoids in man.

Biomed. & Pharmacother 51: 305-310

Hollman, P.C.H., van Trijp, J.M.P., Bysman, N.C.P., v.d. Gaag, M.S., Mengelers, M.J.B., de Vries, J.H.M. und Katan, M.B. 1997

Relative bioavailability of antioxidant flavonoid quercetin from various foods in man.

FEBS Letters 418: 152-156

Ioku, K., Pongpiriyadacha, Y., Konishi, Y., Takei, Y., Nakatani, N. und Terao, J. (1998)

β -Glucosidase activity in the rat small intestine toward quercetin monoglucosides.

Biosci. Biotechnol. Biochem. 62: 1428-1431

Izumi, T., Piskula, M.K., Osawa, S., Obata, A., Tobe, K., Saito, M., Kataoka, S., Kubota, Y. und Kikuchi, M. (2000)

Soy Isoflavone aglycones are absorbed faster and in higher amounts than their glucosides in humans.

J. Nutr. 130: 1695-1699

Janssen, P.L.T.M.K., Mensink, R.P., Cox, F.J.J., Harryvan, J.L., Hovenier R., Hollmann, P.C.H. und Katan, M.B. (1998)

Effect of the flavonoids quercetin and apigenin on hemostasis in healthy volunteers: results from an in vitro and a dietary supplement study.

Am. J. Clin. 67: 255-262

Keli, S.O., Hertog, M.G.L., Feskens, E.J.M. und Kromhout, D. (1996)
Dietary flavonoids, antioxidant vitamins, and incidence of stroke. The Zutphen study.
Arch. Intern. Med.: 154: 637-642

Kneer, R. A. Poulev, A. A. Olesinski, A. und Raskin, I. (1999)
Characterisation of the elicitor-induced biosynthesis and secretion of genestein from
roots of *Lupinus Luteus* L.J.
Exp. Botany. 50: 1553-1559

Knekt, P., Järvinen, R., Reunanen, A. und Maatela, J. (1996)
Flavonoid intake and coronary mortality in Finland: a cohort study.
BMJ 312: 478-481

Kimmich, G.A. und Randles, J. (1978)
Phloretin-like action of bioflavonoids on sugar accumulation capability of isolated
intestinal cells.
Memb. Biochem. 1: 221-237

Kuo, S.-M. (1996)
Antiproliferative potency of structurally distinct dietary flavonoids on human colon
cancer cells.
Cancer lett. 110: 41-48

Kuo, S.-M. (1998)
Transepithelial transport and accumulation of flavone in human intestinal Caco-2
cells.
Life Sci. 63: 2323-2331

Kühnau, J. (1976)
The flavonoids. A class of semi-essential food components: their role in human
nutrition.
Wld Rev. Nutr. Diet. 24: 117-191

Li, C., Homma, M. und Oka, K. (1998)

Characteristics of delayed excretion of flavonoids in human urin after administration of Shosaiko-to, a herbal medicine.

Biol. Pharm. Bull. 21: 1251-1257

Linseisen, J., Radtke, J. und Wolfram, G. (1997).

Flavonoidzufuhr Erwachsener in einem bayerischen Teilkollektiv der Nationalen Verzehrsstudie.

Z. Ernährungswiss. 36: 403-412

Mabe, K., Yamada, M., Oguni, I. und Takahashi, T (1999)

In vitro and *in vivo* activities of tea catechins against *Helicobacter pylori*.

Antimicrob. Agents Chemother. 43: 1788-1791

Manach, C., Morand, C., Demigné, C., Texier, O., Règeat, F. und Rèmešy, C. (1997)

Bioavailability of rutin and quercetin in rats.

FEBS Lett. 409: 12-16

Manach, C., Morand, C., Texier, O., Favier, M.-L., Agullo, G., Règeat, F. und Rèmešy, C. (1995)

Quercetin metabolites in plasma of rats fed diets containing rutin or quercetin.

J. Nutr. 125: 1911-1922

Messina, M.J., Persky, V., Setchell, K.D. und Barnes, S. (1994)

Soy intake and cancer risk: a review of the *in vitro* and *in vivo* data.

Nutr. Cancer 21: 113-131

Middleton, E. Jr. and Kandaswami, C. (1994).

The impact of plant flavonoids on mammalian biology: Implications for immunity, inflammation and cancer.

In: *The Flavonoids: Advances in research since 1986*. (Harborne, J.B., ed.),

pp. 619-652, Chapman and Hall, London

Mizuma, T., Ohta, K., Awazu, S., (1994)

The β -anomeric and glucose preferences of glucose transport carrier for intestinal active absorption of monosaccharide conjugates.

Biochimica Biophysica. Acta 1200: 117-122

Mutschler, E. (1991)

In: *Arzneimittelwirkungen, Lehrbuch der Pharmakologie und Toxikologie*.

pp. 18-30, 6. Auflage, Wissenschaftliche Verlagsgesellschaft, Stuttgart:

Nguyen, T.D., Canada, A.T., Heintz, G.G., Gettys, T.W. und Cohn, J.A. (1991).

Stimulation of secretions by the T₈₄ colonic epithelial cell line with dietary flavonols.

Biochem. Pharmacol. 41: 1879-1886

Nielson, S.E., Breinholt, V., Justesen, U., Cornett, C. und Dragsted, L.O. (1998)

In vitro biotransformation of flavonoids by rat liver mikrosomes

Xenobiotica. 28: 389-401

Noteborn, H.P.J.M., Jansen, E., Benito, S. und Mengelers, M.J.B. (1997)

Oral absorption and metabolism of quercetin and sugar-conjugated derivatives in specific transport systems.

Cancer Lett. 114: 175-177

Ocete, M.A., Galvez, J., Crespo, M.E., Cruz, T., Gonzalez, M., Torres, M.I. und Zarzuelo, A. (1998)

Effects of morin on an experimental model of acute colitis in rats.

Pharmacol. 57: 261-270

Oestmann, A., Südekum, K.-H., Voigt, K. und Stangassinger (1995)

Zur Rolle von lignin und phenolischen Monomeren in Futtermitteln für Wiederkäuer:

1. Vorkommen, Funktionen und Nachweisverfahren.

Übers. Tierernährg. 23: 105-131

Olthof, M.R., Hollmann, P.C.H., Vree, T.B. und Katan, M.B. (1999)
Bioavailabilities of quercetin-3-glucoside and quercetin-4'-glucoside do not differ in humans.

J. Nutr. 130: 1200-1203

Pagana, G. und Rice-Evans, C.A. (1997)
The identification of flavonoids as glycosides in human plasma.

FEBS Lett. 401: 78-82

Piskula M.K., Yamakoshi, J. und Yukihiro, I. (1999)
Diadzein and genestein but not their glucosides are absorbed from the rat stomach.

FEBS Letters 447:287-291

Price, K.R. und Rodes, M.J.C. (1997)
Analysis of the major flavonoid glycosides present in four varieties of onion (*Allium cepa*) and changes in composition resulting from autolysis.

J. Sci. Food Agric. 74: 331-339

Rannelletti, F.O., Ricci, R., Larocca, L.M., Maggiano, N., Capelli, A., Scambia, G., Benedetti-Panici, P., Mancuso, S., Rumi, M. und Piantelli, M. (1992).
Growth-inhibitory effect of quercetin and presence of tyrosine kinase binding sites in human colon-cancer cell lines and primary colorectal tumors.

Int. J. Cancer 50: 486-492

Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Bramley, P.M. und Pridham, J.B. (1995)
The relative antioxidant activities of plant-derived polyphenolic flavonoids.

Free Radical Res. 22: 375-383

Roginsky, V.A., Barsukova, T.K., Remorova, A.A. und Bors, W. (1996)
Moderate antioxidative efficiencies of flavonoids during peroxidation of methyl linoleate in homogeneous and micellar solutions.

JAOCS 73: 777-786

Salunkhe, D.K., Chavan, J.K. und Kadam, S.S. (1990)
Plant phenolics: structure, classification, and biosynthesis.
In: *Dietary Tannins: Consequences and Remedies*. (Kapitel 2), (Salunkhe, D.K., ed.),
pp. 5-25, CRC Press Inc., Boca Ration, Florida, USA

Scharrer, E., Raab, W., Tiemeyer, W. und Amann, B. (1981)
Active transport of hypoxanthine by lamb jejunum *in vitro*.
Pflügers Arch. 391: 41-43

Shimoi, K., Okada, H., Furugori, M., Goda, T., Takase, S., Suzuki, M., Hara, Y.,
Yamamoto, H. und Kinae, N. (1998)
Intestinal absorption of luteolin and luteolin-7-O- β -glucoside in rats and humans.
FEBS Lett. 438: 220-224

Shirley, B.W. (1996)
Flavonoid biosynthesis: 'new' functions of an 'old' pathway.
Trends in Plant Science. 1: 377-382

Spencer, J.P.E., Chowrimootoo, G., Choudhury, R., Debnam, E.S., Srail, S.K. und
Rice-Evans, C. (1999)
The small intestine can both absorb and glucuronidate luminal flavonoids.
FEBS Lett. 458: 224-230

Steinberg, D. (1990).
Arterial metabolism of lipoproteins in relation to atherogenesis.
Ann. New York Acad. Sienc. 598: 125-135

Suzuki, Y., Ishihara, M., Segami, T. und Ito, M. (1998)
Anti-ulcer effect of antioxidants, quercetin, α -tocopherol, nifedipine and tetracycline
in rats.
Jpn. J. Pharmacol. 78: 435-441

Tsuda, T., Horio, F. und Osawa, T (1999)

Absorption and metabolism of cyanidin 3-O- β -D-glucoside in rats

FEBS Letters 449: 179-182

Uddin, S. und Choudhry, M.A. (1995)

Quercetin, a bioflavonoid, inhibits the DNA synthesis of human leukemia cells.

Biochem. Molec. Biol. Int. 36: 545-550

Ueno, I., Nakano, N. und Hirono, I. (1983)

Metabolic fate of [14 C] quercetin in the ACI rat

Jpn. J. Exp. Med. 53: 41-50

Ussing, H.H. und Zerahn, K. (1951)

Active transport of sodium as the source of electric current in the short-circuited isolated frog skin.

Acta Physiol. Scand. 23: 110-127

van Acker, S.A.B.E., de Groot, M.J., van den Berg, D.-J., Tromp, M.N.J.L., Donnè-Op den Kelder, G., van der Vijgh, W.J.F. und Bast, A. (1996a)

Structural aspects of antioxidant activity of flavonoids.

Free Rad. Biol. Med. 20: 331-342.

van Acker, S.A.B.E., de Groot, M.J., van den Berg, D.-J., Tromp, M.N.J.L., Donnè-Op den Kelder, G., van der Vijgh, W.J.F. und Bast, A. (1996b)

A quantum chemical explanation of the antioxidant activity of flavonoids.

Chem. Res. Toxicol. 9: 1305-1312

Walgren, R.A., Karnaky JR., K.J., Lindenmeyer, G.K. und Walle, T. (2000)

Efflux of dietary flavonoid quercetin 4' β -glucoside across human intestinal caco-2 monolayers by apical multidrug resistance-associated protein-2.

JPET 294: 830-836

Walgren, R.A., Lin, J.-T., Kinne, R. K.-H., Walle, T. (2000a)

Cellular uptake of dietary flavonoid quercetin-4'- β -glucosid by sodium-dependent glucose transporter SGLT1.

JPET 294: 837-843

Walgren, R.A., Walle, U.K. und Walle, T (1998)

Transport of quercetin and its glucosides across human intestina caco-2 cells.

Biochem. Pharmacol. 55: 1721-1727

Walle, U.K., French, K.L., Walgren, R.A. und Walle, T. (1999)

Transport of genistein-7-glucoside by human intestinal caco-2 cells: potential role for MRP2.

Res. Comm. Mol. Pathol. Pharmacol. 103: 45-56

Wright, E.W., Hirayama, B.A., Loo, D.D.F., Truk, E. und Hager, K. (1994)

Intestinal sugar transport.

In: *Physiologie of the Gastrointestinal Tract*. Third edition, (Leonard R. Johnson, ed.) pp. 1751-1773, Raven Press, New York