

## 5. Literaturliste

- Abraham H, Losonczy A, Czeh G und Lazar G (2001) Rapid activation of microglial cells by hypoxia, kainic acid, and potassium ions in slice preparations of the rat hippocampus. *Brain.Res.* 906:115-126.
- Acarin L, Vela JM, Gonzalez B und Castellano B (1994) Demonstration of poly-N-acetyl lactosamine residues in ameboid and ramified microglial cells in rat brain by tomato lectin binding. *J.Histochem.Cytochem.* 42:1033-1041.
- Aloisi F, Care A, Borsellino G, Gallo P, Rosa S, Bassani A, Cabibbo A, Testa U, Levi G und Peschle C (1992) Production of hemolymphopoietic cytokines (IL-6, IL-8, colony-stimulating factors) by normal human astrocytes in response to IL-1 beta and tumor necrosis factor-alpha. *J.Immunol.* 149:2358-2366.
- Alt C, Laschinger M und Engelhardt B (2002) Functional expression of the lymphoid chemokines CCL19 (ELC) and CCL21 (SLC) at the blood-brain barrier suggests their involvement in G-protein-dependent lymphocyte recruitment into the central nervous system during experimental autoimmune encephalomyelitis. *Eur.J.Immunol.* 32:2133-2144.
- Altman J (1994) Microglia emerge from the fog. *Trends.Neurosci.* 17:47-49.
- Andersen NE, Gyrding J, Hansen AJ, Laursen H und Siesjo BK (1989) Brain acidosis in experimental pneumococcal meningitis. *J.Cereb.Blood.Flow.Metab.* 9:381-387.
- Araujo DM und Cotman CW (1993) Trophic effect of interleukin-4, -7 and -8 on hippocampal neuronal cultures: potential involvement of glial-derived factors. *Brain.Res.* 600:49-55.
- Aschner M, Allen JW, Kimelberg HK, LoPachin RM und Streit WJ (1999) Glial cells in neurotoxicity development. *Annu.Rev.Pharmacol.Toxicol.* 39:151-173.
- Asensio VC und Campbell IL (1999) Chemokines in the CNS: plurifunctional mediators in diverse states. *Trends Neurosci.* 22:504-512.
- Ashwell K (1991) The distribution of microglia and cell death in the fetal rat forebrain. *Brain.Res.Dev.Brain.Res.* 58:1-12.
- Bacon KB und Harrison JK (2000) Chemokines and their receptors in neurobiology: perspectives in physiology and homeostasis. *J.Neuroimmunol.* 104: 92-97.

- Baggiolini M (1998) Chemokines and leukocyte traffic. *Nature* 392:565-568.
- Banati RB, Gehrman J, Schubert P und Kreutzberg GW (1993) Cytotoxicity of microglia. *Glia* 7:111-118.
- Barnes DA, Huston M, Holmes R, Benveniste EN, Yong VW, Scholz P und Perez HD (1996) Induction of RANTES expression by astrocytes and astrocytoma cell lines. *J.Neuroimmunol.* 71:207-214.
- Bazan JF, Bacon KB, Hardiman G, Wang W, Soo K, Rossi D, Greaves DR, Zlotnik A und Schall TJ (1997) A new class of membrane-bound chemokine with a CX3C motif. *Nature* 385:640-644.
- Bechmann I und Nitsch R (1997) Astrocytes and microglial cells incorporate degenerating fibers following entorhinal lesion: a light, confocal, and electron microscopical study using a phagocytosis-dependent labeling technique. *Glia* 20:145-154.
- Bechmann I und Nitsch R (2000) Involvement of non-neuronal cells in entorhinal-hippocampal reorganization following lesions. *Ann.N.Y.Acad.Sci.* 911:192-206.
- Bechmann I, Peter S, Beyer M, Gimsa U und Nitsch R (2001) Presence of B-7.2 (CD86) and lack of B7-1 (CD80) on myelin-phagocytosing MHC-II positive rat microglia are associated with nondestructive immunity *in vivo*. *FASEB* 15:1086-1088.
- Bell MD, Taub DD und Perry VH (1996) Overriding the brain's intrinsic resistance to leukocyte recruitment with intraparenchymal injections of recombinant chemokines. *Neuroscience* 74:283-292.
- Benveniste EN (1995). Cytokine production. In: Neuroglia (Kettenmann, H. und Ransom, BD., eds). Oxford: Oxford University Press, 700-713.
- Benveniste EN (1997) Role of macrophages/microglia in multiple sclerosis and experimental allergic encephalomyelitis. *J.Mol.Med.* 75:165-173.
- Berger T, Schnitzer J und Kettenmann H. (1991) Developmental changes in the membrane current pattern, K<sup>+</sup> buffer capacity, and morphology of glial cells in the corpus callosum slice. *J.Neurosci* 11:3008-3024.
- Bernhardi R und Ramirez G (2001) Microglia-astrocyte interaction in Alzheimer's disease: friends or foes for the nervous system? *Biol.Res* 34:123-128.

Biber K, Sauter A, Brouwer N, Copray SC und Boddeke HW (2001) Ischemia-induced neuronal expression of the microglia attracting chemokine secondary lymphoid-tissue chemokine (SLC). *Glia* 34:121-133.

Biber K, Dijkstra I, Trebst C, De Groot CJA, Ransohoff RM und Boddeke HWGM (2002) Functional expression of CXCR3 in cultured mouse and human astrocytes and microglia. *Neuroscience* 112:487-497.

Blanpain C, Migeotte I, Lee B, Vakili J, Doranz BJ, Govaerts C, Vassart G, Dorms RW und Panmentier M (1999) CCR5 binds multiple CC-chemokines: MCP-3 acts as a natural antagonist. *Blood* 94:1899-1905.

Blinzinger K und Kreutzberg G (1968). Displacement of synaptic terminals from regenerating motoneurons by microglial cells. *Zellforsch.Microsk.Anat.* 85:145-157.

Bonfoco E, Krainc D, Ankarcrona M, Nicotera P und Lipton SA (1995) Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with N-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc.Natl.Acad.Sci.USA.* 92:7162-7166.

Boucsein C, Kettenmann H und Nolte C (2000) Electrophysiological properties of microglial cells in normal and pathologic rat brain slices. *Eur.J.Neurosci.* 12:2049-2058.

Boya J, Calvo J und Prado A (1979). The origin of microglial cells. *J.Anat.* 129:177-186.

Boztug K, Carson MJ, Pham-Mitchell N, Asensio VC, DeMartino J und Campbell IL (2002) Leukocyte infiltration, but not neurodegeneration, in the CNS of transgenic mice with astrocyte production of the CXC chemokine ligand 10. *J.Immunol.* 169:1505-1515.

Braak H und Braak E (1990) Neurofibrillary changes confined to the entorhinal region and an abundance of cortical amyloid in cases of senile dementia. *Acta Neuropath.* 80:479-486.

Braak H und Braak E (1993) Entorhinal-hippocampal interaction in mnemonic disorders. Hippocampus 3 (Special Issue, Herausgeber: Nitsch, R. und Ohm, T.G.):239-46.

Brockhaus J, Ilschner S, Banati R und Kettenmann H (1993) Membrane properties of ameboid microglial cells in the corpus callosum slice from early postnatal mice. *J.Neurosci.* 13:4412-4421.

Brown KA (2001) Factors modifying the migration of lymphocytes across the blood-brain barrier. *Int.Immunopharmacol.* 1:2043-62.

- Buzsaki G, Penttonen M, Bragin A, Nadasdy Z und Chrobak JJ (1995) Possible physiological role of the perforant path-CA1 projection. *Hippocampus* 5: 141-146.
- Cairns JS und D'Souza MP (1998) Chemokines and HIV-1 second receptors: the therapeutic connection. *Nat.Med.* 4:565-568.
- Campbell JJ, Bowman EP, Murphy K, Youngman KR, Siani MA, Thompson DA, Wu L, Zlotnik A und Butcher EC (1998) 6-C-kine (SLC), a lymphocyte adhesion-triggering chemokine expressed by high endothelium, is an agonist for the MIP-3 $\beta$  receptor CCR7. *J.Cell Biol.* 141:1053-1059.
- Chulyan HE, Schall TJ, Yoshimura T und Issekutz AC (1995) IL-1 activation of endothelium supports VLA-4 (CD49d/CD29)-mediated monocyte transendothelial migration to C5a, MIP-1 alpha, RANTES, and PAF but inhibits migration to MCP-1: a regulatory role for endothelium-derived MCP-1. *J.Leukoc.Biol.* 58:71-79.
- Cocchi F, DeVico AL, Garzino-Demo A, Arya SK, Gallo RC und Lusso P (1995) Identification of RANTES, MIP-1 $\alpha$  and MIP-1 $\beta$  as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 270:1811-1815.
- Corradin SB, Mauel J, Donini SD, Quattrocchi E und Ricciardi-Castagnoli P (1993) Inducible nitric oxide synthase activity of cloned murine microglial cells. *Glia* 7:255-262.
- Damaj BB, McColl SR, Neote K, Hebert CA und Naccache PH (1996) Diverging signal transduction pathways activated by interleukin 8 (IL-8) and related chemokines in human neutrophils. IL-8 and Gro-alpha differentially stimulate calcium influx through IL-8 receptors A and B. *J.Biol.Chem.* 271:20540-20544.
- Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R, Goedert JJ, Buchbinder SP, Vittinghoff E, Gomperts E, Donfield S, Vlahov D, Kaslow R, Saah A, Rinaldo C, Detels R und O'Brien SJ (1996) Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CCR5 structural gene. *Science* 273:1856-1862.
- Devalaraja MN und Richmond A (1999) Multiple chemotactic factors: fine control or redundancy? *Trends.Pharmacol.Sci.* 20:151-156.
- Dohlman HG, Thorner J, Caron MG und Lefkowitz RJ (1991) Model systems for the study of seven-transmembrane-segment receptors. *Annu.Rev.Biochem.* 60:653-688.

Downen M, Amaral TD, Hua LL, Zhao ML und Lee SC (1999) Neuroanl death in cytokine-activated primary human brain cell culture: role of tumor necrosis factor-alpha. *Glia* 28:114-127.

Drejer J, Larsson OM, Kvamme E, Svenneby G, Hertz L und Schousboe A (1985) Ontogenetic development of glutamate metabolizing enzymes in cultured cerebellar granule cells and in cerebellum in vivo. *Neurochem.Res.* 10:49-62.

Eder C, Klee R und Heinemann U (1998) Involvement of stretch-activated Cl<sup>-</sup> channels in ramification of murine microglia. *J.Neurosci.* 18:7127-7137.

Eder C (1998) Ion channels in microglia (brain macrophages). *Am.J.Physiol.* 275:C327-C242.

Fagan AM und Gage FH (1994) Mechanisms of sprouting in the adult central nervous system: cellular responses in areas of terminal degeneration and reinnervation in the rat hippocampus. *Neuroscience* 58:705-725.

Ferrari D, Chiozzi P, Falzoni S, Hanau S und Di VF (1997) Purinergic modulation of interleukin-1 beta release from microglial cells stimulated with bacterial endotoxin. *J.Exp.Med.* 185:579-582.

Fife BT, Huffnagle GB, Kuziel WA und Karpus WJ (2000) CC Chemokine receptor 2 is critical for induction of experimental autoimmune encephalomyelitis. *J.Exp.Med.* 192:899-905.

Fife BT, Kennedy KJ, Paniagua MC, Lukacs NW, Kunkel SL, Luster AD und Karpus WJ (2001) CXCL10 (IFN- $\gamma$ -inducible protein-10) control of encephalitogenic CD4<sup>+</sup> T cell accumulation in the central nervous system during experimental autoimmune encephalomyelitis. *J.Immunol.* 166:7617-7624.

Flügel A, Hager G, Horvat A, Spitzer C, Singer GMA, Graeber MB, Kreutzberg GW und Schwaiger FW (2001) Neuronal MCP-1 expression in response to remote nerve injury. *J.Cerebral Blood Flow and Metabolism.* 21:69-76.

Forster R, Schubel A, Breitfeld D, Kremmer E, Renner-Muller I, Wolf E und Lipp M (1999) CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* 99:23-33.

Frotscher M, Heimrich B und Deller T (1997) Sprouting in the hippocampus is layer-specific. *Trends Neurosci.* 20:218-222.

Gall C, Rose G und Lynch G (1979) Proliferative and migratory activity of glial cells in the partially deafferented hippocampus. *J.Comp.Neurol.* 183:539-549.

Gallin EK (1984) Electrophysiological properties of macrophages. *Fed.Proc.* 43:2385-2389.

Gage FH, Olejniczak P und Armstrong DM (1988) Astrocytes are important for sprouting in the septohippocampal circuit. *Exp.Neurol.* 102:2-13.

Garcia-Zepeda EA, Combadiere C, Rothenberg ME, Sarafi MN, Lavigne F, Hamid Q, Murphy PM und Luster AD (1996) Human monocyte chemoattractant protein (MCP)-4 is a novel CC chemokine with activities on monocytes, eosinophils, and basophils induced in allergic and nonallergic inflammation that signals through the CC chemokine receptors (CCR)-2 and -3. *J.Immunol.* 157:5613-5626.

Gebicke-Haerter PJ, Christoffel F, Timmer J, Northoff H, Berger M und Van-Calker D (1996) Both adenosine A1- and A2-receptors are required to stimulate microglial proliferation. *Neurochem.Int.* 29:37-42.

Gerdes J, Lemke H, Baisch H, Wacker HH, Schwab U und Stein H (1984) Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. *J.Immunol.* 133:1710-1715.

Gehrmann J, Schoen SW und Kreutzberg GW (1991) Lesion of the rat entorhinal cortex leads to a rapid microglial reaction in the dentate gyrus: A light and electron microscopical study. *Acta Neuropathol.* 82:442-455.

Gehrmann J, Mies G, Bonnekoh P, Banati R, Iijima T, Kreutzberg GW und Hossmann KA (1993) Microglial reaction in the rat cerebral cortex induced by cortical spreading depression. *Brain.Pathol.* 3:11-17.

Gehrmann J, Matsumoto Y und Kreutzberg GW (1995) Microglia: intrinsic immune effector cell of the brain. *Brain Res.Rev.* 20:269-287.

Glabinski AR, Tani M, Tuohy VK, Tuthill RJ und Ransohoff RM (1995) Central nervous system chemokine mRNA accumulation follows initial leukocyte entry at the onset of acute murine experimental autoimmune encephalomyelitis. *Brain Behav.Immun* 9:315-330.

- Goldberg SH, van der Meer P, Hesselgesser J, Jaffer S, Kolson DL, Albright AV, Gonzalez-Scarano F und Lavi E (2001) CXCR3 expression in human central nervous system diseases. *Neuropathology and applied Neurobiol.* 27:127-138.
- Gottlieb M und Matute C (1997) Expression of ionotropic glutamate receptor subunits in glial cells of the hippocampal CA1 area following transient forebrain ischemia. *J.Cereb.Blood.Flow.Metab.* 17:290-300.
- Gourmala NG, Buttini M, Limonta S, Sauter A und Boddeke HW (1997) Differential and time-dependent expression of monocyte chemoattractant protein-1 mRNA by astrocytes and macrophages in rat brain: effects of ischemia and peripheral lipopolysaccharide administration. *J.Neuroimmunol.* 74:35-44.
- Graeber MB, Streit WJ und Kreutzberg GW (1988) Axotomy of the rat facial nerve leads to increased CR3 complement receptor expression by activated microglial cells. *J.Neurosci.Res.* 21:18-24.
- Gulyas AI, Miles R, Sik A, Toth K, Tamamaki N und Freund TF (1993) Hippocampal pyramidal cells excite inhibitory neurons through a single release site. *Nature* 366:683-687.
- Gunn MD, Tangemann K, Tam C, Cyster JG, Rosen SD und Williams LT (1998) A chemokine expressed in lymphoid high endothelial venules promotes the adhesion and chemotaxis of naive T lymphocytes. *Proc.Natl.Acad.Sci.USA* 95:258-263.
- Gunn MD, Kyuwa S, Tam C, Kakiuchi T, Matsuzawa A, Williams LT und Nakano H (1999) Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J.Exp.Med.* 189:451-460.
- Hainer NP, Grampp A und Nitsch R (1999) Proliferation of microglia and astrocytes in the dentate gyrus following entorhinal cortex lesion: a quantitative bromodeoxyuridine-labelling study. *Eur.J.Neurosci.* 11:3359-3364.
- Hamill OP, Marty A, Neher E, Sakmann B und Sigworth FJ (1981) Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch.* 391:85-100.
- Hamm HE (1998) The many faces of G protein signaling. *J.Biol.Chem.* 273:669-672.

- Hancock WW, Lu B, Gao W, Csizmadia V, Faia K, King JA, Smiley ST, Ling M, Gerard NP und Gerard C (2000) Requirement of the chemokine receptor CXCR3 for acute allograft rejection. *J.Exp.Med.* 192:1515-1520.
- Hanisch UK, Prinz M, Angstwurm K, Hausler KG, Kann O, Kettenmann H und Weber JR (2001) The protein tyrosine kinase inhibitor AG126 prevents the massive microglial cytokine induction by pneumococcal cell walls. *Eur.J.Immunol.* 31:2104-2115.
- Hao C, Richardson A und Federoff S (1991) Macrophage-like cells originate from neuroepithelium in culture: characterization and properties of the macrophage-like cells. *Int.J.Dev.Neurosci.* 9:1-14.
- Harrison JK, Jiang Y, Chen S, Xia Y, Maciejewski D, McNamara RK, Streit WJ, Salafranca MN, Adhikari S, Thompson DA, Botti P, Bacon P und Feng L (1998) Role for neuronally derived fractalkine in mediating unique interactions between neurons and CX3CR1-expressing microglia. *Proc.Natl.Acad.Sci.USA.* 95:10896-10901.
- Hart MN und Fabry Z (1995) CNS antigen presentation. *Trends.Neurosci.* 18:475-481.
- Hayes GM, Woodroffe MN und Cuzner ML (1987) Microglia are the major cell type expressing MHC class II in human white matter. *J.Neurol.Sci.* 80:25-37.
- He J, Chen Y, Farzan M, Choe H, Ohagen A, Gartner S, Busciglio J, Yang X, Hofmann W, Newman W, Mackay CR, Sodroski J und Gabuzda D (1997) CCR3 and CCR5 are coreceptors for HIV-1 infection of microglia. *Nature* 385:645-649.
- Hedrick JA und Zlotnik A (1997) Identification and characterization of a novel beta chemokine containing six conserved cysteines. *J.Immunol.* 159:1589-1593.
- Hickey WF, Hsu BL und Kimura H (1991) T-lymphocyte entry into the central nervous system. *J.Neurosci.Res.* 28:254-260.
- Holevinsky KO und Nelson DJ (1995) Simultaneous detection of free radical release and membrane current during phagocytosis. *J.Biol.Chem.* 270:8328-8336.
- Honda S, Sasaki Y, Ohsawa K, Imai Y, Nakamura Y, Inoue K und Kohsaka S (2001) Extracellular ATP or ADP induce chemotaxis of cultured microglia through Gi/o-coupled P2Y receptors. *J.Neurosci.* 21:1975-1982.

Hope PL, Cady EB, Delpy DT, Ives NK, Gardiner RM und Reynolds EO (1988) Brain metabolism and intracellular pH during ischaemia: effects of systemic glucose and bicarbonate administration studied by  $^{31}\text{P}$  and  $^1\text{H}$  nuclear magnetic resonance spectroscopy in vivo in the lamb. *J.Neurochem.* 50:1394-1402.

Horuk R, Martin AW, Wang Z, Schweitzer L, Gerassimides A, Guo H, Lu Z, Hesselgesser J, Perez HD, Kim J, Parker J, Hadley TJ und Peiper SC (1997) Expression of chemokine receptors by subsets of neurons in the central nervous system. *J.Immunol.* 158:2882-2890.

Hromas R, Kim CH, Klemsz M, Krathwohl M, Fife K, Cooper S, Schnizlein-Bick C und Broxmeyer HE (1997) Isolation and characterization of Exodus-2, a novel C-C chemokine with a unique 37-amino acid carboxyl-terminal extension. *J.Immunol.* 159:2554-2558.

Hromas R, Cripe L, Hangoc G, Cooper S und Broxmeyer HE (2000) The exodus subfamily of CC chemokines inhibits the proliferation of chronic myelogenous leukemia progenitors. *Blood* 95:1506-1508.

Hyman BT, Van Hoesen GW, Damasio AR und Barnes CL (1984) Alzheimers's disease: cell-specific pathology isolates the hippocampal formation. *Science* 225:1168-1170.

Ince C, Coremans JM, Ypey DL, Leijh PC, Verveen AA und Van Furth R (1988) Phagocytosis by human macrophages is accompanied by changes in ionic channel currents. *J.Cell Biol.* 106:1873-1878.

Jenh CH, Cox MA, Kaminski H, Zhang M, Byrnes H, Fine J, Lundell D, Choa CC, Narula SK und Zavodny PJ (1999) Cutting edge: Species specificity of the CC chemokine 6Ckine signaling through the CXC chemokine receptor CXCR3: human 6Ckine is not a ligand for the human or mouse CXCR3 receptors. *J.Immunol.* 162:3765-3769.

Jensen MB, Gonzalez B, Castellano B und Zimmer J (1994) Microglial and astrogliial reactions to anterograde axonal degeneration: a histochemical and immunocytochemical study of the adult rat fascia dentata after entorhinal perforant path lesions. *Exp.Brain.Res.* 98:245-260.

Johnson LV, Leitner WP, Rivest AJ, Staples MK, Radeke MJ und Anderson DH (2002) The Alzheimer's Abeta -peptide is deposited at sites of complement activation in pathologic deposits associated with aging and age-related macular degeneration. *Proc.Natl.Acad.Sci.USA* 99:11830-11835.

- Kakuta Y, Okayama H, Aikawa T, Kanno T, Ohyama T, Sasaki H, Kato T und Takishima T (1988) K channels of human alveolar macrophages. *J.Allergy Clin.Immunol.* 81:460-468.
- Karpus WJ und Ransohoff RM (1998) Chemokine regulation of experimental autoimmune encephalomyelitis: temporal and spatial expression patterns govern disease pathogenesis. *J.Immunol.* 161:2667-2671.
- Kelner GS, Kennedy J, Bacon KB, Kleyensteuber S, Largaespada DA, Jenkins NA, Copeland NG, Bazan JF, Moore KW und Schall TJ (1994) Lymphotactin: a cytokine that represents a new class of chemokine. *Science* 266:1395-1399.
- Kettenmann H, Hoppe D, Gottmann K, Banati R und Kreutzberg G (1990) Cultured microglial cells have a distinct pattern of membrane channels different from peritoneal macrophages. *J.Neurosci.Res.* 26:278-287.
- Kettenmann H, Banati R und Walz W (1993) Electrophysiological behavior of microglia. *Glia* 7:93-101.
- Kosaka T, Katsumaru H, Hama K, Wu JY und Heizmann CW (1987) GABAergic neurons containing the Ca<sup>2+</sup>-binding protein parvalbumin in the rat hippocampus and dentate gyrus. *Brain Res.* 419:119-130.
- Kovac AD, Kwidzinski E, Bittigau P, Heimrich B, Nitsch R und Bechmann I (2003) Trans-synaptic cell death of dentate granule cells following entorhinal cortex lesion in the mouse. Eingereicht.
- Kreutzberg GW (1996) Microglia: a sensor for pathological events in the CNS. *Trends Neurosci.* 19:312-318.
- Krug M, Brodemann R, Matthies R, Ruthrich H und Wagner M (2001) Activation of the dentate gyrus by stimulation of the contralateral perforant pathway: evoked potentials and long-term potentiation after ipsi- and contralateral induction. *Hippocampus* 11:157-167.
- Lahrtz F, Piali L, Spanaus KS, Seebach J und Fontana A (1998) Chemokines and chemotaxis of leukocytes in infectious meningitis. *J.Neuroimmuno.* 85:33-43.
- Lane TE, Asensio VC, Yu N, Paoletti AD, Campbell IL und Buchmeier MJ (1998) Dynamic regulation of alpha- and beta-chemokine expression in the central nervous system during mouse hepatitis virus-induced demyelinating disease. *J.Immunol.* 160:970-978.

Lassmann H (1997) Basic mechanisms of brain inflammation. *J.Neural.Transm.Suppl.* 50:183-190.

Lavi E, Strizki JM, Ulrich AM, Zhang W, Fu L, Wang Q, O'Connor M, Hoxie JA und Gonzalez-Scarano F (1997) CXCR-4 (Fusin), a co-receptor for the type 1 human immunodeficiency virus (HIV-1), is expressed in the human brain in a variety of cell types, including microglia and neurons. *Am.J.Pathol.* 151:1035-1042.

Lawson LJ, Perry VH und Gordon S (1992) Turnover of resident microglia in the normal adult mouse brain. *Neuroscience* 48:405-415.

Leist M und Nicotera P (1997) The shape of cell death. *Biochem.Biophys.Res.Commun.* 236:1-9.

Leist M und Nicotera P (1998) Apoptosis, excitotoxicity, and neuropathology. *Exp.Cell Res.* 239:183-201.

Ling EA und Wong WC (1993) The origin and nature of ramified and ameboid microglia: a historical review and current concepts. *Glia* 7:9-18.

Liu R, Paxton WA, Choe S, Ceradini D, Martin SR, Horuk R, MacDonald ME, Stuhlmann H, Koup RA und Landau NR (1996) Homozygous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* 86:367-377.

Liu QH, Williams DA, McManus C, Baribaud F, Doms RW, Schols D, De Clercq E, Kotlikoff MI, Collman RG und Freedman BD (2000) HIV-1 gp120 and chemokines activate ion channels in primary macrophages through CCR5 and CXCR4 stimulation. *Proc.Natl.Acad.Sci.USA* 97:4832-4837.

Loetscher M, Gerber B, Loetscher P, Jones SA, Piali L, Clark-Lewis I, Baggolini M und Moser B (1996) Chemokine receptor specific for IP10 and mig: structure, function, and expression in activated T-lymphocytes. *J.Exp.Med.* 184:963-969.

Luster AD (1998) Chemokines-chemotactic cytokines that mediate inflammation. *New Engl.J.Med.* 338:436-445.

Lynch G, Rose G, Gall C und Cotman CW (1975) The response of the dentate gyrus to partial deafferentation. In: Golgi Centennial Symposium Proceedings (ed. Santini, M.), 305-317. Raven Press, New York.

- Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T, Bronson RT and Springer TA (1998) Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc.Natl.Acad.Sci.USA* 95:9448-9453.
- Maghazachi AA (1999) Intracellular signalling pathways induced by chemokines in natural killer cells. *Cell.Signal.* 11:385-390.
- Mantovani A (1999) The chemokine system: redundancy for robust outputs. *Immunol.Toda.* 20:254-257.
- Medawar PB (1948) Immunity to homologous grafted skin. III. The fate of skin homografts transplanted to the brain, to subcutaneous tissue, and to the anterior chamber of the eye. *British J.Exp.Pathol.* 29:58-69.
- Mennicken F, Maki R, de Souza EB und Quirion R (1999) Chemokines and chemokine receptors in the CNS: a possible role in neuroinflammation and patterning. *TiPS* 20:73-78.
- Merrill JE, Ignarro LJ, Sherman MP, Melinek J und Lane TE (1993) Microglial cell cytotoxicity of oligodendrocytes is mediated through nitric oxide. *J.Immunol.* 151:2132-2134.
- Minagar A, Shapshak P, Fujimura R, Ownby R, Heyes M und Eisdorfer C (2002) The role of macrophage/microglia and astrocytes in the pathogenesis of three neurologic disorders: HIV-associated dementia, Alzheimer disease, and multiple sclerosis. *J.Neurol. Sci.* 202:13-23.
- Möller T, Nolte C, Burger R, Verkhratsky A und Kettenmann H (1997) Mechanisms of C5a and C3a complement fragment-induced  $[Ca^{2+}]_i$  signaling in mouse microglia. *J.Neurosci.* 17:615-624.
- Moore S und Thanos S (1996) The concept of microglia in relation to central nervous system disease and regeneration. *Prog.Neurobiol.* 48:441-460.
- Morgan BP (1994) Clinical complementology: recent progress and future trends. *Eur.J.Clin.Invest.* 24:219-228.
- Murphy PM, Baggolini M, Charo IF, Hebert CA, Horuk R, Matsushima K, Miller LH, Oppenheim JJ und Power CA (2000) International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol.Rev.* 52:145-176.

- Nagasawa T, Hirota S, Tachibana K, Takakura N, Nishikawa S, Kitamura Y, Yoshida N, Kikutani H und Kishimoto T (1996) Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* 382:635-638.
- Nagira M, Imai T, Hieshima K, Kusuda J, Ridanpaa M, Takagi S, Nishimura M, Kakizaki M, Nomiyama H und Yoshie O (1997) Molecular cloning of a novel human CC chemokine secondary lymphoid-tissue chemokine that is a potent chemoattractant for lymphocytes and mapped to chromosome 9p13. *J.Biol.Chem.* 272:19518-19524.
- Neher E und Sakmann B (1976) Noise analysis of drug induced voltage clamp currents in denervated frog muscle fibres. *J.Physiol.* 258:705-729.
- Nibbs RJ, Yang J, Landau NR, Mao JH und Graham GJ (1999) LD78beta, a non-allelic variant of human MIP-1alpha (LD78alpha), has enhanced receptor interactions and potent HIV suppressive activity. *J.Biol.Chem.* 274:17478-17483.
- Nitsch R, Soriano E und Frotscher M (1990) The parvalbumin-containing nonpyramidal neurons in the rat hippocampus. *Embryol.* 181:413-425.
- Nitsch R und Frotscher M (1991) Maintenance of peripheral dendites of GABAergic neurons requires specific input. *Brain Res.* 554:304-307.
- Nörenberg W, Gebicke HP und Illes P (1994) Voltage-dependent potassium channels in activated rat microglia. *J.Physiol.Lond.* 475:15-32.
- Numberger M und Draguhn A (1996) Patch-clamp-Technik. Aus der Reihe Labor im Fokus. Spektrum Akademischer Verlag Heidelberg Berlin Oxford.
- Ohm TG, Müller H, Braak H und Bohl J (1995) Close-meshed prevalence rates of different stages as a tool to uncover the rate of Alzheimers's disease-related neurofibrillary changes. *Neuroscience* 64: 209-217.
- Ohomori Y und Hamilton TA (1995) The interferon-stimulated response element and a kappa B site mediate synergistic induction of murine IP-10 gene transcription by IFN-gamma and TNF-alpha. *J.Immunol.* 154:5235-5244.
- Oppermann M, Mack M, Proudfoot AE und Olbrich H (1999) Differential effects of CC chemokines on CC chemokine receptor 5 (CCR5) phosphorylation and identification of phosphorylation sites on the CCR5 carboxyl terminus. *J.Biol.Chem.* 274:8875-8885.

- Pan Y, Lloyd C, Zhou H, Dolich S, Deeds J, Gonzalo JA, Vath J, Gosselin M, Ma J, Dussault B, Woolf E, Alperin G, Culpepper J, Gutierrez-Ramos JC und Gearing D (1997) Neurotactin, a membrane- anchored chemokine upregulated in brain inflammation. *Nature* 387:611-617.
- Pal R, Garzino-Demo A, Markham PD, Burns J, Brown M, Gallo RC und DeVico AL (1997) Inhibition of HIV-1 infection by the beta-chemokine MDC. *Science* 278:695-698.
- Perry VH (1994) Modulation of microglia phenotype. *Neuropathol.Appl.Neurobiol* 20:177.
- Peterson PK, Hu S, Salak-Johnson J, Molitor TW und Chao CC (1997) Differential production of and migratory response to beta chemokines by human microglia and astrocytes. *J.Infect.Dis.* 175:478-481.
- Phillips JW, O'Regan MH und Perkins LM (1993) Adenosine 5'-triphosphate release from the normoxic and hypoxic in vivo rat cerebral cortex. *Neurosci.Lett.* 151:94-96.
- Phipps DJ, Branch DR und Schlichter LC (1996) Chloride-channel block inhibits T lymphocyte activation and signalling. *Cell Signal.* 8:141-149.
- Piali L, Weber C, LaRosa G, Mackay CR, Springer A, Clark-Lewis I und Moser B (1998) The chemokine receptor CXCR3 mediates rapid and shear-resistant adhesion-induction of effector T lymphocytes by the chemokines IP10 and Mig. *Eur.J.Immunol.* 28:961-972.
- Poirier J, Hess M, May PC und Finch CE (1991) Astrocytic apolipoprotein E mRNA and GFAP mRNA in hippocampus after entorhinal cortex lesioning. *Mol.Brain.Res.* 11:97-106.
- Prinz M, Kann O, Draheim HJ, Schumann RR, Kettenmann H, Weber JR und Hanisch UK (1999) Microglial activation by components of gram-positive and -negative bacteria: distinct and common routes to the induction of ion channels and cytokines. *J.Neuropathol.Exp.Neurol.* 58:1078-1089.
- Raivich G, Moreno-Flores MT, Moller JC und Kreutzberg GW (1994) Inhibition of posttraumatic microglial proliferation in a genetic model of macrophage colony-stimulating factor deficiency in the mouse. *Eur.J.Neurosci.* 6:1615-1618.
- Raivich G, Jones LL, Kloss CU, Werner A, Neumann H und Kreutzberg GW (1998) Immune surveillance in the injured nervous system: T-lymphocytes invade the axotomized mouse facial motor nucleus and aggregate around sites of neuronal degeneration. *J.Neurosci.* 18:5804-5816.

Raivich G, Bohatschek M, Kloss CUA, Werner A, Jones LL und Kreutzber GW (1999) Neuroglial activation repertoire in the injured brain: graded response, molecular mechanisms and cues to physiological function. *Brain Res.Rev.* 30:77-105.

Ransohoff RM, Hamilton TA, Tani M, Stoler MH, Shick HE, Major JA, Estes ML, Thomas DM und Tuohy VK (1993) Astrocyte expression of mRNA encoding cytokines IP-10 and JE/MCP-1 in experimental autoimmune encephalomyelitis. *FASEB* 7:592-600.

Rappert A, Biber K, Nolte C, Lipp M, Schubel A, Lu B, Gerard NP, Gerard C, Boddeke HW und Kettenmann H (2002) Secondary lymphoid tissue chemokine (CCL21) activates CXCR3 to trigger a Cl<sup>-</sup> current and chemotaxis in murine microglia. *J.Immunol.* 168:3221-3226.

Rappert, A., Bechmann, I., Pivneva, T., Mahlo, J., Biber, K., Nolte, C., Kovac, A. D., Gerard, C., Boddeke, H. W. G. M., Nitsch, R. and Kettenmann H. (2003) CXCR3-dependent microglial recruitment is essential for dendrite loss after brain lesion. Eingereicht.

Richardson A, Hao C und Federoff S (1993) Microglia progenitor cells: a subpopulation in cultures of mouse neopallial astroglia. *Glia* 7:25-33.

Rio-Hortega PD (1932) Microglia. In: Cytology and Cellular Pathology of the Nervous System (Penfield W. ed), 483-534. New York: Hafner Publishing Company.

Robinson S, Tani M, Skrietre RM, Ransohoff RM und Miller RH (1998) The chemokine growth-regulated oncogene-a promotes spinal cord oligodendrocyte precursor proliferation. *J.Neurosci.* 18:10457-10463.

Rollins B (1997) Chemokines. *Blood* 90:909-928.

Rose G, Lynch G und Cotman CW (1976) Hypertrophy and redistribution of astrocytes in the deafferented dentate gyrus. *Brain Res.Bull.* 1:87-92.

Rottman JB, Ganley KP, Williams K, Lu L, Mackay CR und Ringler DJ (1997) Cellular localization of the chemokine receptor CCR5. Correlation to cellular targets of HIV-1 infection. *Am.J.Pathol.* 151:1341-1350.

Rottman JB, Slavin AJ, Silva R, Weiner HL, Gerard CG und Hancock WW (2000) Leukocyte recruitment during onset of experimental allergic encephalomyelitis is CCR1 dependent. *Eur.J.Immunol.* 30:2372-2377.

Ruth RE, Collier TJ und Routtenberg A (1982) Topography between the entorhinal cortex and the dentate septotemporal axis in rats: I. Medial and intermediate entorhinal projecting cells. *J.Comp.Neurol.* 209:69-78.

Schlichter LC, Sakellaropoulos G, Ballyk B, Pennefather PS und Phipps DJ (1996) Properties of K<sup>+</sup> and Cl<sup>-</sup> channels and their involvement in proliferation of rat microglial cells. *Glia* 17:225-236.

Schmidtmayerova H, Nottet HS, Nuovo G, Raabe T, Flanagan CR, Dubrovsky L, Gendelman HE, Cerami A, Bukrinsky M und Sherry B (1996) Human immunodeficiency virus type 1 infection alters chemokine beta peptide expression in human monocytes: implications for recruitment of leukocytes into brain and lymph nodes. *Proc.Natl.Acad.Sci.USA.* 93:700-704.

Schwaeble WJ, Stover CM, Schall TJ, Dairaghi DJ, Trinder PKE, Linington C, Iglesias A, Schubart A, Lynch NJ, Weihe E und Schafer MKH (1998) Neuronal expression of fractalkine in the presence and absence of inflammation. *FEBS Lett.* 439:203-207.

Shyamala V und Khoja H (1998) Interleukin-8 receptors R1 and R2 activate mitogen-activated protein kinases and induce c-fos, independent of Ras and Raf-1 in Chinese hamster ovary cells. *Biochemistry* 37:15918-15924.

Simpson JE, Newcombe J, Cuzner ML und Woodroffe MN (2000a) Expression of the interferon-gamma-inducible chemokines IP-10 and Mig and their receptor, CXCR3, in multiple sclerosis lesions. *Neuropathol.Appl.Neurobiol.* 26:133-142.

Simpson J, Rezaie P, Newcombe J, Cuzner ML, Male D und Woodroffe MN (2000b) Expression of the beta-chemokine receptors CCR2, CCR3 and CCR5 in multiple sclerosis central nervous system tissue. *J.Neuroimmunol.* 108:192-200.

Slepko N und Levi G (1996) Progressive activation of adult microglial cells in vitro. *Glia* 16:241-246.

Sloviter RS (1989) Calcium-binding protein (calbindin-D28k) and parvalbumin immunocytochemistry: localization in the rat hippocampus with specific reference to the selective vulnerability of hippocampal neurons to seizure activity. *J.Comp.Neuro.* 280:186-196.

Sigworth FJ (1985) Open channel noise. I. Noise in acetylcholine receptor currents suggests conformational fluctuations. *Biophys.J.* 47:709-720.

- Sontheimer H und Waxman SG (1993) Expression of voltage-activated ion channels by astrocytes and oligodendrocytes in the hippocampal slice. *J.Neurophysiol.* 70:1863-1873.
- Sorensen JC, Dalmau I, Zimmer J und Finsen B (1996) Microglial reactions to retrograde degeneration of tracer-identified thalamic neurons after frontal sensorimotor cortex lesions in adult rats. *Exp.Brain Res.* 112:203-212.
- Sorensen TL, Tani M, Jensen J, Pierce V, Lucchinetti C, Folcik VA, Qin S, Rottman J, Sellebjerg F, Strieter RM, Frederiksen JL und Ransohoff RM (1999) Expression of specific chemokines and chemokine receptors in the central nervous system of multiple sclerosis patients. *J.Clin.Invest.* 103:807-815.
- Soto H, Wang W, Strieter RM, Copeland NG, Gilbert DJ, Jenkins NA, Hedrick J und Zlotnik A (1998) The CC chemokine 6Ckine binds the CXC chemokine receptor CXCR3. *Proc.Natl.Acad.Sci.USA* 95:8205-8210.
- Steward O (1976) Topographic organization of the projections from the entorhinal area to the hippocampal formation in the rat. *J.Comp.Neurol.* 167:285-314.
- Steward O und Vinsant SL (1983) The process of reinnervation in the dentate gyrus of the adult rat: a quantitative electron microscopic analysis of terminal proliferation and reactive synaptogenesis. *J.Comp.Neurol.* 214:370-386.
- Streit WJ, Graeber MB und Kreutzberg GW (1988) Functional plasticity of microglia: a review. *Glia* 1:301-307.
- Streit WJ und Graeber MB (1993) Heterogeneity of microglial and perivascular cell populations: insights gained from the facial nucleus paradigm. *Glia* 7:68-74.
- Streit WJ und Kincaid Colton CA (1995) The brain's immune system. *Sci.Am.* 273:54-61.
- Streit WJ, Walter SA und Pennell NA (1999) Reactive microgliosis. *Prog.Neurobiol.* 57:563-81.
- Tanabe S, Heesen M, Berman MA, Fischer MB, Yoshizawa I, Luo Y und Dorf ME (1997a) Murine astrocytes express a functional chemokine receptor. *J.Neurosci.* 17:6522-6528.
- Tanabe S, Heesen M, Yoshizawa I, Berman MA, Luo Y, Bleul CC, Springer T, Okuda K, Gerard N und Dorf ME (1997b) Functional expression of the CXC-chemokine receptor-4/fusin on mouse microglial cells and astrocytes. *J.Immunol.* 159:905-911.

- Taub DD, Lloyd AR, Conlon K, Wang JM, Ortaldo R, Harada A, Matsushima K, Kelvin DJ und Oppenheim JJ (1993) Recombinant human-interferon-inducible protein 10 is a chemoattractant for human monocytes and T lymphocytes and promotes T cell adhesion to endothelial cells. *J.Exp.Med.* 177:1809-1814.
- Thanos S (1991) The Relationship of Microglial Cells to Dying Neurons During Natural Neuronal Cell Death and Axotomy-induced Degeneration of the Rat Retina. *Eur.J.Neurosci.* 3:1189-1207.
- Van Hoesen GW, Hyman BT und Damasio A (1991) Entorhinal cortex pathology in Alzheimer's disease. *Hippocampus* 1:1-8.
- Vassileva G, Soto H, Zlotnik A, Nakano H, Kakiuchi T, Hedrick JA und Lira SA (1999) The reduced expression of 6Ckine in the plt mouse results from the deletion of one of two 6Ckine genes. *J.Exp.Med.* 190:1183-1188.
- Visentin S, Agresti C, Patrizio M und Levi G (1995) Ion channels in rat microglia and their different sensitivity to lipopolysaccharide and interferon-gamma. *J.Neurosci.Res.* 42:439-451.
- Walz W, Ilschner S, Ohlemeyer C, Banati R und Kettenmann H (1993) Extracellular ATP activates a cation conductance and a K<sup>+</sup> conductance in cultured microglial cells from mouse brain. *J.Neurosci.* 13:4403-4411.
- Walz W und Bekar LK (2001) Ion channels in cultured microglia. *Microsc.Res.Tech.* 54:26-33.
- Wekerle H (1995) Antigen presentation by central nervous system glia. In: Neuroglia (Kettenmann, H. and B.R. Ransom eds), 685-699. New York: Oxford University Press.
- Willimann K, Legler DF, Loetscher M, Roos RS, Delgado MB, Clark-Lewis I, Baggiolini M und Moser B (1998) The chemokine SLC is expressed in T cell areas of lymph nodes and mucosal lymphoid tissues and attracts activated T cells via CCR7. *Eur.J.Immunol.* 28:2025-2034.
- Witter MP, Van Hoesen GW und Amaral DG (1989) Topographical organization of the rat entorhinal projection to the dentate gyrus of the monkey. *J.Neurosci.* 9:216218.

- Woodroffe MN, Hayes GM und Cuzner ML (1989) Fc receptor density, MHC antigen expression and superoxide production are increased in interferon-gamma-treated microglia isolated from adult rat brain. *Immunology*. 68:421-426.
- Wu CH, Wen CY, Shieh JY und Ling EA (1994) Down-regulation of membrane glycoprotein in ameboid microglia transforming into ramified microglia in postnatal rat brain. *J.Neurocytol.* 23:258-269.
- Wu DT, Woodman SE, Weiss JM, McManus CM, D'Aversa TG, Hesselgesser J, Major EO, Nath A und Berman JW (2000) Mechanisms of leukocyte trafficking into the CNS. *J.Neurovirol.* 6: S82-85.
- Xia Z, Dickens M, Raingeaud J, Davis RJ und Greenberg ME (1995) Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270:1326-1331.
- Xia MQ und Hyman BT (1999) Chemokines/chemokine receptors in the central nervous system and Alzheimer's disease. *J.Neurovirol.* 5:32-41.
- Xia MQ, Baeski BJ, Knowles RB, Qin SX und Hyman BT (2000) Expression of the chemokine receptor CXCR3 on neurons and the elevated expression of its ligand IP-10 in reactive astrocytes: in vitro ERK1/2 activation and role in Alzheimer's disease. *J.Neuroimmunol.* 108:227-235.
- y Cayal R (1901) Estudios sobre la corteza cerebral humana. IV. Estructura de la corteza cerebral olfativa del hombre y mamíferos. *Trab.Lab.Invest.Biol.Univ.Madrid* 1:1-140.
- Yoshida R, Nagira M, Kitaura M, Imagawa N, Imai T und Yoshie O (1998) Secondary lymphoid-tissue chemokine is a functional ligand for the CC chemokine receptor CCR7. *J.Biol.Chem.* 273:7118-7122.
- Zipp F, Nitsch R, Soriano E und Frotscher M (1989) Entorhinal fibers form synaptic contacts on parvalbumin-immunoreactive neurons in the rat fascia dentate. *Brain Res.* 495:161-166.
- Zou YR, Kottmann AH, Kuroda M, Taniuchi I und Littman DR (1998) Function of the chemokine receptor CXCR4 in hematopoiesis and in cerebellar development. *Nature* 393:595-599.