

# References

- Aceñolaza, F.G. (2000): La Formación Paraná (Miocene medio: estratigrafía, distribución regional y unidades equivalentes). In: Aceñolaza, F.G. and Herbst, R. (Editor), El Neógeno de Argentina. Serie Correlación Geológica, v. 14, p. 9-28.
- Aceñolaza, F. and Sprechmann, P. (2002): The Miocene marine transgression in the meridional Atlantic of South America. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, v. 225, n. 1, p. 75-84.
- Allmendinger, R.W., Jordan, T.E., Kay, S.M., and Isacks, B.L. (1997): The Evolution of the Altiplano-Puna Region of the central Andes. *Annual Review of Earth and Planetary Sciences*, v. 25, p. 139-174.
- Alonso, R.N. (2000): El terciario de la Puna en tiempos de la ingresión marina paranense. In: Aceñolaza, F.G. and Herbst, R. (Editor), El Neógeno de Argentina. Serie Correlación Geológica, v. 14, p. 163-180.
- Baby, P., Rochat, P., Mascle, G., and Héral, G. (1997): Neogene shortening contribution to crustal thickening in the back arc of the Central Andes. *Geology*, v. 25, n. 10, p. 883-886.
- Bachmann, G.H. and Müller, M. (1992): Sedimentary and structural evolution of the German Molasse Basin. *Eclogae Geologicae Helvetiae*, v. 85, p. 519-530.
- Bandel, K. and Hoefs, J. (1975): Die Isotopenzusammensetzung von Molluskenschalen am Beispiel der Gastropoden. *Neues Jahrbuch für Geologie und Paläontologie*, v. 1, p. 1-11.
- Beaumont, C. (1981): Foreland basins. *Geophysical Journal of the Royal Astronomical Society*, v. 65, p. 291-329.
- Bilodeau, W. L. and Keith, S. B. (1986): Lower Jurassic Navajo-Aztec-equivalent sandstones in southern Arizona and their paleogeographic significance. *American Association of Petroleum Geologists Bulletin*, v. 70, n. 6, p. 690-701.
- Blatt, H. (1967): Provenance determinations and recycling of sediments. *Journal of Sedimentary Petrology*, v. 37, p. 1031-1040.
- Blatt, H. and Tracy, R.J. (1995): Petrology. W. H. Freeman and Company, New York, pp. 529.
- Carey, S.W. (1958): A tectonic approach to continental drift. In: Carey, S.W (Editor), Continental drift. A symposium. Hobart, Tasmania. p. 177-355.
- Catuneanu, O., Sweet, A.R., and Miall, A.D. (1997): Reciprocal architecture bearpaw T-P sequences, uppermost Cretaceous, western Canada Sedimentary Basin. *Bulletin of Canadian Petroleum Geology*, v. 45, n. 1, p. 75-94.

- Catuneanu, O., Sweet, A.R., and Miall, A.D. (1999): Concept and styles of reciprocal stratigraphies: Western Canada foreland basin. *Terra Nova*, v. 11, p. 1-8.
- Chivas, A.R., Deckker, P., Cali, J.A., Chapman, A., Kiss, E., and Shelley, J.M.G. (1993): Coupled stable-isotope and trace-element measurements of lacustrine carbonates as paleoclimatic indicator. In: Swart, P.K., Lohmann, K.C., McKenzie, J., and Savin, S. (Editor), *Climate Change in Continental Isotopic Records*. Geophysical Monograph, v. 78, p. 113–122.
- Coira, B., Davidson, J., Mpodozis, C., and Ramos, V. (1982): Tectonic and magmatic evolution of the Andes of northern Chile and Argentina. *Earth-Science Reviews*, v. 18, p. 303-332.
- Coira, B., Kay, S.M., and Viramonte, J. (1993): Upper Cenozoic magmatic evolution of the Argentine Puna—A model for changing subduction geometry. *International Geology Review*, v. 35, p. 677–720.
- Coplen, T.B., Kendall, C., and Hopple, J. (1983): Comparison of stable isotope reference samples. *Nature*, v. 302, p. 236–38.
- Coudert, L., Frappa, M., Viguier, C., and Arias, R. (1995): Tectonic subsidence and crustal flexure in the Neogene Chaco basin of Bolivia. *Tectonophysics*, v. 243, p. 277-292.
- Cox, R. and Lowe, D.R. (1995): A conceptual review of regional-scale controls on the composition of clastic sediment and the co-evolution of the continental blocks and their sedimentary cover. *Journal of Sedimentary Research*, v. A65, n. 1, p. 1-12.
- Craig, H. (1961): Isotopic variations in meteoric waters. *Science*, v. 133, p. 1702–1703.
- Dalrymple, G.B. and Lanphere, M.A. (1974):  $^{40}\text{Ar}/^{39}\text{Ar}$  age spectra of some undisturbed terrestrial samples. *Geochimica et Cosmochimica Acta*, v. 38, n. 5, p. 715-738.
- DeCelles, P.G. and Gilles, K.A. (1996): Foreland basins systems. *Basin Research*, v. 8, p. 105-123.
- DeCelles, P.G. and Horton, B.K. (2003): Early to middle Tertiary foreland basin development and the history of Andean crustal shortening in Bolivia. *Geological Society of American Bulletin*, v. 115, n. 1, p. 58-77.
- Dickinson, W.R. (1970): Interpreting detrital modes of graywacke and arkose. *Journal of Sedimentary Petrology*, v. 40, p. 695-707.
- Dickinson, W.R. and Suczek, C.A. (1979): Plate tectonics and sandstone compositions. *American Association of Petroleum Geologists Bulletin*, v. 63, p. 2164-2182.
- Dickinson, W.R. (1985): Interpreting provenance relations from detrital modes of sandstones. In: Zuffa, G.G. (Editor), *Provenance of arenites*. D. Reidel Publishing Company, Dordrecht, Boston, Lancaster, p. 333-361.
- Dunn, J.F., Hartshorn, K.G., and Hartshorn, P.W. (1995): Structural styles and hydrocarbon potential of the Subandean thrust belt of Southern Bolivia. In: Tankhard, A.J., Suárez-Soruco, R., and Welsink, H.J. (Editor), *Petroleum basins of South America*. American Association of Petroleum Geologists Memoir, v. 62, p. 523-543.

- Echavarria, L., Hernández, R., Allmendinger, R., and Reynolds, J. (2003): Subandean thrust and fold belt of northwestern Argentina: Geometry and timing of the Andean evolution. American Association of Petroleum Geologists Bulletin, v. 87, n. 6, p. 965-985.
- Ege, H. (2004): Exhumations- und Hebungsgeschichte der zentralen Anden in Südbolivien ( $21^{\circ}$ S) durch Spaltspur-Thermochronologie an Apatit. Ph. D. Thesis, Freie Universität Berlin, Germany, pp. 173.
- Ege, H., Sobel, E.R., Scheuber, E., and Jacobshagen, V. (in preparation): AFT-data from the Andean foreland thrust belt of S-Bolivia ( $21^{\circ}$ S).
- Epstein, S., Buchsbaum, R., Lowenstam, H.A., and Urey, H.C. (1953): Revised carbonate water isotopic temperature scale. Geological Society of American Bulletin, v. 64, p. 1315-1326.
- Erdtmann, B.-D., Kley, J., Müller, J., and Jacobshagen, V. (1995): Ordovician basin dynamics and new graptolite data from the Tarija region, Eastern Cordillera, south Bolivia. In: Cooper, J.D., Drosser, M.L., Finney, S.C. (Editors), Ordovician Odyssey. Short papers from the Seventh International symposium on the Ordovician System. Society of Sedimentary Research, Pacific Section, Fullerton, CA, p. 69-73.
- Erikson, J.P. and Kelley, S.A. (1995): Late Oligocene initiation of foreland-basin formation in Bolivia based on newly dated volcanic ash. Congreso Latinoamericano de Geología, Ministerio de Energía y Minas, Caracas, Venezuela.
- Fastovsky, D., Arthur, M., Strater, N., and Foss, A. (1993): Freshwater bivalves (Unionidae), disequilibrium isotopic fractionation, and temperatures. Palaios, v. 8, p. 602-608.
- Faure, G. (1986). Principles of Isotope Geology. John Wiley, New York, p. 66-116.
- Finger, K. L. (1990): Atlas of California Neogene Foraminifera. Cushman Foundation for Foraminiferal Research Special Publications, v. 28, pp. 271.
- Finger, K. L. (1992): Biostratigraphic atlas of Miocene foraminifera from the Monterey and Modelo Formations, Central and Southern California. Cushman Foundation for Foraminiferal Research Special Publications, v. 29, pp. 179.
- Flemings, P.B. and Jordan, T.E. (1989): A synthetic model of foreland basin development. Journal of Geophysical Research, v. 94, n. B4, p. 3851-3866.
- Flemings, P.B. and Jordan, T.E. (1990): Stratigraphic modeling of foreland basins: Interpreting thrust deformation and lithospheric rheology. Geology, v. 18, p. 430-434.
- Folk, R.L. (1951): Textural maturity of sands. Journal of Sedimentary Petrology, v. 21, p. 128.
- Folk, R. L. (1974): Sandstones. Macropedia, v. 16, p. 212-216.
- Fritz, P. and Poplawski, S. (1974):  $^{18}\text{O}$  and  $^{13}\text{C}$  in the shells of freshwater molluscs and their environment. Earth and Planetary Science Letters, v. 24, p. 91-98.

Galli, C.I., Hernández, R.M., and Reynolds, J.H. (1996): Análisis paleoambiental y ubicación geocronológica del Subgrupo Metán (Grupo Orán, Neógeno) en el río Piedras, departamento Metán: Salta, Argentina. Boletín de Informaciones Petroleras, Tercera Epoca, v. 12, n. 46, p. 99-107.

Garrasino, C.A. and Vrba, A.V. (2000): La Formación Paraná: aspectos estratigráficos y estructurales de la región chacoparanense. In: Aceñolaza, F.G. and Herbst, R. (Editor), El Neógeno de Argentina. Serie Correlación Geologica, v. 14, p. 139-146.

Gazzi, P. (1966): Le arenite del flysch sopracretaceo dell' Apenino modense: correlazione con il flysch di Monghidoro. Mineralogiez e Petrografica Acta, v. 16, p. 69-97.

Gingras, M., Räsänen, M., Pemberton, S., and Romero, L. (2002a). The Significance of the Bioturbated Inclined Heterolithic Stratification in the Southern Part of the Miocene Solimoes Formation, Rio Acre, Amazonia, Brazil. *Palaios*, v. 17, p. 591-601.

Gingras, M., Räsänen, M., Ranzi, A. (2002b). Ichnology and Sedimentology reveal depositional characteristics of bay-margin parasequences in the Miocene Amazonian foreland basin. *Journal of Sedimentary Research*, v. 72, n. 6, p. 871-883.

Gubbels, R.W., Isacks, B.L., and Farrar, E. (1993): High-level surfaces, plateau uplift, and foreland development, Bolivian central Andes. *Geology*, v. 21, p. 695-698.

Günther, A. (2001): Strukturgeometrie, Kinematik und Deformationsgeschichte des oberkretazisch-alttertiären magmatischen Bogens (nord-chilenische Prækordillere, 21,7-23°S). Ph. D. Thesis, Freie Universität Berlin, Germany, Berliner Geowissenschaftliche Abhandlungen, A213, pp. 170.

Gupta, S. and Allen, P.A. (2000): Implications of foreland paleotopography for stratigraphic development in the Eocene distal Alpine foreland basin. *Geological Society of American Bulletin*, v. 112, p. 515-530.

Haq, B.U., Hardenbol, J., and Vail, P.R. (1988): Mesozoic and Cenozoic Chronostratigraphy and Cycles of Sea-level Change. In: Wilgus, C.K. (Editor), *Sea-level changes: an integrated approach*. Society of Economic Paleontologists and Mineralogists, Special Publication, v. 42, p. 71-108.

Herbst, R., Zabert, L.L., and Farrar, E. (1987): Microfaunas de la Formacion Paraná (Miocene Superior) de la cuenca Chaco-Paranense (Argentina). Facultad de Ciencias Exactas y Naturales del Litoral, Revista, v. 7, p. 165-206.

Héral, G., Oller, J., Baby, P., Bonhomme, M., and Soler, P. (1996): Strike-slip faulting, thrusting and related basins in the Cenozoic evolution of the southern branch of the Bolivian Orocline. *Tectonophysics*, v. 259, p. 201-212.

Hernández, R. M., Galli, C.I., and Reynolds, J. (1999): Estratigraphía del Terciario en el noroeste Argentino. In: Asociación Geológica Argentina (Editor), *Geología del Noroeste Argentino*, Congreso Geológica Argentina, v. 14, p. 316-328.

- Hinsch, R. (2001): Frontal Akkretion und Verformungsaufteilung eines gebogenen Falten- und Überschiebungsgürtels. Scientific Technical Report STR01/15, GeoForschungsZentrum Potsdam, Germany, pp. 128.
- Hoorn, C. (1993): Marine incursions and the influence of Andean tectonics on the Miocene depositional history of northwestern Amazonia: results of a palynostratigraphic study. *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 105, p. 267-309.
- Hoorn, C. (1994a): An environmental reconstruction of the paleo-Amazon River system (Middle-Late Miocene, NW Amazonia). *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 112, p. 187-238.
- Hoorn, C. (1994b): Fluvial palaeoenvironment in the intracratonic Amazon Basin (Early Miocene-early Middle Miocene), Columbia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 109, p. 1-54.
- Horton, B.K. and DeCelles, P.G. (1997): The modern foreland basin system adjacent to the Central Andes. *Geology*, v. 25, n. 10, p. 895-898.
- Horton, B.K. (1998): Sediment accumulation on top of the Andean orogenic wedge: Oligocene to late Miocene basins of the Eastern Cordillera, southern Bolivia. *Geological Society of America Bulletin*, v. 110, n. 9, p. 1174- 1192.
- Horton, B.K. and DeCelles, P.G. (2001). Modern and ancient fluvial megafans in the foreland basin system of the central Andes, southern Bolivia: Implications for drainage network evolution in the fold-thrust belts. *Basin Research*, v. 13, p. 43-63.
- Horton, B.K., Hampton, B.A., and Waanders, G.L. (2001): Paleogene synorogenic sedimentation in the Altiplano plateau and implications for initial mountain building in the central Andes. *Geological Society of America Bulletin*, v. 113, n. 11, p. 1387-1400.
- Hulka, C., Gräfe, K.-U., Sames, B., Uba C., and Heubeck, C. (in press): Depositional setting of the middle to late Miocene Yecua Formation of the Chaco Foreland basin, southern Bolivia. *Journal of South American Earth Science*.
- Husson, L. and Moretti, I. (2002): Thermal regime of fold and thrust belts - an application to the Bolivian sub Andean zone. *Tectonophysics*, v. 345, p. 253-280.
- Ihering, H. (1927): Die Geschichte des Atlantischen Ozeans. Gustav Fischer Verlag. Jena.
- Ingersoll, R.V., Bullard, T., Ford, R., Grimm, J., Pickle, J., and Sares, S. (1984): The effect of grain size on the detrital modes: a test of the Gazzi-Dickinson point-counting method. *Journal of Sedimentary Petrology*, v. 54, p. 102-116.
- Isacks, B.L. (1988): Uplift of the Central Andean Plateau and Bending of the Bolivian Orocline. *Journal of Geophysical Research*, v. 93, n. B4, p. 3211-3231.
- James, D.E. (1971): Plate tectonic model for the evolution of the Central Andes. *Geological Society of America Bulletin*, v. 82, p. 3325-3346.

Jacobshagen, V., Müller, J., Wemmer, K., Ahrendt, H., and Manutsoglu, E. (2002): Hercynian deformation and metamorphism in the Cordillera Oriental of Southern Bolivia, Central Andes. *Tectonophysics*, v. 345, n. 1-4, p. 119-130.

Johnsson, M. J. (1990): Tectonics versus chemical-weathering controls on the composition of fluvial sands in tropical environments. *Sedimentology*, v. 37, n. 4, p. 713-726.

Johnsson, M.J., Stallard, R.F., and Lundberg, N. (1991): Controls on the composition of fluvial sands from a tropical weathering environment: Sands of the Orinoco River drainage basin, Venezuela and Colombia. *Journal of Geology*, v. 96, p. 263-277.

Jones, J. (1985): The southern border of the Guaporé Shield in western Brazil and Bolivia: An interpretation of its geologic evolution. *Precambrian Research*, v. 85, p. 111-135.

Jordan, T.E. (1981): Thrust loads and foreland basin evolution, Cretaceous, western United States. *American Association of Petroleum Geologists Bulletin*, v. 65, p. 2506-2520.

Jordan, T.E. and Flemings, P.B. (1991): Large-scale stratigraphic architecture, eustatic variations, and unsteady tectonism: a theoretical evaluation. *Journal of Geophysical Research*, v. 96, p. 6681-6699.

Jordan, T.E., Reynolds, J.H., and Erikson, J.P. (1997): Variability in age of initial shortening and uplift in the central Andes, 16-33°30'S. In: Ruddiman, W.F. (Editor), *Tectonic uplift and Climate change*. Plenum Press, p. 41-61.

Kennan, L., Lamb, S., and Rundle, C. (1995): K-Ar dates from the Altiplano and Cordillera Oriental of Bolivia: Implications for Cenozoic stratigraphy and tectonics. *Journal of South American Earth Sciences*, v. 8, p. 163-186.

Kinsman, D.J. (1969): Interpretation of Sr<sup>2+</sup> concentrations in carbonate minerals and rocks. *Journal of Sedimentary Petrology*, v. 39, p. 486-508.

Kley, J. (1993): Der Übergang vom Subandin zur Ostkordillere in Südbolivien (21°15-22° S): Geologische Struktur und Kinematik. Ph. D. Thesis, Freie Universität Berlin, Germany, pp. 88.

Kley, J. and Reinhardt, M. (1994): Geothermal and tectonic evolution of the eastern Cordillera and the Subandean Ranges of southern Bolivia. In: Reutter, K.J., Scheuber, E., and Wigger, P. (Editor), *Tectonics of the South Central Andes*. Springer-Verlag, Berlin, p. 155-170.

Kley, J. (1996): Transition from basement-involved to thin-skinned thrusting in the Cordillera Oriental of southern Bolivia. *Tectonics*, v. 15, n. 4, p. 763-775.

Kley, J., Müller, J., Tawackoli, S., Jacobshagen, V., and Manutsoglu, E. (1997): Pre-Andean and Andean-age Deformation in the Eastern Cordillera of Southern Bolivia. *Journal of South American Earth Sciences*, v. 10, n. 1, p. 1-19.

Kley, J. (1999): Geological and geometric constraints on a kinematic model of the Bolivian orocline. *Journal of South American Earth Sciences*, v. 12, p. 221-235.

- Lamb, S.H. and Hoke, L. (1997): Origin of the high plateau in the Central Andes, Bolivia, Southern America. *Tectonics*, v. 16, n. 4, p. 623-649.
- Lamb, S.H., Hoke, L., Kennan, L., and Dewey, J. (1997): Cenozoic evolution of the central Andes in Bolivia and northern Chile. In: Burg, J.-P. and Ford, M. (Editors), *Orogeny Through Time*. Geological Society of Special Publications, v. 121, p. 237-264.
- Lee, J.K.W., Onstott, T.C., Cashman, K.V., Cumbe, R.J., and Johnson, D. (1991): Incremental heating of hornblende in vacuo; implications for  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology and the interpretation of thermal histories. *Geology*, v. 19, n. 9, p. 872-876.
- Li, L., Keller, G., Adatte, T., and Stennesbeck, W. (2000): Late Cretaceous sea-level changes in Tunisia: a multi-disciplinary approach. *Journal of Geological Society, London*, v. 157, p. 447-457.
- Lo, C.-H. and Onstott, T.C. (1989):  $^{39}\text{Ar}$  recoil of artefacts in chloritized biotite. *Geochemistry and Cosmochimistry*, v. 53, p. 2697-2711.
- Lo, C.-H. and Onstott, T.C. (1995): Rejuvenation of K-Ar for minerals in the Taiwan Mountain Belt. *Earth and Planetary Science Letter*, v. 131, p. 71-98.
- Longinelli, A. and Edmond, J. (1983): Isotope geochemistry of the Amazon Basin: a reconnaissance. *Journal of Geophysical Research*, v. 88, p. 3701-3717.
- Lovejoy, N.R., Bermingham, E., and Martin, A.P. (1998): Marine incursion into South American. *Nature*, v. 396, p. 421-422.
- Mack, G. H. (1984): Exceptions to the relationship between plate tectonics and sandstone composition. *Journal Sedimentary Petrology*, v. 54, p. 212 - 220.
- Marengo, H.G. (2000): Rasgos micropaleontológicos de los depósitos de la transgresión Enterriense-Paranense en la cuenca Chaco-Paranense y Noroeste Argentino. In: Aceñolaza, F.G. and Herbst, R. (Editor), *El Neógeno de Argentina. Serie Correlación Geológica*, v. 14, p. 29-45.
- Marshall, L.G. and Sempere, T. (1991): The Eocene to Pleistocene vertebrates of Bolivia and their stratigraphic context: A review. In: Suárez-Soruco, R. (Editor), *Fósiles y facies de Bolivia. Revista Técnica de Yacimientos Petrolíferos Fiscales Bolivianos*, v. 1, p. 631-652.
- Marshall, L.G., Sempere, T., and Gayet, M. (1993): The Petaca (late Oligocene-middle Miocene) and Yecua (late Miocene) formations of the Subandean Chaco basin, Bolivia and their tectonic significances. *Documents du Laboratoire de Géologie de Lyon*, v. 125, p. 291-301.
- McConaughey, T. (1989):  $^{13}\text{C}$  and  $^{18}\text{O}$  isotopic disequilibrium in biological carbonates: I. Patterns. *Geochimica et Cosmochimica Acta*, v. 53, p. 151-162.
- McQuarrie, N. (2002): The kinematic history of the central Andean fold-thrust belt, Bolivia: Implications for building a high plateau. *Geological Society of America Bulletin*, v. 114, n. 8, p. 950-963.

McQuarrie, N., Horton, B.K., Zandt, G., Beck, S., and DeCelles, P.G. (2005): Lithospheric evolution of the Andean fold-thrust belt, Bolivia, and the origin of the central Andean plateau. *Tectonophysics*, v. 399, p.15–37.

Meisch, C. (2000): Freshwater Ostracoda of Western and Central Europe. In: Schwoerbel, J. and Zwick, P.(Editors), *Suesswasserfauna von Mitteleuropa*. Spektrum Akademischer Verlag, Heidelberg – Berlin, pp. 522.

Meister, A. (1997): Lebenszyklus, Autökologie und Populationsökologie der Körbchenmusche *Corbicula fluminea* and *Corbicula fluminali* (Bivalvia, Corbiculidae) im Inselrhein. *Umweltplanung, Arbeits- und Umweltschutz*, v. 238, pp. 170.

Merrihue, C. (1965): Trace-element determinations and potassium-argon dating by mass spectroscopy of neutronirradiated samples: *Transactions - American Geophysical Union*, v. 46, p. 125.

Merrihue, C. and Turner, G. (1966): Potassium-argon dating by activation with fast neutrons. *Journal of Geophysical Research*, v. 71, p. 2852-2857.

Miall, A.D. (1997): The geology of stratigraphic sequences. Springer-Verlag. Berlin, pp. 433.

Moretti, I., Baby, P., Mendez, E., and Zubieta, D. (1996): Hydrocarbon generation in relation to thrusting in the Sub Andean Zone from 18° to 22°S, Bolivia. *Petroleum Geoscience*, v. 2, p. 17-28.

Müller, J., Kley, J., and Jacobshagen, V. (2002): Structure and Cenozoic kinematics of the eastern Cordillera, southern Bolivia (21°S). *Tectonics*, v. 21, n. 5, p. 1-23.

Muñoz, N. and Charrier, R. (1996): Uplift of the western border of the Altiplano on a west-vergent thrust system, northern Chile. *Journal of South American Earth Sciences*, v. 9, n. 3-4, p. 171-181.

Oliver, P.G. (1992): Bivalved Seashells of the Red Sea. Christa Verlag, Hemmen, Wiesbaden, pp. 330.

Palmer, M.R. and Edmond, J.M. (1992): Controls on the strontium isotope composition of river water. *Geochemistry and Cosmochemistry*, v. 56, p. 2099-2111.

Pareja, J., Vargas, C., Suárez, R., Ballón, R., Carrasco, R., and Villareol, C. (1978): Mapa geológico de Bolivia y memoria explicativa. Yacimientos Petrolíferos Fiscales Bolivianos e Servicio Geológico de Bolivia, 1:1.000.000, pp. 27.

Pettijohn, F. J. (1975): Sedimentary rocks. Harper and Row Publications, New York, pp. 628.

Räsänen, M.E., Linna, A.M., Santos, J.C.R., and Negri, F.R. (1995): Miocene deposits in the Amazonian Foreland Basin. *Sciences*, v. 269, p. 386-389.

Reutter, K.-J., Döbel, R., Bogdanic, T., and Kley, J. (1994): Geological map of the Central Andes between 20° and 26°. In: Reutter, K.-J., Scheuber, E., and Wigger, P.J. (Editor), *Tectonics of the southern Central Andes*. Springer-Verlag, Berlin-Heidelberg, p. 121-139.

- Reynolds, J.H., Idleman, B.D., Hernández, R.M., and Naeser, C.W. (1994): Preliminary chronostratigraphic constraints on Neogene tectonic activity in the Eastern Cordillera and Santa Bárbara System, Salta Province, NW Argentina. Geological Society of America Abstract, v. 26, n. 7, p.A-503.
- Reynolds, J.H., Galli, C.I., Hernández, R.M., Idleman, B.D., Kotila, J.M., Hilliard R.V., and Naeser, C.W. (2000): Middle Miocene tectonic development of the transition zone, Salta Province, northwestern Argentina: magnetic stratigraphy from Métan Subgroup, Sierras González. Geological Society Memoir, v. 10, p. 297-311.
- Rochat, P., Heraïl, G., Baby, P., and Mascle, G. (1999): Bilan crustal et contrôlé de la dynamique érosive et sédimentaire sur les mécanismes de formation de l'Altiplano. Comptes Rendus de l'Academie des Sciences, Serie II. Sciences de la Terre et des Planètes, v. 328, n. 3, p. 189-195.
- Roeder, D. (1988): Andean-age structure of the Eastern Cordillera (Province of La Paz, Bolivia). Tectonics, v. 7, n. 1, p. 23-39.
- Scasso, R.A., McArthur, J.M., del Río, C.J., Martínez, S., and Thirwall, M.S. (2001):  $^{87}\text{Sr}/^{86}\text{Sr}$  Late Miocene age of fossil molluscs in the Entrerriense of the Valdés Peninsula. Journal of South American Earth Sciences, v.14, p. 319-329.
- Scheuber, E. and Reutter, K.-J. (1992): Magmatic arc tectonics in the central andes between 21° and 25°S. Tectonophysics, v. 205, p. 127-140.
- Scheuber, E. and Giese, P. (1999): Architecture of the Central Andes - A compilation of geoscientific data along a transect at 21°S. Journal of South American Earth Sciences, v. 12, p. 103-107.
- Schmidowski, M., Eichmann, R., and Junge, C.E. (1976): Carbon isotope geochemistry of the Precambrian Lomagundi carbonate province, Rhodesia. Geochimistry and Cosmochimistry Acta, v. 40, p. 449-455.
- Schmitz, M. (1994): A balanced model of the southern Central Andes. Tectonics, v.13, n. 2, p. 484-492.
- Schmitz, M. and Kley J. (1997): The geometry of the central Andean backarc crust: joint interpretation of cross-section balancing and seismic refraction data. Journal of South American Earth Sciences, v. 10, p. 99-110.
- Sclater, J.G. and Christie, P.A.F. (1980): Continental stretching: an explanation of the post Mid-Cretaceous subsidence of the central North Sea basin. Journal of Geophysical Research, v. 85, p. 3711-3739.
- Sempere, T., Heraïl, G., Oller, J., and Bonhomme, M.G. (1990): Late Oligocene-early Miocene major tectonic crisis and related basins in Bolivia. Geology, v. 18, p. 946-949.
- Sempere, T., Butler, R.F., Richards, D.R., Marshall, L.G., Sharp, W., Swisher, C.C. (1997): Stratigraphy and chronology of Upper Cretaceous-lower Paleogene strata in Bolivia and northwest Argentina. Geological Society of America Bulletin, v. 109, n. 6, p. 709-727.

- Sheffels, B.M. (1990): Lower bound on the amount of crustal shortening in the central Bolivian Andes. *Geology*, v. 18, p. 812-815.
- Sinclair, H.D., Coakley, B.J., Allen, P.A., Watts, A.B. (1991): Simulation of foreland basin stratigraphy using a diffusion model of mountain belt uplift and erosion: an example from the central Alps, Switzerland. *Tectonics*, v. 10, p. 599-620.
- Sinclair, H.D. and Allen, P.A. (1992): Vertical versus horizontal motions in the Alpine orogenic wedge: stratigraphic response in the foreland basin. *Basin Research*, v. 4, p. 215-232.
- Sinclair, H.D. (1997): Tectonostratigraphic model for underfilled peripheral foreland basins: An Alpine perspective. *Geological Society of America Bulletin*, v. 109, n. 3, p. 324-346.
- Sprechmann, P. (1978): The paleoecology and palaeogeography of the Uruguayan coastal area during the Neogene and Quaternary. *Zitteliana*, v. 4, p. 3-72.
- Sprechmann, P., Ferrando, L.A., and Martínez, S. (2000): Estado actual de los conocimientos sobre la Formación Camacho (Mioceno?-superior?), Uruguay. In: Aceñolaza, F.G. and Herbst, R. (Editor), *El Neógeno de Argentina. Serie Correlación Geologica*, v. 14, p. 47-65.
- Steinmann, M., Hungerbühler, D., Seward, D., and Winkler, W. (1999): Neogene tectonic evolution and exhumation of the southern Andes: a combined stratigraphy and fission-track approach. *Tectonophysics*, v. 307, p. 255-276.
- Stockmal, G. and Beaumont, C. (1987): Geodynamic models of convergent margin tectonics; the southern Canadian Cordillera and the Swiss Alps. In: Beaumont, C. and Tankard, A., (Editor), *Sedimentary Basins and Basin-Forming Mechanisms*. Canadian Society of Petroleum Geologists Memoir, v. 12, p. 393-412.
- Suárez-Soruco, R. (2000): Compendio de Geología de Bolivia. *Revista Técnica de Yacimientos Petrolíferos Fiscales Bolivianos*, v. 18, n. 1-2, p. 1-144.
- Suttner, L. J. and Basu, A. (1981): Climate and the origin of quartz arenites. *Journal of Sedimentary Petrology*, v. 51, p. 1235-1246.
- Tanaka, N., Monaghan, M.C., and Rye, D.M. (1986): Contribution of metabolic carbon to mollusc and barnacle shell carbonate. *Nature*, v. 320, p. 520-523.
- Uba, C.E., Heubeck, C., and Hulka, C. (in revision): Facies analysis and basin architecture of the Neogene Subandean synorogenic wedge, southern Bolivia. *Sedimentary Geology*.
- Uba, C.E. (2005): Late Cenozoic evolution of the Chaco Foreland Basin, Bolivia. Ph. D. Thesis, Freie Universität Berlin, Germany, pp. 112.
- Urey, H.C., Lowenstam, H.A., Epstein, S., and McKinney, C.R. (1951): Measurement of paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark, and the southeastern United States. *Bulletin of the Geological Society of America*, v. 62, p. 399-416.

- Veizer, J., Ala, D., Azmy, K., Bruckschen, P., Bruhn, F., Buhl, D., Carden, G., Diener, A., Ebneth, S., Goddris, Y., Jasper, T., Korte, C., Pawellek, F., Podlaha, O., and Strauss, H. (1999):  $^{87}\text{Sr}/^{86}\text{Sr}$ ,  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  evolution of the Phanerozoic seawater. *Chemical Geology*, v. 161, p. 59-88.
- Vonhof, H.B., Wesselingh, F.P., and Ganssen, G.M. (1998): Reconstruction of the Miocene western Amazonian aquatic system using molluscan isotopic signatures. *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 141, p. 85-93.
- Vonhof, H.B., Wesselingh, F.P., Kaandorp, R.J.G., Davies, G.R., van Hinte, J.E., Guerrero, J., Räsänen, M., Romero-Pitman, L., and Ranzi, A. (2003): Palaeogeography of Miocene Western Amazonia: Isotopic molluscan shells constrain the influence of marine incursions. *Geological Society of American Bulletin*, v. 3, n. 20, p. 983-993.
- Waschbusch, P.J. and Royden L.H. (1992): Episodicity in foredeep basins. *Geology*, v. 20, n. 10, p. 915-918.
- Webb, S.D. (1995): Biological implications of the Middle Miocene Amazon seaway. *Science*, v. 269, p. 361-362.
- Wefer, D. (1985): Die Verteilung stabiler Isotope in Kalkschalen mariner Organismen. *Geologisches Jahrbuch*, v. A82, p. 3-111.
- Welsink, H.J., Franco, M.A., and Oviedo, G.C. (1995): Andean and pre-Andean deformation, Boomerang Hills area, Bolivia. In: Tankard, A.J., Suárez-Soroco, R., and Welsink, H.J. (Editors), *Petroleum Basins of South America*. American Association of Petroleum Geologists Memoir, v. 62, p. 481-499.
- Wesselingh, F.P., Räsänen, M.E., Irion, G., Vonhof, H.B., Kaandorp, R., Renema, W., Romero Pittman, L., and Gingras, M. (2002): Lake Pebas: a palaeoecological reconstruction of a Miocene, long-lived lake complex in western Amazonia. *Cainozoic Research*, v. 1, p. 35-81.
- York, D. (1969): Least squares fitting of a straight line with correlated errors. *Earth and Planetary Science Letters*, v. 5, p. 320-324.