

Literaturverzeichnis

- 1 Hübbel VF, Rhiem K, Warm M, Mallmann P, Schmutzler RK. Aktuelle Endwicklung der Inzidenz und Mortalität beim Mammakarzinom. Gynäkologe. 2003;36:746-752.
- 2 Fink D, Kimmig R, Winter R. Therapy of breast cancer--change from radical to quality-controlled individualized treatment. Gynakol Geburtshilfliche Rundsch. 2005;45(3):125-6.
- 3 Ernst MF, Voogd AC, Coebergh JW, Poortmans PM, Roukema JA. Using loco-regional recurrence as an indicator of the quality of breast cancer treatment. Eur J Cancer. 2004;40(4):487-93.
- 4 Leitch AM, Meek AG, Smith RA, et al. American Cancer Society Lymphedema Workshop. Workgroup I: Treatment of the axilla with surgery and radiation--preoperative and postoperative risk assessment. Cancer. 1998;83(12 Suppl American):2877-9.
- 5 Runowicz CD, Passik SD, Hann D, et al. American Cancer Society Lymphedema Workshop. Workgroup II: Patient education--pre-and posttreatment. Cancer. 1998;83(12 Suppl American):2880-1.
- 6 Halsted WS. The results of operations for the cure of cancer of the breast performed at Johns Hopkins Hospital from June 1889 to January 1894. Ann Surg. 1894; 20:497-555.
- 7 Halsted WS. Developments in skin-grafting operation for cancer of breast. J Am Med Assoc. 1913(60):416-8.
- 8 Jatoi I. Breast cancer: a systemic or local disease? Am J Clin Oncol. 1997;20(5):536-9.
- 9 Hirsch, J. Radiumchirurgie des Brustkrebses. Dtsch Med Wschr. 1927;53:1419-1421.
- 10 Leis HP Jr, Greene FL, Hilfer SE. Bedeutung modifiziert radikaler Mastektomietechniken. In: Bohmert H. Brustkrebs: Organerhaltung und Rekonstruktion. Stuttgart: Thieme, 1989:148-54.
- 11 Berg JW. The significance of axillary node levels in the study of breast carcinoma. Cancer. 1955;8(4):776-8.

-
- 12 Kodama H. Modification of muscle-preserving radical mastectomy. *Cancer*. 1979;44(4):1517-22.
 - 13 Auchincloss H. Modified radical mastectomy: why not? *Am J Surg*. 1970;119(5):506-9.
 - 14 Madden JL. Modified radical mastectomy. *Surg Gynecol Obstet*. 1965;121(6):1221-30.
 - 15 Robinson GN, van Heerden JA, Payne WS, Taylor WF, Gaffey TA. The primary surgical treatment of carcinoma of the breast: a changing trend toward modified radical mastectomy. *Mayo Clin Proc*. 1976;51(7):433-42.
 - 16 Patey DH. A review of 146 cases of carcinoma of the breast operated on between 1930 and 1943. *Br J Cancer*. 1967;21(2):260-9.
 - 17 Handley RS, Thackray AC. Conservative radial mastectomy (Patey's operation). *Ann Surg*. 1969;170(6):880-2.
 - 18 Nemoto T, Vana J, Bedwani RN, Baker HW, McGregor FH, Murphy GP. Management and survival of female breast cancer: results of a national survey by the American College of Surgeons. *Cancer*. 1980;45(12):2917-24.
 - 19 Auchincloss H. Significance of location and number of axillary metastases in carcinoma of the breast. *Ann Surg*. 1963;158:37-46.
 - 20 Urban JA, Baker HW. Radical mastectomy in continuity with en bloc resection of the internal mammary lymph-node chain; a new procedure for primary operable cancer of the breast. *Cancer*. 1952;5(5):992-1008.
 - 21 Urban J.A. Surgical management of palpable breast cancer. *Cancer*. 1980;46(4 Suppl):983-7.
 - 22 Lacour J, Le M, Caceres E, Koszarowski T, Veronesi U, Hill C. Radical mastectomy versus radical mastectomy plus internal mammary dissection. Ten year results of an international cooperative trial in breast cancer. *Cancer*. 1983;51(10):1941-3.
 - 23 Meier P, Ferguson DJ, Karrison T. A controlled trial of extended radical versus radical mastectomy. Ten-year results. *Cancer*. 1989;63(1):188-95.
 - 24 Morimoto T, Monden Y, Takashima S, et al. Five-year results of a randomized clinical trial comparing modified radical mastectomy and extended radical mastectomy for stage II breast cancer. *Surg Today*. 1994;24(3):210-4.

-
- 25 McWhirter R. Treatment of cancer of breast by simple mastectomy and roentgenotherapy. *Arch Surg.* 1949;59:830-42.
 - 26 McWhirter R. Cancer of the breast. *JRadiol Electrol Med Nucl.* 1967;48(11):768-9.
 - 27 Keynes G. Conservative treatment of cancer of the breast. *Br Med J.* 1937; 2:643.
 - 28 Peters MV. Wedge resection with or without radiation in early breast cancer. *Int J Radiat Oncol Biol Phys.* 1977;2(11-12):1151-6.
 - 29 Crile G Jr. Biological observations on the treatment of cancer, *Zentralbl Chir.* 1974;99(13):395-401.
 - 30 Calle R. The role of radiation therapy in the loco-regional treatment of breast cancer. *Recent Results Cancer Res.* 1976;(57):164-75.
 - 31 Pierquin B, Baillet F, Wilson JF. Radiation therapy in the management of primary breast cancer. *AJR Am J Roentgenol.* 1976;127(4):645-8.
 - 32 Spitalier JM, Gambarelli J, Brandone H, et al. Breast-conserving surgery with radiation therapy for operable mammary carcinoma: a 25-year experience. *World J Surg.* 1986;10(6):1014-20.
 - 33 Montague ED. Experience with altered fractionation in radiation therapy of breast cancer. *Radiology.* 1968;90(5):962-6.
 - 34 Hellman S. The place of radiation in the definitive management of carcinoma of the breast. *Int J Radiat Oncol Biol Phys.* 1977;2(3-4):369-70.
 - 35 Veronesi U, Saccozzi R, Del Vecchio M, et al. Comparing radical mastectomy with quadrantectomy, axillary dissection, and radiotherapy in patients with small cancers of the breast. *N Engl J Med.* 1981;305(1):6-11.
 - 36 Fisher B, Montague E, Redmond C, et al. Comparison of radical mastectomy with alternative treatments for primary breast cancer. A first report of results from a prospective randomized clinical trial. *Cancer.* 1977;39(6 Suppl):2827-39.
 - 37 Fisher B. From Halsted to prevention and beyond: advances in the management of breast cancer during the twentieth century. *Eur J Cancer.* 1999;35(14):1963-73.
 - 38 Hayward JL. The Guy's trial of treatments of "early" breast cancer. *World J Surg.* 1977;1(3):314-6.

-
- 39 van Dongen JA, Bartelink H, Fentiman IS, et al. Factors influencing local relapse and survival and results of salvage treatment after breast-conserving therapy in operable breast cancer: EORTC trial 10801, breast conservation compared with mastectomy in TNM stage I and II breast cancer. *Eur J Cancer*. 1992;28A(4-5):801-5.
- 40 Blichert-Toft M, Brincker H, Andersen JA, et al. A Danish randomized trial comparing breast-preserving therapy with mastectomy in mammary carcinoma. Preliminary results. *Acta Oncol*. 1988;27(6A):671-7.
- 41 Lichter AS, Lippman ME, Danforth DN Jr, et al. Mastectomy versus breast-conserving therapy in the treatment of stage I and II carcinoma of the breast: a randomized trial at the National Cancer Institute. *J Clin Oncol*. 1992;10(6):976-83.
- 42 Veronesi U, Cascinelli N, Mariani L, et al. Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med*. 2002;347(16):1227-32.
- 43 Fisher B, Anderson S, Bryant J, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med*. 2002;347(16):1233-41.
- 44 Fisher B, Fisher ER: Barrier function of lymph node to tumor cells and erythrocytes I. Normal nodes. *Cancer*. 1967;20(11):1907-1913.
- 45 Fisher B. Entwicklung, Technik und Ergebnisse der brusterhaltenden Therapie. In: Bohmert H. *Brustkrebs: Organerhaltung und Rekonstruktion*. Stuttgart: Thieme, 1989:42-63.
- 46 Hellman S. Karnofsky Memorial Lecture. Natural history of small breast cancers. *J Clin Oncol*. 1994;12(10):2229-34.
- 47 Sakorafas GH. Breast cancer surgery--historical evolution, current status and future perspectives. *Acta Oncol*. 2001;40(1):5-18.
- 48 Clemons M, Danson S, Hamilton T, Goss P. Locoregionally recurrent breast cancer: incidence, risk factors and survival. *Cancer Treat Rev*. 2001;27(2):67-82.
- 49 Willner J, Kiricuta IC, Kolbl O. Locoregional recurrence of breast cancer following mastectomy: always a fatal event? Results of univariate and multivariate analysis. *Int J Radiat Oncol Biol Phys*. 1997;37(4):853-63.
- 50 Keshtgar MR, Baum M. Axillary dissection over the years: where to from here? *World J Surg*. 2001;25(6):761-6.

-
- 51 Carter CL, Allen C, Henson DE. Relation of tumor size, lymph node status, and survival in 24,740 breast cancer cases. *Cancer*. 1989;63(1):181-7.
- 52 Clodius L. Minimizing secondary arm lymphedema from axillary dissection. *Lymphology*. 2001;34(3):106-10.
- 53 Meek AG. Breast radiotherapy and lymphedema. *Cancer*. 1998;83(12 Suppl American):2788-97.
- 54 Maunsell E, Brisson J, Deschenes L. Arm problems and psychological distress after surgery for breast cancer, *Can J Surg*. 1993;36(4):315-20.
- 55 Stewart FW, Treves N. Lymphangiosarcoma in postmastectomy lymphedema: a report of six cases in elephantiasis chirurgica. *Cancer*. 1948;1:64-81.
- 56 Heitmann C, Ingianni G. Stewart-Treves syndrome: lymphangiosarcoma following mastectomy. *Ann Plast Surg*. 2000;44(1):72-5.
- 57 Harwood CA, Mortimer PS. Causes and clinical manifestations of lymphatic failure. *Clin Dermatol*. 1995;13(5):459-71.
- 58 Petrek JA, Lerner R. Lymphedema. In: Harris JR, Lippman ME, Morrow M, Hellman S. *Diseases of the breast*. Philadelphia: Lippincott-Raven, 1995:896-900.
- 59 Pain SJ, Purushotham AD. Lymphoedema following surgery for breast cancer. *Br J Surg*. 2000;87(9):1128-41.
- 60 Schuchhardt C, Bimler E. Das sekundäre Lymphödem nach operativen gynäkologischen Eingriffen – Stiefkind der onkologischen Nachsorgebehandlung. *Gynäkologe*. 2003;36(6):496-506.
- 61 Földi E, Földi M. Lymphostatische Krankheitsbilder. In Földi M, Kubik S: *Lehrbuch der Lymphologie für Mediziner und Physiotherapeuten*. 4. Aufl. Stuttgart, Jena Lübeck, Ulm: G. Fischer, 1999:275-327.
- 62 Halsted WS. Swelling of arm after operation for cancer of breast – elephantiasis chirurgica – its cause and prevention. *Bull Johns Hopkins Hosp*. 1921;32:309-313.
- 63 Matas R. Surgical treatment of elephantiasis and elephantoid states dependent upon chronic obstruction of lymphatic and venous channels. *Am J Trop Dis*. 1913;1:60-85.
- 64 Földi E, Földi M. Das postmastektomische Armlymphödem. In: Schildberg FW, Kiffner E. *Interdisziplinäre Therapie des Mammakarzinoms*. Erlangen: Perimed Fachbuch-Verlagsgesellschaft, 1985:272-86.

-
- 65 van der Veen P, De Voogdt N, Lievens P, Duquet W, Lamote J, Sacre R. Lymphedema development following breast cancer surgery with full axillary resection. *Lymphology*. 2004;37(4):206-8.
- 66 Kissin MW, Querci della Rovere G, Easton D, Wesbury G. Risk of lymphoedema following the treatment of breast cancer. *Br J Surg*. 1986;73(7):580-4.
- 67 Segerstrom K, Bjerle P, Graffman S, Nystrom A. Factors that influence the incidence of brachial oedema after treatment of breast cancer. *Scand J Plast Reconstr Surg Hand Surg*. 1992;26(2):223-7.
- 68 Casley-Smith JR, Casley-Smith JR. Lymphedema initiated by aircraft flights. *Aviat Space Environ Med*. 1996;67(1):52-6.
- 69 Rockson SG. Precipitating factors in lymphedema: myths and realities. *Cancer*. 1998;83(12 Suppl American):2814-6.
- 70 Földi E, Földi M, Clodius L. The lymphedema chaos: a lancet. *Ann Plast Surg*. 1989;22(6):505-15.
- 71 Bunce IH, Mirolo BR, Hennessy JM, Ward LC, Jones LC. Post-mastectomy lymphoedema treatment and measurement. *Med J Aust*. 1994;161(2):125-8.
- 72 Brown J. A clinically useful method for evaluating lymphedema. *Clin J Oncol Nurs*. 2004;8(1):35-8.
- 73 Cheville AL, McGarvey CL, Petrek JA, Russo SA, Thiadens SR, Taylor ME. The grading of lymphedema in oncology clinical trials. *Semin Radiat Oncol*. 2003;13(3):214-25.
- 74 Harris SR, Hugi MR, Olivotto IA, Levine M; Steering Committee for Clinical Practice Guidelines for the Care and Treatment of Breast Cancer. Clinical practice guidelines for the care and treatment of breast cancer: 11. Lymphedema. *CMAJ*. 2001;164(2):191-9.
- 75 Kaulesar Sukul DM, den Hoed PT, Johannes EJ, van Dolder R, Benda E. Direct and indirect methods for the quantification of leg volume: comparison between water displacement volumetry, the disk model method and the frustum sign model method, using the correlation coefficient and the limits of agreement. *J Biomed Eng*. 1993;15(6):477-80.
- 76 Clodius L, Deak L, Piller NB. A new instrument for the evaluation of tissue tonicity in lymphoedema. *Lymphology*. 1976;9(1):1-5.

-
- 77 Piller NB, Clodius L. The use of a tissue tonometer as a diagnostic aid in extremity lymphoedema: a determination of its conservative treatment with benzopyrones. *Lymphology*. 1976;9(4):127-32.
- 78 Cornish BH, Chapman M, Hirst C, et al. Early diagnosis of lymphedema using multiple frequency bioimpedance. *Lymphology*. 2001;34(1):2-11.
- 79 Duewell S, Hagspiel KD, Zuber J, Schulthess GK, Bollinger A, Fuchs WA. Swollen lower extremity: role of MR imaging. *Radiology*. 1992;184(1):227-31.
- 80 Goltner E, Gass P, Haas JP, Schneider P. The importance of volumetry, lymphoscintigraphy and computer tomography in the diagnosis of brachial edema after mastectomy. *Lymphology*. 1988;21(3):134-43.
- 81 Filippetti M, Santoro E, Graziano F, Petric M, Rinaldi G. Modern therapeutic approaches to postmastectomy brachial lymphedema. *Microsurgery*. 1994;15(8):604-10.
- 82 Bourgeois P, Leduc O, Leduc A. Imaging techniques in the management and prevention of posttherapeutic upper limb edemas. *Cancer*. 1998;83(12 Suppl American):2805-13.
- 83 Stemmer R. A clinical symptom for the early and differential diagnosis of lymphedema. *Vasa*. 1976;5(3):261-2.
- 84 Stemmer R. Stemmer's sign--possibilities and limits of clinical diagnosis of lymphedema. *Wien Med Wochenschr*. 1999;149(2-4):85-6.
- 85 Rockson SG, Miller LT, Senie R, et al. American Cancer Society Lymphedema Workshop. Workgroup III: Diagnosis and management of lymphedema. *Cancer*. 1998;83(12 Suppl American):2882-5.
- 86 Petrek JA, Senie RT, Peters M, Rosen PP. Lymphedema in a cohort of breast carcinoma survivors 20 years after diagnosis. *Cancer*. 2001;92(6):1368-77.
- 87 Norman SA, Miller LT, Erikson HB, Norman MF, McCorkle R. Development and validation of a telephone questionnaire to characterize lymphedema in women treated for breast cancer. *Phys Ther*. 2001;81(6):1192-205.
- 88 Runowicz CD. Lymphedema: patient and provider education: current status and future trends. *Cancer*. 1998;83(12 Suppl American):2874-6.
- 89 Lobb AW, Harkins HN. Postmastectomy swelling of the arm; with a note on the effect of segmental resection of the axillary vein at the time of radical mastectomy. *West J Surg Obstet Gynecol*. 1949;57(11):550-7.

-
- 90 Daland EM. The incidence of swollen arms after radical mastectomy and suggestions for prevention *N Engl J Med*. 1950;242(13):497-502.
- 91 MacDonald I, Osman K. Postmastectomy lymphedema. *Am J Surg*. 1955;90(2):281-6.
- 92 Treves N. An evaluation of the etiological factors of lymphedema following radical mastectomy; an analysis of 1,007 cases. *Cancer*. 1957;10(3):444-59.
- 93 Britton RC, Nelson PA. Causes and treatment of post-mastectomy lymphedema of the arm. Report of 114 cases. *JAMA*. 1962;180:95-102.
- 94 Logan V. Incidence and prevalence of lymphoedema: a literature review. *J Clin Nurs*. 1995;4(4):213-9.
- 95 Petrek JA, Heelan MC. Incidence of breast carcinoma-related lymphedema. *Cancer*. 1998;83(12 Suppl American):2776-81.
- 96 Petrek JA, Pressman PI, Smith RA. Lymphedema: current issues in research and management. *CA Cancer J Clin*. 2000;50(5):292-307.
- 97 Erickson VS, Pearson ML, Ganz PA, et al. Arm edema in breast cancer patients. *J Natl Cancer Inst*. 2001;93(2):96-111.
- 98 Schünemann H, Willich N. Lymphedema after breast carcinoma. A study of 5868 cases. *Dtsch Med Wochenschr*. 1997;122(17):536-41.
- 99 Armer J, Fu MR, Wainstock JM, Zagar E, Jacobs LK. Lymphedema following breast cancer treatment, including sentinel lymph node biopsy. *Lymphology*. 2004;37(2):73-91.
- 100 Werner RS, McCormick B, Petrek J, et al. Arm edema in conservatively managed breast cancer: obesity is a major predictive factor. *Radiology*. 1991;180(1):177-84.
- 101 Rietman JS, Dijkstra PU, Hoekstra HJ, et al. Late morbidity after treatment of breast cancer in relation to daily activities and quality of life: a systematic review. *Eur J Surg Oncol*. 2003;29(3):229-38.
- 102 Swedborg I, Borg G, Sarnelid M. Somatic sensation and discomfort in the arm of post-mastectomy patients. *Scand J Rehabil Med*. 1981;13(1):23-9.
- 103 Hack TF, Cohen L, Katz J, Robson LS, Goss P. Physical and psychological morbidity after axillary lymph node dissection for breast cancer. *J Clin Oncol*. 1999;17(1):143-9.

-
- 104 Sugden EM, Rezvani M, Harrison JM, Hughes LK. Shoulder movement after the treatment of early stage breast cancer. *Clin Oncol (R Coll Radiol)*. 1998;10(3):173-81.
- 105 Gosselink R, Rouffaer L, Vanhelden P, Piot W, Troosters T, Christiaens MR. Recovery of upper limb function after axillary dissection. *J Surg Oncol*. 2003;83(4):204-11.
- 106 Jansen RF, van Geel AN, de Groot HG, Rottier AB, Olthuis GA, van Putten WL. Immediate versus delayed shoulder exercises after axillary lymph node dissection. *Am J Surg*. 1990;160(5):481-4.
- 107 Christodoulakis M, Sanidas E, de Bree E, Michalakis J, Volakakis E, Tsiftsis D. Axillary lymphadenectomy for breast cancer - the influence of shoulder mobilisation on lymphatic drainage. *Eur J Surg Oncol*. 2003;29(4):303-5.
- 108 Wingate L, Croghan I, Natarajan N, Michalek AM, Jordan C. Rehabilitation of the mastectomy patient: a randomized, blind, prospective study. *Arch Phys Med Rehabil*. 1989;70(1):21-4.
- 109 Box RC, Reul-Hirche HM, Bullock-Saxton JE, Furnival CM. Shoulder movement after breast cancer surgery: results of a randomised controlled study of postoperative physiotherapy. *Breast Cancer Res Treat*. 2002;75(1):35-50.
- 110 Tengrup I, Tennvall-Nittby L, Christiansson I, Lavrin M. Arm morbidity after breast-conserving therapy for breast cancer. *Acta Oncol*. 2000;39(3):393-7.
- 111 Ernst MF, Voogd AC, Balder W, Klinkenbijn JH, Roukema JA. Early and late morbidity associated with axillary levels I-III dissection in breast cancer. *J Surg Oncol*. 2002;79(3):151-6.
- 112 Tasmuth T, von Smitten K, Kalso E. Pain and other symptoms during the first year after radical and conservative surgery for breast cancer. *Br J Cancer*. 1996;74(12):2024-31.
- 113 Carpenter JS, Sloan P, Andrykowski MA, et al. Risk factors for pain after mastectomy/lumpectomy. *Cancer Pract*. 1999;7(2):66-70.
- 114 Freeman SR, Washington SJ, Pritchard T, Barr L, Baildam AD, Bundred NJ. Long term results of a randomised prospective study of preservation of the intercostobrachial nerve. *Eur J Surg Oncol*. 2003;29(3):213-5.
- 115 Taylor KO. Morbidity associated with axillary surgery for breast cancer. *ANZ J Surg*. 2004;74(5):314-7.

-
- 116 Torresan RZ, Cabello C, Conde DM, Brenelli HB. Impact of the preservation of the intercostobrachial nerve in axillary lymphadenectomy due to breast cancer. *Breast J.* 2003;9(5):389-92.
- 117 Engel J, Kerr J, Schlesinger-Raab A, Sauer H, Holzel D. Axilla surgery severely affects quality of life: results of a 5-year prospective study in breast cancer patients. *Breast Cancer Res Treat.* 2003;79(1):47-57.
- 118 Kwan W, Jackson J, Weir LM, Dingee C, McGregor G, Olivotto IA. Chronic arm morbidity after curative breast cancer treatment: prevalence and impact on quality of life. *J Clin Oncol.* 2002;20(20):4242-8.
- 119 Voogd AC, Ververs JM, Vingerhoets AJ, Roumen RM, Coebergh JW, Crommelin MA. Lymphoedema and reduced shoulder function as indicators of quality of life after axillary lymph node dissection for invasive breast cancer. *Br J Surg.* 2003;90(1):76-81.
- 120 Herfarth C, Bastert G. Die Revolution in der Onkologie am Beispiel des Mammakarzinoms. *Chirurg.* 2004;75(11):1129-34.
- 121 Pavy JJ, Denekamp J, Letschert J, et al. EORTC Late Effects Working Group. Late effects toxicity scoring: the SOMA scale. *Radiother Oncol.* 1995;35(1):11-5.
- 122 Common Terminology Criteria for Adverse Events (CTCAE) v3.0. National Cancer Institute, 2003. (Assessed Aug 18, 2005, at <http://ctep.cancer.gov/reporting/ctc.html>)
- 123 Strandén E. A comparison between surface measurements and water displacement volumetry for the quantification of leg edema. *J Oslo City Hosp.* 1981;31(12):153-5.
- 124 Pani SP, Vanamail P, Yuvaraj J. Limb circumference measurement for recording edema volume in patients with filarial lymphedema. *Lymphology.* 1995;28(2):57-63.
- 125 Schrenk P, Rieger R, Shamiyeh A, Wayand W. Morbidity following sentinel lymph node biopsy versus axillary lymph node dissection for patients with breast carcinoma. *Cancer.* 2000;88(3):608-14.
- 126 Deutsch M, Flickinger JC. Shoulder and arm problems after radiotherapy for primary breast cancer. *Am J Clin Oncol.* 2001;24(2):172-6.

-
- 127 Johansson S, Svensson H, Larsson LG, Denekamp J. Brachial plexopathy after postoperative radiotherapy of breast cancer patients--a long-term follow-up. *Acta Oncol.* 2000;39(3):373-82.
- 128 Haid A, Kuehn T, Konstantiniuk P, et al. Shoulder-arm morbidity following axillary dissection and sentinel node only biopsy for breast cancer. *Eur J Surg Oncol.* 2002;28(7):705-10.
- 129 Peintinger F, Reitsamer R, Stranzl H, Ralph G. Comparison of quality of life and arm complaints after axillary lymph node dissection vs sentinel lymph node biopsy in breast cancer patients. *Br J Cancer.* 2003;89(4):648-52.
- 130 Mansel RE, Goyal A, Fallowfield L, Newcombe RG, on behalf of the ALMANAC Trialists Group. Sentinel node biopsy versus standard axillary treatment results of the randomized multicenter UK ALMANAC trial. *Breast cancer Res Treat.* 2004;88(1 Suppl):13.
- 131 Veronesi U, Paganelli G, Viale G, et al. A randomized comparison of sentinel-node biopsy with routine axillary dissection in breast cancer. *N Engl J Med.* 2003;349(6):546-53.
- 132 Veronesi U, Galimberti V, Mariani L, et al. Sentinel node biopsy in breast cancer: early results in 953 patients with negative sentinel node biopsy and no axillary dissection. *Eur J Cancer.* 2005;41(2):231-7.
- 133 Cabanas RM. An approach for the treatment of penile carcinoma. *Cancer.* 1977;39(2):456-66.
- 134 Morton DL, Wen DR, Wong JH, et al. Technical details of intraoperative lymphatic mapping for early stage melanoma. *Arch Surg.* 1992;127(4):392-9.
- 135 Krag DN, Weaver DL, Alex JC, Fairbank JT. Surgical resection and radiolocalization of the sentinel lymph node in breast cancer using a gamma probe. *Surg Oncol.* 1993;2(6):335-40.
- 136 Giuliano AE, Kirgan DM, Guenther JM, Morton DL. Lymphatic mapping and sentinel lymphadenectomy for breast cancer. *Ann Surg.* 1994;220(3):391-401.
- 137 Cox CE, Pendas S, Cox JM, et al. Guidelines for sentinel node biopsy and lymphatic mapping of patients with breast cancer. *Ann Surg.* 1998;227(5):645-53.
- 138 Pressman PI. Surgical treatment and lymphedema. *Cancer.* 1998;83(12 Suppl American):2782-7.

-
- 139 Kuhn T, Bembenek A, Buchels H, et al. German Breast Society. Sentinel node biopsy in breast cancer. *Nuklearmedizin*. 2004;43(1):4-9.
- 140 Kuhn T, Bembenek A, Buchels H, et al. Sentinel-Node-Biopsie beim Mammakarzinom. *Geburtshilfe Frauenheilkd* 2003;63(09):835-40.
- 141 Suzanne F, Emering C, Wattiez A, Bournazeau JA, Bruhat MA, Jacquetin B. Axillary lymphadenectomy by lipo-aspiration and endoscopic picking. Apropos of 72 cases. *Chirurgie*. 1997;122(2):138-42.
- 142 Salvat J, Knopf JF, Ayoubi JM, et al. Endoscopic exploration and lymph node sampling of the axilla. Preliminary findings of a randomized pilot study comparing clinical and anatomic-pathologic results of endoscopic axillary lymph node sampling with traditional surgical treatment. *Eur J Obstet Gynecol Reprod Biol*. 1996;70(2):165-73.
- 143 Brunt LM, Jones DB, Wu JS, Brunt EM, Radford DM. Endoscopic axillary lymph node dissection: an experimental study in human cadavers. *J Am Coll Surg*. 1998;187(2):158-63.
- 144 Avrahami R, Nudelman I, Watenberg S, Lando O, Hiss Y, Lelchuk S. Minimally invasive surgery for axillary dissection. Cadaveric feasibility study. *Surg Endosc*. 1998;12(5):466-8.
- 145 Paepke S, Schwarz-Boeger U, Kiechle M, Jacobs VR. Axillary Dissection with Access Minimized (ADAM): a new technique for lymph node dissection in conservative surgery for breast cancer. *Int J Fertil Womens Med*. 2003;48(5):232-7.
- 146 Tsangaris TN, Trad K, Brody FJ, Jacobs LK, Tsangaris NT, Sackier JM. Endoscopic axillary exploration and sentinel lymphadenectomy. *Surg Endosc*. 1999;13(1):43-7.
- 147 Langer I, Kocher T, Guller U, et al.. Long-term outcomes of breast cancer patients after endoscopic axillary lymph node dissection: a prospective analysis of 52 patients. *Breast Cancer Res Treat*. 2005;90(1):85-91.
- 148 Britton KE, Jan H, al-Yasi AR, Biassoni L, Carroll MJ, Granowska M. Efficacy of immunoscintigraphy for detection of lymph node metastases. *Recent Results Cancer Res*. 2000;157:3-11.
- 149 Avril N, Dose J, Ziegler S, Janicke S, Schwaiger M. Diagnosis of breast carcinoma and locoregional lymph nodes with positron emission tomography, *Radiologe*. 1997;37(9):741-8.

-
- 150 Albrecht MR, Zink K, Busch W, Ruhl U. Dissection or irradiation of the axilla in postmenopausal patients with breast cancer? Long-term results and long-term effects in 655 patients. *Strahlenther Onkol.* 2002;178(9):510-6.
- 151 Taneja C, Cady B. Decreasing role of lymphatic system surgery in surgical oncology. *J Surg Oncol.* 2005 Feb 1;89(2):61-6.
- 152 Fisher B, Jeong JH, Anderson S, Bryant J, Fisher ER, Wolmark N. Twenty-five-year follow-up of a randomized trial comparing radical mastectomy, total mastectomy, and total mastectomy followed by irradiation. *N Engl J Med.* 2002;347(8):567-75.
- 153 Luini A, Gatti G, Ballardini B, et al. Development of axillary surgery in breast cancer. *Ann Oncol.* 2005;16(2):259-62.
- 154 Greco M, Agresti R, Cascinelli N, et al. Breast cancer patients treated without axillary surgery: clinical implications and biologic analysis. *Ann Surg.* 2000;232(1):1-7.
- 155 Martelli G, Boracchi P, De Palo M, et al. A randomized trial comparing axillary dissection to no axillary dissection in older patients with T1N0 breast cancer: results after 5 years of follow-up. *Ann Surg.* 2005;242(1):1-9.
- 156 Cady B. A contemporary view of axillary dissection. *Ann Surg.* 2000;232(1):8-9.
- 157 Early Breast Cancer Trialists' Collaborative Group. Polychemotherapy for early breast cancer: an overview of the randomised trials. *Lancet.* 1998;352(9132):930-42.
- 158 Kaufmann M, von Minckwitz G, Eiermann W, Hilfrich J, Jonat W, Kreienberg R. Therapie primärer Mammakarzinome. *Dtsch Ärzteblatt.* 2004;101(4):190-4.
- 159 Lin PP, Allison DC, Wainstock J, et al. Impact of axillary lymph node dissection on the therapy of breast cancer patients. *J Clin Oncol.* 1993;11(8):1536-44.
- 160 Goldhirsch A, Glick JH, Gelber RD, Coates AS, Senn HJ. Meeting highlights: International Consensus Panel on the Treatment of Primary Breast Cancer. Seventh International Conference on Adjuvant Therapy of Primary Breast Cancer. *J Clin Oncol.* 2001;19(18):3817-27.

-
- 161 Menard S, Bufalino R, Rilke F, Cascinelli N, Veronesi U, Colhaghi MI. Prognosis based on primary breast carcinoma instead of pathological nodal status. *Br J Cancer*. 1994;70(4):709-12.
- 162 van de Vijver MJ, He YD, van't Veer LJ, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med*. 2002;347(25):1999-2009.
- 163 Cady B, Stone MD, Schuler JG, Thakur R, Wanner MA, Lavin PT. The new era in breast cancer. Invasion, size, and nodal involvement dramatically decreasing as a result of mammographic screening. *Arch Surg*. 1996;131(3):301-8.
- 164 Velanovich V, Szymanski W. Quality of life of breast cancer patients with lymphedema. *Am J Surg*. 1999;177(3):184-8.
- 165 Walley DR, Augustine E, Saslow D, et al. American Cancer Society Lymphedema Workshop. Workgroup IV: Lymphedema treatment resources-- professional education and availability of patient services. *Cancer*. 1998;83(12 Suppl American):2886-7.