

## 8 Literaturverzeichnis

- Ardila-Osorio, H., B. Clause, et al. (1999). "Evidence of LMP1-TRAF3 interactions in glycosphingolipid-rich complexes of lymphoblastoid und nasopharyngeal carcinoma cells." International Journal of Cancer **81**(4): 645-9.
- Bain, G., E. C. Maandag, et al. (1994). "E2A proteins are required for proper B cell development und initiation of immunoglobulin gene rearrangements." Cell **79**(5): 885-92.
- Bargou, R. C., F. Emmerich, et al. (1997). "Constitutive nuclear factor-kappaB-RelA activation is required for proliferation und survival of Hodgkin's disease tumor cells." The Journal of clinical investigation. **100**(12): 2961-9.
- Bargou, R. C., C. Leng, et al. (1996). "High-level nuclear NF-kappa B und Oct-2 is a common feature of cultured Hodgkin/Reed-Sternberg cells." Blood **87**(10): 4340-7.
- Beg, A. A., W. C. Sha, et al. (1995). "Constitutive NF-kappa B activation, enhanced granulopoiesis, und neonatal lethality in I kappa B alpha-deficient mice." Genes und Development **9**(22): 2736-46.
- Benezra, R., R. L. Davis, et al. (1990). "The protein Id: a negative regulator of helix-loop-helix DNA binding proteins." Cell **61**(1): 49-59.
- Brenner, M. (2000). "To be or notch to be." Nature Medicine **6**(11): 1210-1.
- Brielmeier, M., J. Mautner, et al. (1996). "The latent membrane protein 2 gene of Epstein-Barr virus is important for efficient B cell immortalization." Journal of Genetic Virology **77** (Pt 11): 2807-18.
- Cabannes, E., G. Khan, et al. (1999). "Mutations in the I kappa B alpha gene in Hodgkin's disease suggest a tumour suppressor role for I kappa B alpha." Oncogene **18**(20): 3063-70.
- Caldwell, R. G., R. C. Brown, et al. (2000). "Epstein-Barr virus LMP2A-induced B-cell survival in two unique classes of EmuLMP2A transgenic mice." Journal of Virology **74**(3): 1101-13.
- Caldwell, R. G., J. B. Wilson, et al. (1998). "Epstein-Barr virus LMP2A drives B cell development und survival in the absence of normal B cell receptor signals." Immunity **9**(3): 405-11.
- Casola, S., K. L. Otipoby, et al. (2004). "B cell receptor signal strength determines B cell fate." Nature Immunology **5**(3): 317-27.
- Chen, C. L., N. Singh, et al. (2000). "Lymphocytes lacking I kappa B-alpha develop normally, but have selective defects in proliferation und function." The Journal of Immunology **165**(10): 5418-27.
- Cheng, P., A. Zlobin, et al. (2001). "Notch-1 regulates NF-kappaB activity in hemopoietic progenitor cells." The Journal of Immunology **167**(8): 4458-67.
- Cornell, M., D. A. Evans, et al. (1999). "The Drosophila melanogaster Suppressor of deltex gene, a regulator of the Notch receptor signaling pathway, is an E3 class ubiquitin ligase." Genetics **152**(2): 567-76.
- Day, M. J., M. Kyaw-Tanner, et al. (1999). "T-cell-rich B-cell lymphoma in the cat." Journal of comparative pathology **120**(2): 155-67.
- Drexler, H. G. und J. Minowada (1992). "Hodgkin's disease derived cell lines: a review." Human Cell **5**(1): 42-53.
- Dykstra, M. L., R. Longnecker, et al. (2001). "Epstein-Barr virus coopts lipid rafts to block the signaling und antigen transport functions of the BCR." Immunity **14**(1): 57-67.
- Eliopoulos, A. G., C. W. Dawson, et al. (1996). "CD40-induced growth inhibition in epithelial cells is mimicked by Epstein-Barr Virus-encoded LMP1: involvement of TRAF3 as a common mediator." Oncogene **13**(10): 2243-54.
- Eliopoulos, A. G. und A. B. Rickinson (1998). "Epstein-Barr virus: LMP1 masquerades as an active receptor." Current Biology **8**: R196-R198.

- Emmerich, F., M. Meiser, et al. (1999). "Overexpression of I kappa B alpha without inhibition of NF-kappaB activity und mutations in the I kappa B alpha gene in Reed-Sternberg cells." Blood **94**(9): 3129-34.
- Engels, N., M. Merchant, et al. (2001). "Epstein-Barr virus latent membrane protein 2A (LMP2A) employs the SLP-65 signaling module." The Journal of Experimental Medicine **194**(3): 255-64.
- Fruehling, S. und R. Longnecker (1997). "The immunoreceptor tyrosine-based activation motif of Epstein-Barr virus LMP2A is essential for blocking BCR-mediated signal transduction." Virology **235**(2): 241-51.
- Fruehling, S., R. Swart, et al. (1998). "Tyrosine 112 of latent membrane protein 2A is essential for protein tyrosine kinase loading und regulation of Epstein-Barr virus latency." The Journal of Virology **72**(10): 7796-806.
- Guan, E., J. Wang, et al. (1996). "T cell leukemia-associated human Notch/translocation-associated Notch homologue has I kappa B-like activity und physically interacts with nuclear factor-kappa B proteins in T cells." The Journal of Experimental Medicine **183**(5): 2025-32.
- Hertel, C. B., X. G. Zhou, et al. (2002). "Loss of B cell identity correlates with loss of B cell-specific transcription factors in Hodgkin/Reed-Sternberg cells of classical Hodgkin lymphoma." Oncogene **21**(32): 4908-20.
- Hinz, M., P. Loser, et al. (2001). "Constitutive NF-kappaB maintains high expression of a characteristic gene network, including CD40, CD86, und a set of antiapoptotic genes in Hodgkin/Reed-Sternberg cells." Blood **97**(9): 2798-807.
- Horcher, M., A. Souabni, et al. (2001). "Pax5/BSAP maintains the identity of B cells in late B lymphopoiesis." Immunity **14**(6): 779-90.
- Horwitz, B. H., M. L. Scott, et al. (1997). "Failure of lymphopoiesis after adoptive transfer of NF-kappaB-deficient fetal liver cells." Immunity **6**(6): 765-72.
- Ikeda, A., R. G. Caldwell, et al. (2003). "Itchy, a Nedd4 ubiquitin ligase, downregulates latent membrane protein 2A activity in B-cell signaling." Journal of Virology **77**(9): 5529-34.
- Ikeda, M., A. Ikeda, et al. (2001a). "PY motifs of Epstein-Barr virus LMP2A regulate protein stability und phosphorylation of LMP2A-associated proteins." Journal of Virology **75**(12): 5711-8.
- Ikeda, M., A. Ikeda, et al. (2001b). "PY motifs of Epstein-Barr virus LMP2A regulate protein stability und phosphorylation of LMP2A-associated proteins." The Journal of virology **75**(12): 5711-8.
- Jarriault, S., C. Brou, et al. (1995). "Signalling downstream of activated mammalian Notch." Nature **377**(6547): 355-8.
- Jenne, C. N., L. J. Kennedy, et al. (2003). "A new model of sheep Ig diversification: shifting the emphasis toward combinatorial mechanisms und away from hypermutation." The Journal of Immunology **170**(7): 3739-50.
- Jundt, F., I. Anagnostopoulos, et al. (1999). "Hodgkin/Reed-Sternberg cells induce fibroblasts to secrete eotaxin, a potent chemoattractant for T cells und eosinophils." Blood **94**(6): 2065-71.
- Jundt, F., I. Anagnostopoulos, et al. (2002a). "Activated Notch1 signaling promotes tumor cell proliferation und survival in Hodgkin und anaplastic large cell lymphoma." Blood **99**(9): 3398-403.
- Jundt, F., K. Kley, et al. (2002b). "Loss of PU.1 expression is associated with defective immunoglobulin transcription in Hodgkin und Reed-Sternberg cells of classical Hodgkin disease." Blood **99**(8): 3060-2.
- Jundt, F., K. S. Probsting, et al. (2004). "Jagged1-induced Notch signaling drives proliferation of multiple myeloma cells." Blood **103**(9): 3511-5.
- Jungnickel, B., A. Staratschek-Jox, et al. (2000). "Clonal deleterious mutations in the IkappaBalpha gene in the malignant cells in Hodgkin's lymphoma." The Journal of experimental medicine **191**(2): 395-402.

- Kanzler, H., R. Küppers, et al. (1996). "Hodgkin und Reed-Sternberg cells in Hodgkin's disease represent the outgrowth of a dominant tumor clone derived from (crippled) germinal center B cells." The Journal of Experimental Medicine **184**(4): 1495-505.
- Karin, M. (1999). "How NF-kappaB is activated: the role of the IkappaB kinase (IKK) complex." Oncogene **18**(49): 6867-74.
- Kee, B. L. und C. Murre (1998). "Induction of early B cell factor (EBF) und multiple B lineage genes by the basic helix-loop-helix transcription factor E12." The Journal of Experimental Medicine **188**(4): 699-713.
- Khan, W. N., F. W. Alt, et al. (1995). "Defective B cell development und function in Btk-deficient mice." Immunity **3**(3): 283-99.
- Khan, W. N., A. Nilsson, et al. (1997). "Impaired B cell maturation in mice lacking Bruton's tyrosine kinase (Btk) und CD40." Int Immunol **9**(3): 395-405.
- Kieff, E. und A. Rickinson (2001). Epstein-Barr virus und its replication. Field's Virology. D. M. In Knipe, Howley, P., Griffin, D.E., Martin, M.A., Lamb, R.A., Roizman, B., Straus, S.E. Philadelphia, Lippincott - Williams & Wilkins. **2**: 2511-2573.
- Kitamura, D. und K. Rajewsky (1992). "Targeted disruption of mu chain membrane exon causes loss of heavy-chain allelic exclusion." Nature **356**(6365): 154-6.
- Kitamura, D., J. Roes, et al. (1991). "A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin mu chain gene." Nature **350**(6317): 423-6.
- Klement, J. F., N. R. Rice, et al. (1996). "IkappaBalpha deficiency results in a sustained NF-kappaB response und severe widespread dermatitis in mice." Molecular und cellular biology **16**(5): 2341-9.
- Krappmann, D., F. Emmerich, et al. (1999). "Molecular mechanisms of constitutive NF-kappaB/Rel activation in Hodgkin/Reed-Sternberg cells." Oncogene **18**(4): 943-53.
- Küppers, R. und K. Rajewsky (1998). "The origin of Hodgkin und Reed/Sternberg cells in Hodgkin's disease." Annual review of immunology **16**: 471-93.
- Küppers, R., M. Zhao, et al. (1993). "Detection of clonal B cell populations in paraffin-embedded tissues by polymerase chain reaction." American Journal of Pathology **143**(1): 230-9.
- Lai, E. C. (2002). "Protein degradation: four E3s for the notch pathway." Current Biology **12**(2): R74-8.
- Lanning, D., X. Zhu, et al. (2000). "Development of the antibody repertoire in rabbit: gut-associated lymphoid tissue, microbes, und selection." Immunological Reviews **175**: 214-28.
- Laux, G., M. Perricaudet, et al. (1988). "A spliced Epstein-Barr virus gene expressed in immortalized lymphocytes is created by circularization of the linear viral genome." Embo Journal **7**(3): 769-74.
- Lin, H. und R. Grosschedl (1995). "Failure of B-cell differentiation in mice lacking the transcription factor EBF." Nature **376**(6537): 263-7.
- Longnecker, R. (2000). "Epstein-Barr virus latency: LMP2, a regulator or means for Epstein-Barr virus persistence?" Advanced Cancer Research **79**: 175-200.
- Maeda, H., K. Ozaki, et al. (1993). "Hodgkin's-like lymphoma in a dog." Zentralblatt für Veterinärmedizin. Reihe A **40**(3): 200-4.
- Majeed, S. K. und C. Gopinath (1985). "Hodgkin's disease-like lesion in a rat." Journal of comparative pathology **95**(1): 123-6.
- Marafioti, T., M. Hummel, et al. (2000). "Hodgkin und reed-sternberg cells represent an expansion of a single clone originating from a germinal center B-cell with functional immunoglobulin gene rearrangements but defective immunoglobulin transcription." Blood **95**(4): 1443-50.
- Mathas, S., A. Lietz, et al. (2003). "Inhibition of NF-kappaB essentially contributes to arsenic-induced apoptosis." Blood **102**(3): 1028-34.

- Matskova, L., I. Ernberg, et al. (2001). "C-terminal domain of the Epstein-Barr virus LMP2A membrane protein contains a clustering signal." Journal of Virology **75**(22): 10941-9.
- Merchant, M., R. G. Caldwell, et al. (2000). "The LMP2A ITAM is essential for providing B cells with development und survival signals in vivo." Journal of Virology **74**(19): 9115-24.
- Miller, C. L., A. L. Burkhardt, et al. (1995). "Integral membrane protein 2 of Epstein-Barr virus regulates reactivation from latency through dominant negative effects of protein- tyrosine kinases." Immunity **2**: 155-166.
- Morse, H. C., 3rd, F. W. Shen, et al. (1987). "Genetic nomenclature for loci controlling mouse lymphocyte antigens." Immunogenetics **25**(2): 71-8.
- Nakayama, H., K. Nakanaga, et al. (1984). "Three cases of feline lymphosarcoma with formation of multinucleated giant cells." Nippon juigaku zasshi. The Japanese journal of veterinary science **46**(2): 225-8.
- Nutt, S. L., B. Heavey, et al. (1999). "Commitment to the B-lymphoid lineage depends on the transcription factor Pax5." Nature **401**(6753): 556-62.
- Nutt, S. L., P. Urbanek, et al. (1997). "Essential functions of Pax5 (BSAP) in pro-B cell development: difference between fetal und adult B lymphopoiesis und reduced V-to-DJ recombination at the IgH locus." Genes und Development **11**(4): 476-91.
- O'Riordan, M. und R. Grosschedl (1999). "Coordinate regulation of B cell differentiation by the transcription factors EBF und E2A." Immunity **11**(1): 21-31.
- Pahl, H. L. (1999). "Activators und target genes of Rel/NF-kappaB transcription factors." Oncogene **18**(49): 6853-66.
- Pear, W. S. und F. Radtke (2003). "Notch signaling in lymphopoiesis." Semin Immunol **15**(2): 69-79.
- Portis, T., L. Cooper, et al. (2002). "The LMP2A signalosome--a therapeutic target for Epstein-Barr virus latency und associated disease." Frontiers in Bioscience **7**: d414-26.
- Portis, T., P. Dyck, et al. (2003). "Epstein-Barr Virus (EBV) LMP2A induces alterations in gene transcription similar to those observed in Reed-Sternberg cells of Hodgkin lymphoma." Blood **102**(12): 4166-78.
- Portis, T. und R. Longnecker (2003). "Epstein-Barr virus LMP2A interferes with global transcription factor regulation when expressed during B-lymphocyte development." Journal of Virology **77**(1): 105-14.
- Pui, J. C., D. Allman, et al. (1999). "Notch1 expression in early lymphopoiesis influences B versus T lineage determination." Immunity **11**(3): 299-308.
- Reynaud, C. A., V. Anquez, et al. (1987). "A hyperconversion mechanism generates the chicken light chain preimmune repertoire." Cell **48**(3): 379-88.
- Reynaud, C. A., C. R. Mackay, et al. (1991). "Somatic generation of diversity in a mammalian primary lymphoid organ: the sheep ileal Peyer's patches." Cell **64**(5): 995-1005.
- Roperto, F., S. Damiano, et al. (1983). "Hodgkin's disease in a cat." Zentralblatt für Veterinärmedizin. Reihe A **30**(3): 182-8.
- Saito, T., S. Chiba, et al. (2003). "Notch2 is preferentially expressed in mature B cells und indispensable for marginal zone B lineage development." Immunity **18**(5): 675-85.
- Sigvardsson, M., M. O'Riordan, et al. (1997). "EBF und E2A collaborate to induce expression of the endogenous immunoglobulin surrogate light chain genes." Immunity **7**(1): 25-36.
- Skinninger, B. F. und T. W. Mak (2002). "The role of cytokines in classical Hodgkin lymphoma." Blood **99**(12): 4283-97.
- Smith, D. A. und I. K. Barker (1983). "Four cases of Hodgkin's disease in striped skunks (*Mephitis mephitis*)." Veterinary pathology **20**(2): 223-9.

- Souabni, A., C. Cobaleda, et al. (2002). "Pax5 promotes B lymphopoiesis und blocks T cell development by repressing Notch1." Immunity **17**(6): 781-93.
- Speck, P., K. A. Kline, et al. (1999). "Epstein-Barr virus lacking latent membrane protein 2 immortalizes B cells with efficiency indistinguishable from that of wild-type virus." Journal of Genetic Virology **80 (Pt 8)**: 2193-203.
- Staudt, L. M. (2000). "The molecular und cellular origins of Hodgkin's disease." J Exp Med **191**(2): 207-12.
- Stein, H., T. Marafioti, et al. (2001). "Down-regulation of BOB.1/OBF.1 und Oct2 in classical Hodgkin disease but not in lymphocyte predominant Hodgkin disease correlates with immunoglobulin transcription." Blood **97**(2): 496-501.
- Strobl, L. J., H. Hofelmayr, et al. (2000). "Activated Notch1 modulates gene expression in B cells similarly to Epstein-Barr viral nuclear antigen 2." Journal of Virology **74**(4): 1727-35.
- Thevenin, C., S. L. Nutt, et al. (1998). "Early function of Pax5 (BSAP) before the pre-B cell receptor stage of B lymphopoiesis." The Journal of Experimental Medicine **188**(4): 735-44.
- Thorley-Lawson, D. A. und K. P. Mann (1985). "Early events in Epstein-Barr virus infection provide a model for B cell activation." The Journal of Experimental Medicine **162**: 45-59.
- Tierney, R., H. Kirby, et al. (2000). "The Epstein-Barr virus promoter initiating B-cell transformation is activated by RFX proteins und the B-cell-specific activator protein BSAP/Pax5." The Journal of Virology **74**(22): 10458-67.
- Walton, R. M. und M. J. Hendrick (2001). "Feline Hodgkin's-like lymphoma: 20 cases (1992-1999)." Vet Pathol **38**(5): 504-11.
- Wang, J., L. Shelly, et al. (2001). "Human Notch-1 inhibits NF-kappa B activity in the nucleus through a direct interaction involving a novel domain." The Journal of Immunology **167**(1): 289-95.
- Wells, G. A. (1974). "Hodgkin's disease-like lesions in the dog." The Journal of pathology **112**(1): 5-10.
- Wilson, A., I. Ferrero, et al. (2000). "Cutting edge: an essential role for Notch-1 in the development of both thymus-independent und -dependent T cells in the gut." Journal of Immunology **165**(10): 5397-400.
- Winberg, G., L. Matskova, et al. (2000). "Latent membrane protein 2A of Epstein-Barr virus binds WW domain E3 protein-ubiquitin ligases that ubiquitinate B-cell tyrosine kinases." Molecular und Cellular Biology **20**(22): 8526-35.
- Woisetschlaeger, M., C. N. Yandava, et al. (1990). "Promoter switching in Epstein-Barr virus during the initial stages of infection of B lymphocytes." Proc Natl Acad Sci U S A **87**(5): 1725-9.
- Xu, S., J. E. Tan, et al. (2000). "B cell development und activation defects resulting in xid-like immunodeficiency in BLNK/SLP-65-deficient mice." Int Immunol **12**(3): 397-404.
- Yonezawa, M., H. Nakamine, et al. (1989). "Hodgkin's disease in a killer whale (*Orcinus orca*)." Journal of comparative pathology **100**(2): 203-7.
- Young, L. S., C. W. Dawson, et al. (2000). "The expression und function of Epstein-Barr virus encoded latent genes." Molecular Pathology **53**(5): 238-47.
- Young, L. S. und A. B. Rickinson (2004). "Epstein-Barr virus: 40 years on." Nature Reviews Cancer **4**(10): 757-68.