

## Literaturverzeichnis

- Abbate M, Zoja C, Corna D, Capitanio M, Bertani T and Remuzzi G (1998): In progressive nephropathies, overload of tubular cells with filtered proteins translates glomerular permeability dysfunction into cellular signals of interstitial inflammation. *J Am Soc Nephrol* **9**:1213-1224.
- Abdulkadyrov KM and Bessmeltsev SS (1996): Renal insufficiency in multiple myeloma: basic mechanisms in its development and methods for treatment. *Ren Fail* **18**:139-146.
- Abrass CK (1984): Diabetic proteinuria: glomerular or tubular in origin? *Am J Nephrol* **4**:337-346.
- Addadi L and Weiner S (1985): Interactions between acidic proteins and crystals: stereochemical requirements in biominerilization. *Proc Natl Acad Sci U S A* **82**:4110-4114.
- Adonccecchi L, Marrocco W, Suraci C, Pecora P, Gallinella B, Porra R and Cavina G (1984): Effect of renal and liver failure on blood levels of vitamin A, its precursor (beta-carotene) and its carrier proteins (prealbumin and retinol binding protein). *Boll Soc Ital Biol Sper* **60**:881-886.
- Albers A, Wagner CA, Schmidt C, Lahme S, Bichler KH, Lang F, Zerres K and Eggermann T (2001): [Molecular biology of cystinuria] *Fortschr Med Orig* **119**:49-50.
- Allen T, Jaenke RS and Fettman MJ (1987): A technique for estimation progression of chronic renal failure in the dog. *J Am Vet Med Assoc* **190**:866-868.
- Almy FS, Christopher MM, King DP and Brown SA (2002): Evaluation of cystatin C as an endogenous marker of glomerular filtration rate in dogs. *J Vet Intern Med* **16**:45-51.
- Anderson MS and Recant L (1962): Fine structural alterations in the rat kidney following intraperitoneal bovine albumin. *Am J Pathol* **40**:555-569.
- Ang BS (1995): Urinary tract infections. *Singapore Med J* **36**:314-317.
- Appleman RM, Hallenbeck GA and Shorter RG (1966): Effect of reciprocal allogenic renal transplantation between Dalmatians and non-Dalmatian dogs on urinary excretion of uric acid. *Proc Soc Exp Biol Med* **121**:1094-1102.
- Asplin JR, Arsenault D, Parks JH, Coe FL and Hoyer JR (1998): Contribution of human uropontin to inhibition of calcium oxalate crystallization. *Kidney Int* **53**:194-199.
- Atmani F, Lacour B, Strecker G, Parvy P, Drüeke BT and Daudon M (1993): Molecular characteristics of uronic acid-rich-protein (UAP), a strong inhibitor of calcium oxalate crystallization in vitro. *Biochem Biophys Res Commun* **191**:1158-1165.
- Atmani F, Lacour B, Jungers P, Drüeke BT and Daudon M (1994): Reduced inhibitory activity of uronic-acid-rich protein in urine of stone formers. *Urol Res* **22**:257-260.
- Atmani F, Mizon J and Khan SR (1996a): Identification of uronic-acid-rich protein as urinary bikunin: the light chain of inter-alpha-inhibitor. *European J Biochem* **236**:984-990.
- Atmani F, Opalko FJ and Khan SR (1996b): Association of urinary macromolecules with calcium oxalate crystals induced in vitro in normal human and rat urine. *Urol Res* **24**:45-50.
- Atmani F, Glenton PA and Khan SR (1998): Identification of proteins extracted from calcium oxalate and calcium phosphate crystals induced in the urine of healthy and stone forming subjects. *Urol Res* **26**:201-207.
- Atmani F and Khan SR (1999): Role of bikunin in the inhibition of calcium oxalate crystallization. *J Am Soc Nephrol* **10**:S385-388.
- Atmani F (2001): Characterisation of urinary bikunin and its role in urolithiasis. *Rev Biol Biotech* **1**:17-20.
- Atmani F and Khan SR (2002): Quantification of proteins extracted from calcium oxalate and calcium phosphate crystals induced in vitro in the urine of healthy controls and stone-forming patients. *Urol Int* **68**:54-59.

- Bachmann S, Metzger R and Bunnemann B (1990): Tamm-Horsfall protein-mRNA synthesis is localized to the thick ascending limb of Henle's loop in rat kidney. *Histochemistry* **94**:517-523.
- Bade JJ, Marrink J, Karrenbeld A, van der Weele L and Mensink HJ (1996): Increased urinary levels of Tamm-Horsfall glycoprotein suggest a systemic etiology of interstitial cystitis. *J Urol* **156**:943-946.
- Baker H, Schorr SM, Murphy BD, De Angelis B, Feingold S and Frank O (1986): Blood vitamin and choline concentrations in healthy domestic cats, dogs and horses. *Am J Vet Res* **47**:1468-1471.
- Bakoush O, Torffvit O, Rippe B and Tencer J (2001): High proteinuria selectivity index based upon IgM is a strong predictor of poor renal survival in glomerular diseases. *Nephrol Dial Transplant* **16**:1357-1363.
- Balant L and Fabre J (1979): Clinical relevation of different electrophoretic methods for the analysis of urinary proteins. *Curr Probl Clin Biochem* **9**:216-234.
- Banaszak L, Winter N, Xu Z, Bernlohr DA, Cowan S and Jones TA (1994): Lipid-binding proteins: A family of fatty acid and retinoid transport proteins. *Adv Protein Chem* **45**:89-151.
- Barsanti JA and Finco DR (1979): Protein concentration in urine of normal dogs. *Am J Vet Res* **40**:1583-1588.
- Bartges JW, Osborne CA, Lulich JP, Unger LK, Koehler LA, Bird KA, Clinton CW and Davenport MP (1994): Prevalence of cystine and urate uroliths in bulldogs and urate uroliths in dalmatians. *J Am Vet Med Assoc* **204**:1914-1918.
- Bartges JW, Osborne CA, Lulich JP, Kruger JM, Sanderson SL, Koehler LA and Ulrich LK (1999): Canine urate urolithiasis. Etiopathogenesis, diagnosis, and management. *Vet Clin North Am Small Anim Pract* **29**:161-191, xii-xiii.
- Barth JL and Argraves WS (2001): Cubilin and megalin: partners in lipoprotein and vitamin metabolism. *Trends Cardiovasc Med* **11**:26-31.
- Batuman V, Verroust PJ, Navar GL, Kaysen JH, Goda FO, Campbell WC, Simon E, Pontillon F, Lyles M, Bruno J and Hammond TG (1998): Myeloma light chains are ligands for cubilin (gp280). *Am J Physiol* **275**:F246-254.
- Baumann JM (1988): How reliable are the measurements of crystallization conditions in urine? *Urol Res* **16**:133-135.
- Baumann JM, Affolter B, Caprez U, Lauper D, Maier F, Siegrist HP and Zyssset T (2000): Nucleation and aggregation of calcium oxalate in whole urine; spectrophotometric sedimentation analysis: a new approach to study the aggregation of calcium oxalate dihydrate. *Urol Res* **28**:147-154.
- Bazzi C and D'Amico G (2002): The urinary excretion of IgG and alpha1-microglobulin predicts renal outcome and identifies patients deserving treatment in membranous nephropathy. *Kidney Int* **61**:2276.
- Benkovic J, Furedi-Milhofer H, Hlady V, Cvoriscec D and Stavljenic-Rukavina A (1995): Effect of Tamm-Horsfall protein on calcium oxalate precipitation. *Eur J Clin Chem Clin Biochem* **33**:705-710.
- Bernard AM, Thielemans NO and Lauwers RR (1994): Urinary protein 1 or Clara cell protein: a new sensitive marker of proximal tubular dysfunction. *Kidney Int Suppl* **47**:s34-37.
- Bianchi-Bosisio A, D'Agrosa F, Gabardi F, Gianazza E and Righetti PG (1991): Sodium dodecyl sulphate electrophoresis of urinary proteins. *J Chromatogr* **569**:243-260.
- Biberstein EL (1977): Urinary tract infections in dogs: Microbiological diagnosis. *Calif Vet* **31**:10-17.
- Bichler KH and Kirchner C (1975): Immune histological representation of uromucoid in the kidney of man and rat. *Urologie A* **14**:246-249.
- Bichler KH, Ideler V and Harzmann R (1979): Uromucoid excretion in normal individuals and stone formers. *Curr Probl Clin Biochem* **9**:309-324.
- Bichler KH, Kirchner C, Strohmaier W, Weiser H, Schmitz P and Korn S (1982): Effect of vitamin A deficiency on the excretion of uromucoid and other urine constituents in rats. *Fortschr Urol Nephrol*

20:205-211.

Bichler KH, Kirchner C, Weiser H, Korn S, Strohmaier W, Schmitz-Moormann P, Hanck A and Nelde HJ (1983): Influence of vitamin A deficiency on the excretion of uromucoid and other substances in the urine of rats. *Clin Nephrol* **20**:32-39.

Bichler KH, Henzler B, Strohmaier WL, Stahl C and Korn S (1995): The significance of citrate, uromucoid and GAG for diagnosis of renal tubular acidosis in patients with urinary calculi. *Urologe A* **34**:437-443.

Bichler KH, Mitermuller WL, Strohmaier W, Feil G and Eipper E (1999): Excretion of Tamm-Horsfall protein in patients with uric acid stones. *Urol Int* **62**:87-92.

Biewenga WJ, Gruys E and Hendriks HJ (1982): Urinary protein loss in the dog: nephrological study of 29 dogs without signs of renal disease. *Res Vet Science* **33**:366-374.

Biewenga WJ and Gruys E (1986): Proteinuria in the dog: a clinicopathological study in 51 proteinuric dogs. *Res Vet Science* **41**:257-264.

Birn H, Verroust PJ, Nexo E, Hager H, Jacobsen C, Christensen EI and Moestrup SK (1997): Characterization of an epithelial approximately 460-kDa protein that facilitates endocytosis of intrinsic factor-vitamin B12 and binds receptor-associated protein. *J Biol Chem* **272**:26497-26504.

Birn H, Fyfe JC, Jacobsen C, Mounier F, Verroust PJ, Orskov H, Willnow TE, Moestrup SK and Christensen EI (2000): Cubilin is an albumin binding protein important for renal tubular albumin reabsorption. *J Clin Invest* **105**:1353-1361.

Blaner WS (1989): Retinol-binding protein: the serum transport protein for vitamin A. *Endocr Rev* **10**:308-316.

Blomhoff R, Green MH, Berg T and Norum KR (1990): Transport and storage of vitamin A. *Science* **250**:399-404.

Blomhoff R, Green MH, Green JB, Berg T and Norum KR (1991): Vitamin A metabolism: new perspectives on absorption, transport, and storage. *Physiol Rev* **71**:951-990.

Blomhoff R, Green MH and Norum KR (1992): Vitamin A: physiological and biochemical processing. *Annu Rev Nutr* **12**:37-57.

Blomhoff R (1994a): Transport and metabolism of vitamin A. *Nutr Rev* **52**:S13-23.

Blomhoff R (1994b): Overview of vitamin A metabolism and function. In: *Vitamin A in health and disease*. Blomhoff R (eds.), New York, Basel, Hong Kong, Marcel Dekker Inc. pp 1-36.

Bloom F (1939): Classification and pathology of renal disease in the dog. Comparison with nephritis in man. *Archs Path* **28**:236-245.

Bloom F (1954): Pathology of the dog and cat. The genitourinary system, with clinical considerations. Am Vet Publications, Illinois, Inc Evanston.

Boesken WH, Kopf K and Schollmeyer P (1973): Differentiation of proteinuric diseases by discelectrophoretic molecular weight analysis of urinary proteins. *Clin Nephrol* **1**:311-318.

Boesken WH (1975): Discelectrophoretic molecular weight analysis of urinary proteins. A contribution to the clinical diagnostic differentiation and the pathophysiology of proteinuria. *Contrib Nephrol* **1**:143-155.

Boesken WH, Batsford SR, Schmitz-Hubner U and Schneider G (1976a): Molecular weight analysis of proteinuria in experimental glomerulopathies. *Res Exp Med (Berl)* **167**:23-30.

Boesken WH, Schneider GH und Reuscher C (1976b): Veränderungen des Urinprotein-Musters bei Diabetes mellitus, ein Frühsymptom der renalen Mikroangiopathie? *Diabetologia* **12**:381 (Abstract).

Boesken WH, Schindera F, Billingham M, Hardwicke J, White RH and Williams A (1977): Polymeric albumin in the urine of patients with nephrotic syndrome. *Clin Nephrol* **8**:395-399.

Boesken WH (1985): Bedeutung der Proteinurie-Analyse für die nephrologische Diagnostik. *Mitt Klin*

Nephrologie **XIV**:96-111.

Boesken WH und Mamier A (1985): Molekulargewichtsbezogene Urinprotein-Elektrophorese in der Diagnostik von Nierenkrankheit. Lab med **9**:285-291.

Boeve ER, Cao LC, De Bruijn WC, Robertson WG, Romijn JC and Schroder FH (1994): Zeta potential distribution on calcium oxalate crystal and Tamm-Horsfall protein surface analyzed with Doppler electrophoretic light scattering. J Urol **152**:531-536.

Boie W (1977): Über die Beziehung zwischen Nierenfunktion und Serumkonzentration des Retinol-bindenden Proteins, Präalbumins, Vitamin A und Carotins. A-Promotion, Würzburg, pp 166-177.

Borghi L, Meschi T, Guerra A, Bergamaschi E, Mutti A and Novarini A (1995): Effects of urinary macromolecules on the nucleation of calcium oxalate in idiopathic stone formers and healthy controls. Clin Chim Acta **239**:1-11.

Bosin E and Monji N (1987): Analysis of serum and urinary retinol binding protein in hepato-renal syndrome. Clin Biochem **20**:47-51.

Bovee KC and McGuire T (1984): Qualitative and quantitative analysis of uroliths in dogs: definitive determination of chemical type. J Am Vet Med Assoc **185**:983-987.

Boyce WH, Garvey FK and Norfleet CM (1954): Ion binding properties of electrophoretically homogeneous mucoproteins of urine in normal subjects and in patients with renal calculus disease. J Urol **72**:1019-1031.

Boyce WH and Garvey FK (1956): The amount and nature of the organic matrix in urinary calculi: a review. J Urol **76**:213-227.

Boyce WH and King JS (1959): Crystal-matrix interrelationships in calculi. J Urol **81**:351-355.

Boyce WH, King JS and Fielden ML (1962): Total non-dialyzable solids in human urine XIII. Immunological detection of a component peculiar to renal calculous matrix and to urine of calculous patients. J Clin Invest **41**:1180-1189.

Boyce WH and King JS (1963): Present concepts concerning the origin of matrix in stones. Ann N Y Acad Sci **104**:579-583.

Boyce WH (1968): Organic matrix of human urinary concretions. Am J Med **45**:673-683.

Boyce WH (1970): Proteinuria in kidney calculous disease. In: Proteins in normal and pathological urine., Manuel Y, Revillard JP and Betuel H (eds.), Basel, New York, Karger

Bricker NS, Morrin PAF and Kime SW (1960): The pathologic physiology of chronic Bright's disease. - An exposition of the "intact nephron hypothesis". Am J Med **28**:77-98.

Brown NO, Parks JL and Gremm RW (1977): Canine urolithiasis: Retrospective analysis of 438 cases. J Am Vet Med Assoc **170**:414-418.

Buchborn E und Edel HH (1968): Akutes Nierenversagen. In: Handbuch der Inneren Medizin, Berlin, Springer

Buffington CA, Blaisdell JL and Sako T (1994): Effects of Tamm-Horsfall glycoprotein and albumin on struvite crystal growth in urine of cats. Am J Vet Res **55**:965-971.

Burne MJ, Vyas SV, Smit MF, Pratt LM and Comper WD (1997): Competition between polyanions in glomerular binding and renal clearance. Arch Biochem Biophys **340**:257-264.

Burri BJ, Neidlinger TR and Zwick H (1993): Comparison of the properties and concentrations of the isoforms of retinol-binding protein in animals and human beings. Am J Vet Res **54**:1213-1220.

Butt AJ (1952): Role of protective urinary colloid in prevention of renal lithiasis. J Urol **67**:450-459.

Caliskan S, Hacibekiroglu M, Sever L, Ozbay G and Arisoy N (1996): Urinary N-acetyl-beta-D-glucosaminidase and beta 2-microglobulin excretion in primary nephrotic children. Nephron **74**:401-404.

Cao A, Balant L, Fabre J and Canavèse JC (1975): Physiological proteinuria. Data of acrylamide-SDS gel electrophoresis and other methods of qualitative and quantitative analysis. Schweiz Med Wochenschr **105**:421-429.

Catalano C and Torffvit O (1996): Urinary excretion of Tamm-Horsfall protein in normotensive, normo-albuminuric type 1 diabetic patients. Nephron **72**:436-441.

Center SA, Wilkinson E, Smith CA, Erb H and Lewis RM (1985): 24-Hour urine protein/creatinine ratio in dogs with protein-losing nephropathies. J Am Vet Med Assoc **187**:820-824.

Chen WC, Lin HS, Chen HY, Shih CH and Li CW (2001): Effects of Tamm-Horsfall protein and albumin on calcium oxalate crystallization and importance of sialic acids. Mol Urol **5**:1-5.

Christensen EI, Rennke HG and Carone FA (1983): Renal tubular uptake of protein: effect of molecular charge. Am J Physiol **244**:F436-441.

Christensen EI, Gliemann J and Moestrup SK (1992): Renal tubule gp330 is a calcium binding receptor for endocytic uptake of protein. J Histochem Cytochem **40**:1481-1490.

Christensen EI, Birn H, Verroust P and Moestrup SK (1998): Megalin-mediated endocytosis in renal proximal tubule. Ren Fail **20**:191-199.

Christensen EI, Moskaug JO, Vorum H, Jacobsen C, Gundersen TE, Nykjaer A, Blomhoff R, Willnow TE and Moestrup SK (1999): Evidence for an essential role of megalin in transepithelial transport of retinol. J Am Soc Nephrol **10**:685-695.

Christensen EI and Birn H (2001): Megalin and cubilin: synergistic endocytic receptors in renal proximal tubule. Am J Physiol Renal Physiol **280**:F562-573.

Ciarimboli G, Bokenkamp A, Schurek HJ, Fels LM, Kilian I, Maess B and Stolte H (2001): The "fixed" charge of glomerular capillary wall as determinant of permselectivity. Ren Fail **23**:365-376.

Clausen SW, Baum WS, McCoord AB, Rydeen JO and Bresse BB (1942): The mobilization by alcohols of vitamin A from its stores in the tissue. J Nutr **24**:1-14.

Cleave AJ, Kent PW and Peacocke AR (1972): The binding of hydrogen and calcium ions by Tamm-Horsfall glycoprotein. Biochim Biophys Acta **285**:208-233.

Coe FL, Parks JH and Nakagawa Y (1991a): Protein inhibitors of crystallization. Semin Nephrol **11**:98-109.

Coe FL, Nakagawa Y and Parks JH (1991b): Inhibitors within the nephron. Am J Kidney Dis **17**:407-413.

Coll E, Botey A, Alvarez L, Poch E, Quinto L, Saurina A, Vera M, Piera C and Darnell A (2000): Serum cystatin C as a new marker for noninvasive estimation of glomerular filtration rate and as a marker for early renal impairment. Am J Kidney Dis **36**:29-34.

Comper WD, Lee ASN, Tay M and Adal Y (1993): Anionic charge concentration of rat kidney glomeruli and glomerular basement membrane. Biochem J **289**:647-652.

Constable PD (1991): Clinical assessment of renal tubular function through determination of electrolyte and creatinine concentrations. J Vet Intern Med **5**:357-358.

Corso A, Serricchio G, Zappasodi P, Klersy C, Bosoni T, Moratti R, Castagnola C, Lazzarino M, Pagnucco G and Bernasconi C (1999): Assessment of renal function in patients with multiple myeloma: the role of urinary proteins. Ann Hematol **78**:371-375.

Coye RD, Maude DL, Dibble RF and Yuile CL (1955): Experimental Proteinuria. Arch Pathol **60**:548-555.

Cui S, Verroust PJ, Moestrup SK and Christensen EI (1996): Megalin/gp330 mediates uptake of albumin in renal proximal tubule. Am J Physiol **271**:F900-907.

Cunningham BA, Wang JL, Berggard I and Peterson PA (1973): The complete amino acid sequence of beta 2-microglobulin. Biochemistry **12**:4811-4822.

Dakopatts (1983): General Procedure, quantitation of trace amounts of antigen by two-site micro ELISA. Handbook of Dakopatts (Denmark)

D'Amico G and Bazzi C (2003): Pathophysiology of proteinuria. *Kidney Int* **63**:809-825.

Daudon M, Bader CA and Jungers P (1993): Urinary calculi: review of classification methods and correlations with etiology. *Scanning Microsc* **7**:1081-1104; discussion 1104-1086.

Daudon M, Hennequin C, Bader C, Jungers P, Lacour B and Druecke T (1995): Inhibitors of crystallization. *Adv Nephrol Necker Hosp* **24**:167-216.

Dawnay AB and Cattell WR (1981): Serum Tamm-Horsfall glycoprotein levels in health and in renal disease. *Clin Nephrol* **15**:5-8.

De Schepper J (1977): Nieraandoeningen bij de hond: Duizend klinische Gevallen. Vlaams Diergeneesk. Tijdschr. **46**:18.

Dean C, Kanellos J, Pham H, Gomes M, Oates A, Grover P and Ryall R (2000): Effects of inter-alpha-inhibitor and several of its derivatives on calcium oxalate crystallization in vitro. *Clin Sci (Colch)* **98**:471-480.

DiBartola SP, Chew DJ and Jacobs G (1980): Quantitative urinalysis including 24-hour protein excretion in the dog. *J Am Anim Hosp Assoc* **16**:537-546.

Doetsch K (1972): Determination of total urinary protein by combined gel filtration and automated biuret reaction. *Clin Chem* **18**:296-298.

Donadio C, Lucchesi A, Ardini M and Giordani R (2001): Cystatin C, beta 2-microglobulin, and retinol-binding protein as indicators of glomerular filtration rate: comparison with plasma creatinine. *J Pharm Biomed Anal* **24**:835-842.

Doyle IR, Ryall RL and Marshall VR (1991): Inclusion of proteins into calcium oxalate crystals precipitated from human urine: a highly selective phenomenon. *Clin Chem* **37**:1589-1594.

Doyle S, Conlon P and Royston D (2000): Vitamin A induced stellate cell hyperplasia and fibrosis in renal failure. *Histopathology* **36**:88-93.

Dussol B, Geider S, Lilova A, Leonetti F, Dupuy P, Daudon M, Berland Y, Dagorn JC and Verdier JM (1995): Analysis of the soluble organic matrix of five morphologically different kidney stones. Evidence for a specific role of albumin in the constitution of the stone protein matrix. *Urol Res* **23**:45-51.

Dworkin LD (2001): Serum cystatin C as a marker of glomerular filtration rate. *Curr Opin Nephrol Hypertens* **10**:551-553.

Eddy AA, McCulloch L, Adams J and Liu E (1989): Interstitial nephritis induced by protein-overload proteinuria. *Am J Pathol* **135**:719-733.

Ellis D, Forrest KY, Erbey J and Orchard TJ (1998): Urinary measurement of transforming growth factor-beta and type IV collagen as new markers of renal injury: application in diabetic nephropathy. *Clin Chem* **44**:950-956.

Eppel GA, Osicka TM, Pratt LM, Jablonski P, Howden BO, Glasgow EF and Comper WD (1999): The return of glomerular-filtered albumin to the rat renal vein. *Kidney Int* **55**:1861-1870.

Faro S and Fenner DE (1998): Urinary tract infections. *Clin Obstet Gynecol* **41**:744-754.

Fasth A, Hoyer JR and Seiler MW (1986): Renal tubular immune complex formation in mice immunized with Tamm-Horsfall protein. *Am J Pathol* **125**:555-562.

Fellstrom B, Danielson BG, Ljunghall S and Wikstrom B (1985): The inhibition of calcium oxalate crystal growth by chondroitin sulphates, heparin, pentosan polysulphate and Tamm-Horsfall glycoprotein. In: *Urolithiasis and related clinical research*, Schwille PO, Smith LH, Robertson WG and Vahlensieck W (eds.), New York, Plenum Press, pp 887-890.

Fellstrom B, Danielson BG, Ljunghall S and Wikstrom B (1986): Crystal inhibition: the effects of polyanions on calcium oxalate crystal growth. *Clin Chim Acta* **158**:229-235.

Fex G and Hansson B (1979): Retinol-binding protein from human urine and its interaction with retinol an prealbumin. *Eur J Biochem* **94**:307-313.

Finco DR and Duncan JR (1976): Evaluation of blood urea nitrogen and serum creatinine concentrations as indicators of renal dysfunction: a study of 111 cases and a review of related literature. *J Am Vet Med Assoc* **168**:593-601.

Finco DR and Barsant JA (1982): Mechanism of urinary excretion of creatinine by the cat. *Am J Vet Res* **43**:2207-2209.

Finco DR (1995a): Applied physiology of the kidney. In: *Canine and Feline Nephrology and Urology*, Osborne CA and Finco DR (eds.), Philadelphia, Williams & Wilkins, pp 29-46.

Finco DR (1995b): Evaluation of renal functions. In: *Canine and Feline Nephrology and Urology*, Osborne CA and Finco DR (eds.), Philadelphia, Williams & Wilkins, pp 216-229.

Fleisch H (1978): Inhibitors and promoters of stone formation. *Kidney Int* **13**:361-371.

Fleisch H (1983): Pathophysiologie der Harnsteinbildung. In: *Urologie in Klinik und Praxis*, Hohenfellner R, Zingg E (eds.), Band II, Thieme Verlag, Stuttgart, New York, pp 744-748.

Fletcher AP, Neuberger A and Ratcliffe WA (1970a): Tamm-Horsfall urinary glycoprotein. The chemical composition. *Biochem J* **120**:417-424.

Fletcher AP, Neuberger A and Ratcliffe WA (1970b): Tamm-Horsfall urinary glycoprotein. The subunit structure. *Biochem J* **120**:425-432.

Flynn FV (1990): Assessment of renal function: selected developments. *Clin Biochem* **23**:49-54.

Foreman JW (1984): Renal handling of urate and other organic acids. In: *Canine Nephrology*, Bovee KD (eds.), Media PA, Harwal, pp 135-152.

Fraj BM (1989): Separation and identification of urinary proteins and stone-matrix proteins by mini-slab sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Clin Chem* **35**:658-662.

Franti CE, Ling GV, Ruby AL and Johnson DL (1999): Urolithiasis in dogs V: regional comparisons of breed, age, sex, anatomic location, and mineral type of calculus. *Am J Vet Res* **60**:29-42.

Fuselier HA, Ward DM, Lindberg JS, Allen JM, Husserl FE, Marcucci PA, Cole FE, Turnipseed J, Alam J and Kok DJ (1995): Urinary Tamm-Horsfall protein increased after potassium citrate therapy in calcium stone formers. *Urology* **45**:942-946.

Fyfe JC, Giger U, Hall CA, Jezyk PF, Klumpp SA, Levine JS and Patterson DF (1991): Inherited selective intestinal cobalamin malabsorption and cobalamin deficiency in dogs. *Pediatr Res* **29**:24-31.

Gambaro G, Baggio B, Favaro S, Cicerello E, Marchini F und Borsatti A (1984): Die Rolle von Tamm-Horsfall Mucoprotein in der Calciumoxalat Lithogenese. *Nephrologie* **5**:171-172.

Ganter K, Bongartz D and Hesse A (1999): Tamm-Horsfall protein excretion and its relation to citrate in urine of stone-forming patients. *Urology* **53**:492-495.

Gärtner K (1967): Untersuchungen zur Verbreitung, Differentialdiagnostik, Ätiologie und Pathogenese der spontanen Nephritiden beim Hund unter besonderer Berücksichtigung von Leptospireninfektionen. *Habil, Tierärzt Hochschule Hannover*, Hannover, Schaper, pp 91.

Geider S, Dussol B, Nitsche S, Veesler S, Berthezene P, Dupuy P, Astier JP, Boistelle R, Berland Y, Dagorn JC and Verdier JM (1996): Calcium carbonate crystals promote calcium oxalate crystallization by heterogeneous or epitaxial nucleation: possible involvement in the control of urinary lithogenesis. *Calcif Tissue Int* **59**:33-37.

Gerlach TH and Zile MH (1990): Upregulation of serum retinol in experimental acute renal failure. *Faseb J* **4**:2511-2517.

Gerlach TH and Zile MH (1991a): Effect of retinoic acid and apo-RBP on serum retinol concentration in acute renal failure. *Faseb J* **5**:86-92.

Gerlach TH and Zile MH (1991b): Metabolism and secretion of retinol transport complex in acute renal

failure. *J Lipid Res* **32**:515-520.

Glauser A, Hochreiter W, Jaeger P and Hess B (2000): Determinants of urinary excretion of Tamm-Horsfall protein in non-selected kidney stone formers and healthy subjects. *Nephrol Dial Transplant* **15**:1580-1587.

Glover J (1983): Factors affecting vitamin A transport in animals and man. *Proc Nutr Soc* **42**:19-30.

Gokhale JA, McKee MD and Khan SR (1996): Immunocytochemical localization of Tamm-Horsfall protein in the kidneys of normal and nephrolithic rats. *Urol Res* **24**:201-209.

Gokhale JA, Glenton PA and Khan SR (2001): Characterization of tamm-horsfall protein in a rat nephrolithiasis model. *J Urol* **166**:1492-1497.

Goodyer P, Saadi I, Ong P, Elkas G and Rozen R (1998): Cystinuria subtype and the risk of nephrolithiasis. *Kidney Int* **54**:56-61.

Grant AM, Baker LR and Neuberger A (1973): Urinary Tamm-Horsfall glycoprotein in certain kidney diseases and its content in renal and bladder calculi. *Clin Sci* **44**:377-384.

Grases F and Conte A (1992): Urolithiasis, inhibitors and promoters. *Urol Res* **20**:86-88.

Grases F, Garcia-Ferragut L and Costa-Bauza A (1996): Study of the early stages of renal stone formation: experimental model using urothelium of pig urinary bladder. *Urol Res* **24**:305-311.

Grases F, Garcia-Gonzalez R, Genestar C, Torres JJ and March JG (1998): Vitamin A and urolithiasis. *Clin Chim Acta* **269**:147-157.

Grauer GF, Thomas CB and Eicker SW (1985): Estimation of quantitative proteinuria in the dog, using the urine protein-to-creatinine ratio from a random, voided sample. *Am J Vet Res* **46**:2116-2119.

Gregory H and Preston BM (1977): The primary structure of human urogastone. *Int J Pept Protein Res* **9**:107-118.

Griffith DP and Osborne CA (1987): Infection (urease) stones. *Miner Electrolyte Metab* **13**:278-285.

Grillenberger A, Weninger M and Lubec G (1987): Determination of urinary low molecular weight proteins for the diagnosis of tubular damage. *Padiatr Padol* **22**:229-234.

Grover PK, Ryall RL and Marshall VR (1990): Does Tamm-Horsfall mucoprotein inhibit or promote calcium oxalate crystallization in human urine? *Clin Chim Acta* **190**:223-238.

Grover PK and Resnick MI (1995): Evidence for the presence of abnormal proteins in the urine of recurrent stone formers. *J Urol* **153**:1716-1721.

Grover PK, Moritz RL, Simpson RJ and Ryall RL (1998): Inhibition of growth and aggregation of calcium oxalate crystals in vitro--a comparison of four human proteins. *Eur J Biochem* **253**:637-644.

Grünbaum EG, Freese K und Popp J-P (1991): Praktische Aspekte der Nierendiagnostik. *Kleintierpraxis* **2**:49-60.

Guder WG and Hofmann W (1992): Markers for the diagnosis and monitoring of renal tubular lesions. *Clin Nephrol* **38**, Suppl 1:3-7.

Guder WG and Hofmann W (1993): Differentiation of proteinuria and haematuria by single protein analysis in urine. *Clin Biochem* **26**:277-282.

Guder WG (1995): Niere und ableitende Harnwege. In: Lehrbuch der klinischen Chemie und Pathobiochemie, Greiling H und Gressner AM (eds.), Stuttgart, New York, Schattauer, pp 721-767.

Guy JM, Brammah TB, Holt L, Bernstein RM, McMurray JR, Tieszen K and Cooper RG (1997): Urinary excretion of albumin and retinol binding protein in systemic lupus erythematosus. *Ann Clin Biochem* **34**:668-674.

Hallson PC and Rose GA (1976): Crystalluria in normal subjects and stone formers with and without thiazide and cellulose phosphate treatment. *Br J Urol* **48**:515-524.

- Hallson PC and Rose GA (1979): Uromucoids and urinary stone formation. *Lancet* **1**:1000-1002.
- Hallson PC, Rose GA and Samuell CT (1981): The role of uromucoids in urinary stone formation. In: Urinary calculus., Brockis JG and Finlayson B (eds.), Littleton, PSG Publishing Co., pp 285-295.
- Hallson PC, Choong SK, Kasidas GP and Samuell CT (1997): Effects of Tamm-Horsfall protein with normal and reduced sialic acid content upon the crystallization of calcium phosphate and calcium oxalate in human urine. *Br J Urol* **80**:533-538.
- Hammad SM, Stefansson S, Twal WO, Drake CJ, Fleming P, Remaley A, Brewe, HB Jr. and Argraves WS (1999): Cubilin, the endocytic receptor for intrinsic factor-vitamin B(12) complex, mediates high-density lipoprotein holoparticle endocytosis. *Proc Natl Acad Sci USA* **96**:10158-10163.
- Hampel DJ, Sansome C, Sha M, Brodsky S, Lawson WE and Goligorsky MS (2001): Toward proteomics in uroscopy: urinary protein profiles after radiocontrast medium administration. *J Am Soc Nephrol* **12**:1026-1035.
- Harvey DG and Hoe CM (1966): The use of paper electrophoresis for the routine identification of urinary proteins in the dog. *J Small Anim Pract* **7**:431-440.
- Hess B (1989): Neue pathophysiologische Aspekte der Nephrolithiasis. *Schweiz Med Wschr* **119**:929-934.
- Hess B, Nakagawa Y and Coe FL (1989): Inhibition of calcium oxalate monohydrate crystal aggregation by urine proteins. *Am J Physiol* **257**:F99-106.
- Hess B (1991): The role of Tamm-Horsfall glycoprotein and Nephrocalcins in calcium oxalate monohydrate crystallization processes. *Scanning Microsc* **5**:689-695; discussion 696.
- Hess B, Nakagawa Y, Parks JH and Coe FL (1991): Molecular abnormality of Tamm-Horsfall glycoprotein in calcium oxalate nephrolithiasis. *Am J Physiol* **260**:F569-578.
- Hess B, Zipperle L and Jaeger P (1993): Citrate and calcium effects on Tamm-Horsfall glycoprotein as a modifier of calcium oxalate crystal aggregation. *Am J Physiol* **265**:F784-791.
- Hess B (1994): Tamm-Horsfall glycoprotein and calcium nephrolithiasis. *Miner Electrolyte Metab* **20**:393-398.
- Hess B, Jordi S, Zipperle L, Ettinger E and Giovanoli R (2000): Citrate determines calcium oxalate crystallization kinetics and crystal morphology-studies in the presence of Tamm-Horsfall protein of a healthy subject and a severely recurrent calcium stone former. *Nephrol Dial Transplant* **15**:366-374.
- Hesse A und Brühl M (1988): Comparative aspects of urolithiasis in man and animals (dogs, cats). Proceedings of the III. Annual Symposium, WSAVA, ESVNU, Barcelona:183-194.
- Hesse A (1990): Canine urolithiasis: epidemiological data and findings from analysis of urinary calculi. *J Small Anim Pract* **31**:599-604.
- Hesse A, Steffes H-J, Graf C, Bongartz D und Albrecht F (1997): Aktuelle Daten zur Zusammensetzung und Rasseverteilung der Harnsteine von Hunden. *Berl Münch Tierarztl Wochenschr* **110**:436-439.
- Hession C, Decker JM, Sherblom AP, Kumar S, Yue CC, Mattaliano RJ, Tizard R, Kawashima E, Schmeissner U and Heletky S (1987): Uromodulin (Tamm-Horsfall glycoprotein): a renal ligand for lymphokines. *Science* **237**:1479-1484.
- Heukeshoven J and Dernick R (1988): Improved silver staining procedure for fast staining in PhastSystem Development Unit. I. Staining of sodium dodecyl sulfate gels. *Electrophoresis* **9**:28-32.
- Holmquist P, Torffvit O, Jorgensen PE, Torring N, Nexo E and Sjöblad S (2001): Early urinary changes in Tamm-Horsfall protein and epidermal growth factor in diabetic children. *Pediatr Nephrol* **16**:488-492.
- Hörauf A, Reusch C und Minkus G (1989): Vergleich von Nierenbiopsie und SDS-Polyacrylamidgel-Elektrophorese (SDS-PAGE) bei der Katze. *Tierärztl Prax Suppl* **5**:29-32.

Hörauf A, Reusch C und Minkus G (1990): Aussagekraft des Protein-Kreatinin-Verhältnisses im Urin zur Differenzierung feliner Nephropathien. Tierärztl Prax **18**:423-425.

Howie AJ, Lote CJ, Cunningham AA, Zaccone G and Fasulo S (1993): Distribution of immunoreactive Tamm-Horsfall protein in various species in the vertebrate classes. Cell Tissue Res **274**:15-19.

Howie AJ (1994): Morphometric studies of acute renal failure using anti-brush-border and other antisera. Nephrol Dial Transplant **9**:37-39.

Hoyer JR (1992): Uropontin is selectively incorporated into the organic matrix of urinary calcium oxalate monohydrate and brushite stones. In: VIIth International Symposium on Urolithiasis, Cairns, Australia, pp 118.

Iida S, Peck AB, Byer KJ and Khan SR (1999): Expression of bikunin mRNA in renal epithelial cells after oxalate exposure. J Urol **162**:1480-1486.

Israele V, Darabi A and McCracken GH Jr. (1987): The role of bacterial virulence factors and Tamm-Horsfall protein in the pathogenesis of Escherichia coli urinary tract infection in infants. Am J Dis Child **141**:1230-1234.

Iwata H, Kamei O, Abe Y, Nishio S, Wakatsuki A, Ochi K and Takeuchi M (1988): The organic matrix of urinary uric acid crystals. J Urol **139**:607-610.

Iwata H, Iio S, Nishio S and Takeuchi M (1992): Architecture of mixed calcium oxalate dihydrate and monohydrate stones. Scanning Microsc **6**:231-237; discussion 237-238.

Jaconi S, Rose K, Hughes GJ, Saurat JH and Siegenthaler G (1995): Characterization of two post-translationally processed forms of human serum retinol-binding protein: altered ratios in chronic renal failure. J Lipid Res **36**:1247-1253.

Jaconi S, Saurat JH and Siegenthaler G (1996): Analysis of normal and truncated holo- and apo-retinol-binding protein (RBP) in human serum: altered ratios in chronic renal failure. Eur J Endocrinol **134**:576-582.

Jergens AE, McCaw DL and Hewett JE (1987): Effects of collection time and food consumption on the urine protein/creatinine ratio in the dog. Am J Vet Res **48**:1106-1109.

Joachim GR, Cameron JS, Schwartz M and Becker L (1964): Selectivity of protein excretion in patients with the nephrotic syndrome. J Clin Invest **43**:2332-2346.

Kanalas JJ and Hopfer U (1997): Effect of TGF-beta 1 and TNF-alpha on the plasminogen system of rat proximal tubular epithelial cells. J Am Soc Nephrol **8**:184-192.

Kazumi T, Hozumi T, Ishida Y, Ikeda Y, Kishi K, Haykawa M and Yoshino G (1999): Increased urinary transferrin excretion predicts microalbuminuria in patients with type 2 diabetes. Diabetes Care **22**:1176-1180.

Kelleher J, Humphrey CS, Homer D, Davison AM, Giles GR and Losowsky MS (1983): Vitamin A and its transport proteins in patients with chronic renal failure receiving maintenance haemodialysis and after renal transplantation. Clin Sci **66**:619-626.

Kerjaschki D and Farquhar MG (1982): The pathogenic antigen of Heymann nephritis is a membrane glycoprotein of the renal proximal tubule brush border. Proc Natl Acad Sci USA **79**:5557-5581.

Kerjaschki D and Farquhar MG (1983): Immunocytochemical localization of the Heymann nephritis antigen (GP330) in glomerular epithelial cells of normal Lewis rats. J Exp Med **157**:667-686.

Keutel HJ and King JS (1964): Further studies of matrix substance A. Invest Urol **2**:115-122.

Khan SR, Finlayson B and Hackett RL (1983): Stone matrix as proteins adsorbed on crystal surfaces: a microscopic study. Scan Electron Microsc **87**:379-385.

Khan SR and Hackett RL (1993): Role of organic matrix in urinary stone formation: an ultrastructural study of crystal matrix interface of calcium oxalate monohydrate stones. J Urol **150**:239-245.

King JS, Fielden ML and Boyce WH (1961): Total nondialyzable solids in human urine. XI. chemical

analysis of purified normal uromucoid. *Arch Biochem Biophys* **95**:424f.

King JS and Boyce WH (1963): Immunological studies on serum and urinary proteins in urolith matrix in man. *Ann N Y Acad Sci* **104**:579f.

Kitamura T and Pak CY (1982): Tamm and Horsfall glycoprotein does not promote spontaneous precipitation and crystal growth of calcium oxalate in vitro. *J Urol* **127**:1024-1026.

Klausner JS and Osborne CA (1980): Urinary tract infection and urolithiasis. *Vet Clin North Am Small Anim Pract* **9**:701-711.

Klausner JS, Osborne CA, O'Leary TP, Muscoplat CM and Griffith DP (1980): Experimental induction of struvite uroliths in miniature schnauzer and beagle dogs. *Invest Urol* **18**:127-132.

Knorle R, Schnierle P, Koch A, Buchholz NP, Hering F, Seiler H, Ackermann T and Rutishauser G (1994): Tamm-Horsfall glycoprotein: role in inhibition and promotion of renal calcium oxalate stone formation studied with Fourier-transform infrared spectroscopy. *Clin Chem* **40**:1739-1743.

Kozyraki R, Fyfe J, Kristiansen M, Gerdes C, Jacobsen C, Cui S, Christensen EI, Aminoff M, de la Chapelle A, Krahe R, Verroust PJ and Moestrup SK (1999): The intrinsic factor-vitamin B12 receptor, cubilin, is a high-affinity apolipoprotein A-I receptor facilitating endocytosis of high-density lipoprotein. *Nat Med* **5**:656-661.

Kozyraki R, Fyfe J, Verroust PJ, Jacobsen C, Dautry-Varsat A, Gburek J, Willnow TE, Christensen EI and Moestrup SK (2001): Megalin-dependent cubilin-mediated endocytosis is a major pathway for the apical uptake of transferrin in polarized epithelia. *Proc Natl Acad Sci USA* **98**:12491-12496.

Krohn K, Mero M and Oksanen A (1971): Immunologic observations in canine interstitial nephritis. *Am J Pathol* **65**:157-162.

Kruger JM and Osborne CA (1986): Etiopathogenesis of uric acid and ammonium urate uroliths in non-Dalmatian dogs. *Vet Clin North Am Small Anim Pract* **16**:87-126.

Kruger JM, Osborne CA, Nachreiner RF and Refsal KR (1996): Hypercalcemia and renal failure. Etiology, pathophysiology, diagnosis, and treatment. *Vet Clin North Am Small Anim Pract* **26**:1417-1445.

Kruger JM, Osborne CA and Lulich JP (1999): Canine calcium phosphate uroliths. Etiopathogenesis, diagnosis, and management. *Vet Clin North Am Small Anim Pract* **29**:141-159, xii.

Kumar S and Muchmore A (1990): Tamm-Horsfall protein--uromodulin (1950-1990). *Kidney Int* **37**:1395-1401.

Kunin CM, Chesney RW, Craig WA, England AC and Deangelis C (1978): Enzymuria as a marker of renal injury and disease: Studies of N-acetyl- $\beta$ -glucosaminidase in the general population and in patients with renal disease. *Pediatrics* **62**:751-760.

Kuster G, Shorter RG, Dawson B and Hallenbeck GA (1972): Uric acid metabolism in dalmatians and other dogs. Role of the liver. *Arch Intern Med* **129**:492-496.

Laemmli UK (1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680-685.

Laine J, Etelamaki P, Holmberg C and Dunkel L (1997): Apoptotic cell death in human chronic renal allograft rejection. *Transplantation* **63**:101-105.

Leal JJ and Finlayson B (1977): Adsorption of naturally occurring polymers onto calcium oxalate crystal surfaces. *Invest Urol* **14**:278-283.

Leeker A, Kreft B, Sandmann J, Bates J, Wasenauer G, Muller H, Sack K and Kumar S (1997): Tamm-Horsfall protein inhibits binding of S- and P-fimbriated Escherichia coli to human renal tubular epithelial cells. *Exp Nephrol* **5**:38-46.

Lees GE (1995): Fundamentals of the practice of veterinary nephrology and urology. In: *Canine and feline nephrology and urology*, Osborne CA and Finco DR (eds.), Baltimore, Williams and Williams, pp 49-67.

Leheste JR, Rolinski B, Vorum H, Hilpert J, Nykjaer A, Jacobsen C, Aucouturier P, Moskaug JO, Otto A, Christensen EI and Willnow TE (1999): Megalin knockout mice as an animal model of low molecular weight proteinuria. *Am J Pathol* **155**:1361-1370.

Leheste JR, Melsen F, Wellner M, Jansen P, Schlichting U, Renner-Muller I, Andreassen TT, Wolf E, Bachmann S, Nykjaer A and Willnow TE (2003): Hypocalcemia and osteopathy in mice with kidney-specific megalin gene defect. *Faseb J* **17**:247-249.

Leopold-Temmler B und Nolte I (1995): Verlaufsuntersuchung von nierenkranken Hunden mit der Sodium Dodecyl Sulfat-Polyacrylamid-Gradientengel Urinelektrophorese. *Kleintierpraxis* **40**:103-113.

Liebich H-G (1993): Harnorgane. In: Funktionelle Histologie, 2. Auflage., Stuttgart, New York, Schattauer Verlag, pp 218-232.

Lieske JC, Hammes MS, Hoyer JR and Toback FG (1997): Renal cell osteopontin production is stimulated by calcium oxalate monohydrate crystals. *Kidney Int* **51**:679-686.

Ling GV, Biberstein EL and Hirsh DC (1979): Bacterial pathogens associated with urinary tract infections. *Vet Clin North Am* **9**:617-630.

Ling GV and Rudby AL (1986): Canine uroliths: analysis of data derived from 813 specimens. *Vet Clin North Am* **16**:303-316.

Ling GV, Franti CE, Johnson DL and Ruby AL (1998a): Urolithiasis in dogs. IV: Survey of interrelations among breed, mineral composition, and anatomic location of calculi, and presence of urinary tract infection. *Am J Vet Res* **59**:650-660.

Ling GV, Franti CE, Ruby AL, Johnson DL and Thurmond M (1998b): Urolithiasis in dogs. I: Mineral prevalence and interrelations of mineral composition, age, and sex. *Am J Vet Res* **59**:624-629.

Lloyd SE, Gunther W, Pearce SH, Thomson A, Bianchi ML, Bosio M, Craig IW, Fisher SE, Scheinman SJ, Wrong O, Jentsch TJ and Thakker RV (1997): Characterisation of renal chloride channel, CLCN5, mutations in hypercalciuric nephrolithiasis (kidney stones) disorders. *Hum Mol Genet* **6**:1233-1239.

Lubec D, Grillenberger A, Weninger M and Lubec G (1987): Value of SDS-polyacrylamide gel electrophoresis, small molecular weight proteins and alpha-1-acid glycoprotein for the diagnosis of tubular damage. *Padiatr Padol* **22**:319-324.

Lulich JP and Osborne DC (1990): Interpretation of urine protein-creatinine ratios in dogs with glomerular and nonglomerular disorders. *Compendium* **12**:59-73.

Lulich JP, Osborne CA, Nagode LA, Polzin DJ and Parke ML (1991): Evaluation of urine and serum metabolites in miniature schnauzers with calcium oxalate urolithiasis. *Am J Vet Res* **52**:1583-1590.

Lulich JP, Osborne CA, Thumchai R, Lekcharoensuk C, Ulrich LK, Koehler LA, Bird KA, Swanson LL and Nakagawa Y (1999): Epidemiology of canine calcium oxalate uroliths. Identifying risk factors. *Vet Clin North Am Small Anim Pract* **29**:113-122, xi.

Lynn K (1982): Tamm-Horsfall glycoprotein--what relevance to health and disease? *N Z Med J* **95**:457-458.

Lynn KL, Shenkin A and Marshall RD (1982): Factors affecting excretion of human urinary Tamm-Horsfall glycoprotein. *Clin Sci* **62**:21-26.

Lynn KL and Marshall RD (1984): Excretion of Tamm-Horsfall glycoprotein in renal disease. *Clin Nephrol* **22**:253-257.

MacDougall DF, Cook T, Steward AP and Cattell V (1986): Canine chronic renal disease: Prevalence and types of glomerulonephritis in the dog. *Kidney Int* **29**:1144-1151.

MacGillivray RT, Mendez E, Sinha SK, Sutton MR, Lineback-Zins J and Brew K (1982): The complete amino acid sequence of human serum transferrin. *Proc Natl Acad Sci USA* **79**:2504-2508.

Marino M, Andrews D, Brown D and McCluskey RT (2001): Transcytosis of retinol-binding protein across renal proximal tubule cells after megalin (gp 330)-mediated endocytosis. *J Am Soc Nephrol* **12**:637-648.

Markwell PJ, Robertson WG and Stevenson AE (2000): Urolithiasis: a comparison of humans, cats and dogs. Proceedings of 9th International Symposium on Urolithiasis, pp 785-788.

Mastroianni Kirsztajn G, Nishida SK, Silva MS, Ajzen H and Pereira AB (2000): Urinary retinol-binding protein as a prognostic marker in the treatment of nephrotic syndrome. *Nephron* **86**:109-114.

Maxfield M (1961): Molecular forms of human urinary mucoprotein present under physiological conditions. *Biochim Biophys Acta* **49**:548-558.

McKenzie JK and McQueen EG (1969): Immunofluorescent localization of Tamm-Horsfall mucoprotein in human kidney. *J Clin Pathol* **22**:334-339.

McQueen EG (1962): The nature of urinary casts. *J Clin Pathol* **15**:367.

McQueen EG and Engel GB (1966): Factors determining the aggregation of urinary mucoprotein. *J Clin Pathol* **19**:392-396.

Medetognon-Benissan J, Tardivel S, Hennequin C, Daudon M, Drueke T and Lacour B (1999): Inhibitory effect of bikunin on calcium oxalate crystallization in vitro and urinary bikunin decrease in renal stone formers. *Urol Res* **27**:69-75.

Melick RA, Quelch KJ and Rhodes M (1980): The demonstration of sialic acid in kidney stone matrix. *Clin Sci (Colch)* **59**:401-404.

Meyer-Lindenberg A, Wohlsein P, Trautwein G und Nolte I (1997): Urinproteinanalyse mit der Sodium-Dodecyl-Sulfat-Polyacrylamid-Gradienten-Elektrophorese (SDS-PAGE) bei gesunden und nierenkranken Katzen. *J Vet Med* **44**:39-54.

Meyrier A (1995): Upper and lower urinary tract infections. Epidemiology, etiology, physiopathology, diagnosis, development, treatment. *Rev Prat* **45**:1951-1961.

Miltenyi M (1979): Urinary protein excretion in healthy children. *Clin Nephrol* **12**:216-221.

Min W, Shiraga H, Chalko C, Goldfarb S, Krishna GG and Hoyer JR (1998): Quantitative studies of human urinary excretion of uropontin. *Kidney Int* **53**:189-193.

Minic Z, Hranisavljevic J and Vucelic D (1997): Isolation and characterization of isoforms of retinol binding protein by isoelectrofocusing. *Biochem Molecular Biology International* **41**:1057-1066.

Mitch WE, Collier VU and Walser M (1980): Creatinine metabolism in chronic renal failure. *Clin Sci* **58**:327-335.

Moestrup SK, Cui S, Vorum H, Bregengard C, Bjorn SE, Norris K, Gliemann J and Christensen EI (1995): Evidence that epithelial glycoprotein 330/megalin mediates uptake of polybasic drugs. *J Clin Invest* **96**:1404-1413.

Moestrup SK, Birn H, Fischer PB, Petersen CM, Verroust PJ, Sim RB, Christensen EI and Nexo E (1996): Megalin-mediated endocytosis of transcobalamin-vitamin-B12 complexes suggests a role of the receptor in vitamin-B12 homeostasis. *Proc Natl Acad Sci USA* **93**:8612-8617.

Moestrup SK, Kozyraki R, Kristiansen M, Kayser JH, Rasmussen HH, Brault D, Pontillon F, Goda FO, Christensen EI, Hammond TG and Verroust PJ (1998a): The intrinsic factor-vitamin B12 receptor and target of teratogenic antibodies is a megalin-binding peripheral membrane protein with homology to developmental proteins. *J Biol Chem* **273**:5235-5242.

Moestrup SK, Schousboe I, Jacobsen C, Leheste JR, Christensen EI and Willnow TE (1998b): Beta2-glycoprotein-I (apolipoprotein H) and beta2-glycoprotein-I-phospholipid complex harbor a recognition site for the endocytic receptor megalin. *J Clin Invest* **102**:902-909.

Moestrup SK and Kozyraki R (2000): Cubilin, a high-density lipoprotein receptor. *Curr Opin Lipidol* **11**:133-140.

Monaco HL and Zanotti G (1992): Three-dimensional structure and active site of three hydrophobic molecule-binding proteins with significant amino acid sequence similarity. *Biopolymers* **32**:457-465.

Monroe WE, Davenport DJ and Saunders GK (1989): Twenty-four hour urinary protein loss in healthy

cats and the urinary protein-creatinine ratio as an estimate. Am J Vet Res **50**:1906-1909.

Moore S and Gowland GC (1975): The immunological integrity of matrix substanz A and its possible detection an quantitation in urine. Br J Urol **47**:489-494.

Moorhead JF, Chan MK and Varghese Z (1986): The role of abnormalities of lipid metabolism in the progression of renal disease. In: The progressive nature of renal disease, Mitch WE (ed), New York, Churchill Livingstone, pp 133-148

Morse RM and Resnick MI (1988): A new approach to the study of urinary macromolecules as a participant in calcium oxalate crystallization. J Urol **139**:869-873.

Morse RM and Resnick MI (1989): A study of the incorporation of urinary macromolecules onto crystals of different mineral compositions. J Urol **141**:641-644.

Müller LF (1966): Die Diagnostik der Nierenkrankheiten des Hundes. WTM **53**:740-749.

Müller-Peddinghaus R, Schaefer B, Greiffenhagen U, Wirth W und Trautwein G (1976): Beitrag zur Pathologie, Klinik und Ätiologie der Glomerulonephritis und interstitiellen Nephritis des Hundes. Berl Münch Tierärztl Wschr **89**:7-11.

Müller-Peddinghaus R und Trautwein G (1977a): Harnanalyse mittels SDS-Polyacrylamidgelektrophorese beim Hund. Zbl Vet Med A **24**:731-755.

Müller-Peddinghaus R and Trautwein G (1977b): Spontaneus glomerulonephritis in dogs. Vet Pathol **14**:1-13.

Müller-Peddinghaus R und Trautwein G (1978): Differenzierung von Proteinurien mittels SDS-Polyacrylamidgelektrophorese. Fortschritte der Veterinärmedizin **28**:292-301.

Mustafi D and Nakagawa H (1996): Characterization of Ca 2+ - binding sites in the kidney stone inhibitor glycoprotein nephrocalcin using vanadyl ions: different metal binding properties in strong and weak inhibitor proteins revealed by EPR and ENDOR. Biochemistry **35**:14703-14709.

Muto Y, Smith FR and Goodman DS (1973): Comparative studies of retinol transport in plasma. J Lipid Res **14**:525-532.

Nakagawa H, Kaiser ET and Coe FL (1978): Isolation and characterization of calcium oxalate crystal growth inhibitors form human urine. Biochem Biophys Res Commun **84**:1038-1044.

Nakagawa H, Parks JH, Kézdy FJ and Coe FL (1985): Molecular abnormality of urinary glycoprotein crystal growth inhibitor in calcium nephrolithiasis. Trans Assoc Am Physicians **98**:281-289.

Nakagawa Y (1987): Urinary Tamm-Horsfall glycoprotein in patients with various renal diseases. Nippon Jinzo Gakkai Shi **29**:529-534.

Nakagawa Y, Ahmed M, Hall SL, Deganello S and Coe FL (1987): Isolation from human calcium oxalate renal stones of nephrocalcin, a glycoprotein inhibitor of calcium oxalate crystal growth. Evidence that nephrocalcin from patients with calcium oxalate nephrolithiasis is deficient in gamma-carboxyglutamic acid. J Clin Invest **79**:1782-1787.

Narayanan SH and Appleton HS (1980): Creatinine: a review. Clin Chem **26**:1119-1126.

Narita T, Fujita H, Koshimura J, Meguro H, Kitazato H, Shimotomai T, Kagaya E, Suzuki K, Murata M, Usami A and Ito S (2001): Glycemic control reverses increases in urinary excretions of immunoglobulin G and coeruloplasmin in type 2 diabetic patients with normoalbuminuria. Horm Metab Res **33**:370-378.

Nedelkov D and Nelson RW (2001): Analysis of human urine protein biomarkers via biomolecular interaction analysis mass spectrometry. Am J Kidney Dis **38**:481-487.

Newman DJ, Thakkar H and Gallagher H (2000): Progressive renal disease: does the quality of the proteinuria matter or only the quantity? Clin Chim Acta **297**:43-54.

Nishio S, Abe Y, Wakatsuki A, Iwata H, Ochi K, Takeuchi M and Matsumoto A (1985): Matrix glycosaminoglycan in urinary stones. J Urol **134**:503-505.

Nortier JL, Deschodt-Lanckman MM, Simon S, Thielemans NO, de Prez EE, Depierreux MF, Thielemans CL, Richard C, Lquwerys RR, Bernard AM and Vanherweghem JL (1997): Proximal tubular injury in Chinese herbs nephropathy: monitoring by neutral endopeptidase enzymuria. *Kidney Int* **51**:288-293.

Noseda G, Reiner M und Scali G (1980): Haarzellleukemie und Myelom. *Schweiz Med Wochenschr* **110**:1494-1498.

Nykjaer A, Dragun D, Walther D, Vorum H, Jacobsen C, Herz J, Melsen F, Christensen EI and Willnow TE (1999): An endocytic pathway essential for renal uptake and activation of the steroid 25-(OH) vitamin D3. *Cell* **96**:507-515.

Oddoze C, Morange S, Portugal H, Berland Y and Dussol B (2001): Cystatin C is not more sensitive than creatinine for detecting early renal impairment in patients with diabetes. *Am J Kidney Dis* **38**:310-316.

Orlando RA, Rader K, Authier F, Yamazaki H, Posner BI, Bergeron JJ and Farquhar MG (1998): Megalin is an endocytic receptor for insulin. *J Am Soc Nephrol* **9**:1759-1766.

Osborne CA, Johnson KH, Perman V and Schall WD (1968): Renal amyloidosis in the dog. *J Am Vet Med Assoc* **153**:669-688.

Osborne CA, Johnson KH and Perman V (1969): Amyloid nephrotic syndrome in the dog. *J Am Vet Med Assoc* **154**:1545-1560.

Osborne CA, Johnson KH, Perman V, Fangmann GM and Riis RC (1970): Clinicopathologic progression of renal amyloidosis in a dog. *J Am Vet Med Assoc* **157**:203-219.

Osborne CA, Hammer RF, Stevens JB, Resnick JS and Michael AF (1977): The glomerulus in health and disease: a comparative review of domestic animals and man. *Adv Vet Sci Comp Med* **21**:207-285.

Osborne CA, Polzin DJ, Abdullahi SU, Leininger JR, Clinton CW and Griffith DP (1985): Struvite urolithiasis in animals and man: formation, detection, and dissolution. *Adv Vet Sci Comp Med* **29**:1-101.

Osborne CA, Clinton CW, Bamman LK, Moran HC, Coston BR and Frost AP (1986a): Prevalence of canine uroliths. Minnesota Urolith Center. *Vet Clin North Am Small Anim Pract* **16**:27-44.

Osborne CA, Klausner JS, Polzin DJ and Griffith DP (1986b): Etiopathogenesis of canine struvite urolithiasis. *Vet Clin North Am Small Anim Pract* **16**:67-86.

Osborne CA, Poffenbarger EM, Klausner JS, Johnston SD and Griffith DP (1986c): Etiopathogenesis, clinical manifestations, and management of canine calcium oxalate urolithiasis. *Vet Clin North Am Small Anim Pract* **16**:133-170.

Osborne CA, Polzin DJ, Lulich JP, Kruger JM, Johnston GR and Felice LJ (1989): Relationship of nutritional factors to the cause, dissolution, and prevention of canine uroliths. *Vet Clin North Am Small Anim Pract* **19**:583-619.

Osborne CA, Kruger JP, Lulich JP, Bartges JW, Polzin D, Molitor T, Beauclair KD and Onffroy J (1992): Feline matrix-crystalline urethral plugs: a unifying hypothesis of causes. *J Small Anim Pract* **33**:172-177.

Osborne CA, Stevens JB, Lulich JP, Ulrich LK, Bird KA, Koehler LA and Swanson LL (1995): A clinician's analysis of urinalysis. In: *Canine and Feline Nephrology and Urology*, Osborne CA and Finco DR (eds.), Philadelphia, Williams & Wilkins, pp 136-205.

Osborne CA, Lulich JP, Polzin DJ, Allen TA, Kruger JM, Bartges JW, Koehler LA, Ulrich LK, Bird KA and Swanson LL (1999a): Medical dissolution and prevention of canine struvite urolithiasis. Twenty years of experience. *Vet Clin North Am Small Anim Pract* **29**:73-111, xi.

Osborne CA, Lulich JP, Polzin DJ, Sanderson SL, Koehler LA, Ulrich LK, Bird KA, Swanson LL, Pederson LA and Sudo SZ (1999b): Analysis of 77,000 canine uroliths. Perspectives from the Minnesota Urolith Center. *Vet Clin North Am Small Anim Pract* **29**:17-38, ix-x.

Osborne CA, Sanderson SL, Lulich JP, Bartges JW, Ulrich LK, Koehler LA, Bird KA and Swanson LL

(1999c): Canine cystine urolithiasis. Cause, detection, treatment, and prevention. *Vet Clin North Am Small Anim Pract* **29**:193-211, xiii.

Osicka TM and Comper WD (1998): Tubular inhibition destroys charge selectivity for anionic and neutral horseradish peroxidase. *Biochim Biophys Acta* **1381**:170-178.

Pak CY (1987): Citrate and renal calculi. *Miner Electrolyte Metab* **13**:257-266.

Payne RB (1986): Creatinine clearance: a redundant clinical investigation. *Ann Clin Biochem* **23**:243-250.

Peach RJ, Day WA, Ellingsen PJ and McGiven AR (1988): Ultrastructural localization of Tamm-Horsfall protein in human kidney using immunogold electron microscopy. *Histochem J* **20**:156-164.

Peraldi MN (1992): Le point sur la protéine de Tamm-Horsfall. *Néphrologie* **13**:7-11.

Pesce AJ, Hsu A, Kornhauser C, Sethi K, Ooi BS and Pollak VE (1976): Method for measuring the concentration of urinary proteins according to their molecular size category. *Clin Chem* **22**:667-672.

Polzin DJ, Osborne CA and Bell FW (1986): Canine distal renal tubular acidosis and urolithiasis. *Vet Clin North Am Small Anim Pract* **16**:241-250.

Polzin DJ and Osborne CA (1995): Pathophysiology of renal failure and uremia. In: *Canine and Feline Nephrology and Urology*, Osborne CA and Finco DR (eds.), Philadelphia, Williams & Wilkins, pp 335-367.

Polzin DJ, Osborne DC, Jacob F and Ross S (2000): Chronic renal failure. In: *Textbook of veterinary internal medicine*, Ettinger SJ and Feldman EC (eds.), Philadelphia, WB Saunders, pp 1634-1662.

Popper R, Steigmann F and Dyniewicz HA (1954): Plasma vitamin a level in renal disease. *Amer J Clin Pathol* **15**:272-276.

Porile JL, Asplin JR, Nakagawa H and Coe FL (1993): Evidence for elevated urinary Tamm-Horsfall protein (THP) but not albumin excretion among patients with accelerated calcium oxalate nephrolithiasis (abstract). *J Am Soc Nephrol* **4**:713.

Poulik MD, Farrah D, Malek GH, Shinnick CJ and Smithies O (1975): Low molecular weight urinary proteins - partial amino acid sequences of the retinol-binding proteins of man and dog. *Biochimica et Biophysica Acta* **412**:326-334.

Praga M and Morales E (2002): Renal damage associated with proteinuria. *Kidney Int* **62**:S42-S46.

Rabiet MJ, Blashill A, Furie B and Furie BC (1986): Prothrombin fragment 1 X 2 X 3, a major product of prothrombin activation in human plasma. *J Biol Chem* **261**:13210-13215.

Raila J (1999): Untersuchungen zur Ausscheidung von Vitamin A im Harn von Kaniden. Diss vet med, Leipzig, Berlin: Logos-Verlag, pp 129.

Raila J, Buchholz I, Aupperle H, Raila G, Schoon HA and Schweigert FJ (2000): The distribution of vitamin A and retinol-binding protein in the blood plasma, urine, liver and kidneys of carnivores. *Vet Res* **31**:541-551.

Raila J und Schweigert FJ (2001): Zur Bedeutung der Nieren im Vitamin-Stoffwechsel. *Berl Münch Tierärztl Wschr* **114**:257-266.

Raila J, Radon R, Trupschuch A and Schweigert FJ (2002): Retinol and retinyl ester responses in the blood plasma and urine of dogs after a single oral dose of vitamin A. *J Nutr* **132**:1673S-1675S.

Raila J, Neumann U and Schweigert FJ (2003): Immunochemical localization of megalin, retinol-binding protein and Tamm-Horsfall glycoprotein in the kidneys of dogs. *Vet Res Com* **27**:125-135.

Rask L, Anundi H, Bohme J, Eriksson U, Ronne H, Sege K and Peterson PA (1981): Structural and functional studies of vitamin A-binding proteins. *Ann N Y Acad Sci* **359**:79-90.

Remuzzi G, Ruggenenti P and Benigni A (1997): Understanding the nature of renal disease progression. *Kidney Int* **51**:2-15.

Resnick MI, Munday D and Boyce WH (1980): Urinary B2-microglobulin excretion and renal lithiasis. *Invest Urol* **18**:137-138.

Rhodes DC, Hinsman EJ, Rhodes JA and Hawkins EC (1992): Urinary Tamm-Horsfall glycoprotein concentrations in normal and urolithiasis-affected male cats determined by an ELISA. *Zentralbl Veterinarmed [A]* **39**:621-634.

Rhodes DC, Hinsman EJ and Rhodes JA (1993): Cation-induced aggregation of cat Tamm-Horsfall glycoprotein and its possible role in feline urolithiasis. *Zentralbl Veterinarmed [A]* **40**:283-291.

Ribaya-Mercado JD, Blanco MC, Fox JG and Russell RM (1994): High concentrations of vitamin A esters circulate primarily as retinyl stearate and are stored primarily as retinyl palmitate in ferret tissues. *J Am Coll Nutr* **13**:83-86.

Rigas DA and Heller CG (1951): The amount and nature of urinary proteins in normal human subjects. *J Clin Invest* **30**:853-861.

Robertson WG (1993): Urinary tract calculi. In: *Metabolic bone and stone disease*, Nording BEC, Need AG and Morris HA (eds.), Edinburgh, Churchill Livingstone, pp 249-311.

Robertson WG, Jones JS, Heaton MA, Stevenson AE and Markwell PJ (2002): Predicting the crystallization potential of urine from cats and dogs with respect to calcium oxalate and magnesium ammonium phosphate (struvite). *J Nutr* **32**:1637S-1641S.

Romero MC, Nocera S and Nesse AB (1997): Decreased Tamm-Horsfall protein in lithiasic patients. *Clin Biochem* **30**:63-67.

Rose GA and Sulaiman S (1982): Tamm-Horsfall mucoproteins promote calcium oxalate crystal formation in urine: quantitative studies. *J Urol* **127**:177-179.

Ryall RL, Harnett RM, Hibberd CM, Edyvane KA and Marshall VR (1991): Effects of chondroitin sulphate, human serum albumin and Tamm-Horsfall mucoprotein on calcium oxalate crystallization in undiluted human urine. *Urol Res* **19**:181-188.

Ryall RL, Grover PK, Stapleton AM, Barrell DK, Tang Y, Moritz RL and Simpson RJ (1995): The urinary F1 activation peptide of human prothrombin is a potent inhibitor of calcium oxalate crystallization in undiluted human urine in vitro. *Clin Sci (Colch)* **89**:533-541.

Ryall RL (1996): Glycosaminoglycans, proteins, and stone formation: adult themes and child's play. *Pediatr Nephrol* **10**:656-666.

Sahali D, Mulliez N, Chatelet F, Laurent-Winter C, Citadelle D, Sabourin JC, Roux C, Ronco P and Verroust P (1993): Comparative immunochemistry and ontogeny of two closely related coated pit proteins. The 280-kd target of teratogenic antibodies and the 330-kd target of nephritogenic antibodies. *Am J Pathol* **142**:1654-1667.

Samuell CT (1979): Uromucoid excretion in normal subjects, calcium stone formers and in patients with chronic renal failure. *Urol Res* **7**:5-12.

Sarre H (1958): *Nierenkrankheiten: Physiologie, Pathophysiologie, Klinik und Therapie.*, 1. Auflage, Seiten 540.

Savory J, Pu PH and Sunderman FW Jr. (1968): A biuret method for determination of protein in normal urine. *Clin Chem* **14**:1160-1171.

Scarpioni L, Dall'Aglio PP, Poisetti PG and Buzio C (1976): Retinol-binding protein in serum and in urine of glomerular and tubular nephropathies. *Clin Chim Acta* **68**:107-113.

Scheinman SJ, Cox JP, Lloyd SE, Pearce SH, Salenger PV, Hoopes RR, Bushinsky DA, Wrong O, Asplin JR, Langman CB, Norden AG and Thakker RV (2000): Isolated hypercalciuria with mutation in CLCN5: relevance to idiopathic hypercalciuria. *Kidney Int* **57**:232-239.

Schiwara HW und Spiller A (1989): Differenzierung von renalen und postrenalen Blutungen und Proteinurie mittels SDS Polyacrylamidegelektrophoresis und Immunoblotting. *Klin Wochenschr* **67**:14-18.

Schmitz C, Hilpert J, Jacobsen C, Boensch C, Christensen EI, Luft FC and Willnow TE (2002): Megalin deficiency offers protection from renal aminoglycoside accumulation. *J Biol Chem* **277**:618-622.

Schnierle P, Hering F and Seiler H (1996): Isoelectric focusing of Tamm-Horsfall glycoproteins: a simple tool for recognizing recurrent calcium oxalate renal stone formers. *Urol Res* **24**:79-82.

Schöne S, Stein G, Stelzner A, Geinitz S, Scholz S, Keil E und Schöne F (1984): Untersuchungen über den Vitamin-A-Status von Patienten mit chronischer Niereninsuffizienz. *Z Klin Med* **40**:645-648.

Schultz CJ, Dalton RN, Neil HA, Konopelska-Bahu T and Dunger DB (2001): Markers of renal tubular dysfunction measured annually do not predict risk of microalbuminuria in the first few years after diagnosis of Type I diabetes. *Diabetologia* **44**:224-229.

Schultze AE and Jensen RK (1989): Sodium dodecyl sulfate polyacrylamid gel electrophoresis of canine urinary proteins for the analysis and differentiation of tubular and glomerular diseases. *Vet Clin Path* **18**:93-97.

Schurek HJ, Panzer J, Wiemeyer A, Kuhn K, Aeikens B and Brod J (1982): Effects of aging on the glomerular capillaries, blood pressure, and renal function in rats. *Contrib Nephrol* **30**:157-162.

Schweigert FJ (1988): Insensitivity of dogs to the effects of nonspecific bound vitamin A in plasma. *Int J Vitam Nutr Res* **58**:23-25.

Schweigert FJ und Uehlein-Harrell S (1989): Diagnostische Bedeutung Lipoprotein-gebundener Vitamin-A-Ester im Blutplasma von Karnivoren. *Tierärztl Prax Suppl* **5**:78-80.

Schweigert FJ, Uehlein-Harrell S and Zucker H (1990): Effect of feeding on vitamin A concentrations in blood plasma of dogs. *Zentralbl Veterinarmed A* **37**:605-609.

Schweigert FJ, Thomann E and Zucker H (1991): Vitamin A in the urine of carnivores. *Int J Vitam Nutr Res* **61**:110-113.

Schweigert FJ (1998): Vitamin A: Stoffwechsel, Genexpression und embryonale Entwicklung. Übers Tierernährg **26**:1-24.

Schweigert FJ, Buchholz I and Bonitz K (1998): Effect of age on the levels of retinol and retinyl esters in blood plasma, liver and kidney of dogs. *Int J Vitam Nutr Res* **68**:237-241.

Schweigert FJ and Bok V (2000): Vitamin A in blood plasma and urine of dogs is affected by the dietary level of vitamin A. *Int J Vitam Nutr Res* **70**:84-91.

Schweigert FJ, Raila J and Haebel S (2002): Vitamin A excreted in the urine of canines is associated with a Tamm-Horsfall like protein. *Vet Res* **33**:299-311.

Scurr DS and Robertson WG (1986a): Modifiers of calcium oxalate crystallization found in urine. II. Studies on their mode of action in an artificial urine. *J Urol* **136**:128-131.

Scurr DS and Robertson WG (1986b): Modifiers of calcium oxalate crystallization found in urine. III. Studies on the role of Tamm-Horsfall mucoprotein and of ionic strength. *J Urol* **136**:505-507.

See WA and Williams RD (1992): Urothelial injury and clotting cascade activation: common denominators in particulate adherence to urothelial surfaces. *J Urol* **147**:541-548.

Selting KA, Ogilvie GK, Lana SE, Fettman MJ, Mitchener KL, Hansen RA, Richardson KL, Walton JA and Scherk MA (2000): Serum alpha 1-acid glycoprotein concentrations in healthy and tumor-bearing cats. *J Vet Intern Med* **14**:503-506.

Shiraga H, Min W, Vandusen WJ, Claymen MD, Miner D, Terrel CH, Sherbotie JR, Foreman JW, Przysiecki C, Neilson EG and Hoyer JR (1992): Inhibition of calcium oxalate crystal growth in vitro by uropontin: another member of the aspartic acid-rich protein superfamily. *Proc Natl Acad Sci U S A* **89**:426-430.

Siddiqui AA, Sultana T, Buchholz NP, Waqar MA and Talati J (1998): Proteins in renal stones and urine of stone formers. *Urol Res* **26**:383-388.

Sikri KL, Foster CL, MacHugh N and Marshall RD (1981): Localization of Tamm-Horsfall glycoprotein in the human kidney using immuno-fluorescence and immuno-electron microscopical techniques. *J Anat* **132**:597-605.

Smith FR and Goodman DS (1971): The effects of diseases of the liver, thyroid, and kidneys on the transport of vitamin A in human plasma. *J Clin Invest* **50**:2426-2436.

Smith FR and Goodman DS (1976): Vitamin A transport in human vitamin A toxicity. *N Engl J Med* **294**:805-808.

Smith LH (1987): Pathogenesis of renal stones. *Miner Electrolyte Metab* **13**:214-219.

Smith MF and Comper WD (1996): A codiffusing system as a novel model for capillary wall charge selectivity. *Biophys Chem* **62**:73-80.

Sophasan S, Chatasingh S, Thanphaichitr P and Dhanamitta S (1980): Tamm Horsfall mucoprotein in urine of potential bladder stone formers. *J Urol* **124**:522-524.

Sousa MM, Norden AG, Jacobsen C, Willnow TE, Christensen EI, Thakker RV, Verroust PJ, Moestrup SK and Saraiva MJ (2000): Evidence for the role of megalin in renal uptake of transthyretin. *J Biol Chem* **275**:38176-38181.

Spector AR, Gray A and Prien EL Jr. (1976): Kidney stone matrix. Differences in acidic protein composition. *Invest Urol* **13**:387-389.

Spitzauer S, Schweiger C, Sperr WR, Pandjaitan B, Valent P, Muhl S, Ebner C, Scheiner O, Kraft D, and Rumpold H (1994): Molecular characterization of dog albumin as a cross-reactive allergen. *J Allergy Clin Immunol* **93**:614-627.

Stapleton AM, Dawson CJ, Grover PK, Hohmann A, Comacchio R, Boswarva V, Tang Y and Ryall RL (1996): Further evidence linking urolithiasis and blood coagulation: urinary prothrombin fragment 1 is present in stone matrix. *Kidney Int* **49**:880-888.

Stephensen CB, Alvarez JO, Kohatsu J, Hardmeier R, Kennedy JI Jr. and Gammon RB Jr. (1994): Vitamin A is excreted in the urine during acute infection. *Am J Clin Nutr* **60**:388-392.

Stewart WK and Fleming LW (1982): Plasma retinol and retinol binding protein concentrations in patients on maintenance haemodialysis with and without vitamin A supplements. *Nephron* **30**:15-21.

Storch S, Saggi S, Megyesi J, Price PM and Safirstein R (1992): Ureteral obstruction decreases renal prepro-epidermal growth factor and Tamm-Horsfall expression. *Kidney Int* **42**:89-94.

Sugimoto T, Funae Y, Rubben H, Nishio S, Hautmann R and Lutzeyer W (1985): Resolution of proteins in the kidney stone matrix using high-performance liquid chromatography. *Eur Urol* **11**:334-340.

Suzuki K, Moriyama M, Nakajima C, Kawamura K, Miyazawa K, Tsugawa R, Kikuchi N and Nagata K (1994): Isolation and partial characterization of crystal aggregation: Evidence of activation peptide of human prothrombin. *Urol Res* **22**:45-50.

Tamaki K, Okuda S, Ando T, Iwamoto T, Nakayama M and Fujishima M (1994): TGF- $\beta$  1 in glomerulosclerosis and interstitial fibrosis of adriamycin nephropathy. *Kidney Int* **45**:525-536.

Tamm I and Horsfall FL (1950): Characterization and Separation of an Inhibitor of viral hemagglutination present in urine. *Proc Soc Etp Biol Med* **74**:108-114.

Tang Y, Grover PK, Moritz RL, Simpson RJ and Ryall RL (1995): Is nephrocalcin related to the urinary derivative (bikunin) of inter-alpha-trypsin inhibitor? *Br J Urol* **76**:425-430.

Tardivel S, Medetognon J, Randoux C, Kebede M, Drueke T, Daudon M, Hennequin C and Lacour B (1999): Alpha-1-microglobulin: inhibitory effect on calcium oxalate crystallization in vitro and decreased urinary concentration in calcium oxalate stone formers. *Urol Res* **27**:243-249.

Tencer J, Torffvit O, Thysell H, Rippe B and Grubb A (1998): Proteinuria selectivity index based upon alpha 2-macroglobulin or IgM is superior to the IgG based index in differentiating glomerular diseases. Technical note. *Kidney Int* **54**:2098-2105.

- Tencer J, Bakoush O and Torffvit O (2000): Diagnostic and prognostic significance of proteinuria selectivity index in glomerular diseases. *Clin Chim Acta* **297**:73-83.
- Thomas DB, Davies M and Williams JD (1993a): Release of gelatinase and superoxide from human mononuclear phagocytes in response to particulate Tamm Horsfall protein. *Am J Pathol* **142**:249-260.
- Thomas DB, Davies M and Williams JD (1993b): Tamm-Horsfall protein: an aetiological agent in tubulointerstitial disease? *Exp Nephrol* **1**:281-284.
- Thomas ME, Brunskill NJ, Harris KP, Bailey E, Pringle JH, Furness PN and Walls J (1999): Proteinuria induces tubular cell turnover: A potential mechanism for tubular atrophy. *Kidney Int* **55**:890-898.
- Thornley C, Dawnay A and Cattell WR (1985): Human Tamm-Horsfall glycoprotein: urinary and plasma levels in normal subjects and patients with renal disease determined by a fully validated radioimmunoassay. *Clin Sci* **68**:529-535.
- Tishkov I, Razboynikov S, Tchoukanov C and Shipkova P (1978): Biochemical methods for the assessment of the selectivity of proteinuria. *Int Urol Nephrol* **10**:245-249.
- Tomlinson PA, Dalton RN, Turner C and Chantler C (1990): Measurement of  $\beta_2$ -microglobulin, retinol-binding protein, alpha1-microglobulin and urine protein 1 in healthy children using enzyme-linked immunosorbent assay. *Clin Chim Acta* **192**:99-106.
- Tomlinson PA, Dalton RN, Hartley B, Haycock GB and Chantler C (1997): Low molecular weight protein excretion in glomerular disease: a comparative analysis. *Pediatr Nephrol* **11**:285-290.
- Topping MD, Forster HW, Dolman C, Luczynska CM and Bernard AM (1986): Measurement of urinary retinol-binding protein by enzyme-linked immunosorbent assay, and its application to detection of tubular proteinuria. *Clin Chim Acta* **32**:1863-1866.
- Torffvit O, Agardh CD, Kjellsson B and Wieslander J (1992): Tubular secretion of Tamm-Horsfall protein in type 1 (insulin- dependent) diabetes mellitus using a simplified enzyme linked immunoassay. *Clin Chim Acta* **205**:31-41.
- Torffvit O, Jorgensen PE, Kamper AL, Holstein-Rathlou NH, Leyssac PP, Poulsen SS and Strandgaard S (1998): Urinary excretion of Tamm-Horsfall protein and epidermal growth factor in chronic nephropathy. *Nephron* **79**:167-172.
- Trautwein G und Müller-Peddinghaus R (1978): Klassifikation und Pathogenese von Glomerulopathien der Tiere. *Dtsch Tierärztl Wschr* **85**:205-212.
- Truong LD, Petruska S, Yang G, Gurpinar T, Shappell S, Lechago J, Rouse D and Suki WN (1996): Cell apoptosis and proliferation in experimental chronic obstructive uropathy. *Kidney Int* **50**:200-207.
- Umekawa T, Kohri K, Amasaki N, Yamate T, Yoshida K, Yamamoto K, Suzuki Y, Sinohara H and Kurita T (1993): Sequencing of a urinary stone protein identical to alpha-1-antitrypsin. *Biochem Biophys Res Commun* **193**:1049-1053.
- Vahlquist A, Johnsson A and Nygren K (1979): Metabolism of the vitamin A transporting protein-complex. *Amer J Clin Nutrit* **32**:1433-1438.
- Vahlquist A, Berne B and Berne C (1982): Skin content and plasma transport of vitamin A and beta-carotene in chronic renal failure. *Eur J Clin Invest* **12**:63-67.
- Vasiliades J (1976): Reaction of alkaline sodium picrate with creatinine: I. Kinetics and mechanism of formation of the mono-creatinine picric acid complex. *Clin Chem* **22**:1664-1671.
- Verdier JM, Dussol B, Casanova P, Daudon M, Dupuy P, Berthezenne P, Boistelle R, Berland Y and Dagorn JC (1992): Evidence that human kidney produces a protein similar to lithostathin, the pancreatic inhibitor of CaCO<sub>3</sub> crystal growth. *Eur J Clin Invest* **22**:469-474.
- Vermeulen CW and Lyon ES (1968): Mechanism of genesis and growth of calculi. *Am J Med* **45**:684-692.
- Verroust PJ and Kozyraki R (2001): The roles of cubilin and megalin, two multiligand receptors, in

proximal tubule function: possible implication in the progression of renal disease. *Curr Opin Nephrol Hypertens* **10**:33-38.

Vetr H and Gebhard W (1990): Structure of the human alpha 1-microglobulin-bikunin gene. *Biol Chem Hoppe Seyler* **371**:1185-1196.

Vlahou A, Schellhammer PF, Medrinos S, Kondylis FI, Gong L, Nasim S and Wright GL (2001): Development of a novel proteomic approach for the detection of transitional cell carcinoma of the bladder in urine. *Am J Pathol* **158**:1491-1501.

Volpert A, Nolte I und Käufer-Weiß I (1989): Vertikale Sodium-Dodecyl-Sulfat-Polyacrylamid-Gradientengel-Elektrophorese im Microsystem zur Differenzierung von Urinproteinen in der Diagnostik von Nierenerkrankungen beim Hund. *Tierärztl Prax Suppl* **5**:24-28.

Wang Y, Chen J, Chen L, Tay YC, Rangan GK and Harris DC (1997): Induction of monocyte chemoattractant protein-1 in proximal tubule cell by urinary protein. *J Am Soc Nephrol* **8**:1537-1545.

Wangsiripaisan A, Gengaro PE, Edelstein CL and Schrier RW (2001): Role of polymeric Tamm-Horsfall protein in cast formation: oligosaccharide and tubular fluid ions. *Kidney Int* **59**:932-940.

Weaver AD (1970): Canine urolithiasis, chemical composition and outcome of 100 cases. *J Small Anim Pract* **11**:93-107.

Weber MH und Scholz P (1985): Mikro-Flachgelelektrophorese und alpha1-Mikroglobulin - Neue Begriffe in der Proteinuriediagnostik. *Mitt Klin Nephrologie* **XIV**:117-122.

Weber MH, Cheong KS, Schott KJ and Neuhoff V (1986): Microelectrophoresis in the differential diagnosis of proteinuric diseases. *Electrophoresis* **7**:134-141.

Wenkel R, Berg W und Prange H (1998): Harnsteine bei Kleintieren und anderen Tieren - Eine retrospektive Studie von 1980-1989. *Dtsch Tierärztl Wschr* **105**:182-186.

White JV, Olivier NB, Reimann K and Johnson C (1984): Use of protein-to-creatinine ratio in a single urine specimen for quantitative estimation of canine proteinuria. *J Am Vet Med Assoc* **185**:882-885.

White RHR (1981): Clinical application of selectivity of proteinuria. *Contr Nephrol* **24**:63-71.

Wikstrom B and Wieslander J (1981): Excretion of Tamm-Horsfall urinary glycoprotein (uromucoid) in renal calcium stone formers. In: *Urolithiasis, clinical and basic research.*, Smith LH (eds.), New York, Plenum Press, pp 685-688.

Williams JD and Coles GA (1994): Proteinuria - a direct cause of renal morbidity? *Kidney Int* **45**:443-450.

Wilson DE, Chan IF, Cheung AK, Dutz W and Buchi KN (1985): Retinyl ester retention in chronic renal failure. Further evidence for a defect in chylomicron remnant metabolism. *Atherosclerosis* **57**:189-197.

Wilson DE, Hejazi J, Elstad NL, Chan IF, Gleeson JM and Iverius PH (1987): Novel aspects of vitamin A metabolism in the dog: distribution of lipoprotein retinyl esters in vitamin A-deprived and cholesterol-fed animals. *Biochim Biophys Acta* **922**:247-258.

Worcester EM, Nakagawa Y, Wabner CL, Kumar S and Coe FL (1988): Crystal adsorption and growth slowing by nephrocalcin, albumin, and Tamm-Horsfall protein. *Am J Physiol* **255**:F1197-1205.

Worcester EM and Beshensky AM (1995): Osteopontin inhibits nucleation of calcium oxalate crystals. *Ann N Y Acad Sci* **760**:375-377.

Wright GL, Cazares LH, Leung SM, Nasim S, Adam B, Yip T, Schellhammer PF, Gong L and Vlahou A (2000): Proteinchip-surface enhanced laser desorption/ionization (SELDI) mass spectrometry: a novel protein biochip technology for detection of prostate cancer biomarkers in complex protein mixtures. *Prostate Cancer Prostatic Dis* **2**:264-276.

Yatzidis H, Digenis P and Fountas P (1975): Hypervitaminosis A accompanying advanced chronic renal failure. *Br Med J* **3**:352-353.

Yoshioka T, Koide T, Utsunomiya M, Itatani H, Oka T and Sonoda T (1989): Possible role of Tamm-

Horsfall glycoprotein in calcium oxalate crystallisation. Br J Urol **64**:463-467.

Zamparo O and Comper WD (1990): Model anionic polysaccharide matrices exhibit lower charge selectivity than is normally associated with kidney ultrafiltration. Biophys Chem **38**:167-178.

Zimmerhackl LB (1993): Evaluation of nephrotoxicity with renal antigens in children: role of Tamm-Horsfall protein. Eur J Clin Pharmacol **44 Suppl 1**:S39-42.

Zoja C, Morigi M and Figliuzzi M (1995): Proximal tubular cell synthesis and secretion of endothelin-1 on challenge with albumin and other proteins. Am J Kidney Dis **26**:934-941.

Zollinger HU (1971): [Morphological nosology of glomerulonephritides] Beitr Pathol **143**:395-406.