

8. Anhang

8.1 Literaturverzeichnis

- Abbas, A. K., Lichtman, A. H., Pober, J. S. (2000). Cellular and Molecular Immunology. 4th edition, Saunders Company 2000
- Ahn, K., Gruhler, A., Galocha, B., Jones, T. R., Wiertz, E. J., Ploegh, H. L., Peterson, P. A., Yang, Y., and Fruh, K. (1997). The ER-luminal domain of the HCMV glycoprotein US6 inhibits peptide translocation by TAP. *Immunity* 6, 613-21.
- Alexander, J., Payne, J. A., Murray, R., Frelinger, J. A., and Cresswell, P. (1989). Differential transport requirements of HLA and H-2 class I glycoproteins. *Immunogenetics* 29, 380-8.
- Allen, R. L., O'Callaghan, C. A., McMichael, A. J., and Bowness, P. (1999). Cutting edge: HLA-B27 can form a novel beta 2-microglobulin-free heavy chain homodimer structure. *J Immunol* 162, 5045-8.
- Androlewicz, M. J., Anderson, K. S., and Cresswell, P. (1993). Evidence that transporters associated with antigen processing translocate a major histocompatibility complex class I-binding peptide into the endoplasmic reticulum in an ATP-dependent manner. *Proc Natl Acad Sci U S A* 90, 9130-4.
- Androlewicz, M. J., Ortman, B., van Endert, P. M., Spies, T., and Cresswell, P. (1994). Characteristics of peptide and major histocompatibility complex class I/beta 2-microglobulin binding to the transporters associated with antigen processing (TAP1 and TAP2). *Proc Natl Acad Sci U S A* 91, 12716-20.
- Androlewicz, M. J., and Cresswell, P. (1994). Human transporters associated with antigen processing possess a promiscuous peptide-binding site. *Immunity* 1, 7-14.
- Armandola, E. A., Momburg, F., Nijenhuis, M., Bulbuc, N., Fruh, K., and Hammerling, G. J. (1996). A point mutation in the human transporter associated with antigen processing (TAP2) alters the peptide transport specificity. *Eur J Immunol* 26, 1748-55.
- Arnett, K. L., Huang, W., Valiante, N. M., Barber, L. D., and Parham, P. (1998). The Bw4/Bw6 difference between HLA-B*0802 and HLA-B*0801 changes the peptides endogenously bound and the stimulation of alloreactive T cells. *Immunogenetics* 48, 56-61.
- Balow, J. P., Weissman, J. D., and Kearse, K. P. (1995). Unique expression of major histocompatibility complex class I proteins in the absence of glucose trimming and calnexin association. *J Biol Chem* 270, 29025-9.
- Bangia, N., Lehner, P. J., Hughes, E. A., Surman, M., and Cresswell, P. (1999). The N-terminal region of tapasin is required to stabilize the MHC class I loading complex. *Eur J Immunol* 29, 1858-70.
- Barnden, M. J., Purcell, A. W., Gorman, J. J., and McCluskey, J. (2000). Tapasin-mediated retention and optimization of peptide ligands during the assembly of class I molecules. *J Immunol* 165, 322-30.
- Barnstable, C. J., Bodmer, W. F., Brown, G., Galfre, G., Milstein, C., Williams, A. F., and Ziegler, A. (1978). Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens-new tools for genetic analysis. *Cell* 14, 9-20.
- Baumeister, W., Walz, J., Zuhl, F., and Seemuller, E. (1998). The proteasome: paradigm of a self-compartmentalizing protease. *Cell* 92, 367-80.

- Beersma, M. F., Bijlmakers, M. J., and Ploegh, H. L. (1993). Human cytomegalovirus down-regulates HLA class I expression by reducing the stability of class I H chains. *J Immunol* 151, 4455-64.
- Beissbarth, T., Sun, J., Kavathas, P. B., and Ortmann, B. (2000). Increased efficiency of folding and peptide loading of mutant MHC class I molecules. *Eur J Immunol* 30, 1203-13.
- Benjamin, R. J., Madrigal, J. A., and Parham, P. (1991). Peptide binding to empty HLA-B27 molecules of viable human cells. *Nature* 351, 74-7.
- Bjorkman, P. J., Saper, M. A., Samraoui, B., Bennett, W. S., Strominger, J. L., and Wiley, D. C. (1987). Structure of the human class I histocompatibility antigen, HLA-A2. *Nature* 329, 506-12.
- Bluestone, J. A., Kaliyaperumal, A., Jameson, S., Miller, S., and Dick, R., 2nd (1993). Peptide-induced changes in class I heavy chains alter allorecognition. *J Immunol* 151, 3943-53.
- Brodsky, F. M., Bodmer, W. F., and Parham, P. (1979). Characterization of a monoclonal anti-beta 2-microglobulin antibody and its use in the genetic and biochemical analysis of major histocompatibility antigens. *Eur J Immunol* 9, 536-45.
- Burgert, H. G., and Kvist, S. (1985). An adenovirus type 2 glycoprotein blocks cell surface expression of human histocompatibility class I antigens. *Cell* 41, 987-97.
- Carreno, B. M., Solheim, J. C., Harris, M., Stroynowski, I., Connolly, J. M., and Hansen, T. H. (1995). TAP associates with a unique class I conformation, whereas calnexin associates with multiple class I forms in mouse and man. *J Immunol* 155, 4726-33.
- Chen, C. C., Kurokawa, T., Shaw, S. Y., Tillotson, L. G., Kalled, S., and Isselbacher, K. J. (1986). Human erythrocyte glucose transporter: normal asymmetric orientation and function in liposomes. *Proc Natl Acad Sci U S A* 83, 2652-6.
- Copeman, J., Bangia, N., Cross, J. C., and Cresswell, P. (1998). Elucidation of the genetic basis of the antigen presentation defects in the mutant cell line .220 reveals polymorphism and alternative splicing of the tapasin gene. *Eur J Immunol* 28, 3783-91.
- Copeman, J., Han, R. N., Caniggia, I., McMaster, M., Fisher, S. J., and Cross, J. C. (2000). Posttranscriptional regulation of human leukocyte antigen G during human extravillous cytotrophoblast differentiation. *Biol Reprod* 62, 1543-50.
- Corbett, E. F., Oikawa, K., Francois, P., Tessier, D. C., Kay, C., Bergeron, J. J., Thomas, D. Y., Krause, K. H., and Michalak, M. (1999). Ca²⁺ regulation of interactions between endoplasmic reticulum chaperones. *J Biol Chem* 274, 6203-11.
- Corbett, E. F., and Michalak, M. (2000). Calcium, a signaling molecule in the endoplasmic reticulum? *Trends Biochem Sci* 25, 307-11.
- Cotten, J. F., Ostedgaard, L. S., Carson, M. R., and Welsh, M. J. (1996). Effect of cystic fibrosis-associated mutations in the fourth intracellular loop of cystic fibrosis transmembrane conductance regulator. *J Biol Chem* 271, 21279-84.
- Cresswell, P., Bangia, N., Dick, T., and Diedrich, G. (1999). The nature of the MHC class I peptide loading complex. *Immunol Rev* 172, 21-8.
- Danilczyk, U. G., Cohen-Doyle, M. F., and Williams, D. B. (2000). Functional relationship between calreticulin, calnexin, and the endoplasmic reticulum luminal domain of calnexin. *J Biol Chem* 275, 13089-97.
- David, V., Hochstenbach, F., Rajagopalan, S., and Brenner, M. B. (1993). Interaction with newly synthesized and retained proteins in the endoplasmic reticulum

- suggests a chaperone function for human integral membrane protein IP90 (calnexin). *J Biol Chem* 268, 9585-92.
- Day, P. M., Esquivel, F., Lukszo, J., Bennink, J. R., and Yewdell, J. W. (1995). Effect of TAP on the generation and intracellular trafficking of peptide-receptive major histocompatibility complex class I molecules. *Immunity* 2, 137-47.
- De la Salle, H., Hanau, D., Fricker, D., Urlacher, A., Kelly, A., Salamero, J., Powis, S. H., Donato, L., Bausinger, H., Laforet, M., and et al. (1994). Homozygous human TAP peptide transporter mutation in HLA class I deficiency. *Science* 265, 237-41.
- Degen, E., and Williams, D. B. (1991). Participation of a novel 88-kD protein in the biogenesis of murine class I histocompatibility molecules. *J Cell Biol* 112, 1099-115.
- Degen, E., Cohen-Doyle, M. F., and Williams, D. B. (1992). Efficient dissociation of the p88 chaperone from major histocompatibility complex class I molecules requires both beta 2- microglobulin and peptide. *J Exp Med* 175, 1653-61.
- Delp, K., Momburg, F., Hilmes, C., Huber, C., and Seliger, B. (2000). Functional deficiencies of components of the MHC class I antigen pathway in human tumors of epithelial origin. *Bone Marrow Transplant* 25 Suppl 2, S88-95.
- DeMars, R., Rudersdorf, R., Chang, C., Petersen, J., Strandtmann, J., Korn, N., Sidwell, B., and Orr, H. T. (1985). Mutations that impair a posttranscriptional step in expression of HLA-A and -B antigens. *Proc Natl Acad Sci U S A* 82, 8183-7.
- Deverson, E. V., Gow, I. R., Coadwell, W. J., Monaco, J. J., Butcher, G. W., and Howard, J. C. (1990). MHC class II region encoding proteins related to the multidrug resistance family of transmembrane transporters. *Nature* 348, 738-41.
- DiBrino, M., Parker, K. C., Margulies, D. H., Shiloach, J., Turner, R. V., Biddison, W. E., and Coligan, J. E. (1995). Identification of the peptide binding motif for HLA-B44, one of the most common HLA-B alleles in the Caucasian population. *Biochemistry* 34, 10130-8.
- Diedrich, G., Bangia, N., Pan, M., and Cresswell, P. (2001). A role for calnexin in the assembly of the MHC class I loading complex in the endoplasmic reticulum. *J Immunol* 166, 1703-9.
- Ehrlich, R. (1997). Modulation of antigen processing and presentation by persistent virus infections and in tumors. *Hum Immunol* 54, 104-16.
- Elliott, T. (1991). How do peptides associate with MHC class I molecules? *Immunol Today* 12, 386-8.
- Elliott, T., Elvin, J., Cerundolo, V., Allen, H., and Townsend, A. (1992). Structural requirements for the peptide-induced conformational change of free major histocompatibility complex class I heavy chains. *Eur J Immunol* 22, 2085-91.
- Fahnestock, M. L., Johnson, J. L., Feldman, R. M., Tsomides, T. J., Mayer, J., Narhi, L. O., and Bjorkman, P. J. (1994). Effects of peptide length and composition on binding to an empty class I MHC heterodimer. *Biochemistry* 33, 8149-58.
- Falk, K., Rotzschke, O., and Rammensee, H. G. (1990). Cellular peptide composition governed by major histocompatibility complex class I molecules. *Nature* 348, 248-51.
- Farmery, M. R., Allen, S., Allen, A. J., and Bulleid, N. J. (2000). The role of ERp57 in disulfide bond formation during the assembly of major histocompatibility complex class I in a synchronized semipermeabilized cell translation system. *J Biol Chem* 275, 14933-8.
- Fleischhauer, K., Avila, D., Vilbois, F., Traversari, C., Bordignon, C., and Wallny, H. J. (1994). Characterization of natural peptide ligands for HLA-B*4402 and -B*4403: implications for peptide involvement in allorecognition of a single amino acid

- change in the HLA-B44 heavy chain. *Tissue Antigens* 44, 311-7.
- Frangoulis, B., Park, I., Guillemot, F., Severac, V., Auffray, C., and Zoorob, R. (1999). Identification of the Tapasin gene in the chicken major histocompatibility complex. *Immunogenetics* 49, 328-37.
- Fruh, K., Ahn, K., Djaballah, H., Sempe, P., van Endert, P. M., Tampe, R., Peterson, P. A., and Yang, Y. (1995). A viral inhibitor of peptide transporters for antigen presentation. *Nature* 375, 415-8.
- Fruh, K., Gruhler, A., Krishna, R. M., and Schoenhals, G. J. (1999). A comparison of viral immune escape strategies targeting the MHC class I assembly pathway. *Immunol Rev* 168, 157-66.
- Gakamsky, D. M., Bjorkman, P. J., and Pecht, I. (1996). Peptide interaction with a class I major histocompatibility complex- encoded molecule: allosteric control of the ternary complex stability. *Biochemistry* 35, 14841-8.
- Garbi, N., Tan, P., Diehl, A. D., Chambers, B. J., Ljunggren, H. G., Momburg, F., and Hammerling, G. J. (2000). Impaired immune responses and altered peptide repertoire in tapasin- deficient mice. 1, 234-238.
- Garcia, K. C. (1999). Molecular interactions between extracellular components of the T-cell receptor signaling complex. *Immunol Rev* 172, 73-85.
- Glynne, R., Powis, S. H., Beck, S., Kelly, A., Kerr, L. A., and Trowsdale, J. (1991). A proteasome-related gene between the two ABC transporter loci in the class II region of the human MHC. *Nature* 353, 357-60.
- Goldberg, A. L., and Rock, K. L. (1992). Proteolysis, proteasomes and antigen presentation. *Nature* 357, 375-9.
- Gottesman, M. M., and Pastan, I. (1993). Biochemistry of multidrug resistance mediated by the multidrug transporter. *Annu Rev Biochem* 62, 385-427.
- Grande, A. G., Androlewicz, M. J., Athwal, R. S., Geraghty, D. E., and Spies, T. (1995). Dependence of peptide binding by MHC class I molecules on their interaction with TAP. *Science* 270, 105-8.
- Grande, A. G., Lehner, P. J., Cresswell, P., and Spies, T. (1997). Regulation of MHC class I heterodimer stability and interaction with TAP by tapasin. *Immunogenetics* 46, 477-83.
- Grande, A. G., Golovina, T. N., Hamilton, S. E., Sriram, V., Spies, T., Brutkiewicz, R. R., Harty, J. T., Eisenlohr, L. C., and Van Kaer, L. (2000). Impaired assembly yet normal trafficking of MHC class I molecules in Tapasin mutant mice. *Immunity* 13, 213-22.
- Greenwood, R., Shimizu, Y., Sekhon, G. S., and DeMars, R. (1994). Novel allele-specific, post-translational reduction in HLA class I surface expression in a mutant human B cell line. *J Immunol* 153, 5525-36.
- Groettrup, M., Soza, A., Eggers, M., Kuehn, L., Dick, T. P., Schild, H., Rammensee, H. G., Koszinowski, U. H., and Kloetzel, P. M. (1996). A role for the proteasome regulator PA28alpha in antigen presentation. *Nature* 381, 166-8.
- Gumperz, J. E., Litwin, V., Phillips, J. H., Lanier, L. L., and Parham, P. (1995). The Bw4 public epitope of HLA-B molecules confers reactivity with natural killer cell clones that express NKB1, a putative HLA receptor. *J Exp Med* 181, 1133-44.
- Hauri, H., Appenzeller, C., Kuhn, F., and Nufer, O. (2000). Lectins and traffic in the secretory pathway. *FEBS Lett* 476, 32-7.
- Hebert, D. N., Foellmer, B., and Helenius, A. (1995). Glucose trimming and reglucosylation determine glycoprotein association with calnexin in the endoplasmic reticulum. *Cell* 81, 425-33.
- Hebert, D. N., Foellmer, B., and Helenius, A. (1996). Calnexin and calreticulin promote

- folding, delay oligomerization and suppress degradation of influenza hemagglutinin in microsomes. *EMBO J* 15, 2961-8.
- Hebert, D. N., Zhang, J. X., Chen, W., Foellmer, B., and Helenius, A. (1997). The number and location of glycans on influenza hemagglutinin determine folding and association with calnexin and calreticulin. *J Cell Biol* 139, 613-23.
- Heemels, M. T., and Ploegh, H. L. (1994). Substrate specificity of allelic variants of the TAP peptide transporter. *Immunity* 1, 775-84.
- Hengel, H., Koopmann, J. O., Flohr, T., Muranyi, W., Goulmy, E., Hammerling, G. J., Koszinowski, U. H., and Momburg, F. (1997). A viral ER-resident glycoprotein inactivates the MHC-encoded peptide transporter. *Immunity* 6, 623-32.
- Herberg, J. A., Beck, S., and Trowsdale, J. (1998). TAPASIN, DAXX, RGL2, HKE2 and four new genes (BING 1, 3 to 5) form a dense cluster at the centromeric end of the MHC. *J Mol Biol* 277, 839-57.
- Herberg, J. A., Sgouros, J., Jones, T., Copeman, J., Humphray, S. J., Sheer, D., Cresswell, P., Beck, S., and Trowsdale, J. (1998). Genomic analysis of the Tapasin gene, located close to the TAP loci in the MHC. *Eur J Immunol* 28, 459-67.
- Hiles, I. D., Gallagher, M. P., Jamieson, D. J., and Higgins, C. F. (1987). Molecular characterization of the oligopeptide permease of *Salmonella typhimurium*. *J Mol Biol* 195, 125-42.
- Hill, A., Jugovic, P., York, I., Russ, G., Bennink, J., Yewdell, J., Ploegh, H., and Johnson, D. (1995). Herpes simplex virus turns off the TAP to evade host immunity. *Nature* 375, 411-5.
- Hilt, W., and Wolf, D. H. (1996). Proteasomes: destruction as a programme. *Trends Biochem Sci* 21, 96-102.
- Hirano, N., Shibasaki, F., Sakai, R., Tanaka, T., Nishida, J., Yazaki, Y., Takenawa, T., and Hirai, H. (1995). Molecular cloning of the human glucose-regulated protein ERp57/GRP58, a thiol-dependent reductase. Identification of its secretory form and inducible expression by the oncogenic transformation. *Eur J Biochem* 234, 336-42.
- Hochstenbach, F., David, V., Watkins, S., and Brenner, M. B. (1992). Endoplasmic reticulum resident protein of 90 kilodaltons associates with the T- and B-cell antigen receptors and major histocompatibility complex antigens during their assembly. *Proc Natl Acad Sci U S A* 89, 4734-8.
- Hughes, E. A., and Cresswell, P. (1998). The thiol oxidoreductase ERp57 is a component of the MHC class I peptide-loading complex. *Curr Biol* 8, 709-12.
- Ihara, Y., Cohen-Doyle, M. F., Saito, Y., and Williams, D. B. (1999). Calnexin discriminates between protein conformational states and functions as a molecular chaperone in vitro. *Mol Cell* 4, 331-41.
- Jones, B., and Janeway, C. A., Jr. (1981). Functional activities of antibodies against brain-associated T cell antigens. I. Induction of T cell proliferation. *Eur J Immunol* 11, 584-92.
- Jones, T. R., and Sun, L. (1997). Human cytomegalovirus US2 destabilizes major histocompatibility complex class I heavy chains. *J Virol* 71, 2970-9.
- Kelly, A., Powis, S. H., Glynne, R., Radley, E., Beck, S., and Trowsdale, J. (1991). Second proteasome-related gene in the human MHC class II region. *Nature* 353, 667-8.
- Knittler, M. R., Gulow, K., Seelig, A., and Howard, J. C. (1998). MHC class I molecules compete in the endoplasmic reticulum for access to transporter associated with antigen processing. *J Immunol* 161, 5967-77.

- Koopmann, J. O., Post, M., Neefjes, J. J., Hammerling, G. J., and Momburg, F. (1996). Translocation of long peptides by transporters associated with antigen processing (TAP). *Eur J Immunol* 26, 1720-8.
- Krause, K. H., and Michalak, M. (1997). Calreticulin. *Cell* 88, 439-43.
- Kropshofer, H., Hammerling, G. J., and Vogt, A. B. (1997). How HLA-DM edits the MHC class II peptide repertoire: survival of the fittest? *Immunol Today* 18, 77-82.
- Lanier, L. L. (1998). NK cell receptors. *Annu Rev Immunol* 16, 359-93.
- Lankat-Buttgereit, B., and Tampe, R. (1999). The transporter associated with antigen processing TAP: structure and function. *FEBS Lett* 464, 108-12.
- Lauvau, G., Gubler, B., Cohen, H., Daniel, S., Caillat-Zucman, S., and van Endert, P. M. (1999). Tapasin enhances assembly of transporters associated with antigen processing-dependent and -independent peptides with HLA-A2 and HLA-B27 expressed in insect cells. *J Biol Chem* 274, 31349-58.
- Le Gall, S., Buseyne, F., Trocha, A., Walker, B. D., Heard, J. M., and Schwartz, O. (2000). Distinct trafficking pathways mediate Nef-induced and clathrin-dependent major histocompatibility complex class I down-regulation. *J Virol* 74, 9256-66.
- Lehner, P. J., and Cresswell, P. (1996). Processing and delivery of peptides presented by MHC class I molecules. *Curr Opin Immunol* 8, 59-67.
- Lehner, P. J., Surman, M. J., and Cresswell, P. (1998). Soluble tapasin restores MHC class I expression and function in the tapasin-negative cell line .220. *Immunity* 8, 221-31.
- Lewis, J. W., Neisig, A., Neefjes, J., and Elliott, T. (1996). Point mutations in the alpha 2 domain of HLA-A2.1 define a functionally relevant interaction with TAP. *Curr Biol* 6, 873-83.
- Lewis, J. W., Sewell, A., Price, D., and Elliott, T. (1998). HLA-A*0201 presents TAP-dependent peptide epitopes to cytotoxic T lymphocytes in the absence of tapasin. *Eur J Immunol* 28, 3214-20.
- Lindquist, J. A., Jensen, O. N., Mann, M., and Hammerling, G. J. (1998). ER-60, a chaperone with thiol-dependent reductase activity involved in MHC class I assembly. *EMBO J* 17, 2186-95.
- Ljunggren, H. G., Stam, N. J., Ohlen, C., Neefjes, J. J., Hoglund, P., Heemels, M. T., Bastin, J., Schumacher, T. N., Townsend, A., Karre, K., and et al. (1990). Empty MHC class I molecules come out in the cold. *Nature* 346, 476-80.
- Luckey, C. J., King, G. M., Marto, J. A., Venketeswaran, S., Maier, B. F., Crotzer, V. L., Colella, T. A., Shabanowitz, J., Hunt, D. F., and Engelhard, V. H. (1998). Proteasomes can either generate or destroy MHC class I epitopes: evidence for nonproteasomal epitope generation in the cytosol. *J Immunol* 161, 112-21.
- Lutz, C. T., Smith, K. D., Greazel, N. S., Mace, B. E., Jensen, D. A., McCutcheon, J. A., and Goeken, N. E. (1994). Bw4-reactive and Bw6-reactive antibodies recognize multiple distinct HLA structures that partially overlap in the alpha-1 helix. *J Immunol* 153, 4099-110.
- Momburg, F., Ortiz-Navarrete, V., Neefjes, J., Goulmy, E., van de Wal, Y., Spits, H., Powis, S. J., Butcher, G. W., Howard, J. C., Walden, P., and et al. (1992). Proteasome subunits encoded by the major histocompatibility complex are not essential for antigen presentation. *Nature* 360, 174-7.
- Momburg, F., Roelse, J., Hammerling, G. J., and Neefjes, J. J. (1994a). Peptide size selection by the major histocompatibility complex-encoded peptide transporter. *J Exp Med* 179, 1613-23.
- Momburg, F., Roelse, J., Howard, J. C., Butcher, G. W., Hammerling, G. J., and

- Neefjes, J. J. (1994b). Selectivity of MHC-encoded peptide transporters from human, mouse and rat. *Nature* 367, 648-51.
- Momburg, F., Armandola, E. A., Post, M., and Hammerling, G. J. (1996). Residues in TAP2 peptide transporters controlling substrate specificity. *J Immunol* 156, 1756-63.
- Morel, S., Levy, F., Burlet-Schiltz, O., Brasseur, F., Probst-Kepper, M., Peitrequin, A. L., Monsarrat, B., Van Velthoven, R., Cerottini, J. C., Boon, T., Gairin, J. E., and Van den Eynde, B. J. (2000). Processing of some antigens by the standard proteasome but not by the immunoproteasome results in poor presentation by dendritic cells. *Immunity* 12, 107-17.
- Morrice, N. A., and Powis, S. J. (1998). A role for the thiol-dependent reductase ERp57 in the assembly of MHC class I molecules. *Curr Biol* 8, 713-6.
- Muller, C., Herbst, H., Uchanska-Ziegler, B., Ziegler, A., Schunter, F., Steiert, I., and Wernet, P. (1985). Characterization of a monoclonal anti-Bw4 antibody (Tu109): evidence for similar epitopes on the Bw4 and Bw6 antigens. *Hum Immunol* 14, 333-49.
- Muller, C. A., Engler-Blum, G., Gekeler, V., Steiert, I., Weiss, E., and Schmidt, H. (1989). Genetic and serological heterogeneity of the supertypic HLA-B locus specificities Bw4 and Bw6. *Immunogenetics* 30, 200-7.
- Muller, K. M., Ebensperger, C., and Tampe, R. (1994). Nucleotide binding to the hydrophilic C-terminal domain of the transporter associated with antigen processing (TAP). *J Biol Chem* 269, 14032-7.
- Neefjes, J. J., and Ploegh, H. L. (1988). Allele and locus-specific differences in cell surface expression and the association of HLA class I heavy chain with beta 2-microglobulin: differential effects of inhibition of glycosylation on class I subunit association. *Eur J Immunol* 18, 801-10.
- Neefjes, J. J., Smit, L., Gehrman, M., and Ploegh, H. L. (1992). The fate of the three subunits of major histocompatibility complex class I molecules. *Eur J Immunol* 22, 1609-14.
- Neefjes, J. J., Hammerling, G. J., and Momburg, F. (1993a). Folding and assembly of major histocompatibility complex class I heterodimers in the endoplasmic reticulum of intact cells precedes the binding of peptide. *J Exp Med* 178, 1971-80.
- Neefjes, J. J., Momburg, F., and Hammerling, G. J. (1993b). Selective and ATP-dependent translocation of peptides by the MHC- encoded transporter. *Science* 261, 769-71.
- Neefjes, J. J., Dierx, J., and Ploegh, H. L. (1993c). The effect of anchor residue modifications on the stability of major histocompatibility complex class I-peptide interactions. *Eur J Immunol* 23, 840-5.
- Neefjes, J., Gottfried, E., Roelse, J., Gromme, M., Obst, R., Hammerling, G. J., and Momburg, F. (1995). Analysis of the fine specificity of rat, mouse and human TAP peptide transporters. *Eur J Immunol* 25, 1133-6.
- Neisig, A., Roelse, J., Sijts, A. J., Ossendorp, F., Feltkamp, M. C., Kast, W. M., Melief, C. J., and Neefjes, J. J. (1995). Major differences in transporter associated with antigen presentation (TAP)-dependent translocation of MHC class I-presentable peptides and the effect of flanking sequences. *J Immunol* 154, 1273-9.
- Neisig, A., Melief, C. J., and Neefjes, J. (1998). Reduced cell surface expression of HLA-C molecules correlates with restricted peptide binding and stable TAP interaction. *J Immunol* 160, 171-9.
- Nickerson, C. L., Hogen, K. L., Luthra, H. S., and David, C. S. (1990). Effect of H-2

- genes on expression of HLA-B27 and Yersinia-induced arthritis. *Scand J Rheumatol Suppl* 87, 85-90.
- Niedermann, G., Geier, E., Lucchiari-Hartz, M., Hitziger, N., Ramsperger, A., and Eichmann, K. (1999). The specificity of proteasomes: impact on MHC class I processing and presentation of antigens. *Immunol Rev* 172, 29-48.
- Nijenhuis, M., and Hammerling, G. J. (1996). Multiple regions of the transporter associated with antigen processing (TAP) contribute to its peptide binding site. *J Immunol* 157, 5467-77.
- Nijenhuis, M., Schmitt, S., Armandola, E. A., Obst, R., Brunner, J., and Hammerling, G. J. (1996). Identification of a contact region for peptide on the TAP1 chain of the transporter associated with antigen processing. *J Immunol* 156, 2186-95.
- Nilsson, T., and Warren, G. (1994). Retention and retrieval in the endoplasmic reticulum and the Golgi apparatus. *Curr Opin Cell Biol* 6, 517-21.
- Nossal, Sir G. J. V (1999). Das Immunsystem. Spektrum der Wissenschaft Spezial 1999, S.9
- Nossner, E., and Parham, P. (1995). Species-specific differences in chaperone interaction of human and mouse major histocompatibility complex class I molecules. *J Exp Med* 181, 327-37.
- Nuchtern, J. G., Bonifacino, J. S., Biddison, W. E., and Klausner, R. D. (1989). Brefeldin A implicates egress from endoplasmic reticulum in class I restricted antigen presentation. *Nature* 339, 223-6.
- Oliver, J. D., Roderick, H. L., Llewellyn, D. H., and High, S. (1999). ERp57 functions as a subunit of specific complexes formed with the ER lectins calreticulin and calnexin. *Mol Biol Cell* 10, 2573-82.
- Ortiz-Navarrete, V., Seelig, A., Gernold, M., Frentzel, S., Kloetzel, P. M., and Hammerling, G. J. (1991). Subunit of the '20S' proteasome (multicatalytic proteinase) encoded by the major histocompatibility complex. *Nature* 353, 662-4.
- Ortmann, B., Androlewicz, M. J., and Cresswell, P. (1994). MHC class I/beta 2-microglobulin complexes associate with TAP transporters before peptide binding. *Nature* 368, 864-7.
- Ortmann, B., Copeman, J., Lehner, P. J., Sadasivan, B., Herberg, J. A., Grandea, A. G., Riddell, S. R., Tampe, R., Spies, T., Trowsdale, J., and Cresswell, P. (1997). A critical role for tapasin in the assembly and function of multimeric MHC class I-TAP complexes. *Science* 277, 1306-9.
- Otteken, A., and Moss, B. (1996). Calreticulin interacts with newly synthesized human immunodeficiency virus type 1 envelope glycoprotein, suggesting a chaperone function similar to that of calnexin. *J Biol Chem* 271, 97-103.
- Ou, W. J., Cameron, P. H., Thomas, D. Y., and Bergeron, J. J. (1993). Association of folding intermediates of glycoproteins with calnexin during protein maturation. *Nature* 364, 771-6.
- Owen, B. A., and Pease, L. R. (1999). TAP association influences the conformation of nascent MHC class I molecules. *J Immunol* 162, 4677-84.
- Paabo, S., Bhat, B. M., Wold, W. S., and Peterson, P. A. (1987). A short sequence in the COOH-terminus makes an adenovirus membrane glycoprotein a resident of the endoplasmic reticulum. *Cell* 50, 311-7.
- Parker, K. C., DiBrino, M., Hull, L., and Coligan, J. E. (1992). The beta 2-microglobulin dissociation rate is an accurate measure of the stability of MHC class I heterotrimers and depends on which peptide is bound. *J Immunol* 149, 1896-904.
- Paul, W. E. (1999). Fundamental Immunology I. 4th edition. Lippincott-Raven 1999.
- Paz, P., Brouwenstijn, N., Perry, R., and Shastri, N. (1999). Discrete proteolytic

- intermediates in the MHC class I antigen processing pathway and MHC I-dependent peptide trimming in the ER. *Immunity* 11, 241-51.
- Peace-Brewer, A. L., Tussey, L. G., Matsui, M., Li, G., Quinn, D. G., and Frelinger, J. A. (1996). A point mutation in HLA-A*0201 results in failure to bind the TAP complex and to present virus-derived peptides to CTL. *Immunity* 4, 505-14.
- Peh, C. A., Burrows, S. R., Barnden, M., Khanna, R., Cresswell, P., Moss, D. J., and McCluskey, J. (1998). HLA-B27-restricted antigen presentation in the absence of tapasin reveals polymorphism in mechanisms of HLA class I peptide loading. *Immunity* 8, 531-42.
- Peh, C. A., Laham, N., Burrows, S. R., Zhu, Y., and McCluskey, J. (2000). Distinct functions of tapasin revealed by polymorphism in MHC class I peptide loading. *J Immunol* 164, 292-9.
- Perarnau, B., Siegrist, C. A., Gillet, A., Vincent, C., Kimura, S., and Lemonnier, F. A. (1990). Beta 2-microglobulin restriction of antigen presentation. *Nature* 346, 751-4.
- Peruzzi, M., Wagtmann, N., and Long, E. O. (1996). A p70 killer cell inhibitory receptor specific for several HLA-B allotypes discriminates among peptides bound to HLA-B*2705. *J Exp Med* 184, 1585-90.
- Peterson, J. R., Ora, A., Van, P. N., and Helenius, A. (1995). Transient, lectin-like association of calreticulin with folding intermediates of cellular and viral glycoproteins. *Mol Biol Cell* 6, 1173-84.
- Plempner, R. K., Bohmler, S., Bordallo, J., Sommer, T., and Wolf, D. H. (1997). Mutant analysis links the translocon and BiP to retrograde protein transport for ER degradation. *Nature* 388, 891-5.
- Ploegh, H. L. (1997). Destruction of proteins as a creative force. *Immunol Today* 18, 269-71.
- Ploegh, H. L. (1998). Viral strategies of immune evasion. *Science* 280, 248-53.
- Powis, S. J., Townsend, A. R., Deverson, E. V., Bastin, J., Butcher, G. W., and Howard, J. C. (1991). Restoration of antigen presentation to the mutant cell line RMA-S by an MHC-linked transporter. *Nature* 354, 528-31.
- Powis, S. J., Young, L. L., Joly, E., Barker, P. J., Richardson, L., Brandt, R. P., Melief, C. J., Howard, J. C., and Butcher, G. W. (1996). The rat cim effect: TAP allele-dependent changes in a class I MHC anchor motif and evidence against C-terminal trimming of peptides in the ER. *Immunity* 4, 159-65.
- Purcell, A. W., Gorman, J. J., Garcia-Peydro, M., Paradela, A., Burrows, S. R., Talbo, G. H., Laham, N., Peh, C. A., Reynolds, E. C., Lopez De Castro, J. A., and McCluskey, J. (2001). Quantitative and qualitative influences of tapasin on the class I peptide repertoire. *J Immunol* 166, 1016-27.
- Radka, S. F., Kostyu, D. D., and Amos, D. B. (1982). A monoclonal antibody directed against the HLA-Bw6 epitope. *J Immunol* 128, 2804-6.
- Rammensee, H. G. (1995). Chemistry of peptides associated with MHC class I and class II molecules. *Curr Opin Immunol* 7, 85-96.
- Rammensee, H., Bachmann, J., Emmerich, N. P., Bachor, O. A., and Stevanovic, S. (1999). SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 50, 213-9.
- Reits, E. A., Griekspoor, A. C., and Neefjes, J. (2000). How does TAP pump peptides? insights from DNA repair and traffic ATPases. *Immunol Today* 21, 598-600.
- Riordan, J. R., Rommens, J. M., Kerem, B., Alon, N., Rozmahel, R., Grzelczak, Z., Zielenski, J., Lok, S., Plavsic, N., Chou, J. L., and et al. (1989). Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA.

- Science* 245, 1066-73.
- Rodan, A. R., Simons, J. F., Trombetta, E. S., and Helenius, A. (1996). N-linked oligosaccharides are necessary and sufficient for association of glycosylated forms of bovine RNase with calnexin and calreticulin. *EMBO J* 15, 6921-30.
- Roelse, J., Gromme, M., Momburg, F., Hammerling, G., and Neefjes, J. (1994). Trimming of TAP-translocated peptides in the endoplasmic reticulum and in the cytosol during recycling. *J Exp Med* 180, 1591-7.
- Rohren, E. M., McCormick, D. J., and Pease, L. R. (1994). Peptide-induced conformational changes in class I molecules. Direct detection by flow cytometry. *J Immunol* 152, 5337-43.
- Rotem-Yehudar, R., Winograd, S., Sela, S., Coligan, J. E., and Ehrlich, R. (1994). Downregulation of peptide transporter genes in cell lines transformed with the highly oncogenic adenovirus 12. *J Exp Med* 180, 477-88.
- Rotem-Yehudar, R., Shechter, H., and Ehrlich, R. (1994). Transcriptional regulation of class-I major histocompatibility complex genes transformed in murine cells is mediated by positive and negative regulatory elements. *Gene* 144, 265-70.
- Russ, G., Esquivel, F., Yewdell, J. W., Cresswell, P., Spies, T., and Bennink, J. R. (1995). Assembly, intracellular localization, and nucleotide binding properties of the human peptide transporters TAP1 and TAP2 expressed by recombinant vaccinia viruses. *J Biol Chem* 270, 21312-8.
- Sadasivan, B., Lehner, P. J., Ortmann, B., Spies, T., and Cresswell, P. (1996). Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* 5, 103-14.
- Saito, Y., Ihara, Y., Leach, M. R., Cohen-Doyle, M. F., and Williams, D. B. (1999). Calreticulin functions in vitro as a molecular chaperone for both glycosylated and non-glycosylated proteins. *EMBO J* 18, 6718-29.
- Salter, R. D., Howell, D. N., and Cresswell, P. (1985). Genes regulating HLA class I antigen expression in T-B lymphoblast hybrids. *Immunogenetics* 21, 235-46.
- Scaffidi, C., Kirchhoff, S., Krammer, P. H., and Peter, M. E. (1999). Apoptosis signaling in lymphocytes. *Curr Opin Immunol* 11, 277-85.
- Schoenhals, G. J., Krishna, R. M., Grandea, A. G., Spies, T., Peterson, P. A., Yang, Y., and Fruh, K. (1999). Retention of empty MHC class I molecules by tapasin is essential to reconstitute antigen presentation in invertebrate cells. *EMBO J* 18, 743-53.
- Schumacher, T. N., Kantesaria, D. V., Heemels, M. T., Ashton-Rickardt, P. G., Shepherd, J. C., Fruh, K., Yang, Y., Peterson, P. A., Tonegawa, S., and Ploegh, H. L. (1994). Peptide length and sequence specificity of the mouse TAP1/TAP2 translocator. *J Exp Med* 179, 533-40.
- Schiebert, E. M. (1999). ABC transporter-facilitated ATP conductive transport. *Am J Physiol* 276, C1-8.
- Scott, J. E., and Dawson, J. R. (1995). MHC class I expression and transport in a calnexin-deficient cell line. *J Immunol* 155, 143-8.
- Seliger, B., Wollscheid, U., Momburg, F., Blankenstein, T., and Huber, C. (2000). Coordinate downregulation of multiple MHC class I antigen processing genes in chemical-induced murine tumor cell lines of distinct origin. *Tissue Antigens* 56, 327-36.
- Seliger, B., Maeurer, M. J., and Ferrone, S. (2000). Antigen-processing machinery breakdown and tumor growth. *Immunol Today* 21, 455-64.
- Seliger, B., Wollscheid, U., Momburg, F., Blankenstein, T., and Huber, C. (2001). Characterization of the major histocompatibility complex class I deficiencies in

- B16 melanoma cells. *Cancer Res* 61, 1095-9.
- Seliger, B., Schreiber, K., Delp, K., Meissner, M., Hammers, S., Reichert, T., Pawlischko, K., Tampe, R., and Huber, C. (2001). Downregulation of the constitutive tapasin expression in human tumor cells of distinct origin and its transcriptional upregulation by cytokines. *Tissue Antigens* 57, 39-45.
- Shields, M. J., Moffat, L. E., and Ribaldo, R. K. (1998). Functional comparison of bovine, murine, and human beta2-microglobulin: interactions with murine MHC I molecules. *Mol Immunol* 35, 919-28.
- Shields, M. J., Hodgson, W., and Ribaldo, R. K. (1999). Differential association of beta2-microglobulin mutants with MHC class I heavy chains and structural analysis demonstrate allele-specific interactions. *Mol Immunol* 36, 561-73.
- Shimizu, Y., and DeMars, R. (1989). Production of human cells expressing individual transferred HLA-A,-B,-C genes using an HLA-A,-B,-C null human cell line. *J Immunol* 142, 3320-8.
- Shresta, S., Graubert, T. A., Thomas, D. A., Raptis, S. Z., and Ley, T. J. (1999). Granzyme A initiates an alternative pathway for granule-mediated apoptosis. *Immunity* 10, 595-605.
- Sijts, A. J., and Pamer, E. G. (1997). Enhanced intracellular dissociation of major histocompatibility complex class I-associated peptides: a mechanism for optimizing the spectrum of cell surface-presented cytotoxic T lymphocyte epitopes. *J Exp Med* 185, 1403-11.
- Smith, J. D., Myers, N. B., Gorka, J., and Hansen, T. H. (1993). Model for the in vivo assembly of nascent Ld class I molecules and for the expression of unfolded Ld molecules at the cell surface. *J Exp Med* 178, 2035-46.
- Smith, J. D., Solheim, J. C., Carreno, B. M., and Hansen, T. H. (1995). Characterization of class I MHC folding intermediates and their disparate interactions with peptide and beta 2-microglobulin. *Mol Immunol* 32, 531-40.
- Smith, K. D., Kurago, Z. B., and Lutz, C. T. (1997). Conformational changes in MHC class I molecules. Antibody, T-cell receptor, and NK cell recognition in an HLA-B7 model system. *Immunol Res* 16, 243-59.
- Solheim, J. C., Harris, M. R., Kindle, C. S., and Hansen, T. H. (1997). Prominence of beta 2-microglobulin, class I heavy chain conformation, and tapasin in the interactions of class I heavy chain with calreticulin and the transporter associated with antigen processing. *J Immunol* 158, 2236-41.
- Solheim, J. C. (1999). Class I MHC molecules: assembly and antigen presentation. *Immunol Rev* 172, 11-9.
- Spies, T., Bresnahan, M., Bahram, S., Arnold, D., Blanck, G., Mellins, E., Pious, D., and DeMars, R. (1990). A gene in the human major histocompatibility complex class II region controlling the class I antigen presentation pathway. *Nature* 348, 744-7.
- Springer, S., Doring, K., Skipper, J. C., Townsend, A. R., and Cerundolo, V. (1998). Fast association rates suggest a conformational change in the MHC class I molecule H-2Db upon peptide binding. *Biochemistry* 37, 3001-12.
- Srivastava, P. K., Udono, H., Blachere, N. E., and Li, Z. (1994). Heat shock proteins transfer peptides during antigen processing and CTL priming. *Immunogenetics* 39, 93-8.
- Stam, N. J., Spits, H., and Ploegh, H. L. (1986). Monoclonal antibodies raised against denatured HLA-B locus heavy chains permit biochemical characterization of certain HLA-C locus products. *J Immunol* 137, 2299-306.
- Stam, N. J., Vroom, T. M., Peters, P. J., Pastoors, E. B., and Ploegh, H. L. (1990). HLA-A- and HLA-B-specific monoclonal antibodies reactive with free heavy chains in

- western blots, in formalin-fixed, paraffin-embedded tissue sections and in cryo-immuno-electron microscopy. *Int Immunol* 2, 113-25.
- Suh, W. K., Cohen-Doyle, M. F., Fruh, K., Wang, K., Peterson, P. A., and Williams, D. B. (1994). Interaction of MHC class I molecules with the transporter associated with antigen processing. *Science* 264, 1322-6.
- Suh, W. K., Mitchell, E. K., Yang, Y., Peterson, P. A., Waneck, G. L., and Williams, D. B. (1996). MHC class I molecules form ternary complexes with calnexin and TAP and undergo peptide-regulated interaction with TAP via their extracellular domains. *J Exp Med* 184, 337-48.
- Suh, W. K., Derby, M. A., Cohen-Doyle, M. F., Schoenhals, G. J., Fruh, K., Berzofsky, J. A., and Williams, D. B. (1999). Interaction of murine MHC class I molecules with tapasin and TAP enhances peptide loading and involves the heavy chain alpha3 domain. *J Immunol* 162, 1530-40.
- Tahara, T., Yang, S. Y., Khan, R., Abish, S., Hammerling, G. J., and Hammerling, U. (1990). HLA antibody responses in HLA class I transgenic mice. *Immunogenetics* 32, 351-60.
- Takamiya, Y., Sakaguchi, T., Miwa, K., and Takiguchi, M. (1996). Role of HLA-B*5101 binding nonamer peptides in formation of the HLA-Bw4 public epitope. *Int Immunol* 8, 1027-34.
- Tomazin, R., Hill, A. B., Jugovic, P., York, I., van Endert, P., Ploegh, H. L., Andrews, D. W., and Johnson, D. C. (1996). Stable binding of the herpes simplex virus ICP47 protein to the peptide binding site of TAP. *EMBO J* 15, 3256-66.
- Tormo, J., Natarajan, K., Margulies, D. H., and Mariuzza, R. A. (1999). Crystal structure of a lectin-like natural killer cell receptor bound to its MHC class I ligand. *Nature* 402, 623-31.
- Trowsdale, J., Hanson, I., Mockridge, I., Beck, S., Townsend, A., and Kelly, A. (1990). Sequences encoded in the class II region of the MHC related to the 'ABC' superfamily of transporters. *Nature* 348, 741-4.
- Uger, R. A., Chan, S. M., and Barber, B. H. (1999). Covalent linkage to beta2-microglobulin enhances the MHC stability and antigenicity of suboptimal CTL epitopes. *J Immunol* 162, 6024-8.
- Urade, R., and Kito, M. (1992). Inhibition by acidic phospholipids of protein degradation by ER-60 protease, a novel cysteine protease, of endoplasmic reticulum. *FEBS Lett* 312, 83-6.
- Urban, R. G., Chicz, R. M., Lane, W. S., Strominger, J. L., Rehm, A., Kenter, M. J., UytdeHaag, F. G., Ploegh, H., Uchanska-Ziegler, B., and Ziegler, A. (1994). A subset of HLA-B27 molecules contains peptides much longer than nonamers. *Proc Natl Acad Sci U S A* 91, 1534-8.
- Vales-Gomez, M., Reyburn, H., and Strominger, J. (2000). Molecular analyses of the interactions between human NK receptors and their HLA ligands. *Hum Immunol* 61, 28-38.
- van Endert, P. M., Tampe, R., Meyer, T. H., Tisch, R., Bach, J. F., and McDevitt, H. O. (1994). A sequential model for peptide binding and transport by the transporters associated with antigen processing. *Immunity* 1, 491-500.
- Van Kaer, L., Ashton-Rickardt, P. G., Ploegh, H. L., and Tonegawa, S. (1992). TAP1 mutant mice are deficient in antigen presentation, surface class I molecules, and CD4-8+ T cells. *Cell* 71, 1205-14.
- Vassilakos, A., Cohen-Doyle, M. F., Peterson, P. A., Jackson, M. R., and Williams, D. B. (1996). The molecular chaperone calnexin facilitates folding and assembly of class I histocompatibility molecules. *EMBO J* 15, 1495-506.

- Vassilakos, A., Michalak, M., Lehrman, M. A., and Williams, D. B. (1998). Oligosaccharide binding characteristics of the molecular chaperones calnexin and calreticulin. *Biochemistry* 37, 3480-90.
- Vogt, A. B., and Kropshofer, H. (1999). HLA-DM - an endosomal and lysosomal chaperone for the immune system. *Trends Biochem Sci* 24, 150-4.
- Vos, J. C., Spee, P., Momburg, F., and Neefjes, J. (1999). Membrane topology and dimerization of the two subunits of the transporter associated with antigen processing reveal a three-domain structure. *J Immunol* 163, 6679-85.
- Wada, I., Imai, S., Kai, M., Sakane, F., and Kanoh, H. (1995). Chaperone function of calreticulin when expressed in the endoplasmic reticulum as the membrane-anchored and soluble forms. *J Biol Chem* 270, 20298-304.
- Wang, K., Fruh, K., Peterson, P. A., and Yang, Y. (1994). Nucleotide binding of the C-terminal domains of the major histocompatibility complex-encoded transporter expressed in *Drosophila melanogaster* cells. *FEBS Lett* 350, 337-41.
- Ware, F. E., Vassilakos, A., Peterson, P. A., Jackson, M. R., Lehrman, M. A., and Williams, D. B. (1995). The molecular chaperone calnexin binds Glc1Man9GlcNAc2 oligosaccharide as an initial step in recognizing unfolded glycoproteins. *J Biol Chem* 270, 4697-704.
- York, I. A., Goldberg, A. L., Mo, X. Y., and Rock, K. L. (1999). Proteolysis and class I major histocompatibility complex antigen presentation. *Immunol Rev* 172, 49-66.
- Yu, Y. Y., Turnquist, H. R., Myers, N. B., Balendiran, G. K., Hansen, T. H., and Solheim, J. C. (1999). An extensive region of an MHC class I alpha 2 domain loop influences interaction with the assembly complex. *J Immunol* 163, 4427-33.
- Zapun, A., Darby, N. J., Tessier, D. C., Michalak, M., Bergeron, J. J., and Thomas, D. Y. (1998). Enhanced catalysis of ribonuclease B folding by the interaction of calnexin or calreticulin with ERp57. *J Biol Chem* 273, 6009-12.
- Zhang, Q. J., Gavioli, R., Klein, G., and Masucci, M. G. (1993). An HLA-A11-specific motif in nonamer peptides derived from viral and cellular proteins. *Proc Natl Acad Sci U S A* 90, 2217-21.
- Zhang, Q., Tector, M., and Salter, R. D. (1995). Calnexin recognizes carbohydrate and protein determinants of class I major histocompatibility complex molecules. *J Biol Chem* 270, 3944-8.
- Zhang, J. X., Braakman, I., Matlack, K. E., and Helenius, A. (1997). Quality control in the secretory pathway: the role of calreticulin, calnexin and BiP in the retention of glycoproteins with C-terminal truncations. *Mol Biol Cell* 8, 1943-54.
- Zhang, Q., and Salter, R. D. (1998). Distinct patterns of folding and interactions with calnexin and calreticulin in human class I MHC proteins with altered N-glycosylation. *J Immunol* 160, 831-7.
- Ziegler, A., Uchanska-Ziegler, B., Zeuthen, J., and Wernet, P. (1982). HLA antigen expression at the single cell level on a K562 X B cell hybrid: an analysis with monoclonal antibodies using bacterial binding assays. *Somatic Cell Genet* 8, 775-89.
- Zijlstra, M., Bix, M., Simister, N. E., Loring, J. M., Raulet, D. H., and Jaenisch, R. (1990). Beta 2-microglobulin deficient mice lack CD4-8+ cytolytic T cells. *Nature* 344, 742-6.

8.2 Abkürzungsverzeichnis

β_2 m:	β_2 -Mikroglobulin
Δ C:	Carboxyterminale Verkürzung
Δ N:	Aminoternale Verkürzung
ABC:	ATP-Bindungskassette
APC:	Antigen präsentierende Zelle
BZR:	B-Zellrezeptor
BiP:	Binding protein
B-LCL:	Lymphatische B-Zelllinie
BSA:	Rinderserumalbumin
CD:	Clusters of differentiation
CXN:	Calnexin
CRT:	Calretikulin
CTL:	Zytotoxischer T-Lymphozyt
DMSO:	Dimethylsulfoxid
DNA:	Desoxyribonukleinsäure
DTT:	Dithiothreitol
EDTA:	Ethylendiamintetraessigsäure
ER:	Endoplasmatisches Retikulum
ER60:	ER-Protein mit einem Molekulargewicht von etwa 60kD
FACS:	Fluoreszenz aktivierte Zellsortierung
FCS:	Fötale Kälberserum
FL:	voller Länge
HLA:	Humaner Histokompatibilitätskomplex
Hsp:	Hitze-Schock Protein
HSV:	Herpes simplex Virus
hTpn:	humanes Tapasin
Ig:	Immunoglobulin
IL:	Interleukin
INF:	Interferon
LLnL:	N-Acety-L-leucyl-L-leucyl-Norleucinal
LMP:	Low molecular mass proteins
MCF:	Mittlerer Fluoreszenzkanal
MHC:	Haupthistokompatibilitätskomplex
mTpn:	Maus-Tapasin voller Länge
NBD:	Nukleotidbindungsdomäne
NP40:	Nonidet-P40
NK Zelle:	Natürliche Killerzelle
PBS:	Phosphate-gepufferte Salzlösung
PDI:	Protein disulfide isomerase
SDS:	Natriumdodecylsulfat
TAP:	Transporter associated with antigen processing
Tpn:	Tapasin (von "Tapas Inn" - mexikanisches Restaurant)
TZR:	T-Zell Receptor

8.3 Danksagungen

Zunächst möchte ich PD Dr. Frank Momburg für die Betreuung meiner Dissertation und für die Art und Weise seiner Anleitung danken, die mir ein unbedrängtes Arbeiten ermöglicht hat. Insbesondere danke ich ihm dafür, daß ich die Unterstützung von TAs genießen konnte. Wegen seiner erfolgreichen Bemühungen, aus aller Herren Länder Reagenzien für die Laborarbeit zu beschaffen, hat es mir zudem nie an notwendigen Arbeitsmitteln gefehlt. Beindruckt hat mich seine Präzision und Sorgfalt, von der sich manch' einer eine Scheibe abschneiden könnte...

Prof. Dr. Günter J. Hämmerling danke ich für die Bereitstellung aller notwendigen Arbeitsmittel und möchte insbesondere für die Art der Zusammenarbeit im Tapasin-Maus-Projekt loben.

Mein herzlicher Dank gilt allen, mir bei vielen Experimenten mit Hand und Fuß hilfreich gewesen, netten Menschen: DANKE besonders Nadja Bulbuc sowie Ulrike Rothermel, Michelle L. Harris, Markus Post und Beate Plischka.

Thanks to Dr. immunol. aqua. nat. Mac. Garcia for the collaboration in the Tapasin mice project and helpfull discussions (alla guud!).

Dr. Gerd Moldenhauer gilt mein Dank für viele tausend Antikörper und andere Reagenzien, die er mir stets mit seiner großzügigen Art überlassen hat.

Dr. Harald Kropshofer möchte ich sehr für die Erstellung der Peptidprofile danken. Nadine Anders danke ich für die lustige und erfolgreiche Zusammenarbeit bei der Peptidisolierung.

Klaus Hexel danke ich für die erfolgreichen Zellsorts und die kurpfälzische Laune, mit der er mich, trotz häufigen Knuffer auf die Arme, erfreut hat.

Herrn Pipkorn danke ich für die Bereitstellung des Massenspektrometers und die Herstellung mehrerer Peptide sowie Dr. J. McCluskey für die Überlassung der .220.B44-Zellen, Dr. T. Elliott für die .220.B8-Zellen, Dr. T. Hansen für das Antiserum gegen Maus-Tapasin und Dr. J. Neefjes für das Antiserum gegen TAP1.

Jörn Albring danke ich ganz besonders für die netten Späße, und bei Eva Hüter bedanke ich mich, weil sie sich von meinem ersten, strengen Rat nicht hat abschrecken lassen, mir weiterhin Fragen zu stellen ;)

Für das sehr angenehme Arbeitsklima bin ich allen tapferen Mensaessern und den vernünftigen Menschen der Abteilung äußerst dankbar.

Allen, die dabei nicht angesprochen worden sind, bei unserer Weihnachtsfeier aber unermüdlich mitgearbeitet und gelacht haben, danke ich herzlich (Hi, Roland)!

Meinem Bruder Fabian danke für alle Hinweise zur Behebung von Schwierigkeiten am Computer und meinen Eltern möchte zu guter letzt für die geistige Unterstützung mit vielen aufmunternden Worten und den schönen Stunden der Ablenkung in Berlin herzlich danken.