

V LITERATURVERZEICHNIS

- Abate, C.; Smith, J.A.; Joh, T.H. (1988) Characterization of the catalytic domain of bovine adrenal tyrosine hydroxylase. *Biochem. Biophys. Res. Commun.* **151**: 1446-1453.
- Akiskal, H.S.; McKinney, W.T. Jr. (1973) Depressive disorders: toward a unified hypothesis. *Science* **182**(107): 20-29.
- Albert, P.R.; Zhou, Q.Y.; Van Tol, H.H.; Bunzow, J.R.; Civelli, O. (1990) Cloning, functional expression and mRNA tissue distribution of the rat 5-HT1A receptor gene. *J. Biol. Chem.* **265**: 5825-5832.
- Albinsson, A.; Andersson, G.; Andersson, K.; Vega-Matuszczyk, J.; Larsson, K. (1996) The effects of lesions in the mesencephalic raphe systems on male rat sexual behavior and locomotor activity. *Behav. Brain Res.* **80**(1-2): 57-63.
- Antithrombotic Trialists' Collaboration. (2002) Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ.* **324**(7329): 71-86.
- Artigas, F.; Sunol, C.; Tusell, J.M.; Martinez, E.; Gelpi, E. (1985) Comparative ontogenesis of brain tryptamine, serotonin, and tryptophan. *J. Neurochem.* **44**: 31-37.
- Aune, T.M.; McGarth, K.M.; Sarr, T.; Bombara, M.P.; Kelley, K.A. (1993) Expression of 5HT1a receptors on activated human T cells. Regulation of cyclic AMP levels and T cell proliferation by 5-hydroxytryptamine. *J. Immunol.* **151**: 1175-1183.
- Austin, M.C.; O'Donnell, S.M. (1999) Regional distribution and cellular expression of tryptophan hydroxylase messenger RNA in postmortem human brainstem and pineal gland. *J. Neurochem.* **72**: 2065-2073.
- Azmitia, E.C. (2000) Serotonin. *Nat. Encycl. Life Sci.* <http://www.els.net/>
- Azmitia, E.C. (2001) Modern views on an ancient chemical: serotonin effects on cell proliferation, maturation, and apoptosis. *Brain Res. Bull.* **56**(5): 413-424.
- Badawy, A.A. (1999) Tryptophan metabolism in alcoholism. *Adv. Exp. Med. Biol.* **467**: 265-274.
- Baigent, C. (2005) Aspirin for everyone older than 50? Against. *BMJ.* **330**(7505): 1442-1443.
- Baker G.B., Martin I.L. and Mtichel P.R., (1977) The effects of some indolalkylamines on the uptake and release of 5-hydroxytryptamine in rat striatum [proceedings]. *Br. J. Pharmacol.* **61**: 151-152.
- Banik, U.; Wang, G.; Wagner, P.D.; Kaufman, S. (1997) Interaction of phosphorylated tryptophan hydroxylase with 14-3-3 protein. *J. Biol. Chem.* **42**: 219-226.

- Barbosa, R.M.; Silva, A.M.; Tome, A.R.; Stamford, J.A.; Santos, R.M.; Rosario, L.M. (1998) Control of pulsatile 5-HT/insulin secretion from single mouse pancreatic islets by intracellular calcium dynamics. *J. Physiol.* **510** (Pt 1): 135-143.
- Barden, N.; Reul, J.M.; Holsboer, F. (1995) Do antidepressants stabilize mood through actions on the hypothalamic-pituitary-adrenocortical system? *Trends Neurosci.* **18**(1): 6-11.
- Baumgarten, H.G.; Klemm, H.P.; Sievers, J.; Schlossberger, H.G. (1982) Dihydroxytryptamines as tools to study the neurobiology of serotonin. *Brain Res. Bull.* **9**(1-6): 131-150.
- Belanoff, J.K.; Kalehzan, M.; Sund, B.; Fleming-Ficek, S.K.; Schatzberg, A.F. (2001) Cortisol activity and cognitive changes in psychotic major depression. *Am. J. Psychiatry* **158**(10): 1612-1616.
- Bell, C.J.; Nutt, D.J. (1998) Serotonin and panic. *Br. J. Psychiatry* **172**: 465-471.
- Bender, N. (2004) Gene therapy: a concept for the future in medical practice? The current state. *Hamostaseologie* **24**(3): 173-178.
- Bengel, D.; Murphy, D.L.; Andrews, A.M.; Wicherms, C.H.; Feltner, D.; Heils, A.; Mossner, R.; Westphal, H.; Lesch, K.P. (1998) Altered brain serotonin homeostasis and locomotor insensitivity to 3,4-methylenedioxymethamphetamine ("Ecstasy") in serotonin transporter-deficient mice. *Mol. Pharmacol.* **53**: 649-655.
- Bertrand, E.; Pictet, R.; Grange, T. (1994) Can hammerhead ribozymes be efficient tools to inactivate gene function? *Nucl. Acids Res.* **22**(3): 293-300.
- Birnboim, H.C.; Doly, J. (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids Res.* **7**: 1513-1523.
- Boadle-Biber, M.C. (1978) Activation of tryptophan hydroxylase from central serotonergic neurons by calcium and depolarization. *Biochem. Pharmacol.* **27**(7): 1069-1079.
- Boadle-Biber, M.C. (1993) Regulation of serotonin synthesis. *Prog. Biophys. Mol. Biol.* **60**: 1-15.
- Boadle-Biber, M.C.; Phan, T.H. (1986) AGN 2979 [3(3-methoxyphenyl)-3-(3-dimethylaminopropyl)-4, 4-dimethylpiperidine-2,6-dione]. An inhibitor of the activation of tryptophan hydroxylase. *Biochem. Pharmacol.* **35**(9): 1521-1526.
- Bodner, R.A.; Lynch, T.; Lewis, L.; Kahn, D. (1995) Serotonin syndrome. *Neurology* **45**(2): 219-223.
- Borman, R.A.; Burleigh, D.E. (1993) Evidence for the involvement of a 5-HT₄ receptor in the secretory response of human small intestine to 5-HT. *Br. J. Pharmacol.* **110**(3): 927-928.
- Borowsky, B., Hoffman, B.J. (1995) Neurotransmitter transporters: Molecular biology, function, and regulation. *Int. Rev. Neurobiol.* **38**: 139-199
- Bosin, T.R.; Hixson, E.J.; Maickel, R.P. (1976) Pressor effects of tryptamine analogues. *Br. J. Pharmacol.* **56**: 25-27.
- Boston, P.F.; Jackson, P.; Kynoch, P.A.; Thompson, R.J. (1982) Purification and immunohistochemical localization of human brain 14-3-3. *J. Neurochem.* **38**: 1466-1474.

- Bottlander, R.; Dobmeier, P.; Moller, H.J. (1998) Der Einfluß von selektiven Serotonin-Wiederaufnahmehemmern auf die Blutgerinnung. *Fortschr. Neurol. Psychiatr.*, **66**: 32-35.
- Bou-Flores, C.; Lajard, A.M., Monteau, R.; De Maeyer, E.; Seif, I.; Lanoir, J.; Hilaire, G. (2000) Abnormal phrenic motoneuron activity and morphology in neonatal monoamine oxidase A-deficient transgenic mice: possible role of a serotonin excess. *J. Neurosci.* **20**(12): 4646-4656.
- Boularand, S.; Biguet, N.F.; Vidal, B.; Vernon, M.; Mallet, J.; Vincent, J.-D.; Dufour, S.; Vernier, P. (1998) Tyrosine hydroxylase in the european eel (*Anguilla anguilla*): cDNA cloning, brain distribution, and phylogenetic analysis. *J. Neurochem.* **71**: 460-470.
- Boularand, S.; Darmon, M. C.; Mallet, J. (1995a) The human tryptophan hydroxylase gene. An unusual splicing complexity in the 5'-untranslated region. *J. Biol. Chem.* **270**: 3748-3756.
- Boularand, S.; Darmon, M. C.; Ravassard, P.; Mallet, J. (1995b) Characterization of the human tryptophan hydroxylase gene promoter. Transcriptional regulation by cAMP requires a new motif distinct from the cAMP-responsive element. *J. Biol. Chem.* **270**: 3757-3764.
- Boularand, S.; Darmon, M.C.; Ganem, Y.; Launay, J.-M.; Mallet, J. (1990) Complete coding sequence of human tryptophan hydroxylase. *Nucl. Acids Res.* **18**: 4257.
- Boutrel, B.; Monaca, C.; Hen, R.; Hamon, M.; Adrien, J. (2002) Involvement of 5-HT1A receptors in homeostatic and stress-induced adaptive regulations of paradoxical sleep: studies in 5-HT1A knock-out mice. *J. Neurosci.* **22**(11): 4686-4692.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248-254.
- Briley, M.; Chopin, P.; Moret, C. (1990) Effect of serotonergic lesion on "anxious" behaviour measured in the elevated plus-maze test in the rat. *Psychopharmacology (Berl.)* **101**(2): 187-189.
- Brinck, U.; Meyer, T. (2000) Tryptophan hydroxylase and 5-hydroxytryptamine in carcinoids of the gastrointestinal tract. *Anticancer Res.* **20**(4): 2325-2328.
- Brodsky, L.I.; Ivanov, V.V.; Kalaydzidis, Y.L.; Leontovich, A.M., Nikolaev, V.K., Feranchuk, S.I.; Drachev, V.A. (1995) GeneBee-NET: Internet-Based Server for Analyzing Biopolymers Structure. *Biochemistry* **60**(8): 923-928
- Brown, S.M.; Peet, E.; Manuck, S.B.; Williamson, D.E.; Dahl, R.E.; Ferrell, R.E.; Hariri, A.R. (2005) A regulatory variant of the human tryptophan hydroxylase-2 gene biases amygdala reactivity. *Mol. Psychiatry* [Epub ahead of print]
- Brunner, D.; Buhot, M.C.; Hen, R.; Hofer, M. (1999) Anxiety, motor activation, and maternal-infant interactions in 5HT1B knockout mice. *Behav. Neurosci.* **113**(3): 587-601.
- Cabrera-Vera, T.M.; Garcia, F.; Pinto, W.; Battaglia, G. (1997) Effect of Prenatal Fluoxetine (Prozac) Exposure on Brain Serotonin Neurons in Prepubescent and Adult Male Rat Offspring. *J. Pharmacol. Exp. Ther.* **280**(1): 138-145.

- Cady, S.G.; Sono, M. (1991) 1-Methyl-DL-tryptophan, beta-(3-benzofuranyl)-DL-alanine (the oxygen analog of tryptophan), and beta-[3-benzo(b)thienyl]-DL-alanine (the sulfur analog of tryptophan) are competitive inhibitors for indoleamine 2,3-dioxygenase. *Arch. Biochem. Biophys.* **291**: 326–333
- Carlsson, A.; Lindqvist, M. (1972) The effect of L-tryptophan and some psychotropic drugs on the formation of 5-hydroxytryptophan in the mouse brain in vivo. *J. Neural Transm.* **33**: 23-42.
- Carrasco, G.; Cruz, M.A.; Dominguez, A., Gallardo, V.; Miguel, P., Gonzalez, C. (2000b) The expression and activity of monoamine oxidase A, but not of the serotonin transporter, is decreased in human placenta from pre-eclamptic pregnancies. *Life Sci.* **67**(24): 2961-2969.
- Carrasco, G.; Cruz, M.A.; Gallardo, V.; Miguel, P.; Dominguez, A.; Gonzalez, C. (2000a) Transport and metabolism of serotonin in the human placenta from normal and severely pre-eclamptic pregnancies. *Gynecol. Obstet. Invest.* **49**(3): 150-155.
- Cases, O.; Seif, I.; Grimsby, J.; Gaspar, P.; Chen, C.; Pournin, S.; Müller, U.; Aguet, M.; Babinet, C.; Shih, J.C.; De Maeyer, E. (1995) Aggressive behavior and altered amounts of brain serotonin and norepinephrine in mice lacking MAOA. *Science* **268**: 1763-1766.
- Cases, O.; Vitalis, T.; Seif, I.; De Maeyer, E., Sotelo, C.; Gaspar, P. (1996) Lack of barrels in the somatosensory cortex of monoamine oxidase A-deficient mice: role of a serotonin excess during the critical period. *Neuron* **16**(2): 297-307.
- Cash, C.D. (1998) Why tryptophan hydroxylase is difficult to purify: A reactive oxygen-derived species-mediated phenomenon that may be implicated in human pathology. *Gen. Pharmacol.* **30**: 569-574.
- Cash, C.D.; Vayer, P.; Mandel, P.; Maitre, M. (1985) Tryptophan hydroxylase. Rapid purification from whole rat brain and production of a specific antiserum. *Eur. J. Biochem.*, **149**: 239-245.
- Caspi, A.; Sugden, K.; Moffitt, T.E.; Taylor, A.; Craig, I.W.; Harrington, H.; McClay, J.; Mill, J.; Martin, J.; Braithwaite, A.; Poulton, R. (2003) Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* **301**(5631): 386-389.
- Cattaneo, M.G.; Codignola, A.; Vicentini, L.M.; Clementi, F.; Sher, E. (1993) Nicotine stimulates a serotonergic autocrine loop in human small-cell lung carcinoma. *Cancer Res.* **53**(22): 5566-5568.
- Cerda, J.; Reich, G.; Wallace, R.A.; Selman, K. (1998) Serotonin inhibition of steroid-induced meiotic maturation in the teleost Fundulus heteroclitus: role of cyclic AMP and protein kinases. *Mol. Reprod. Dev.* **49**(3): 333-341.
- Chamberlain, B.; Ervin, F.R.; Pihl, R.O.; Young, S.N. (1987) The effect of raising or lowering tryptophan levels on aggression in rhesus monkeys. *Pharmacol. Biochem. Behav.* **28**(4): 503-510.

- Champier, J.; Claustrat, B.; Besncon, R.; Eymin, C.; Killer, C.; Jouvet, A.; Chamba, G.; Fèvre-Montange, M. (1997) Evidence for tryptophan hydroxylase and hydroxy-indol-O-methyl-transferase mRNAs in human blood platelets. *Life Sci.* **60**: 2191-2197.
- Chiang, M.Y.; Chan, H.; Zounes, M.A.; Freier, S.M.; Lima, W.F.; Bennett, C.F. (1991) Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms. *J. Biol. Chem.* **266**(27): 18162-18171.
- Cho, S.H.; Huh, S.O.; Park, D.H.; Joh, T.H.; Son, J.H. (1993) Mouse tryptophan hydroxylase (TPH) promotor regions mediating basal and regulated expression of TPH gene in vitro. *Neurosci. Abst.* **19**: 35.5.
- Chugani, D.C. (2002) Role of altered brain serotonin mechanisms in autism. *Mol. Psychiatry* **7** Suppl. 2: S16-17.
- Clark, J.A.; Pai, L.Y.; Flick, R.B.; Rohrer, S.P. (2005) Differential hormonal regulation of tryptophan hydroxylase-2 mRNA in the murine dorsal raphe nucleus. *Biol. Psychiatry*. **57**(8): 943-946.
- Cohen, R.A.; Shepherd, J.T.; Vanhoutte, P.M. (1983) Inhibitory role of the endothelium in the response of isolated coronary arteries to platelets. *Science* **221**(4607): 273-274.
- Colas, J.F.; Launay, J.M.; Maroteaux, L. (1999a) Maternal and zygotic control of serotonin biosynthesis are both necessary for Drosophila germband extension. *Mech. Dev.* **87**(1-2): 67-76.
- Colas, J.F.; Launay, J.M.; Vonesch, J.L.; Hickel, P.; Maroteaux, L. (1999b) Serotonin synchronises convergent extension of ectoderm with morphogenetic gastrulation movements in Drosophila. *Mech. Dev.* **87**(1-2): 77-91.
- Coligan, J. E.; Kruisbeek, M. A.; Margulies, D. H.; Shevach, E. M.; Strober, W., (1991) *Current Protocols in Immunology*, Wiley: New York
- Compan, V.; Zhou, M.; Grailhe, R.; Gazzara, R.A.; Martin, R.; Gingrich, J.; Dumuis, A.; Brunner, D.; Bockaert, J.; Hen, R. (2004) Attenuated response to stress and novelty and hypersensitivity to seizures in 5-HT4 receptor knock-out mice. *J. Neurosci.* **24**(2): 412-419.
- Conaty, J.; Hendry, P.; Lockett, T. (1999) Selected classes of minimized hammerhead ribozyme have very high cleavage rates at low Mg²⁺ concentration. *Nucl. Acids. Res.* **27** (11): 2400-2407
- Coon, H.; Dunn, D.; Lainhart, J.; Miller, J.; Hamil, C.; Battaglia, A.; Tancredi, R.; Leppert, M.F.; Weiss, R.; McMahon, W. (2005) Possible association between autism and variants in the brain-expressed tryptophan hydroxylase gene (TPH2). *Am. J. Med. Genet. B. Neuropsychiatr. Genet.* **135**(1): 42-46.
- Coon, S.L.; Mazuruk, K.; Bernard, M.; Roseboom, P.H.; Klein, D.C.; Rodriguez, I.R. (1996) The human serotonin N-acetyltransferase (EC 2.3.1.87) gene (AANAT): structure, chromosomal localization, and tissue expression. *Genomics* **34**: 76-84.

- Couture, L.A.; Stinchcomb, D.T. (1996) Anti-gene therapy: the use of ribozymes to inhibit gene function. *Trends Genet.* **12**(12): 510-515.
- D'Amato, R.J.; Blue, M.E.; Largent, B.L.; Lynch, D.R.; Ledbetter, D.J.; Molliver, M.E.; Snyder, S.H. (1987) Ontogeny of the serotonergic projection to rat neocortex: transient expression of a dense innervation to primary sensory areas. *Proc. Natl. Acad. Sci. U. S. A.* **84**: 4322-4326.
- Dagert, M.; Ehrlich, S.D. (1979) Prolonged incubation in calcium chloride improves the competence of Escherichia coli cells. *Gene* **6**: 23-28.
- Dahlström, A.; Fuxe, K. (1964) Evidence for the existence of monoamine-containing neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta Physiol. Scand.* **62**: 1-55.
- Dale, G.L.; Friese, P.; Batar, P.; Hamilton, S.F.; Reed, G.L.; Jackson, K.W.; Clemetson, K.J.; Alberio L. (2002) Stimulated platelets use serotonin to enhance their retention of procoagulant proteins on the cell surface. *Nature* **415**(6868): 175-179.
- Danos, O.; Mulligan, R.C. (1988) Safe and efficient generation of recombinant retroviruses with amphotropic and ecotropic host ranges. *Proc. Natl. Acad. Sci. U. S. A.* **85**(17): 6460-6464.
- Daubner, S.C.; Lohse, D.L.; Fitzpatrick, P.F. (1993) Expression and characterization of catalytic and regulatory domains of rat tyrosine hydroxylase. *Protein Sci.* **2**: 1452-1460.
- De Luca, V.; Mueller, D.J.; Tharmalingam, S.; King, N.; Kennedy, J.L. (2004) Analysis of the novel TPH2 gene in bipolar disorder and suicidality. *Mol. Psychiatry* **9**(10): 896-897.
- Dean, B.; Hayes, W. (1996) Decreased frontal cortical serotonin2A receptors in schizophrenia. *Schizophr. Res.* **21**(3): 133-139.
- Delarue, C.; Becquet, D.; Idres, S.; Hery, F.; Vaudry, H. (1992) Serotonin synthesis in adrenochromaffin cells. *Neurosci.* **46**: 495-500.
- Dewhurst, W.G.; Pare, C.M. (1961) A clinical and biochemical study of monoamine oxidase inhibition in depressed patients. II. 5-Hydroxytryptamine tolerance before and after Nialamide. *J. Ment. Sci.* **107**: 244-249.
- Dumas, S.; Darmon, M.C.; Delort, J.; Mallet, J. (1989) Differential control of tryptophan hydroxylase expression in raphe and in pineal gland: evidence for a role of translation efficiency. *J. Neurosci. Res.* **24**: 537-547.
- Edelman, J.; Edelman, J.; Nemunaitis, J. (2003) Adenoviral p53 gene therapy in squamous cell cancer of the head and neck region. *Curr. Opin. Mol. Ther.* **5**(6): 611-617.
- Efrat, S.; Leiser, M.; Wu, Y.J.; Fusco-DeMane, D.; Emran, O.A.; Surana, M.; Jetton, T.L.; Magnuson, M.A.; Weir, G.; Fleischer, N. (1994) Ribozyme-mediated attenuation of pancreatic beta-cell glucokinase expression in transgenic mice results in impaired glucose-induced insulin secretion. *Proc. Natl. Acad. Sci. U. S. A.* **91**(6): 2051-2055.

- Ehret, M.; Cash, C.D.; Hamon, M.; Maitre, M. (1989) Formal demonstration of the phosphorylation of rat brain tryptophan hydroxylase by a Ca²⁺/calmodulin-dependent protein kinase. *J. Neurochem.* **52**: 1886-1891.
- Ek, A.; Witkop, B. (1954) The Synthesis of Labile Hydroxytryptophan Metabolites. *J. Am. Chem. Soc.* **76**(22): 5579-5588.
- Elbashir, S.M.; Harborth, J.; Lendeckel, W.; Yalcin, A.; Weber, K.; Tuschl, T. (2001b) Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* **411**(6836): 494-498.
- Elbashir, S.M.; Lendeckel, W.; Tuschl, T. (2001a) RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev.* **15**(2): 188-200.
- Elwood, P.; Morgan, G.; Brown, G.; Pickering, J. (2005) Aspirin for everyone older than 50? *For. BMJ.* **330**(7505): 1440-1441.
- Ener, R.A.; Meglathery, S.B.; Van Decker, W.A.; Gallagher, R.M. (2003) Serotonin syndrome and other serotonergic disorders. *Pain Med.* **4**(1): 63-74.
- Farrer, T.; Roller, A.B.; Kent, W.J.; Zahler, A.M. (2002) Analysis of the role of *Caenorhabditis elegans* GC-AG introns in regulated splicing. *Nucl. Acids Res.* **30**(15): 3360-3367.
- Fedor, M.J.; Uhlenbeck, O.C. (1992) Kinetics of intermolecular cleavage by hammerhead ribozymes. *Biochemistry* **31**: 12042-12054
- Feinberg, A.P.; Vogelstein, B. (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**: 6-13.
- Fenton, H.J.H. (1854) Oxidation of tartaric acid in presence of iron. *J. Chem. Soc.* **65**: 899-910.
- Finocchiaro, L.M.; Arzt, E.S.; Fernandez-Castelo, S.; Criscuolo, M.; Finkelman, S.; Nahmod, V.E. (1988) Serotonin and melatonin synthesis in peripheral blood mononuclear cells: stimulation by interferon-gamma as part of an immunomodulatory pathway. *J. Interferon Res.* **8**: 705-716.
- Fiorica-Howells, E.; Hen, R.; Gingrich, J.; Li, Z.; Gershon, M.D. (2002) 5-HT(2A) receptors: location and functional analysis in intestines of wild-type and 5-HT(2A) knockout mice. *Am. J. Physiol. Gastrointest. Liver Physiol.* **282**(5): G877-893.
- Fisher, D.B.; Kaufman, S. (1972) The stimulation of rat liver phenylalanine hydroxylase by phospholipids. *J. Biol. Chem.* **247**: 2250-2252.
- Flatman, P.W. (1991) Mechanisms of Magnesium Transport. *Annu. Rev. Physiol.* **53**: 259-271.
- Fon, E.A.; Pothos, E.N.; Sun, B.C.; Killeen, N.; Sulzer, D.; Edwards, R.H. (1997) Vesicular transport regulates monoamine storage and release but is not essential for amphetamine action. *Neuron* **19**(6): 1271-1283.
- Friedman, P.A.; Kappelman, A.H.; Kaufman, S. (1972) Partial purification and characterization of tryptophan hydroxylase from rabbit hindbrain. *J. Biol. Chem.* **247**: 4165-4173.

- Frungieri, M.B.; Gonzalez-Calvar, S.I.; Rubio, M.; Ozu, M.; Lustig, L.; Calandra, R.S. (1999) Serotonin in golden hamster testes: testicular levels, immunolocalization and role during sexual development and photoperiodic regression-recrudescence transition. *Neuroendocrinology* **69**(4): 299-308.
- Fujimoto, S.; Ueda, H.; Kagawa, H. (1987) Immunocytochemistry on the localization of 5-hydroxytryptamine in monkey and rabbit taste buds. *Acta Anat. (Basel)* **128**(1): 80-83.
- Fujisawa, H.; Nakata, H. (1987) Tryptophan 5-monooxygenase from mouse mastocytoma clone P815. *Meth. Enzymol.* **142**: 93-96.
- Furukawa, Y.; Ikuta, N.; Omata, S.; Yamauchi, T.; Isobe, T.; Ichimura, T. (1993) Demonstration of the phosphorylation-dependent interaction of tryptophan hydroxylase with the 14-3-3 protein. *Biochem. Biophys. Res. Commun.* **194**: 144-149.
- Gal, E.M.; Patterson, K. (1973) Rapid non-isotopic assay of tryptophan-5-hydroxylase activity in tissues. *Anal. Biochem.* **52**: 625-629.
- Gal, E.M.; Sherman, A.D. (1980) L-kynurenone: its synthesis and possible regulatory function in brain. *Neurochem. Res.* **5**(3): 223-239.
- Garriock, H.A.; Allen, J.J.; Delgado, P.; Nahaz, Z.; Kling, M.A.; Carpenter, L.; Burke, M.; Burke, W.; Schwartz, T.; Marangell, L.B.; Husain, M.; Erickson, R.P.; Moreno, F.A. (2005) Lack of association of TPH2 exon XI polymorphisms with major depression and treatment resistance. *Mol. Psychiatry*. [Epub ahead of print].
- Gaughan, D.J.; Whitehead, A.S. (1997) Efficient in vitro cleavage of mouse acute phase serum amyloid A mRNA mediated by a synthetic hammerhead ribozyme. *Scand. J. Immunol.* **46**(1): 51-58.
- Geba, G.P.; Ptak, W.; Anderson, G.M.; Paliwal, V.; Ratzlaff, R.E.; Levin, J.; Askenase, P.W. (1996) Delayed-type hypersensitivity in mast cell-deficient mice. *J. Immunol.* **157**: 557-565.
- Gershon, R.K.; Askenase, P.W.; Gershon, M.D. (1975) Requirement for vasoactive amines for production of delayed-type hypersensitivity skin reactions. *J. Exp. Med.* **142**: 732-747.
- Gholson, R.K.; Hankes, L.V.; Henderson, L.M. (1960) 3-Hydroxyanthranilic acid as an intermediate in the oxidation of the indole nucleus of tryptophan. *J. Biol. Chem.* **235**: 132-135.
- Gibbons, J.L. (1964) Cortisol Secretion Rate In Depressive Illness. *Arch. Gen. Psychiatry* **10**: 572-575.
- Gilbert, J.A.; Frederick, L.M.; Ames, M.M. (2000) The aromatic-L-amino acid decarboxylase inhibitor carbidopa is selectively cytotoxic to human pulmonary carcinoid and small cell lung carcinoma cells. *Clin. Cancer Res.* **6**: 4365-4372.
- Gingrich, J.A.; Hen, R. (2001) Dissecting the role of the serotonin system in neuropsychiatric disorders using knockout mice. *Psychopharmacology (Berl.)* **155**(1): 1-10.

- Goldberg, R.J. (1998) Selective serotonin reuptake inhibitors: infrequent medical adverse effects. *Arch. Fam. Med.* **7**: 78-84.
- Goodman, M.; New, A. (2000) Impulsive aggression in borderline personality disorder. *Curr. Psychiatry Rep.* **2**(1): 56-61.
- Goodwill, K.E.; Sabatier, C.; Marks, C.; Raag, R.; Fitzpatrick, P.F.; Stevens, R.C. (1997) Crystal structure of tyrosine hydroxylase at 2.3 Å and its implications for inherited neurodegenerative diseases. *Nature Struct. Biol.* **4**: 578-585.
- Gordon, J.W.; Scangos, G.A.; Plotkin, D.J.; Barbosa, J.A.; Ruddle, F.H. (1980) Genetic transformation of mouse embryos by microinjection of purified DNA. *Proc. Natl. Acad. Sci. U. S. A.* **77**(12): 7380-7384.
- Grailhe, R.; Waeber, C.; Dulawa, S.C.; Hornung, J.P.; Zhuang, X.; Brunner, D.; Geyer, M.A.; Hen, R.; (1999) Increased exploratory activity and altered response to LSD in mice lacking the 5-HT(5A) receptor. *Neuron* **22**(3): 581-591.
- Green, C.B.; Beharse, J.C. (1994) Tryptophan hydroxylase expression is regulated by a circadian clock in *Xenopus laevis* retina. *J. Neurochem.* **62**: 2420-2428.
- Grenett, H.E.; Ledley, F.D.; Reed, L.L.; Woo, S.L.C. (1987) Full-length cDNA for rabbit tryptophan hydroxylase: functional domains and evolution of aromatic amino acid hydroxylases. *Proc. Natl. Acad. Sci. U. S. A.* **84**: 5530-5534.
- Grimbsby, J.; Toth, M.; Chen, K.; Kumazawa, T.; Klaidman, L.; Adams, J.D.; Karoum, F.; Gal, J.; Shih, J.C. (1997) Increased stress response and beta-phenylethylamine in MAOB-deficient mice. *Nature Gen.* **17**: 206-210.
- Gross C; Santarelli L, Brunner D; Zhuang X, Hen R. (2000) Altered fear circuits in 5-HT(1A) receptor KO mice. *Biol. Psychiatry* **48**(12): 1157-1163.
- Gross C; Zhuang X, Stark K; Ramboz S; Oosting R; Kirby L; Santarelli L; Beck S; Hen R. (2002) Serotonin1A receptor acts during development to establish normal anxiety-like behaviour in the adult. *Nature* **416**(6879): 396-400.
- Guerrier-Takada C; Gardiner K; Marsh T; Pace N; Altman S. (1983) The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* **35**(3 Pt 2): 849-857.
- Hammann, C.; Hormes, R.; Szakiel, G.; Tabler, M. (1997) A spermidine-induced conformational change of long-armed hammerhead ribozymes: ionic requirements for fast cleavage kinetics. *Nucleic Acids Res.* **25**(23): 4715-4722.
- Hamon, M.; Bourgoin, S.; Hery, F.; Simonnet, G. (1978) Activation of tryptophan hydroxylase by adenosine triphosphate, magnesium and calcium. *Mol. Pharmacol.* **14**: 99-110.
- Hanahan, D. (1983) Studies on transformation of *E. coli* with plasmids. *J. Mol. Biol.* **166**: 557-563.

- Hanahan, D.; Jessee, J., Bloom, F.R. (1991) Plasmid transformation of escherichia coli and other bacteria. *Meth. Enzymol.* **204**: 63-113.
- Hansen, M.B. (1995) Serotonin--an intestinal secretagogue--receptor subtypes and intracellular mediators. *Pharmacol. Toxicol.* **77** Suppl 1: 3-39.
- Harlow, E.; Lane, D. (1988) Immunoblotting. In: Antibodies: A Laboratory Manual. CSH Laboratory, Cold Spring Harbor, N. Y.: 471-510.
- Harris, A.L.; Smith, I.E. (1983) Tryptophan in the treatment of carcinoid crisis. *Cancer Chemother. Pharmacol.* **10**(2): 137-139.
- Hart, R.P.; Yang, R.; Riley, L.A.; Green, T.L. (1991) Post-transcriptional control of tryptophan hydroxylase gene expression in rat brain stem and pineal gland. *Mol. Cell. Neurosci.* **2**: 71-77.
- Harvey M; Shink E; Tremblay M; Gagne B; Raymond C; Labbe M; Walther DJ; Bader M; Barden N. (2004) Support for the involvement of TPH2 gene in affective disorders. *Mol. Psychiatry* **9**(11): 980-981.
- Hasegawa, H.; Ichiyama, A. (1987) Tryptophan 5-Monooxygenase from mouse mastocytoma: High-performance liquid chromatography assay. *Meth. Enzymol.* **142**: 88-92.
- Hedlund, P.B.; Danielson, P.E.; Thomas, E.A.; Slanina, K.; Carson, M.J.; Sutcliffe, J.G. (2003) No hypothermic response to serotonin in 5-HT7 receptor knockout mice. *Proc. Natl. Acad. Sci. U. S. A.* **100**(3): 1375-1380.
- Hedner, J.; Lundell, K.H.; Breese, G.R.; Mueller, R.A.; Hedner, T. (1986) Developmental variations in CSF monoamine metabolites during childhood. *Biol. Neonate* **49**: 190-197.
- Heidenreich, O.; Xu, X.; Swiderski, P.; Rossi, J.J.; Nerenberg, M. (1996) Correlation of activity with stability of chemically modified ribozymes in nuclei suspension. *Antisense Nucleic Acid Drug Dev.* **6**(2): 111-118.
- Heisler LK; Chu HM; Tecott LH. (1998a) Epilepsy and obesity in serotonin 5-HT2C receptor mutant mice. *Ann. N. Y. Acad. Sci.* **861**: 74-78.
- Hellstrand K; Hermodsson S. (1993) Serotonergic 5-HT1A receptors regulate a cell contact-mediated interaction between natural killer cells and monocytes. *Scand. J. Immunol.* **37**(1): 7-18.
- Hendricks TJ; Fyodorov DV; Wegman LJ; Lelutiu NB; Pehek EA; Yamamoto B; Silver J; Weeber EJ; Sweatt JD; Deneris ES. (2003) Pet-1 ETS gene plays a critical role in 5-HT neuron development and is required for normal anxiety-like and aggressive behavior. *Neuron* **37**(2): 233-247.
- Herschlag; D. (1991) Implications of ribozyme kinetics for targeting the cleavage of specific RNA molecules in vivo: more isn't always better. *Proc. Natl. Acad. Sci. U S A.* **88**(16): 6921-6925.
- Hertel, K.J.; Herschlag, D.; Uhlenbeck, O.C. (1994) A Kinetic and Thermodynamic Framework for the Hammerhead Ribozyme Reaktion. *Biochemistry* **33**: 3374-3385

- Higley, J.D.; Linnoila, M. (1997) Low central nervous system serotonergic activity is traitlike and correlates with impulsive behavior. A nonhuman primate model investigating genetic and environmental influences on neurotransmission. *Ann. N. Y. Acad. Sci.* **836**: 39-56.
- Hixson, E.J.; Lehrmann, G.V.; Maickel, R.P. (1977) Contractile responses to tryptamine analogues in isolated smooth muscle. *Arch. Int. Pharmacodyn.* **229**: 4-14.
- Hogan, B.; Costantini, F.; Lacy, E. (1986) Manipulating the mouse embryo - A laboratory manual. 1-332.
- Holland, J.M. (1976) Serotonin deficiency and prolonged bleeding in beige mice. *Proc. Soc. Exp. Biol. Med.* **151**: 32-39.
- Holmes, A.; Murphy, D.L.; Crawley, J.N. (2002a) Reduced aggression in mice lacking the serotonin transporter. *Psychopharmacology (Berl.)* **161**(2): 160-167.
- Holmes, A.; Yang, R.J.; Lesch, K.P.; Crawley, J.N.; Murphy, D.L. (2003) Mice lacking the serotonin transporter exhibit 5-HT(1A) receptor-mediated abnormalities in tests for anxiety-like behavior. *Neuropsychopharmacology* **28**(12): 2077-2088.
- Holmes, A.; Yang, R.J.; Murphy, D.L.; Crawley, J.N. (2002b) Evaluation of antidepressant-related behavioral responses in mice lacking the serotonin transporter. *Neuropsychopharmacology* **27**(6): 914-923.
- Holsboer, F.; Barden, N. (1996) Antidepressants and hypothalamic-pituitary-adrenocortical regulation. *Endocr. Rev.* **17**(2): 187-205.
- Houston, D.S.; Vanhouette, P.M. (1986) Serotonin and the vascular system. Role in health and disease, and implications for therapy. *Drugs* **31**: 149-163.
- Hoyer, D. and Martin, G.R. (1997) 5-HT receptor classification and nomenclature: towards a harmonization with the human genome. *Neuropharmacol.* **36**: 419-428.
- Hoyer, D.; Clarke, D.E.; Fozard, J.R.; Hartig, P.R.; Martin, G.R.; Mylecharane, E.J.; Saxena, P.R.; Humphrey, P.P.A. (1994) VII. International union of pharmacology classification of receptors for 5-hydroxytryptamine (serotonin). *Pharmacol. Rev.* **46**: 157-204.
- Huether, G.; Schmidt, S.; Rüther, E. (1998) Essen, Serotonin und Psyche. *Deutsches Ärzteblatt* **95**: C-377-379.
- Huh, S.O.; Park, D.H.; Cho, J.Y.; Joh, T.H.; Son, J.H. (1994) A 6.1 kb 5' upstream region of the mouse tryptophan hydroxylase gene directs expression of *E. coli* lacZ to major serotonergic brain regions and pineal gland in transgenic mice. *Mol. Brain. Res.* **24**: 145-152.
- Humphrey, P.P.; Hartig, P.; Hoyer, D. (1993) A proposed new nomenclature for 5-HT receptors. *Trends Pharmacol. Sci.* **14**(6): 233-236.
- Hutson, P.H.; Sarna, G.S.; Sahakian, B.J.; Dourish, C.T.; Curzon, G. (1986) Monitoring 5HT metabolism in the brain of the freely moving rat. *Ann. N. Y. Acad. Sci.* **473**: 321-336.

- Hwang, S.L.; Chung, N.P.; Chan, J.K.; Lin, C.L. (2005) Indoleamine 2, 3-dioxygenase (IDO) is essential for dendritic cell activation and chemotactic responsiveness to chemokines. *Cell Res.* **15**(3): 167-175.
- Ichimura, T.; Isobe, T.; Okuyama, T.; Takahashi, N.; Araki, K.; Kuwano, R. (1988) Molecular cloning of cDNA coding for brain specific 14-3-3 protein, a protein kinase-dependent activator of tyrosine and tryptophan hydroxylases. *Proc. Natl. Acad. Sci. U. S. A.* **85**: 7084-7088.
- Ichimura, T.; Isobe, T.; Okuyama, T.; Yamauchi, T.; Fujisawa, H. (1987) Brain 14-3-3 protein is an activator protein that activates tryptophan 5-monooxygenase and tyrosine 3-monooxygenase in the presence of Ca²⁺, calmodulin-dependent protein kinase II. *FEBS Lett.* **219**: 79-82.
- Ichimura, T.; Uchiyama, J.; Kunihiro, O.; Ito, M.; Horigome, T.; Omata, S. (1995) Identification of the site of interaction of the 14-3-3 protein with phosphorylated tryptophan hydroxylase. *J. Biol. Chem.* **270**: 515-518.
- Ichinose, H.; Ohye, T.; Fujita, K.; Yoshida, M.; Ueda, S.; Nagatsu, T. (1993) Increased heterogeneity of tyrosine hydroxylase in humans. *Biochem. Biophys. Res. Commun.* **195**: 158-165.
- Imai, Y.; Yamauchi, T.; Fujisawa, H. (1989) Modulation of tryptophan hydroxylase activity by phospholipids: stimulation followed by inactivation. *J. Neurochem.* **53**(4): 1293-1299.
- Ishizuka, J.; Beauchamp, R.D.; Townsend, C.M.Jr.; Greeley, G.H.Jr.; Thompson, J.C. (1992) Receptor-mediated autocrine growth-stimulatory effect of 5-hydroxytryptamine on cultured human pancreatic carcinoid cells. *J. Cell. Physiol.* **150**(1): 1-7
- Isobe, T.; Ichimura, T.; Sunaya, T.; Okuyama, T.; Takahashi, N.; Kuwano, R. (1991) Distinct forms of the protein kinase-dependent activator of tyrosine and tryptophan hydroxylases. *J. Mol. Biol.* **217**: 125-132.
- Iwaki, M.; Phillips, R.S.; Kaufman, S. (1986) Proteolytic modification of the amino-terminal and carbonyl-terminal regions of rat hepatic phenylalanine hydroxylase. *J. Biol. Chem.* **261**: 2051-2056.
- Jackson, A.L.; Bartz, S.R.; Schelter, J.; Kobayashi, S.V.; Burchard, J.; Mao, M.; Li, B.; Cavet, G.; Linsley, P.S. (2003) Expression profiling reveals off-target gene regulation by RNAi. *Nat. Biotechnol.* **21**(6): 635-637.
- Jackson, A.L.; Linsley, P.S. (2004) Noise amidst the silence: off-target effects of siRNAs? *Trends Genet.* **20**(11): 521-524.
- Jann, M.W.; Kurtz, N.M. (1987) Treatment of panic and phobic disorders. *Clin. Pharm.* **6**(12): 947-962.
- Jequier, E.; Robinson, D.S.; Lovenberg, W.; Sjoerdsma, A. (1969) Further studies on tryptophan hydroxylase in rat brainstem and beef pineal. *Biochem. Pharmacol.* **18**(5): 1071-1081.
- Joh, T.H. (1997) Tryptophan hydroxylase: molecular biology and regulation. *Serotonergic Neurons and 5-HT Receptors in the CNS* 117-130.

- Jones, B.J.; Blackburn, T.P. (2002) The medical benefit of 5-HT research. *Pharmacol. Biochem. Behav.* **71**(4): 555-568.
- Jones, R.S. (1982) Tryptamine: a neuromodulator or neurotransmitter in mammalian brain? *Prog. Neurobiol.* **19**: 117–139.
- Jonsson, G. (1980) Chemical neurotoxins as denervation tools in neurobiology. *Annu. Rev. Neurosci.* **3**: 169-187.
- Kaneda, N.; Kobayashi, K.; Ichinose, H.; Kishi, F.; Nakazawa, A.; Kurosawa, Y.; Fujita, K.; Nagatsu, T. (1987) Isolation of a novel cDNA clone for human tyrosine hydroxylase: alternative RNA splicing produces four kinds of mRNA from a single gene. *Biochem. Biophys. Res. Commun.* **146**: 971-975.
- Kantak, K.M.; Hegstrand, L.R.; Eichelman, B. (1980) Dietary tryptophan modulation and aggressive behavior in mice. *Pharmacol Biochem. Behav.* **12**(5): 675-679.
- Kato, Y.; Kuwabara, T.; Warashina, M.; Toda, H.; Taira, K. (2001) Relationships between the activities in vitro and in vivo of various kinds of ribozyme and their intracellular localization in mammalian cells. *J. Biol. Chem.* **276**(18): 15378-15385.
- Kaufman, S. (1959) Studies on the mechanism of the enzymatic conversion of phenylalanine to tyrosine. *J. Biol. Chem.* **234**: 2677-2682.
- Kaufman, S. (1993) The phenylalanine hydroxylating system. *Adv. Enzy. Related Areas Mol. Biol.* **67**: 77-264.
- Kaufman, S.; Fisher, D.B. (1970) Purification and some physical properties of phenylalanine hydroxylase from rat liver. *J. Biol. Chem.*, **245**: 4745-4750.
- Kawasaki, H.; Taira, K. (2004) Induction of DNA methylation and gene silencing by short interfering RNAs in human cells. *Nature* **431**(7005): 211-217.
- Khozhai, L.I.; Puchkov, V.F.; Otellin, V.A. (1995) The effect of a serotonin deficiency on mammalian embryonic development. *Ontogenet* **26**: 350-355.
- Kiehntopf, M.; Esquivel, E.L.; Brach, M.A.; Herrmann, F. (1995) Clinical applications of ribozymes. *Lancet* **345**(8956): 1027-1031.
- Kijima, H.; Ishida, H.; Ohkawa, T.; Kashani-Sabet, M.; Scanlon, K.J. (1995) Therapeutic applications of ribozymes. *Pharmacol. Ther.* **68**(2): 247-267.
- Killam, A.L.; Cohen, M.L. (1991) Characterization of rat platelet serotonin receptors with tryptamine agonists and the antagonists: ketanserin and SCH 23390. *Thromb. Res.* **64**: 331–339.
- Kobayashi, H.; Dorai, T.; Holland, J.F.; Ohnuma, T. (1993) Cleavage of human MDR1 mRNA by a hammerhead ribozyme. *FEBS Lett.* **319**(1-2): 71-74.
- Kobayashi, H.; Dorai, T.; Holland, J.F.; Ohnuma, T. (1994) Reversal of drug sensitivity in multidrug-resistant tumor cells by an MDR1 (PGY1) ribozyme. *Cancer Res.* **54**(5): 1271-1275.

- Kore, A.R.; Vaish, N.K.; Kutzke, U.; Eckstein, F. (1998) Sequence specificity of the hammerhead ribozyme revisited; the NHH rule. *Nucleic Acids Res.* **26**(18): 4116-4120.
- Kore, A.R.; Vaish, N.K.; Kutzke, U.; Eckstein, F. (1998) Sequence specificity of the hammerhead ribozyme revisited; the NHH rule. *Nucl. Acids Res.* **26** (18): 4116-4120.
- Kowlessur, D.; Kaufman, S. (1999) Cloning and expression of recombinant human pineal tryptophan hydroxylase in Escherichia coli: purification and characterization of the cloned enzyme. *Biochim. Biophys. Acta* **1434**(2): 317-330.
- Krantz, D.E.; Peter, D.; Liu, Y.; Edwards, R.H. (1997) Phosphorylation of a vesicular monoamine transporter by casein kinase II. *J. Biol. Chem.* **272**(10): 6752-6759.
- Kruger, K.; Grabowski, P.J.; Zaug, A.J.; Sands, J.; Gottschling, D.E.; Cech, T.R. (1982) Self-splicing RNA: autoexcision and autocyclization of the ribosomal RNA intervening sequence of Tetrahymena. *Cell* **31**(1): 147-157.
- Kudo, Y.; Boyd, C.A.; Sargent, I.L.; Redman, C.W. (2001) Tryptophan degradation by human placental indoleamine 2,3-dioxygenase regulates lymphocyte proliferation. *J. Physiol.* **535**(Pt 1): 207-215.
- Kuhn, D.M.; Arthur, R.; States, J.C. (1997) Phosphorylation and activation of brain tryptophan hydroxylase: identification of serine-58 as a substrate site for protein kinase A. *J. Neurochem.* **68**: 2220-2223.
- Kuhn, D.M.; Arthur, R.E.Jr. (1996) Inactivation of brain tryptophan hydroxylase by nitric oxide. *J. Neurochem.* **67**(3): 1072-1077.
- Kuhn, D.M.; Lovenberg, W. (1982) Role of calmodulin in the activation of tryptophan hydroxylase. *Fed. Proc.* **41**: 2258-2264.
- Kuhn, D.M.; Meyer, M.A.; Lovenberg, W. (1980) Comparision of tryptophan hydroxylase from a malignant murine mast cell tumor and rat mesencephalic tegmentum. *Arch. Biochem. Biophys.* **199**: 355-361.
- Kuhn, D.M.; Vogel, R.L.; Lovenberg, W. (1978) Calcium-dependent activation of tryptophan hydroxylase by ATP and magnesium. *Biochem. Biophys. Res. Commun.* **82**: 759-766.
- Kumar, R.; Dammai, V.; Yadava, P.K.; Kleinau, S. (2005) Gene targeting by ribozyme against TNF-alpha mRNA inhibits autoimmune arthritis. *Gene Ther.* [Epub ahead of print].
- Kumer, S.C.; Mockus, S.M.; Rucker, P.J.; Vrana, K.E. (1997) Amino terminal deletion analysis of tryptophan hydroxylase: PKA phosphorylation occurs at serine-58. *J. Neurosci.* **69**: 1738-1745.
- Laberge, S.; Cruikshank, W.W.; Beer, D.J.; Center, D.M. (1996) Secretion of IL-16 (lymphocyte chemoattractant factor) form serotonin-stimulated CD8+ T cells in vitro. *J. Immunol.* **156**: 310-315.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680-685.

- Lange, W.; Cantin, E.M.; Finke, J.; Dolken, G. (1993) In vitro and in vivo effects of synthetic ribozymes targeted against BCR/ABL mRNA. *Leukemia* **7**(11): 1786-1794.
- Larsson, S.; Hotchkiss, G.; Andang, M.; Nyholm, T.; Inzunza, J.; Jansson, I.; Ahrlund-Richter, L. (1994) Reduced beta 2-microglobulin mRNA levels in transgenic mice expressing a designed hammerhead ribozyme. *Nucl. Acids Res.* **22**(12): 2242-2248.
- Lauder, J.M. (1990) Ontogeny of the serotonergic system in the rat: serotonin as a developmental signal. *Ann. N. Y. Acad. Sci.* **600**: 297-314.
- Lauder, J.M.; Krebs, H. (1978) Serotonin as a differentiation signal in early neurogenesis. *Dev. Neurosci.* **1**(1): 15-30.
- Lauder, J.M.; Towle, A.C.; Patrick, K.; Henderson, P.; Krebs, H. (1985) Decreased serotonin content of embryonic raphe neurons following maternal administration of p-chlorophenylalanine: a quantitative immunocytochemical study. *Brain Res.* **352**: 107-114.
- Lauder, J.M.; Wallace, J.A.; Krebs, H. (1981) Roles for serotonin in neuroembryogenesis. *Adv. Exp. Med. Biol.* **133**: 477-506.
- Lauder, J.M.; Wallace, J.A.; Krebs, H.; Petrusz, P.; McCarthy, K. (1982) In vivo and in vitro development of serotonergic neurons. *Brain Res. Bull.* **9**: 605-625.
- Leathwood, P.D. (1987) Tryptophan availability and serotonin synthesis. *Proc. Nutr. Soc.* **46**: 143-156.
- Lebrand, C.; Cases, O.; Adelbrecht, C.; Doye, A.; Alvarez, C.; El Mestikawy, S.; Seif, I.; Gaspar, P. (1996) Transient uptake and storage of serotonin in developing thalamic neurons. *Neuron* **17**: 823-835.
- Leclerc, C.; Guerrier, P.; Moreau, M. (2000) Role of dihydropyridine-sensitive calcium channels in meiosis and fertilization in the bivalve molluscs *Ruditapes philippinarum* and *Crassostrea gigas*. *Biol. Cell.* **92**(3-4): 285-299.
- Ledley, F.D.; Grenett, H.E.; Dunbar, B.S., Woo, S.L.C. (1990) Mouse Phenylalanine hydroxylase. Homology and divergence from human phenylalanine hydroxylase. *Biochem. J.* **267**: 399-406.
- Ledley, F.D.; van Tuinen, P.; Ledbetter, D.; Gerhardt, T.; Jones, C.; Woo, S.L.C. (1987) Assignment of tryptophan hydroxylase to human chromosome 11p: Duplication and rearrangement in aromatic amino acid hydroxylase evolution. Human gene mapping. *Cytogenet. Cell Genet.* **46**: 646.
- Leibowitz, S.F.; Alexander, J.T. (1998) Hypothalamic serotonin in control of eating behavior, meal size, and body weight. *Biol. Psychiatry* **44**(9): 851-864
- Lesch, K.P.; Beckmann, H. (1990) The serotonin hypothesis of depression. *Fortschr. Neurol. Psychiatr.* **58**(11): 427-438.
- Lesch, K.P.; Zeng, Y.; Reif, A.; Gutknecht, L. (2003) Anxiety-related traits in mice with modified genes of the serotonergic pathway. *Eur. J. Pharmacol.* **480**: 185-204.

- Leung, R.K.; Whittaker, P.A. (2005) RNA interference: from gene silencing to gene-specific therapeutics. *Pharmacol. Ther.* **107**(2): 222-239.
- Levine, R.A.; Kuhn, D.M.; Lovenberg, W. (1979) The regional distribution of hydroxylase cofactor in rat brain. *J. Neurochem.* **32**: 1575-1578.
- Levine, R.A.; Miller, L.P.; Lovenberg, W. (1981) Tetrahydrobiopterin in striatum. *Science* **214**: 919-921.
- Levitt, P.; Rakic, P. (1982) The time of genesis, embryonic origin and differentiation of the brain stem monoamine neurons in the rhesus monkey. *Brain Res.* **256**(1): 35-57.
- L'Huillier, P.J.; Soulier, S.; Stinnakre, M.G.; Lepourry, L.; Davis, S.R.; Mercier, J.C.; Villette, J.L. (1996) Efficient and specific ribozyme-mediated reduction of bovine alpha-lactalbumin expression in double transgenic mice. *Proc. Natl. Acad. Sci. U. S. A.* **93**(13): 6698-6703.
- Lidov, H.G.W.; Molliver, M.E. (1982a) An immunohistochemical study of serotonin neuron development in the rat: ascending pathways and terminal fields. *Brain Res. Bull.*, 8: 389-430.
- Lidov, H.G.W.; Molliver, M.E. (1982b) Immunohistochemical study of the development of serotonergic neurons in the rat CNS. *Brain Res. Bull.* **9**: 559-604.
- Lieber, A.; Strauss, M. (1995) Selection of efficient cleavage sites in target RNAs by using a ribozyme expression library. *Mol. Cell. Biol.* **15**(1): 540-551.
- Lin, X.; Ruan, X.; Anderson, M.G.; McDowell, J.A.; Kroeger, P.E.; Fesik, S.W.; Shen, Y. (2005) siRNA-mediated off-target gene silencing triggered by a 7 nt complementation. *Nucleic Acids Res.* **33**(14): 4527-4535.
- Lipton, R.B.; Scher, A.I.; Kolodner, K.; Liberman, J.; Steiner, T.J.; Stewart, W.F. (2002) Migraine in the United States: epidemiology and patterns of health care use. *Neurology* **58**(6): 885-894.
- Lira, A.; Zhou, M.; Castanon, N.; Ansorge, M.S.; Gordon, J.A.; Francis, J.H.; Bradley-Moore, M.; Lira, J.; Underwood, M.D.; Arango, V.; Kung, H.F.; Hofer, M.A.; Hen, R.; Gingrich, J.A. (2003) Altered depression-related behaviors and functional changes in the dorsal raphe nucleus of serotonin transporter-deficient mice. *Biol. Psychiatry* **54**(10): 960-971.
- Liu, X; Vrana, K.E. (1991) Leucine zippers and coiled-coils in the aromatic amino acid hydroxylases. *Neurochem. Int.* **18**: 27-31.
- Lovenberg, W.; Jequier, E.; Sjoerdsma, A. (1967) Tryptophan hydroxylation: measurement in pineal gland, brainstem, and carcinoid tumor. *Science* **155**(759): 217-219.
- Lovenberg, W.; Jequier, E.; Sjoerdsma, A. (1968) Tryptophan hydroxylation in mammalian systems. *Advan. Pharmacol.* **6A**: 21-25.
- Lovenberg, W.; Victor, S.J. (1974) Regulation of tryptophan and tyrosine hydroxylase. *Life Sci.* **14**: 2337-2353.

- Luca, V.D.; Likhodi, O.; Van Tol, H.H.; Kennedy, J.L.; Wong, A.H. (2005) Gene expression of tryptophan hydroxylase 2 in post-mortem brain of suicide subjects. *Int. J. Neuropsychopharmacol.* **1-5** [Epub ahead of print].
- Ludecke, B.; Dworniczak, B.; Bartholome, K. (1996) Recessively inherited L-DOPA responsive parkinsonism in infancy caused by a point mutation (L205P) in the tyrosine hydroxylase gene. *Hum. Mol. Gen.* **7**: 1023-1028.
- Luyckx, V.A.; Leclercq, B.; Dowland, L.K.; Yu, A.S. (1999) Diet-dependent hypercalciuria in transgenic mice with reduced CLC5 chloride channel expression. *Proc. Natl. Acad. Sci. U. S. A.* **96**(21): 12174-12179.
- Lyngstadaas, S.P.; Risnes, S.; Sproat, B.S.; Thrane, P.S.; Prydz, H.P. (1995) A synthetic, chemically modified ribozyme eliminates amelogenin, the major translation product in developing mouse enamel in vivo. *EMBO J.* **14**(21): 5224-5229.
- Mach, T. (2004) The brain-gut axis in irritable bowel syndrome--clinical aspects. *Med. Sci. Monit.* **10**(6): RA125-131.
- Majzoub, J.A.; Muglia, L.J. (1996) Molecular medicine. Knockout mice. *New Eng. J. Med.* 904-907.
- Makita, Y.; Okuno, S.; Fujisawa, H. (1990) Involvement of activator protein in the activation of tryptophan hydroxylase by cAMP-dependent protein kinase. *FEBS Lett.* **268**: 185-188.
- Malek, Z.S.; Dardente, H.; Pevet, P.; Raison, S. (2005) Tissue-specific expression of tryptophan hydroxylase mRNAs in the rat midbrain: anatomical evidence and daily profiles. *Eur. J. Neurosci.* **22**(4): 895-901.
- Malyszko, J.; Malyszko, J.S.; Pawlak, K.; Buczko, W.; Mysliwiec, M. (1996b) Hemostasis, platelet function and serotonin in acute and chronic renal failure. *Thromb. Res.* **83**: 351-361.
- Malyszko, J.; Malyszko, J.S.; Pawlak, K.; Rydzewski, A.; Mysliwiec, M.; Buczko, W. (1996a) Comprehensive study on platelet function, hemostasis, fibrinolysis, peripheral serotonergic system and serum lipids in nephrotic syndrome. *Pol. J. Pharmacol.* **48**: 191-195.
- Mandel, M.; Higa, A. (1979) Calcium dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**: 154-162.
- Mann, R.; Mulligan, R.C.; Baltimore, D. (1983) Construction of a retrovirus packaging mutant and its use to produce helper-free defective retrovirus. *Cell* **33**(1): 153-159.
- Markey, K.A.; Kondo, S.; Shenkman, L.; Goldstein, M. (1980) Purification and characterization of tyrosine hydroxylase from a clonal pheochromocytoma cell line. *Mol. Pharmacol.* **17**: 79-85.
- Maskos, Z.; Rush, J.D.; Koppenol, W.H. (1992) The hydroxylation of tryptophan. *Arch. Biochem. Biophys.* **296**(2): 514-520.

- Mathews, D.H.; Sabina, J.; Zuker, M.; Turner, D.H. (1999) Expanded Sequence Dependence of Thermodynamic Parameters Improves Prediction of RNA Secondary Structure. *J. Mol. Biol.* **288**: 911-940
- Matsuda, M.; Imaoka, T.; Vomachka, A.J.; Gudelsky, G.A.; Hou, Z.; Mistry, M.; Bailey, J.P.; Nieport, K.M.; Walther, D.J.; Bader, M.; Horseman, N.D. (2004) Serotonin regulates mammary gland development via an autocrine-paracrine loop. *Dev. Cell* **6**(2): 193-203.
- McKinney, J.; Knappskog, P.M.; Haavik, J. (2005) Different properties of the central and peripheral forms of human tryptophan hydroxylase. *J. Neurochem.* **92**(2): 311-320.
- Meek, J.L.; Neckers, L.M. (1975) Measurement of tryptophan hydroxylase in single brain nuclei by high pressure liquid chromatography. *Brain Res.* **91**: 336-340.
- Meeusen, R.; Thorre, K.; Chaouloff, F.; Sarre, S.; De Meirlier, K.; Ebinger, G.; Michotte, Y. (1996) Effects of tryptophan and/or acute running on extracellular 5-HT and 5-HIAA levels in the hippocampus of food-deprived rats. *Brain Res.* **740**(1-2): 245-252.
- Melo, L.G.; Pachori, A.S.; Kong, D.; Gnechi, M.; Wang, K.; Pratt, R.E.; Dzau, V.J. (2004) Molecular and cell-based therapies for protection, rescue, and repair of ischemic myocardium: reasons for cautious optimism. *Circulation* **109**(20): 2386-2393.
- Mena, F.; Enjalbert, A.; Carbonell, L.; Priam, M.; Kordan, C. (1976) Effect of suckling on plasma prolactin and hypothalamic monoamine levels in the rat. *Endocrinology* **99**(2): 445-451.
- Miller, A.D.; Buttimore, C. (1986) Redesign of retrovirus packaging cell lines to avoid recombination leading to helper virus production. *Mol. Cell Biol.* **6**(8): 2895-2902.
- Mistry, S.J.; Benham, C.J.; Atweh, G.F. (2001) Development of ribozymes that target stathmin, a major regulator of the mitotic spindle. *Antisense Nucleic Acid Drug Dev.* **11**(1): 41-49.
- Miwa, S.; Watanabe, Y.; Hayaishi, O. (1985) 6R-L-erythro-5,6,7,8-tetrahydrobiopterin as a regulator of dopamine and serotonin biosynthesis in the rat brain. *Arch. Biochem. Biophys.* **239**: 234-241.
- Mockus, S.M.; Vrana, K.E. (1998) Advances in the molecular characterization of tryptophan hydroxylase. *J. Mol. Neurosci.* **10**: 163-179.
- Modlin, I.M.; Sandor, A. (1997) An analysis of 8305 cases of carcinoid tumors. *Cancer* **79**(4): 813-829.
- Molliver, M.E. (1987) Serotonergic neuronal systems: what their anatomic organization tells us about function. *J. Clin. Psychopharmacol.* **7**(6 Suppl.): 3S-23S.
- Moran, G.R.; Daubner, S.C.; Fitzpatrick, P.F. (1998) Expression and characterization of the catalytic core of tryptophan hydroxylase. *J. Biol. Chem.* **273**: 12259-12266.
- Moskowitz, D.S.; Pinard, G.; Zuroff, D.C.; Annable, L.; Young, S.N. (2001) The effect of tryptophan on social interaction in everyday life: a placebo-controlled study. *Neuropsychopharmacology* **25**(2): 277-289.

- Mossner, R.; Lesch, K.P. (1998) Role of serotonin in the immune system and in neuroimmuno interactions. *Brain Behav. Immun.* **12**(4): 249-271.
- Munn, D.H.; Sharma, M.D.; Lee, J.R.; Jhaver, K.G.; Johnson, T.S.; Keskin, D.B.; Marshall, B.; Chandler, P.; Antonia, S.J.; Burgess, R.; Slingluff, C.L.Jr.; Mellor, A.L. (2002) Potential regulatory function of human dendritic cells expressing indoleamine 2,3-dioxygenase. *Science* **297**(5588): 1867-1870.
- Murphy, D.L.; Zohar, J.; Benkelfat, C.; Pato, M.T.; Pigott, T.A.; Insel, T.R. (1989) Obsessive-compulsive disorder as a 5-HT subsystem-related behavioural disorder. *Br. J. Psychiatry Suppl.* **8**: 15-24.
- Nakata, H.; Fujisawa, H. (1982a) Purification and properties of tryptophan 5-monooxygenase from rat brain stem. *Eur. J. Biochem.* **122**: 41-47.
- Nakata, H.; Fujisawa, H. (1982b) Tryptophan 5-monooxygenase from mouse mastocytoma P815. A simple purification and general properties. *Eur. J. Biochem.* **124**: 595-601.
- Narlikar, G.J.; Herschlag, D. (1997) Mechanistic aspects of enzymatic catalysis: lessons from comparison of RNA and protein enzymes. *Annu. Rev. Biochem.* **66**: 19-59.
- Nathwani, A.C.; Benjamin, R.; Nienhuis, A.W.; Davidoff, A.M. (2004) Current status and prospects for gene therapy. *Vox Sang.* **87**(2): 73-81.
- Nazar, M.; Siemiatkowski, M.; Bidzinski, A.; Czlonkowska, A.; Sienkiewicz-Jarosz, H.; Plaznik, A. (1999) The influence of serotonin depletion on rat behavior in the Vogel test and brain 3H-zolpidem binding. *J. Neural. Transm.* **106**(5-6): 355-368.
- Neary, P.C.; Redmond, P.H.; Houghton, T.; Watson, G.R.K.; Bouchier-Hayes, D. (1997) Carcinoid disease: review of the literature. *Dis. Colon Rectum* **40**: 349-362.
- Nebigil, C.G.; Choi, D.S.; Dierich, A.; Hickel, P.; Le Meur, M.; Messaddeq, N.; Launay, J.M.; Maroteaux, L. (2000) Serotonin 2B receptor is required for heart development. *Proc. Natl. Acad. Sci. U. S. A.* **97**(17): 9508-9513.
- Newman, C.; Wang, D.; Cutz, E. (1993) Serotonin (5-hydroxytryptamine) expression in pulmonary neuroendocrine cells (NE) and a netumor cell line. *Adv. Exp. Med. Biol.* **337**: 73-78
- Nonogaki, K.; Strack, A.M.; Dallman, M.F.; Tecott, L.H. (1998) Leptin-independent hyperphagia and type 2 diabetes in mice with a mutated serotonin 5-HT2C receptor gene. *Nat. Med.* **4**(10): 1152-1156.
- Numano, H. (1953) Studies on the fertilization substance. IV. Presence of acetylcholine-like substance and cholinesterase in echinoderm germ cells during fertilization. *Sci. Papers Coll. Gen. Educ. Univ. Tokyo* **3**: 193-200.
- Oberg, K. (1994) Endocrine tumors of the gastrointestinal tract: systemic treatment. *Anticancer Drugs* **5**: 503-519.

- Offermanns, S.; Toombs, C.F.; Hu, Y.H.; Simon, M.I. (1997) Defective platelet activation in G alpha(q)-deficient mice. *Nature* **389**: 183–186
- O'Kane, R.L.; Vina, J.R.; Simpson, I.; Hawkins, R.A. (2004) Na⁺-dependent neutral amino acid transporters A, ASC, and N of the blood-brain barrier: mechanisms for neutral amino acid removal. *Am. J. Physiol. Endocrinol. Metab.* **287**(4): E622-629.
- Olson, J.M. (1970) Evolution of Photosynthesis. *Science* **168**: 438-446
- Onishi, A.; Liotta, L.J.; Benkovic, S.J. (1991) Cloning and expression of Chromobacterium violaceum phenylalanine hydroxylase in Escherichia coli and comparison of amino acid sequence with mammalian aromatic amino acid hydroxylases. *J. Biol. Chem.* **266**: 18454-18459.
- Ormsbee III, H.S.; Fondacaro, J.D. (1985) Action of serotonin on the gastrointestinal tract. *Proc. Soc. Exp. Biol. Med.* **178**: 333-338.
- Paddison, P.J.; Caudy, A.A.; Hannon, G.J. (2002) Stable suppression of gene expression by RNAi in mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* **99**(3): 1443-1448.
- Palmer, D.S.; Aye, M.T.; Ganz, P.R.; Halpenny, M.; Hashemi, S. (1994) Adenosine nucleotides and serotonin stimulate von Willebrand factor release from cultured human endothelial cells. *Thromb. Haemost.* **72**: 132-139.
- Pannell, D.; Osborne, C.S.; Yao, S.; Sukonnik, T.; Pasceri, P.; Karaiskakis, A.; Okano, M.; Li, E.; Lipshitz, H.D.; Ellis, J. (2000) Retrovirus vector silencing is de novo methylase independent and marked by a repressive histone code. *EMBO J.* **19**(21): 5884-5894.
- Parsons, B.; Roxas, A.Jr.; Huang, Y.Y.; Dwork, A.; Stanley, M. (1992) Regional studies of serotonin and dopamine metabolism and quantification of serotonin uptake sites in human cerebral cortex. *J. Neural. Transm. Gen. Sect.* **87**(1): 63-75.
- Patel, P.D.; Pontrello, C.; Burke, S. (2004) Robust and tissue-specific expression of TPH2 versus TPH1 in rat raphe and pineal gland. *Biol. Psychiatry* **55**(4): 428-433.
- Pelley, R.J.; Bukowski, R.M. (1997) Recent advances in diagnosis and therapy of neuroendocrine tumors of the gastrointestinal tract. *Curr. Opin. Oncol.* **9**: 68-74.
- Persico, A.M.; Baldi, A.; Dell'Acqua, M.L.; Moessner, R.; Murphy, D.L.; Lesch, K.P.; Keller, F. (2003) Reduced programmed cell death in brains of serotonin transporter knockout mice. *Neuroreport* **14**(3): 341-344.
- Persico, A.M.; Mengual, E.; Moessner, R.; Hall, F.S.; Revay, R.S.; Sora, I.; Arellano, J.; DeFelipe, J.; Gimenez-Amaya, J.M.; Conciatori, M.; Marino, R.; Baldi, A.; Cabib, S.; Pascucci, T.; Uhl, G.R.; Murphy, D.L.; Lesch, K.P.; Keller, F. (2001) Barrel pattern formation requires serotonin uptake by thalamocortical afferents, and not vesicular monoamine release. *J. Neurosci.* **21**(17): 6862-6873.

- Peter, D.; Jimenez, J.; Liu, Y.; Kim, J.; Edwards, R.H. (1994) The chromaffin granule and synaptic vesicle amine transporters differ in substrate recognition and sensitivity to inhibitors. *J. Biol. Chem.* **269**(10): 7231-7237.
- Peter, D.; Liu, Y.; Sternini, C.; de Giorgio, R.; Brecha, N.; Edwards, R.H. (1995) Differential expression of two vesicular monoamine transporters. *J. Neurosci.* **15**(9): 6179-6188.
- Pezawas, L.; Meyer-Lindenberg, A.; Drabant, E.M.; Verchinski, B.A.; Munoz, K.E.; Kolachana, B.S.; Egan, M.F.; Mattay, V.S.; Hariri, A.R.; Weinberger, D.R. (2005) 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nat. Neurosci.* **8**(6): 828-834.
- Pratt, H.L.; Carroll, R.C.; Jones, J.B.; Lothrop, C.D. (1991) Platelet aggregation, storage pool deficiency, and protein phosphorylation in mice with Chediak-Higashi syndrome. *Am. J. Vet. Res.* **52**: 945-950.
- Preisig, M.; Ferrero, F.; Malafosse, A. (2005) Monoamine oxidase a and tryptophan hydroxylase gene polymorphisms: are they associated with bipolar disorder? *Am. J. Pharmacogenomics* **5**(1): 45-52.
- Pyle, A.M.; Green, J.B. (1995) RNA folding. *Curr. Opin. Struct. Biol.* **5**(3): 303-310.
- Raleigh, M.J.; Brammer, G.L.; Yuwiler, A.; Flannery, J.W.; McGuire, M.T.; Geller, E. (1980) Serotonergic influences on the social behavior of vervet monkeys (*Cercopithecus aethiops sabaeus*). *Exp. Neurol.* **68**(2): 322-334.
- Raleigh, M.J.; McGuire, M.T. (1991) Bidirectional relationships between tryptophan and social behavior in vervet monkeys. *Adv. Exp. Med. Biol.* **294**: 289-298.
- Raleigh, M.J.; McGuire, M.T.; Brammer, G.L.; Pollack, D.B.; Yuwiler, A. (1991) Serotonergic mechanisms promote dominance acquisition in adult male vervet monkeys. *Brain Res.* **559**(2): 181-190.
- Rapport, M.M.; Green, A.A.; Page, I.H. (1947) Purification of the substance which is responsible for the vasoconstrictor activity of serum. *Fed. Proc.* **6**: 184.
- Rapport, M.M.; Green, A.A.; Page, I.H. (1948) Serum vasoconstrictor (serotonin). III. Chemical inactivation. *J. Biol. Chem.* **176**: 1237.
- Raymond, J.R.; Mukhin, Y.V.; Gelasco, A.; Turner, J.; Collinsworth, G.; Gettys, T.W.; Grewal, J.S.; Garnovskaya, M.N. (2001) Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacol. Ther.* **92**(2-3): 179-212.
- Reader, T.A.; Le Marec, N.; Ase, A.R.; Lalonde, R. (1999) Effects of L-tryptophan on indoleamines and catecholamines in discrete brain regions of wild type and Lurcher mutant mice. *Neurochem. Res.* **24**(9): 1125-1134.

- Rebsam, A.; Seif, I.; Gaspar, P. (2002) Refinement of thalamocortical arbors and emergence of barrel domains in the primary somatosensory cortex: a study of normal and monoamine oxidase a knock-out mice. *J. Neurosci.* **22**(19): 8541-8552.
- Reed, G.E.; Kirchner, J.E.; Carr, L.G. (1995) NF-Y activates mouse tryptophan hydroxylase transcription. *Brain Res.* **682**: 1-12.
- Renaud, F.; Parisi, E.; Capasso, A.; De Prisco, P. (1983) On the role of serotonin and 5-methoxytryptamine in the regulation of cell division in sea urchin eggs. *Dev. Biol.* **98**(1): 37-46.
- Rhoades, R.W.; Mooney, R.D.; Chiaia, N.L.; Bennett-Clarke, C.A. (1990) Development and plasticity of the serotonergic projection to the hamster's superior colliculus. *J. Comp. Neurol.* **299**: 151-166.
- Ribeiro, P.; Wang, Y.; Citron, B.A.; Kaufman, S. (1993) Deletion mutagenesis of rat PC12 tyrosine hydroxylase regulatory and catalytic domains. *J. Mol. Neurosci.* **4**: 125-139.
- Robertson, G.; Garrick, D.; Wu, W.; Kearns, M.; Martin, D. (1995) Position-dependent variegation of globin transgene expression in mice. *Proc. Natl. Acad. Sci. U. S. A.* **92**: 5371-5375.
- Römpf Online Lexikon (2004) Serotonin. *Georg Thieme Verlag*, Version 2.2.
- Rosa-Neto, P.; Diksic, M.; Okazawa, H.; Leyton, M.; Ghadirian, N.; Mzengeza, S.; Nakai, A.; Debonnel, G.; Blier, P.; Benkelfat, C. (2004) Measurement of brain regional alpha-[11C]methyl-L-tryptophan trapping as a measure of serotonin synthesis in medication-free patients with major depression. *Arch. Gen. Psychiatry* **61**(6): 556-563.
- Rotman, A. (1977) The mechanism of action of neurocytotoxic compounds. *Life Sci.* **21**(7): 891-900
- Salter, M.; Knowles, R.G.; Pogson, C.I. (1989) How does displacement of albumin-bound tryptophan cause sustained increases in the free tryptophan concentration in the plasma and 5-hydroxytryptamine synthesis in brain? *Biochem. J.* **262**: 365-368.
- Sambrook, J.; Fritsch, E.F.; Maniatis, T. (1989) Molecular cloning: a laboratory manual. 2nd edn.: 771pp.
- Sanchez, R.L.; Reddy, A.P.; Centeno, M.L.; Henderson, J.A.; Bethea, C.L. (2005) A second tryptophan hydroxylase Isoform, TPH-2 mRNA, is increased by ovarian steroids in the raphe region of macaques. *Mol. Brain Res.* **135**(1-2): 194-203.
- Santarelli, L.; Saxe, M.; Gross, C.; Surget, A.; Battaglia, F.; Dulawa, S.; Weisstaub, N.; Lee, J.; Duman, R.; Arancio, O.; Belzung, C.; Hen, R. (2003) Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* **301**(5634): 805-809.
- Santoso, D.I.; Rogers, P.; Wallace, E.M.; Manelpillai, U.; Walker, D.; Subakir, S.B. (2002) Localization of indoleamine 2,3-dioxygenase and 4-hydroxynonenal in normal and pre-eclamptic placentae. *Placenta* **23**(5): 373-379.

- Sarna, G.S.; Hutson, P.H.; Tricklbank, M.D.; Curzon, G. (1983) Determination of brain 5-hydroxytryptamine turnover in freely moving rats using repeated sampling of cerebrovascular fluid. *J. Neurochem.* **40**(2): 383-388.
- Saudou, F.; Amara, D.A.; Dierich, A.; LeMeur, M.; Ramboz, S.; Segu, L.; Buhot, M.C.; Hen, R. (1994) Enhanced aggressive behavior in mice lacking 5-HT1B receptor. *Science* **265**(5180): 1875-1878.
- Sawada, M.; Sugimoto, T.; Matsuura, S.; Nagatsu, T. (1986) (6R)-tetrahydrobiopterin increases the activity of tryptophan hydroxylase in rat raphe slices. *J. Neurochem.* **47**: 1544-1547.
- Saxena, S.; Jonsson, Z.O.; Dutta, A. (2003) Small RNAs with imperfect match to endogenous mRNA repress translation: implications for off-target activity of siRNA in mammalian cells. *J. Biol. Chem.* **278**(45): 44312-44319.
- Scacheri, P.C.; Rozenblatt-Rosen, O.; Caplen, N.J.; Wolfsberg, T.G.; Umayam, L.; Lee, J.C.; Hughes, C.M.; Shanmugam, K.S.; Bhattacharjee, A.; Meyerson, M.; Collins, F.S. (2004) Short interfering RNAs can induce unexpected and divergent changes in the levels of untargeted proteins in mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* **101**(7): 1892-1897.
- Scherer, L.J.; Rossi, J.J. (2003) Approaches for the sequence-specific knockdown of mRNA. *Nat. Biotechnol.* **21**(12): 1457-1465.
- Schikowski, A.; Thewissen, M.; Mathis, C.; Ross, H.G.; Enck, P. (2002) Serotonin type-4 receptors modulate the sensitivity of intramural mechanoreceptive afferents of the cat rectum. *Neurogastroenterol. Motil.* **14**(3): 221-227.
- Schuller, H.M.; Hegedus, T.J. (1989) Effects of endogenous and tobacco-related amines and nitrosamines on cell growth and morphology of a cell line derived from a human neuroendocrine lung cancer. *Toxicol. In Vitro* **31**: 37-43.
- Shamir, A.; Shaltiel, G.; Levi, I.; Belmaker, R.H.; Agam, G. (2005) Postmortem Parietal Cortex TPH2 Expression Is Not Altered in Schizophrenic, Unipolar-Depressed, and Bipolar Patients vs Control Subjects. *J. Mol. Neurosci.* **26**(1): 33-38.
- Sheehan, K.; Lowe, N.; Kirley, A.; Mullins, C.; Fitzgerald, M.; Gill, M.; Hawi, Z. (2005) Tryptophan hydroxylase 2 (TPH2) gene variants associated with ADHD. *Mol. Psychiatry* [Epub ahead of print].
- Shelburne, C.P.; Huff, T.F. (1999) Inhibition of kit expression in P815 mouse mastocytoma cells by a hammerhead ribozyme. *Clin. Immunol.* **93**(1): 46-58.
- Shulkin, B.L., Betz, A.L., Koeppen, R.A., Agranoff, B.W. (1995) Inhibition of neutral amino acid transport across the human blood-brain barrier by phenylalanine. *J. Neurochem.* **64**(3): 1252-1257.
- Siegel, G.J.; Agranoff, B.W.; Fisher, S.K.; Albers, R.W.; Uhler, M.D. (1999) Basic Neurochemistry: Molecular, Cellular and Medical Aspects. Sixth Edition. *American Society for Neurochemistry*.

- Sioud, M.; Natvig, J.B.; Forre, O. (1992) Preformed ribozyme destroys tumour necrosis factor mRNA in human cells. *J. Mol. Biol.* **223**(4): 831-835.
- Snyder, S.H. (2002) Neurobiology: serotonin sustains serenity. *Nature* **416**(6879): 377-380.
- Sole, M.J.; Madapallimattam, A.; Baines, A.D. (1986) An active pathway for serotonin synthesis by renal proximal tubules. *Kidney Int.* **29**: 689-694.
- Soriano, P.; Jaenisch, R. (1986) Retroviruses as probes for mammalian development: allocation of cells to the somatic and germ cell lineages. *Cell* **46**(1): 19-29.
- Soslau, G.; Silverberg, M.; Brodsky, I.; McCarty, P.P. (1997) ATP modification of serotonin-induced contraction of the rat pulmonary artery. *Proc. Soc. Exp. Biol. Med.* **214**: 233-241.
- Spink, J.; Geddes, D. (2004) Gene therapy progress and prospects: bringing gene therapy into medical practice: the evolution of international ethics and the regulatory environment. *Gene Ther.* **11**(22): 1611-1616.
- Statham, S.; Morgan, R.A. (1999) Gene therapy clinical trials for HIV. *Curr. Opin. Mol. Ther.* **1**(4): 430-436.
- Steinman, J.L.; Carlton, S.M.; Haber, B.; Willis, W.D. (1987) Differential effects of p-chlorophenylalanine on indoleamines in brainstem nuclei and spinal cord of rats. I. Biochemical and behavioral analysis. *Brain Res.* **426**(2): 297-309.
- Stoll, J.; Goldman, D. (1991) Isolation and structural characterization of the murine tryptophan hydroxylase gene. *J. Neurosci. Res.* **28**: 457-465.
- Stoll, J.; Kozak, C.A.; Goldman, D. (1990) Characterization and chromosomal mapping of a cDNA encoding tryptophan hydroxylase from a mouse mastocytoma cell line. *Genomics* **7**(1): 88-96.
- Suzuki, O.; Hattori, H.; Sawada, M.; Nagatsu, T.; Miki, N.; Higashida, H. (1983) Serotonin in neuroblastoma x glioma NG108-15 hybrid cells. *Neurochem. Int.* **5**: 599-601.
- Swindle, C.S.; Kim, H.G.; Klug, C.A. (2004) Mutation of CpGs in the murine stem cell virus retroviral vector long terminal repeat represses silencing in embryonic stem cells. *J. Biol. Chem.* **279**(1): 34-41.
- Szasz, R., Dale, G.L. (2002) Thrombospondin and fibrinogen bind serotonin-derivatized proteins on COAT-platelets. *Blood* **100**(8): 2827-2831.
- Tabatabaie, T.; Goyal, R.N.; Blank, C.L.; Dryhurst, G. (1993) Further insights into the molecular mechanisms of action of the serotonergic neurotoxin 5,7-dihydroxytryptamine. *J. Med. Chem.* **36**: 229-236.
- Tecott, L.H.; Sun, L.M.; Akana, S.F.; Strack, A.M.; Lowenstein, D.H.; Dallman, M.F.; Julius, D. (1995) Eating disorder and epilepsy in mice lacking 5-HT_{2C} serotonin receptors. *Nature* **374**: 542-546.

- Thompson, J.D.; Ayers, D.F.; Malmstrom, T.A.; McKenzie, T.L.; Ganousis, L.; Chowrira, B.M.; Couture, L.; Stinchcomb, D.T. (1995) Improved accumulation and activity of ribozymes expressed from a tRNA-based RNA polymerase III promoter. *Nucl. Acids Res.* **23**(12): 2259-2268.
- Tong, J.H.; Kaufman, S. (1975) Tryptophan hydroxylase. Purification and some properties of the enzyme from rabbit hindbrain. *J. Biol. Chem.* **250**: 4152-4158.
- Törk, I. (1990) Anatomy of the serotonergic system. *Ann. N. Y. Acad. Sci.* **600**: 9-35
- Toth, G.; Fekete, M. (1986) 5-Hydroxyindole acetic acid excretion in newborns, infants and children. *Acta Paediatr. Hung.* **27**: 221-226.
- Tuschl, T.; Eckstein, F. (1993) Hammerhead ribozymes: importance of stem-loop II for activity. *Proc. Natl. Acad. Sci. U. S. A.* **90**(15): 6991-6994.
- Tutton, P.J.; Barkla, D.H. (1978) Evaluation of the cytotoxicity of dihydroxytryptamines and 5-hydroxytryptamine antagonists as cytotoxic agents in dimethylhydrazine-induced adenocarcinomata. *Cancer Chemother. Pharmacol.* **1**(4): 209-213.
- Twarog, B.M.; Page, I.H. (1953) Serotonin content of some mammalian tissues and urine and a method for its determination. *Am. J. Physiol.* **175**(1): 157-161.
- Udenfriend, S.; Clark, C.T.; Axelrod, J.; Brodie, B.B. (1954) Ascorbic acid in aromatic hydroxylation. I. A model system for aromatic hydroxylation. *J. Biol. Chem.* **208**(2): 731-739.
- Ummadi, M.; Weimer, B.C. (2001) Tryptophan Catabolism in *Brevibacterium linens* as a Potential Cheese Flavor Adjunct. *J. Dairy Sci.* **84**: 1773-1782
- Upton, A.L.; Ravary, A.; Salichon, N.; Moessner, R.; Lesch, K.P.; Hen, R.; Seif, I.; Gaspar, P. (2002) Lack of 5-HT(1B) receptor and of serotonin transporter have different effects on the segregation of retinal axons in the lateral geniculate nucleus compared to the superior colliculus. *Neuroscience* **111**(3): 597-610.
- Upton, A.L.; Salichon, N.; Lebrand, C.; Ravary, A.; Blakely, R.; Seif, I.; Gaspar, P. (1999) Excess of serotonin (5-HT) alters the segregation of ipsilateral and contralateral retinal projections in monoamine oxidase A knock-out mice: possible role of 5-HT uptake in retinal ganglion cells during development. *J. Neurosci.* **19**(16): 7007-7024.
- Vicaut, E.; Laemmel, E.; Stucker, O. (2000) Impact of serotonin on tumour growth. *Ann. Med.* **32**(3): 187-194.
- Villalon, C.M.; Centurion, D.; Valdivia, L.F.; de Vries, P.; Saxena, P.R. (2003) Migraine: pathophysiology, pharmacology, treatment and future trends. *Curr. Vasc. Pharmacol.* **1**(1): 71-84.
- Vitalis, T.; Cases, O.; Callebert, J.; Launay, J.M.; Price, D.J.; Seif, I.; Gaspar, P. (1998) Effects of monoamine oxidase A inhibition on barrel formation in the mouse somatosensory cortex: Determination of a sensitive developmental period. *J. Comp. Neurol.* **393**: 169-184.

- Vitalis, T.; Parnavelas, J.G. (2003) The role of serotonin in early cortical development. *Dev. Neurosci.* **25**(2-4): 245-256.
- Voog, L.; Eriksson, T. (1982) Is rat brain content of large neutral aminiacids (LNAs) a reflection of plasma LNAA concentrations? *J. Neural. Transm. Gen. Sect.* **87**(2) 133-143.
- Vrana, K.E.; Rucker, P.J.; Kumer, S.C. (1994) Recombinant rabbit tryptophan hydroxylase is a substrate for cAMP-dependent protein kinase. *Life Sci.* **55**: 1045-1052.
- Wald, N.J.; Law, M.R. (2003) A strategy to reduce cardiovascular disease by more than 80%. *BMJ.* **326**(7404): 1419.
- Walitza, S.; Renner, T.J.; Dempfle, A.; Konrad, K.; Wewetzer, C.; Halbach, A.; Herpertz-Dahlmann, B.; Remschmidt, H.; Smidt, J.; Linder, M.; Flierl, L.; Knolker, U.; Friedel, S.; Schafer, H.; Gross, C.; Hebebrand, J.; Warnke, A.; Lesch, K.P. (2005) Transmission disequilibrium of polymorphic variants in the tryptophan hydroxylase-2 gene in attention-deficit/hyperactivity disorder. *Mol. Psychiatry* [Epub ahead of print].
- Walker, S.J.; Liu, X.; Roskoski, Jr. R.; Vrana, K.E. (1994) Catalytic core of rat tyrosine hydroxylase: Terminal deletion analysis of bacterially-expressed enzyme. *Biochim. Biophys. Acta* **1206**: 113-119.
- Wall, R.J. (1996) Transgenic livestock: progress and prospects for the future. *Theriogenology* **45**: 57-68.
- Walsh, H.A.; Daya, S. (1997) Inhibition of hepatic tryptophan-2,3-dioxygenase: Superior potency of melatonin over serotonin. *J. Pineal Res.* **23**: 20-23.
- Walther, D.J.; Bader, M. (1998) Tryptophan Hydroxylase expression and serotonin synthesis in murine embryonic stem cells. *Biol. Chem.* **379**: S35.
- Walther, D.J.; Bader, M. (1999) Serotonin synthesis in murine embryonic stem cells. *Mol. Brain Res.* **68**: 55-63.
- Walther, D.J.; Kusserow, H.; Hörtnagl, H.; Theuring, F.; Bader, M. (1998) Serotonin 5-HT1A receptor expression and tryptophan hydroxylase activity in neuroblastoma x glioma hybrid NG108-15. *Eur. J. Neurosci.* **10**: 173.
- Walther, D.J.; Peter, J.U.; Bader, M. (2002) 7-Hydroxytryptophan, a novel, specific, cytotoxic agent for carcinoids and other serotonin-producing tumors. *Cancer* **94**(12): 3135-3140.
- Walther, D.J.; Peter, J.U.; Bashammakh, S.; Hortnagl, H.; Voits, M.; Fink, H.; Bader, M. (2003a) Synthesis of serotonin by a second tryptophan hydroxylase isoform. *Science* **299**(5603): 76.
- Walther, D.J.; Peter, J.U.; Winter, S.; Holtje, M.; Paulmann, N.; Grohmann, M.; Vowinckel, J.; Alamo-Bethencourt, V.; Wilhelm, C.S.; Ahnert-Hilger, G.; Bader, M. (2003b) Serotonylation of small GTPases is a signal transduction pathway that triggers platelet alpha-granule release. *Cell* **115**(7): 851-862.

- Wang, Q.; Liu, M.; Kozasa, T.; Rothestein, J.D.; Sternweis, P.C.; Neubig, R.R. (2004) Ribozyme- and siRNA-mediated suppression of RGS-containing RhoGEF proteins. *Methods Enzymol.* **389**: 244-265.
- Wang, Y.M.; Gainetdinov, R.R.; Fumagalli, F.; Xu, F.; Jones, S.R.; Bock, C.B.; Miller, G.W.; Wightman, R.M.; Caron, M.G. (1997) Knockout of the vesicular monoamine transporter 2 gene results in neonatal death and supersensitivity to cocaine and amphetamine. *Neuron* **19**(6): 1285-1296.
- Webb, A.; Cunningham, D.; Cotter, F.; Clarke, P.A.; di Stefano, F.; Ross, P.; Corbo, M.; Dziewanowska, Z. (1997) BCL-2 antisense therapy in patients with non-Hodgkin lymphoma. *Lancet* **349**(9059): 1137-1141.
- Weber, L.J.; Horita, A. (1965) A study of 5-hydroxytryptamine formation from L-tryptophan in the brain and other tissues. *Biochem. Pharmacol.* **14**: 1141-1149.
- Weihe, E.; Schafer, M.K.; Erickson, J.D.; Eiden, L.E. (1994) Localization of vesicular monoamine transporter isoforms (VMAT1 and VMAT2) to endocrine cells and neurons in rat. *J. Mol. Neurosci.* **5**(3): 149-164.
- Weiner, C.P. (1987) The role of serotonin in the genesis of hypertension in preeclampsia. *Am. J. Obstet. Gynecol.* **156**(4): 885-888.
- Whitaker-Azmitia, P.M. (2001) Serotonin and brain development: role in human developmental diseases. *Brain Res. Bull.* **56**(5): 479-485.
- Williams, J.; Spurlock, G.; McGuffin, P.; Mallet, J.; Nothen, M.M.; Gill, M.; Aschauer, H.; Nylander, P.O.; Macchiardi, F.; Owen, M.J. (1996) Association between schizophrenia and T102C polymorphism of the 5-hydroxytryptamine type 2a-receptor gene. European Multicentre Association Study of Schizophrenia (EMASS) Group. *Lancet* **347**(9011): 1294-1296.
- Winchel, R.M.; Stanley, M. (1991) Self-injurious behavior: a review of the behavior and biology of self-mutilation. *Am. J. Psychiatry* **148**(3): 306-317.
- Wisor, J.P. (2002) Disorders of the circadian clock: etiology and possible therapeutic targets. *Curr. Drug Targets CNS Neurol. Disord.* **1**(6): 555-566.
- Wolf, W.A.; Anastasiadis, P.Z.; Kuhn, D.M.; Levine, R.A. (1990) Influence of tetrahydrobiopterin on serotonin synthesis, metabolism and release in synaptosomes. *Neurochem. Int.* **16**: 335-340.
- Wolf, W.A.; Ziaja, E.; Arthur, Jr. R.A.; Anastasiadis, P.Z.; Levine, R.A.; Kuhn, D.M. (1991) Effect of tetrahydrobiopterin on serotonin synthesis, release, and metabolism in superfused hippocampal slices. *J. Neurochem.* **57**: 1191-1197.
- Wurtman, R.J.; Hefti, F.; Melamed, E. (1981) Precursor control of neurotransmitter synthesis. *Pharmacol. Rev.* **32**: 315-335.
- Yamaguchi, T.; Sawada, M.; Kato, T.; Nagatsu, T. (1981) Demonstration of tryptophan 5-monooxygenase activity in human brain by highly sensitive high-performance liquid chromatography with fluorometric detection. *Biochem. Int.* **2**: 295-303.

- Yamauchi, T.; Fujisawa, H. (1979) Activation of tryptophan 5-monooxygenase by calcium-dependent regulator protein. *Biochem. Biophys. Res. Commun.* **90**(1): 28-35.
- Yamauchi, T.; Fujisawa, H. (1979) Regulation of rat brainstem tryptophan 5-monooxygenase. Calcium dependent reversible activation by ATP and magnesium. *Arch. Biochem. Biophys.* **198**: 219-226.
- Yamauchi, T.; Fujisawa, H. (1983) Purification and characterization of the brain calmodulin-dependent protein kinase (kinase II), which is involved in the activation of tryptophan 5-monooxygenase. *Eur. J. Biochem.* **132**(1): 15-21.
- Yang, X.J.; Kaufman, S. (1994) High-level expression and deletion mutagenesis of human tryptophan hydroxylase. *Proc. Natl. Acad. Sci. U. S. A.* **91**: 6659-6663.
- YohrlingIV, G.J.; Jiang, G.C.; DeJohn, M.M.; Robertson, D.J.; Vrana, K.E.; Cha; J.H. (2002) Inhibition of tryptophan hydroxylase activity and decreased 5-HT1A receptor binding in a mouse model of Huntington's disease. *J. Neurochem.* **82**(6): 1416-1423.
- Yokoi, N.; Kitada, K.; Kuramoto, T.; Maihara, T.; Andoh, Y.; Takami, S.; Higaki, J.; Miki, T.; Ogihara, T.; Lathorp, G.M.; Serikawa, T. (1996) Mapping of 20 polymorphic DNA markers in the rat by somatic hybrid and linkage analysis. *Mamm. Genome* **7**: 71-73.
- Yokoyama, Y.; Morishita, S.; Takahashi, Y.; Hashimoto, M.; Tamaya T. (1997) Modulation of c-fms proto-oncogene in an ovarian carcinoma cell line by a hammerhead ribozyme. *Br. J. Cancer* **76**(8): 977-982.
- Young, M.R.I.; Kut, J.L.; Coogan, M.P.; Wright, M.A.; Young, M.E.; Matthews, J. (1993) Stimulation of splenic T-lymphocyte function by endogenous serotonin and by low-dose exogenous serotonin. *Immunol.* **80**: 395-400.
- Young, S.N.; Leyton, M. (2002) The role of serotonin in human mood and social interaction Insight from altered tryptophan levels. *Pharmacol. Biochem. Behav.* **71**: 857-865.
- Zeitz, K.P.; Guy, N.; Malmberg, A.B.; Dirajlal, S.; Martin, W.J.; Sun, L.; Bonhaus, D.W.; Stucky, C.L.; Julius, D.; Basbaum, A.I. (2002) The 5-HT3 subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. *J. Neurosci.* **22**(3): 1010-1019.
- Zhang, S.; Pohnert, G.; Kongsaeree, P.; Wilson, D.B.; Clardy, J.; Ganem, B. (1998) Chorismate mutase-prephenate dehydratase from Escherichia coli. Study of catalytic and regulatory domains using genetically engeneered proteins. *J. Biol. Chem.* **273**: 6248-6253.
- Zhang, X.; Beaulieu, J.M.; Sotnikova, T.D.; Gainetdinov, R.R.; Caron, M.G. (2004) Tryptophan hydroxylase-2 controls brain serotonin synthesis. *Science* **305**(5681): 217.
- Zhang, X.; Gainetdinov, R.R.; Beaulieu, J.M.; Sotnikova, T.D.; Burch, L.H.; Williams, R.B.; Schwartz, D.A.; Krishnan, K.R.; Caron, M.G. (2005) Loss-of-function mutation in tryptophan hydroxylase-2 identified in unipolar major depression. *Neuron* **45**(1): 11-16.

- Zhao, G.; Xia, T.; Song, J.; Jensen, R.A. (1994) *Pseudomonas aeruginosa* possesses homologues of mammalian phenylalanine hydroxylase and 4alpha-carbinolamine dehydratase/DCoH as part of a three-component gene cluster. *Proc. Natl. Acad. Sci. U. S. A.* **91**: 1366-1370.
- Zhao, R.C.; McIvor, R.S.; Griffin, J.D.; Verfaillie, C.M. (1997) Gene therapy for chronic myelogenous leukemia (CML): a retroviral vector that renders hematopoietic progenitors methotrexate-resistant and CML progenitors functionally normal and nontumorigenic in vivo. *Blood* **90**(12): 4687-4698.
- Zhou, J.M.; Zhou, D.M.; Takagi, Y.; Kasai, Y.; Inoue, A.; Baba, T.; Taira, K. (2002) Existence of efficient divalent metal ion-catalyzed and inefficient divalent metal ion-independent channels in reactions catalyzed by a hammerhead ribozyme. *Nucl. Acids Res.* **30**(11): 2374-82.
- Zill, P.; Baghai, T.C.; Zwanzger, P.; Schule, C.; Eser, D.; Rupprecht, R.; Moller, H.J.; Bondy, B.; Ackenheil, M. (2004a) SNP and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene provide evidence for association with major depression. *Mol. Psychiatry* **9**(11): 1030-1036.
- Zill, P.; Buttner, A.; Eisenmenger, W; Bondy, B; Ackenheil, M. (2004b) Regional mRNA expression of a second tryptophan hydroxylase isoform in postmortem tissue samples of two human brains. *Eur. Neuropsychopharmacol.* **14**(4): 282-284.
- Zill, P.; Buttner, A.; Eisenmenger, W.; Moller, H.J.; Ackenheil, M.; Bondy, B. (2005) Analysis of tryptophan hydroxylase I and II mRNA expression in the human brain: A post-mortem study. *J. Psychiatr. Res.* [Epub ahead of print].
- Zill, P.; Buttner, A.; Eisenmenger, W.; Moller, H.J.; Bondy, B.; Ackenheil, M. (2004c) Single nucleotide polymorphism and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene in suicide victims. *Biol. Psychiatry* **56**(8): 581-586.
- Zuker, M. (2003) Mfold web server for nucleic acid folding and hybridization prediction. *Nucl. Acids Res.* **31**(13): 3406-3415