

## 7 Literaturverzeichnis

- Alberts, B.** (1998) The cell as a collection of protein machines: preparing the next generation of molecular biologists. *Cell*, **92**, 291-294.
- Alliotte, T., Tire, C., Engler, G., Peleman, J., Caplan, A., Van Montagu, M. und Inze, D.** (1989) An auxin-regulated gene of *Arabidopsis thaliana* encodes a DNA-binding protein. *Plant Phys.*, **89**, 743-752.
- Anantharaman, V. und Aravind, L.** (2001) The CHASE domain: a predicted ligand-binding module in plant cytokinin receptors and other eukaryotic and bacterial receptors. *Trends Biochem. Sci.*, **10**, 579-582.
- Asakura, Y., Hagino, T., Ohta, Y., Aoki, K., Yonekura-Sakakibara, K., Deji, A., Yamaya, T., Sugiyama, T. und Sakakibara, H.** (2003) Molecular characterization of His-Asp phosphorelay signalling factors in maize leaves: implication of the signal divergence by cytokinin-inducible response regulators in the cytosol and the nuclei. *Plant Mol. Biol.*, **52**, 331-341.
- Auerbach, D., Thaminy, S., Hottiger, M.O. und Stagljar, I.** (2002) The post-genomic era of interactive proteomics: Facts and perspectives. *Proteomics*, **2**, 611-623.
- Bader, G.D., Heilbut, A., Andrews, B., Tyers, M., Hughes, T. und Boone, C.** (2003) Functional genomics and proteomics: charting a multidimensional map of the yeast cell. *Trends Cell Biol.*, **13**, 344-356.
- Bader, G.D. und Hogue, D.W.V.** (2002) Analysing yeast protein-protein interaction data obtained from different sources. *Natur Biotechnology*, **20**, 991-997.
- Barylko, B., Gerber, S.H., Binnst, D.D., Grichine, N., Khvotchev, M., Südhof, T.C. und Albanesi, J.P.** (2001) A Novel Family of Phosphatidylinositol 4-Kinases Conserved from Yeast to Humans. *The Journal of Biological Chemistry*, **276**, 7705-7708.
- Bauer, A. und Kuster, B.** (2003) Affinity purification-mass spectrometry. *Eur. J. Biochem.*, **270**, 570-578.
- Bernard, P. und Couturier, M.** (1992) Cell killing by the F plasmid CcdB protein involves poisoning of DNA-topoisomerase II complexes. *J. Mol. Biol.*, **226**, 735-745.
- Bleecker, A.B. und Kende, H.** (2000) Ethylene: a gaseous signal molecule in plants. *Annu. Rev. Cell Dev. Biol.*, **16**, 1-18.
- Birnboim, H.C. und Doly, J.** (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids. Res.*, **7**, 1513-1523.
- Bracha-Drori, K., Shichrur, K., Katz, A., Oliva, M., Angelovici, R., Yalovsky, S. und Ohad, N.** (2004) Detection of protein-protein interactions in plants using bimolecular fluorescence complementation. *The Plant Journal*, **40**, 419-427.
- Brandstatter, I. und Kieber, J.J.** (1998) Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell*, **10**, 1009-1020.

- Brenner, W.G., Romanov, G.A., Kollmer, I., Bürkle, L. und Schmölling, T.** (2005) Immediate-early and delayed cytokinin response genes of *Arabidopsis thaliana* identified by genome-wide expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades. *Plant J.*, **44**, 314-333.
- Bürkle, L., Meyer, S., Dortay, H., Lehrach, H. und Heyl, A.** (2005) In vitro recombination cloning of entire cDNA libraries in *Arabidopsis thaliana* and its application to the yeast two-hybrid system. *Funct. Integr. Genomics.*, **5**, 175-183.
- Chang, C. und Stewart, R. C.** (1998) The two-component system. *Plant Physiol.*, **117**, 723-731.
- Chang, C., Kwok, S.F., Bleeker, A.B. und Meyerowitz, E.M.** (1993) *Arabidopsis* ethylene response gene ETR1: similarity of product to two-component regulators. *Science*, **262**, 539-544.
- Calvin, N.M. und Hanawalt, P.C.** (1988) High-efficiency transformation of bacterial cells by electroporation. *J. Bacteriol.*, **170**, 2796-2801.
- Claire, R.W., David, G.L. und Roger, J.D.** (2002) MAP Kinase signaling specificity. *Science*, **296**, 2345-2356.
- Cohen, S.N., Chang, A.C.Y. und Hsu, L.** (1972) Nonchromosomal antibiotic resistance in bacteria: Genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci.*, **70**, 1293-1297.
- Conrad, P.A. und Heppler, P.K.** (1988) The effect of 1,4-dihydropyridines on the initiation and development of gametophore buds in the moss *Funaria*. *Plant Physiol.*, **86**, 684-687.
- D'Agostino, I.B., Deruere, J. und Kieber, J.J.** (2000) Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiol.*, **124**, 1706-1717.
- D'Agostino, I.B. und Kieber, J.J.** (1999) Phosphorelay signal transduction: the emerging family of plant response regulators. *Trends Biochem. Sci.*, **24**, 452-456.
- de Folter, S., Immink, R.G., Kieffer, M., Parenicova, L., Henz, S.R., Weigel, D., Busscher, M., Kooiker, M., Colombo, L., Kater, M.M. et al.** (2005) Comprehensive interaction map of the *Arabidopsis* MADS box transcription factors. *Plant Cell*, **17**, 1424-1433.
- Drobak, B.K.** (1993) Plant Phosphoinositides and intracellular signaling. *Plant Physiol.*, **102**, 705-709.
- Elion, E.A., Qi, M. und Chen, W.** (2005) Signal transduction. Signaling specificity in yeast. *Science*, **307**, 687-688.
- Fabret, C., Feher, V.A. und Hoch, J.A.** (1999) Two-component signal transduction in *Bacillus subtilis*: how one organism sees its world. *J. Bacteriol.*, **181**, 1975-1983.
- Ferreira, F.J. und Kieber, J.J.** (2005) Cytokinin signaling. *Curr. Opin. Plant Biol.*, **8**, 518-525.
- Fleischmann, R.D., Adams, M.D., White, O., Clayton, R.A., Kirkness, E.F., Kerlavage, A.R., Bult, C.J., Tomb, J.F., Dougherty, B.A., Merrick, J.M. et al.** (1995) Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science*, **269**, 496-512.

- Fraser, C.M., Gocayne, J.D., White, O., Adams, M.D., Clayton, R.A., Fleischmann, R.D., Bult, C.J., Kerlavage, A.R., Sutton, G., Kelley, J.M. et al.** (1995) The minimal gene complement of *Mycoplasma genitalium*. *Science*, **270**, 397-403.
- Gavin, A.C., Bosche, M., Krause, R., Grandi, P., Marzioch, M., Bauer, A., Schultz, J., Rick, J.M., Michon, A.M., Cruciat, C.M. et al.** (2002) Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature*, **415**, 141-147.
- Gietz, R.D. und Woods, R.A.** (2002) Transformation of yeast by the Liac/SS carrier DNA/PEG method. *Methods in Enzymology*, **350**, 87-96.
- Giote, L., Bader, J.S., Brouwer, C., Chaudhuri, A., Kuang, B., Li, Y., Hao, Y.L., Ooi, C.E., Godwin, B., Vitols, E. et al.** (2003) A protein interaction map of *Drosophila melanogaster*. *Science*, **302**, 1727-1736.
- Gobom, J., Schuerenberg, M., Mueller, M., Theiss, D., Lehrach, H. und Nordhoff, E.** (2001)  $\alpha$ -Cyano-4-hydroxycinnamic acid affinity sample preparation. A protocol for MALDI-MS peptide analysis in proteomics. *Anal. Chem.*, **73**, 434-438.
- Goehler, H., Lalowski, M., Stelzl, U., Waelter, S., Stroedicke, M., Worm, U., Droege, A., Lindenberg, K.S., Knoblich, M., Haenig, C. et al.** (2004): A Protein Interaction Network Links GIT1, an Enhancer of Huntingtin Aggregation, to Huntington's Disease. *Molecular Cell*, **15**, 853-865.
- Golemis, E.** (2002) Protein-Protein-Interactions. A molecular cloning manual. *Cold Spring Harbor Laboratory Press*.
- Grefen, C. und Harter, K.** (2004) Plant two-component systems: principles, functions, complexity and cross talk. *Planta*, **219**, 733-742.
- Grunberg-Manago, M.** (1999) Messenger RNA stability and its role in control of gene expression in bacteria and phages. *Annu. Rev. Gen.*, **33**, 193-227.
- Hanahan, D.** (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.*, **166**, 557-580.
- Hass, C., Lohrmann, J., Albrecht, V., Sweere, U., Hummel, F., Yoo, S.D., Hwang, I., Zhu, T., Schafer, E., Kudla, J. und Harter, K.** (2004) The response regulator 2 mediates ethylene signalling and hormone signal integration in Arabidopsis. *Embo J.*, **23**, 3290-3302.
- Heilmeyer, Jr. L.M.G., Vereb, Jr. G., Vereb, G., Kakuk, A. und Szivak, I.** (2003) Mammalian Phosphatidylinositol 4-Kinases. *IUBMB Life*, **55**, 59-65.
- Heyl, A., Werner, T. und Schmülling, T.** (2006) *Annual Plant Reviews in press*.
- Heyl, A. und Schmülling, T.** (2003) Cytokinin signal perception and transduction. *Curr. Opin. Plant Biol.*, **6**, 480-488.
- Higuchi, M., Pischke, M.S., Mähönen, A.P., Miyawaki, K., Hashimoto, Y., Seki, M., Kobayashi, M., Shinozaki, K., Kato, T., Tabata, S. et al.** (2004) *In planta* functions of the *Arabidopsis* cytokinin receptor family. *Proc. Natl. Acad. Sci.*, **101**, 8821-8826.
- Hilton, J.L., Kearny, P. und Ames, B.N.** (1965) Mode of action of the herbicide 3-amino-1,2,4-triazole (amitrole): inhibition of an enzyme of histidine biosynthesis. *Arch. Biochem. Biophys.*, **112**, 544-547.

- Ho, Y., Gruhler, A., Heilbutt, A., Bader, G.D., Morre, L., Adams, S.-L., Millar, A., Taylor, P., Bennett, K., Boutilier, K. et al.** (2002) Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature*, **415**, 180-183.
- Hoffman, W.L. und Jump, A.A.** (1986) Tween 20 removes antibodies and other proteins from nitrocellulose. *J. Immunol. Methods*, **94**, 191-196.
- Hosoda, K., Imamura, A., Katoh, E., Hatta, T., Tachiki, M., Yamada, H., Mizuno, T. und Yamazaki, T.** (2002) Molecular structure of the GARP family of plant Myb-related DNA binding motifs of the *Arabidopsis* response regulators. *Plant Cell*, **14**, 2015-2020.
- Hughes, T.R., Marton, M.J., Jones, A.R., Roberts, C.J., Stoughton, R., Armour, C.D., Bennett, H.A., Coffey, E., Dai, H., He, Y.D. et al.** (2000) Functional discovery via a compendium of expression profiles. *Cell*, **102**, 109-126.
- Hwang, I., Chen, H.C. und Sheen, J.** (2002) Two-component signal transduction pathways in *Arabidopsis*. *Plant Physiol.*, **129**, 500-515.
- Hwang, I. und Sheen, J.** (2001) Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature*, **413**, 383-389.
- Imamura, A., Kiba, T., Tajima, Y., Yamashino, T. und Mizuno, T.** (2003) *In vivo* and *in vitro* characterization of the ARR11 response regulator implicated in the His-Asp phosphorelay signal transduction in *Arabidopsis thaliana*. *Plant Cell Physiol.*, **44**, 122-131.
- Imamura, A., Yoshino, Y. und Mizuno, T.** (2001) Cellular localization of the signalling components of *Arabidopsis* His-to-Asp phosphorelay. *Biosci. Biotechnol. Biochem.*, **65**, 2113-2117.
- Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Kiba, T., Ueguchi, C., Sugiyama, T. und Mizuno, T.** (1999) Compilation and characterization of *Arabidopsis thaliana* response regulators implicated in His-Asp phosphorelay signal transduction. *Plant Cell Physiol.*, **40**, 733-742.
- Imamura, A., Hanaki, N., Umeda, H., Nakamura, A., Suzuki, T., Ueguchi, C. und Mizuno, T.** (1998) Response regulators implicated in His-to-Asp phosphotransfer signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci.*, **95**, 2691-2696.
- Immink, R.G. und Angenent, G.C.** (2002) Transcription factors do it together: the hows and whys of studying protein-protein interactions. *Trends Plant Sci.*, **7**, 531-534.
- Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., Tabata, S., Shinozaki, K. und Kakimoto, T.** (2001) Identification of CRE1 as a cytokinin receptor from *Arabidopsis*. *Nature*, **409**, 1060-1063.
- Inouye, M.** (2003) Histidine kinases in signal transduction. *Academic Press*.
- Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M. und Sakai, Y.** (2001) A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc. Natl. Acad. Sci.*, **98**, 4569-4574.
- Jeong, H., Mason, S.P., Barabasi, A.L. und Oltvai, Z.N.** (2001) Lethality and centrality in protein networks. *Nature*, **411**, 41-42.
- Jessee, J.** (1984) *FOCUS*, **6**, 5.

- Jones, D.R., Paneda, C., Villar, A.V., Alonso, A., Goni, F.M., Bütikofer, P., Brodbeck, U., Shepherd, P.R. und Varela-Nieto, I.** (2005) Phosphorylation of glycosyl-phosphatidylinositol by phosphatidylinositol 3-kinase changes its properties as a substrate for phospholipases. *FEBS Letters*, **579**, 59-65.
- Jung, J.-Y., Kim, Y.-W., Kwak, J.M., Hwang, J.-U., Young, J., Schroeder, J.I., Hwang, I. und Lee, Y.** (2002) Phosphatidylinositol 3- and 4-phosphate are required for normal stomatal movements. *The Plant Cell*, **14**, 2399-2412.
- Kafri, R., Bar-Even, A. und Pilpel, Y.** (2005) Transcription control reprogramming in genetic backup circuits. *Nat. Genet.*, **37**, 295-299.
- Kakimoto, T.** (2003) Perception and signal transduction of cytokinins. *Annu. Rev. Plant Biol.*, **54**, 605-627.
- Kakimoto, T.** (1996) CKI1, a histidine kinase homolog implicated in cytokinin signal transduction. *Science*, **274**, 982-985.
- Kanaho, Y. und Suzuki, T.** (2002) Phosphoinositide kinases as enzymes that produce versatile signaling lipids, phosphoinositides. *J. Biochem.*, **131**, 503-509.
- Kane, J.F.** (1995) Effects of rare codon clusters on high-level expression of heterologous proteins in *Escherichia coli*. *Curr. Opin. Biotechnol.*, **6**, 494-500.
- Kiba, T., Naitou, T., Koizumi, N., Yamashino, T., Sakakibara, H. und Mizuno, T.** (2005) Combinatorial microarray analysis revealing Arabidopsis genes implicated in Cytokinin responses through the His > Asp phosphorelay circuitry. *Plant Cell Physiol.*, **46**, 339-355.
- Kiba, T., Aoki, K., Sakakibara, H. und Mizuno, T.** (2004) Arabidopsis response regulator, ARR22, ectopic expression of which results in phenotypes similar to the *wol* cytokinin-receptor mutant. *Plant Cell Physiol.*, **45**, 1063-1077.
- Kiba, T., Yamada, H. und Mizuno, T.** (2002) Characterization of the ARR15 and ARR16 response regulators with special reference to the cytokinin signaling pathway mediated by the AHK4 histidine kinase in roots of *Arabidopsis thaliana*. *Plant Cell Physiol.*, **43**, 1059-1066.
- Kiba, T., Taniguchi, M., Imamura, A., Ueguchi, C., Mizuno, T. und Sugiyama, T.** (1999) Differential expression of genes for response regulators in response to cytokinins and nitrate in *Arabidopsis thaliana*. *Plant Cell Physiol.*, **40**, 767-771.
- Kido, M., Yamanaka, K., Mitani, T., Niki, H., Ogura, T. und Hiraga, S.** (1996) RNase E Polypeptides Lacking a Carboxyl-terminal Half Suppress a *mukB* mutation in *Escherichia coli*. *J. Bacteriol.*, **178**, 3917-3925.
- King, P.V. und Blakesley, R.** (1986) *FOCUS*, **8**, 1.
- Kim, H.J., Ryu, H., Hong, S.H., Woo, H.R., Lim, P.O., Lee, I.C., Sheen, J., Nam, H.G. und Hwang, I.** (2006) Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of ARR2 in *Arabidopsis*. *Proc. Natl. Acad. Sci.*, **103**, 814-819.
- Klopotowski, T. und Wiater, A.** (1965) Synergism of aminotriazole and phosphate on the inhibition of yeast imidazole glycerol phosphate dehydratase. *Arch. Biochem. Biophys.*, **112**, 562-566.
- Koretke, K.K., Lupas, A.N., Warren, P.V., Rosenberg, M. und Brown, J.R.** (2000) Evolution of two-component signal transduction. *Mol. Biol. Evol.*, **17**, 1956-1970.

- Kumar, A., Agaward, S., Heyman, J.A., Matson, S., Heidtman, M., Piccirillo, S., Umansky, L., Drawid, A., Jansen, R., Cheung, K.H. et al.** (2002) Subcellular localization of the yeast proteome. *Genes Dev.*, **16**, 707-719.
- Kurland, C. und Gallant, J.** (1996) Errors of heterologous protein expression. *Curr. Opin. Biotechnol.*, **7**, 489-493.
- Landy, A.** (1989) Dynamic, Structural, and Regulatory Aspects of Lambda Site specific Recombination. *Ann. Rev. Biochem.* **58**, 913-949.
- Letunic, I., Copley, R.R., Schmidt, S., Ciccarelli, F.D., Doerks, T., Schultz, J., Ponting, C.P. und Bork, P.** (2004) SMART 4.0: towards genomic data integration. *Nucleic Acids Res.*, **32**, 142-144.
- Li, J., Krichevsky, A., Vaidya, M., Tzfira, T. und Citovsky, V.** (2005) Uncoupling of the functions of *Arabidopsis* Vip1 protein in transient and stable plant genetic transformation by *Agrobacterium*. *Proc. Natl. Acad. Sci.*, **102**, 5733-5738.
- Li, S., Armstrong, C.M., Bertin, N., Ge, H., Milstein, S., Boxem, M., Vidalain, P.O., Han, J.D., Chesneau, A., Hao, T., et al.** (2004) A map of the interactome network of the metazoan *C. elegans*. *Science*, **303**, 540-543.
- Lin, W.H., Ye, R., Ma, H., Xu, Z.H. und Xue, H.W.** (2004) DNA chip-based expression profile analysis indicates involvement of the phosphatidylinositol signaling pathway in multiple plant responses to hormone and abiotic treatments. *Cell Research*, **14**, 34-45.
- Lopez, P.J., Marchand, I., Joyce, S.A. und Dreyfus, M.** (1999) The C-terminal Half of RNase E, Which Organizes the Escherichia coli Degradosome, Participates in mRNA Degradation but not rRNA Processing *in vivo*. *Mol. Microbiol.*, **33**, 188-199.
- Lohrmann, J. und Harter, K.** (2002) Plant two-component signaling systems and the role of response regulators. *Plant Phys.*, **128**, 363-369.
- Lohrmann, J., Sweere, U., Zabaleta, E., Baeurle, I., Keitel, C., Kozma, B.L., Brennicke, A., Schaefer, E., Kudla, J. und Harter, K.** (2001) The response regulator ARR2: A pollen-specific transcription factor involved in the expression of nuclear genes for components of mitochondrial complex I in *Arabidopsis*. *Mol. Gen. Genet.*, **265**, 2-13.
- Lohrmann, J., Buchholz, G., Keitel, C., Sweere, C., Kircher, S., Bäurle, I., Kudla, J. und Harter, K.** (1999) Differentially-expressed and nuclear-localized response regulator-like proteins from *Arabidopsis thaliana* with transcription factor properties. *Plant Biol.*, **1**, 495-506.
- Luo, D., Carpenter, R., Copsey, L., Vincent, C., Clark, J. und Coen, E.** (1999) Control of organ asymmetry in flowers of *Antirrhinum*. *Cell*, **99**, 367-376.
- Ma, Q.H. und Tian, B.** (2005) Characterization of a wheat histidine-containing phosphotransfer protein (HP) that is regulated by cytokinin-mediated inhibition of leaf senescence. *Plant Sci.*, **168**, 1507-1514.
- Maciver, S.K.** (2003) Phosphatidylinositol Signalling. *The Dynamic Cell*, **32**, 1-6.
- MacNicol, M., Jefferson, A.B. und Schulman, H.** (1990) Ca<sup>2+</sup>/Calmodulin Kinase is activated by the phosphatidyl inositol signaling pathway and becomes Ca<sup>2+</sup>-independent in PC12 Cell. *The Journal of Biological Chemistry*, **265**, 18055-18058.

- Maeda, T., Wurgler-Murphy, S.M. und Saito, H.** (1994) A two-component system that regulates an osmosensing MAP kinase cascade in yeast. *Nature*, **369**, 242-245.
- Mähönen, A.P., Bishopp, A., Higuchi, M., Nieminen, K.M., Kinoshita, K., Törmäkangas, K., Ikeda, Y., Oka, A., Kakimoto, T. und Helariutta, Y.** (2006) Cytokinin Signaling and its inhibitor AHP6 regulate cell fate during vascular development. *Science*, **311**, 94-98.
- Mähönen, A.P., Bonke, M., Kaupinnen, L., Riikonen, M., Benfey, P.N. und Helariutta, Y.** (2000) A novel two-component hybrid molecule regulates vascular morphogenesis of the *Arabidopsis* root. *Genes Dev.*, **14**, 2938-2943.
- Makino, S., Kiba, T., Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Ueguchi, C., Sugiyama, T. und Mizuno, T.** (2000) Genes encoding pseudo-response regulators: insight into His-to-Asp phosphorelay and circadian rhythm in *Arabidopsis thaliana*. *Plant Cell Physiol.*, **41**, 791-803.
- Marina, A., Waldburger, C.D. und Hendrickson, W.A.** (2005) Structure of the entire cytoplasmic portion of a sensor histidine-kinase protein. *The EMBO Journal*, **24**, 4247-4259.
- Mason, M.G., Mathews, D.E., Argyros, D.A., Maxwell, B.B., Kieber, J.J., Alonso, J.M., Ecker, J.R. und Schaller, G.E.** (2005) Multiple type-B response regulators mediate cytokinin signal transduction in *Arabidopsis*. *Plant Cell*, **17**, 3007-3018.
- Mason, M.G., Li, J., Mathews, D.E., Kieber, J.J. und Schaller, G.E.** (2004) Type-B response regulators display overlapping expression patterns in *Arabidopsis*. *Plant Physiol.*, **135**, 927-937.
- Matsubayashi, Y.** (2003) Ligand-receptor pairs in plant peptide signaling. *J. Cell. Sci.*, **116**, 3863-3870.
- Minet, M., Dufour, M.-E. und Francois, M.E.** (1992) Complementation of *Saccharomyces cerevisiae* auxotrophic mutants by *Arabidopsis thaliana* cDNAs. *The Plant Journal*, **2**, 417-422.
- Mirgorodskaya, E., Braeuer, C., Fucini, P., Lehrach, H. und Gobom, J.** (2005) Nanoflow liquid chromatography coupled to matrix-assisted laser desorption/ionization mass-spectrometry: Sample preparation, data analysis, and application to the analysis of complex peptide mixtures. *Proteomics*, **5**, 399-408.
- Miyata, S., Urao, T., Yamaguchi-Shinozaki, K. und Shinozaki, K.** (1998) Characterization of genes for two-component phosphorelay mediators with a single HPT domain in *Arabidopsis thaliana*. *FEBS Letters*, **437**, 11-14.
- Mizuno, T.** (2005) Two-Component Phosphorelay Signal Transduction Systems in Plants: from Hormone Responses to Circadian Rhythms. *Biosci. Biotechnol. Biochem.*, **69**, 2263-2276.
- Mizuno, T.** (1997) Compilation of all genes encoding two-component phosphotransfer signal transducers in the genome of *Escherichia coli*. *DNA Res.*, **4**, 161-168.
- Mizuno, T., Kaneko, T. und Tabata, S.** (1996) Compilation of all genes encoding bacterial two-component signal transducers in the genome of the cyanobacterium, *Synechocystis sp.* strain PCC 6803. *DNA Res.*, **3**, 407-414.
- Mougel, C. und Zhulin, I.B.** (2001) CHASE: an extracellular sensing domain common to transmembrane receptors from prokaryotes, lower eukaryotes and plants. *Trends Biochem. Sci.*, **26**, 582-584.

- Müller-Röber, B. und Pical, C.** (2002) Inositol phospholipid metabolism in *Arabidopsis*. Characterized and putative isoforms of inositol phospholipids kinase and phosphoinositide-specific phospholipase c. *Plant Physiology*, **130**, 22-46.
- Mullis, K.B. und Faloona, F.A.** (1987) Specific synthesis of DNA *in vitro* via a polymerase catalyzed chain reaction. *Methods Enzymol.*, **155**, 335-350.
- Murashige, T. und Skoog, F.** (1962) A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Physiologia Plantarum*, **15**, 473-497.
- Munnik, T., Irvine, R.F. und Musgrave, A.** (1998) Phospholipid signalling in plants. *Biochimica et Biophysica Acta*, **1389**, 222-272.
- Napier, R.** (2004) Plant Hormone Binding Sites. *Annals of Botany*, **93**, 227-233.
- Nishimura, C., Ohashi, Y., Sato, S., Kato, T., Tabata, S. und Ueguchi, C.** (2004) Histidine kinase homologs that act as cytokinin receptors possess overlapping functions in the regulation of shoot and root growth in *Arabidopsis*. *Plant Cell*, **16**, 1365-1377.
- Orosz, A., Boros, I. und Venetianer, P.** (1991) Analysis of the Complex Transcription Termination Region of the *Escherichia coli* *rrnB* Gene. *Eur. J. Biochem.*, **201**, 653-659.
- Osakabe, Y., Miyata, S., Urao, T., Seki, M., Shinozaki, K. und Yamaguchi-Shinozaki, K.** (2002) Overexpression of *Arabidopsis* response regulators, ARR4/ATRR1/IBC7 and ARR8/ATRR3, alters cytokinin responses differentially in the shoot and in callus formation. *Biochem. Biophys. Res. Commun.*, **293**, 806-815.
- Ota, I.M. und Varshavsky, A.** (1993) A yeast protein similar to bacterial two-component regulator. *Science*, **262**, 566-569.
- Palkova, Z., Pospisek, M. und Zikanova, B.** (1996) Techniques for Colony DNA Hybridization and Protein Immunoassays for a Broad Spectrum of Yeast Genera. *Biotechniques*, **21**, 982-986.
- Pawson, T. und Nash, P.** (2003) Assembly of cell regulatory systems through protein interaction domains. *Science*, **300**, 445-452.
- Pluskal, M.G., Przekop, M.B., Kavonian, M.R., Vecoli, C. und Hicks, D.A.** (1986) Immobilon PVDF transfer membrane – A new membrane substrate for western blotting of proteins. *Biotechniques*, **4**, 272-283.
- Prescott, S.M.** (1999) A thematic series on kinases and phosphatases that regulate lipid signalling. *The Journal of Biological Chemistry*, **274**, 8345.
- Pukahainen, T., Hess, M.W., Makela, P., Svensson, J., Heino, P. und Palva, E.T.** (2004) Overexpression of multiple dehydrin genes enhances tolerance to freezing stress in *Arabidopsis*. *Plant Mol. Biol.*, **54**, 743-753.
- Rabilloud, T.** (1990) Mechanism of protein silver staining in polyacrylamide gels: A 10-year synthesis. *Electrophoresis*, **11**, 785-794.
- Raleigh, E.A., Murray, N.E., Revel, H., Blumenthal, R.M., Westaway, D., Reith, A.D., Rigby, P.W., Elhai, J. und Hanahan, D.** (1988) McrA and McrB restriction phenotypes of some *E. coli* strains and implications for gene cloning. *Nucl. Acids Research*, **16**, 1523.
- Raleigh, E.A., Lech, K. und Brent, R.** (1989) Selected topics from classical bacterial genetics. *In: Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith,*



- J.A., Struhl, K., Albright, C.M., Coen, D.M. und Verki, A. (Hrsg.)** Current protocols in molecular biology. *Wiley*, New York, Vol. 1a, pp. 1.4.1-1.4.16.
- Repp, A., Mikami, K., Mittmann, F. und Hartmann, E. (2004)** Phosphoinositide-specific phospholipase C is involved in cytokinin and gravity responses in the moss *Physcomitrella patens*. *The Plant Journal*, **40**, 250-259.
- Riechmann, J.L., Heard, J., Martin, G., Reuber, L., Jiang, C., Keddie, J., Adam, L., Pineda, O., Ratcliffe, O.J., Samaha, R.R. et al. (2000)** Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science*, **290**, 2105-2110.
- Riefler, M., Novak, O., Strnad, M. und Schmülling, T. (2006)** *Arabidopsis* cytokinin receptor mutants reveal functions in shoot growth, leaf senescence, seed size, germination, root development and cytokinin metabolism. *The Plant Cell*, *in press*.
- Rohila, J.S., Chen, M., Chen, S., Chen, J., Cerny, R., Dardick, C., Canlas, P., Xu, X., Gribskov, M., Kanrar, S. et al. (2006)** Protein-protein interactions of tandem affinity purification-tagged protein kinases in rice. *The Plant Journal*, **46**, 1-13.
- Russinova, E., Borst, J.W., Kwaaitaal, M., Cano-Delgado, A., Yin, Y., Chory, J. und de Vries, S.C. (2004)** Heterodimerization and endocytosis of Arabidopsis brassinosteroid receptors BRI1 and AtSERK3 (BAK1). *Plant Cell*, **16**, 3216-3229.
- Saito, H. (2001)** Histidine phosphorylation and two-component signaling in eukaryotic cells. *Chem. Rev.*, **101**, 2497-2509.
- Sakai, H., Honma, T., Aoyama, T., Sato, S., Kato, T., Tabata, S. und Oka, A. (2001)** ARR1, a transcription factor for genes immediately responsive to cytokinins. *Science*, **294**, 1519-1521.
- Sakai, H., Aoyama, T. und Oka, A. (2000)** Arabidopsis ARR1 and ARR2 response regulators operate as transcriptional activators. *Plant J.*, **24**, 703-711.
- Sakai, H., Aoyama, T., Bono, H. und Oka, A. (1998)** Two-component Response Regulators from *Arabidopsis thaliana* Contain a Putative DNA-binding Motif. *ICR Annual Report*, **5**, 50-51.
- Sambrook, J., Fritsch, E.F. und Maniatis, T. (1989)** Molecular Cloning: A Laboratory Manual. *Coldspring Harbor Laboratory*, New York.
- Schaller, G.E., Mathews, D.E., Gribskov, M. und Walker, J.C. (2002)** Two-component signalling elements and histidyl-aspartyl phosphorelays. In: The Arabidopsis Book. (eds. Somerville, C., Meyerowitz, E., Rockville, M.D), American Society of Plant Biologists, URL: <http://www.aspb.org/publications/Arabidopsis>.
- Schaller, G.E., Ladd, A.N., Lanahan, M.B., Spanbauer, J.M. und Bleecker, A.B. (1995)** The ethylene response mediator *ETR1* from *Arabidopsis* forms a disulfide-linked dimer. *J. Biol. Chem.*, **270**, 12526-12530.
- Schlessinger, J. (2002)** Ligand-induced, receptor-mediated dimerization and activation of EGF receptor. *Cell*, **110**, 669-672.
- Schmülling, T. (2002)** New insights into the functions of cytokinins in plant development. *J. Plant Growth Reg.*, **21**, 40-49.

- Schuster, S.C., Noegel, A.A., Oehme, F., Gerisch, G. und Simon, M.I.** (1996) The hybrid histidine kinase DokA is part of the osmotic response system of Dictyostelium. *The EMBO Journal*, **15**, 3880-3889.
- Schwikowski, B., Uetz, P. und Fields, S.** (2000) A network of protein-protein interactions in yeast. *Nature Biotechnology*, **18**, 1257-1261.
- Sheen, J.** (2002) Phosphorelay and transcription control in cytokinin signal transduction. *Science*, **296**, 1650-1652.
- Singleton, C.K., Zinda, M.J., Mykytka, B. und Yang, P.** (1998) The histidine kinase dhkC regulates the choice between migrating slugs and terminal differentiation in Dictyostelium discoideum. *Dev. Biol.*, **203**, 345-357.
- Spichal, L., Rakova, N.Y., Riefler, M., Mizuno, T., Romanov, G.A., Strnad, M. und Schmülling, T.** (2004) Two Cytokinin Receptors of *Arabidopsis thaliana*, CRE1/AHK4 and AHK3, Differ in their Ligand Specificity in a Bacterial Assay. *Plant Cell Physiol.*, **45**, 1299-1305.
- Stepanova, A.N. und Ecker, J.R.** (2000) Ethylene signaling: from mutants to molecules. *Curr. Opin. Plant Biol.*, **3**, 353-360.
- Stock, A.M., Robinson, V.L. und Goudreau, P.N.** (2000) Two-component signal transduction. *Annu. Rev. Biochem.*, **69**, 183-215.
- Stover, C.K., Pham, X.Q., Erwin, A.L., Mizoguchi, S.D., Warrenner, P., Hickey, M.J., Brinkman, F.S., Hufnagel, W.O., Kowalik, D.J., Lagrou, M. et al.** (2000) Complete genome sequence of *Pseudomonas aeruginosa* PA01, an opportunistic pathogen. *Nature*, **406**, 959-964.
- Subramaniam, R., Desveaux, D., Spickler, C., Michnick, S.W. und Brisson, N.** (2001) Direct visualization of protein interactions in plant cells. *Nature Biotechnology*, **19**, 769-772.
- Suzuki, T., Ishikawa, K., Yamashino, T. und Mizuno, T.** (2002) An Arabidopsis histidine-containing phosphotransfer (Hpt) factor implicated in phosphorelay signal transduction: overexpression of AHP2 in plants results in hypersensitiveness to cytokinin. *Plant Cell Physiol.*, **43**, 123-129.
- Suzuki, T., Miwa, K., Ishikawa, K., Yamada, H., Aiba, H. und Mizuno, T.** (2001a) The Arabidopsis sensor His-kinase, AHk4, can respond to cytokinins. *Plant Cell Physiol.*, **42**, 107-113.
- Suzuki, T., Sakurai, K., Ueguchi, C. und Mizuno, T.** (2001b) Two types of putative nuclear factors that physically interact with histidine-containing phosphotransfer (Hpt) domains, signaling mediators in His-to-Asp phosphorelay, in Arabidopsis thaliana. *Plant Cell Physiol.*, **42**, 37-45.
- Suzuki, T., Sakurai, K., Imamura, A., Nakamura, A., Ueguchi, C. und Mizuno, T.** (2000) Compilation and characterization of histidine-containing phosphotransmitters implicated in His-to-Asp phosphorelay in plants: AHP signal transducers of *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.*, **64**, 2486-2489.
- Suzuki, T., Imamura, A., Ueguchi, C. und Mizuno, T.** (1998) Histidine-containing phosphotransfer (HPT) signal transducers implicated in His-to-Asp phosphorelay in *Arabidopsis*. *Plant Cell Physiol.*, **39**, 1258-1268.

- Sweere, U., Eichenberg, K., Lohrmann, J., Mira-Rodado, V., Baurle, I., Kudla, J., Nagy, F., Schäfer, E. und Harter, K.** (2001) Interaction of the response regulator ARR4 with the photoreceptor phytochrome B in modulating red light signaling. *Science*, **294**, 1108-1111.
- Tajima, Y., Imamura, A., Kiba, T., Amano, Y., Yamashino, T. und Mizuno, T.** (2004) Comparative studies on the type-B response regulators revealing their distinctive properties in the His-to-Asp phosphorelay signal transduction of *Arabidopsis thaliana*. *Plant Cell Physiol.*, **45**, 28-39.
- Tanaka, Y., Suzuki, T., Yamashino, T. und Mizuno, T.** (2004) Comparative studies of the AHP histidine-containing phosphotransmitters implicated in His-to-Asp phosphorelay in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.*, **68**, 462-465.
- Taniguchi, M., Kiba, T., Sakakibara, H., Ueguchi, C., Mizuno, T. und Sugiyama, T.** (1998) Expression of *Arabidopsis* response regulator homolog is induced by cytokinins and nitrate. *FEBS Letters*, **429**, 259-262.
- To, J.P., Haberer, G., Ferreira, F.J., Deruere, J., Mason, M.G., Schaller, G.E., Alonso, J.M., Ecker, J.R. und Kieber, J.J.** (2004) Type-A *Arabidopsis* response regulators are partially redundant negative regulators of cytokinin signaling. *Plant Cell*, **16**, 658-671.
- Towbin, H., Staehelin, T. und Gordon, J.** (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. *Proc. Natl. Acad. Sci.*, **76**, 4350-4354.
- Ueguchi, C., Koizumi, H., Suzuki, T. und Mizuno, T.** (2001) Novel family of sensor histidine kinase genes in *Arabidopsis thaliana*. *Plant Cell Physiol.*, **42**, 231-235.
- Uetz, P., Russel, L. und Finley, Jr.** (2005) From protein networks to biological systems. *FEBS Letters*, **579**, 1821-1827.
- Uetz, P. und Pohl, E.** (2004) Protein-Protein- und Protein-DNA-Interaktionen. In: *Wink et al., Molekulare Biotechnologie, Wiley-VCH*.
- Uetz, P., Giot, L., Cagney, G., Mansfield, T.A., Judson, R.S., Knight, J.R., Lockshon, D., Narayan, V., Srinivasan, M., Pochart, P. et al.** (2000) A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature*, **403**, 623-627.
- Urao, T., Yamaguchi-Shinozaki, K. und Shinozaki, K.** (2000) Two-component systems in plant signal transduction. *Trends Plant Sci.*, **5**, 67-74.
- Urao, T., Yakubov, B., Satoh, R., Yamaguchi-Shinozaki, K., Seki, M., Hirayama, T. und Shinozaki, K.** (1999) A transmembrane hybrid-type histidine kinase in *Arabidopsis* functions as an osmosensor. *Plant Cell*, **11**, 1743-1754.
- Urao, T., Yakubov, B., Yamaguchi-Shinozaki, K. und Shinozaki, K.** (1998) Stress responsive expression of genes for two-component response regulator-like proteins in *Arabidopsis thaliana*. *FEBS Letters*, **427**, 175-178.
- VanOss, C.J., Good, R.J. und Chaudhury, M.K.** (1987) Mechanisms of DNA (Southern) and protein (Western) blotting on cellulose nitrate and other membranes. *J. Chromatogr.*, **391**, 53-65.
- von Mering, C., Krause, R., Snel, B., Cornell, M., Oliver, S.G., Fields, S. und Bork, P.** (2002) Comparative assessment of large-scale data sets of protein-protein interactions. *Nature*, **417**, 399-403.

- Walhout, A.J.M. und Vidal, M.** (2001) Protein interaction maps for model organisms. *Nature Reviews*, **2**, 55-62.
- Walhout, A.J.M., Sordella, R., Lu, X., Hartley, J.L., Temple, G.F., Brasch, M.A., Thierry-Mieg, N. und Vidal, M.** (2000) Protein interaction mapping in *C. elegans* using proteins involved in vulval development. *Science*, **287**, 116-122.
- Walter, M., Chaban, C., Schütze, K., Bastitc, O., Weckermann, K., Näke, C., Blazevic, D., Grefen, C., Schumacher, K., Oecking, C. et al.** (2004) Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *The Plant Journal*, **40**, 428-438.
- Wang, X., Goshe, M.B., Soderblom, E.J., Phinney, B.S., Kuchar, J.A., Li, J., Asami, T., Yoshida, S., Huber, S.C. und Clouse, S.D.** (2005) Identification and functional analysis of *in vivo* phosphorylation sites of the Arabidopsis BRASSINOSTEROID-INSENSITIVE1 receptor kinase. *Plant Cell*, **17**, 1685-1703.
- Wang, N., Soderbom, F., Anjard, C., Shaulsky, G. und Loomis, W.F.** (1999) SDF-2 induction of terminal differentiation in *Dictyostelium discoideum* is mediated by the membrane-spanning sensor kinase DhkA. *Mol. Cell Biol.*, **19**, 4750-4756.
- West, A. H. und Stock, A.M.** (2001) Histidine kinases and response regulator proteins in two-component signaling systems. *Trends in Biochemical Sciences*, **26**, 369-376.
- Westergren, T., Ekblad, L., Jergil, B. und Sommarin, M.** (1999) Phosphatidylinositol 4-kinase associated with spinach plasma membranes. Isolation and characterization of two distinct forms. *Plant Phys.*, **121**, 507-516.
- Wiater, A., Hulanicka, D. und Klotowski, T.** (1971) Structural requirements for inhibition of yeast imidazoleglycerol phosphate dehydratase by triazole and anion inhibitors. *Acta Biochim. Pol.*, **18**, 289-297.
- Wolanin, P.M., Thomason, P.A. und Stock, J.B.** (2002) Histidine protein kinases: key signal transducers outside the animal kingdom. *Genome Biology*, **3**, 3013.1-3013.8.
- Xenarios, I. und Eisenberg, D.** (2001) Protein interaction databases. *Current Opinion in Biotechnology*, **12**, 334-339.
- Xue, H.W., Pical, C., Brearley, C., Elge, S. und Müller-Röber, B.** (1999) A plant 126-kDa phosphatidylinositol 4-kinase with a novel repeat structure. *The Journal of Biol. Chem.*, **274**, 5738-5745.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K. und Mizuno, T.** (2001) The *Arabidopsis* AHK4 histidine kinase is a cytokinin-binding receptor that transduces cytokinin signals across the membrane. *Plant Cell Physiol.*, **42**, 1017-1023.
- Yamada, H., Hanaki, N., Imamura, A., Ueguchi, C. und Mizuno, T.** (1998) An *Arabidopsis* protein that interacts with the cytokinin-inducible response regulator, ARR4, implicated in the His-Asp phosphorelay signal transduction. *FEBS Letters*, **436**, 76-80.
- Yang, Y. und Inouye, M.** (1991) Intermolecular complementation between two defective mutants of *Escherichia coli* signal transducing receptors. *Proc. Natl. Acad. Sci.*, **88**, 11057-11061.
- Yates, J.R.** (2000) Mass spectrometry from genomics to proteomics. *TIG*, **16**, 5-8.

- Yeh, K.C., Murphy, J.T. und Lagarias, J.C.** (1998) Eukaryotic phytochromes: light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc. Natl. Acad. Sci.*, **12**, 659-667.
- Yonekura-Sakakibara, K., Kojima, M., Yamaya, T. und Sakakibara, H.** (2004) Molecular characterization of cytokinin-responsive histidine kinases in maize. Differential ligand preferences and response to cis-zeatin. *Plant Physiol.*, **134**, 1654-1661.
- Zhao, L., Nakazawa, M., Takase, T., Manabe, K., Kobayashi, M., Seki, M., Shinozaki, K. und Matsui, M.** (2004) Overexpression of LSH1, a member of an uncharacterized gene family, causes enhanced light regulation of seedling development. *The Plant Journal*, **37**, 694-706.
- Zhao, X.H., Bondeva, T. und Balla, T.** (2000) Characterization of recombinant phosphatidylinositol 4-Kinase  $\beta$  reveals auto- and heterophosphorylation of the enzyme. *The Journal of Biological Chemistry*, **275**, 14642-14648.
- Zhu, Y. und Inouye, M.** (2004) The HAMP linker in histidine kinase dimeric receptors is critical for symmetric transmembrane signal transduction. *J. Biol. Chem.*, **279**, 48152-48158.
- Zoller, M.J. und Smith, M.** (1982) Oligonucleotid-directed mutagenesis using M13-derived vectors: An efficient and generell procedure for the production of point mutations in any fragment of DNA. *Nucl. Acids Res.*, **10**, 6487-6500.