

## 8 Literatur

**Aerssens, J., Boonen, S., Lowet, G., Dequeker, J.** (1998): Interspecies differences in bone composition, density, and quality: potential implications for in vivo bone research. *Endocrinology*, 139(2): 663-70.

**Amling, M., Delling, G.** (1996): Cell biology of osteoclasts and molecular mechanisms of bone resorption. *Pathologe*, 17(5): 358-67.

**Arnold, U., Lindenhayn, K., Perka, C.** (2002): In vitro-cultivation of human periosteum derived cells in bioresorbable polymer-TCP-composites. *Biomaterials*, 23(11): 2303-10.

**Aro, H. T., Wippermann, B. W., Hodgson, S. F., Chao, E. Y.** (1990): Internal remodeling of periosteal new bone during fracture healing. *J Orthop Res*, 8(2): 238-46.

**Arrington, E. D., Smith, W. J., Chambers, H. G., Bucknell, A. L., Davino, N. A.** (1996): Complications of iliac crest bone graft harvesting. *Clin Orthop Relat Res*, (329): 300-9.

**Audige, L., Griffin, D., Bhandari, M., Kellam, J., Ruedi, T. P.** (2005): Path analysis of factors for delayed healing and nonunion in 416 operatively treated tibial shaft fractures. *Clin Orthop Relat Res*, 438: 221-32.

**Babhulkar, S., Pande, K., Babhulkar, S.** (2005): Nonunion of the diaphysis of long bones. *Clin Orthop Relat Res*, (431): 50-6.

**Baksh, D., Song, L., Tuan, R. S.** (2004): Adult mesenchymal stem cells: characterization, differentiation, and application in cell and gene therapy. *J Cell Mol Med*, 8(3): 301-16.

**Barry, F., Boynton, R. E., Liu, B., Murphy, J. M.** (2001): Chondrogenic differentiation of mesenchymal stem cells from bone marrow: differentiation-dependent gene expression of matrix components. *Exp Cell Res*, 268(2): 189-200.

**Bauer, T. W., Muschler, G. F.** (2000): Bone graft materials. An overview of the basic science. *Clin Orthop Relat Res*, (371): 10-27.

**Bennett, J. H., Joyner, C. J., Triffitt, J. T., Owen, M. E.** (1991): Adipocytic cells cultured from marrow have osteogenic potential. *J Cell Sci*, 99 (Pt 1): 131-9.

**Bianchi, G., Muraglia, A., Daga, A., Corte, G., Cancedda, R., Quarto, R.** (2001): Microenvironment and stem properties of bone marrow-derived mesenchymal cells. *Wound Repair Regen*, 9(6): 460-6.

**Bianco, P., Riminucci, M., Gronthos, S., Robey, P. G.** (2001): Bone marrow stromal stem cells: nature, biology, and potential applications. *Stem Cells*, 19(3): 180-92.

**Bonewald, L. F.** (2002): Osteocytes: a proposed multifunctional bone cell. *J Musculoskelet Neuronal Interact*, 2(3): 239-41.

**Bosch, U., Skutek, M., Kasperczyk, W. J., Tscherne, H.** (1999): Nonunion of the humeral diaphysis - operative and nonoperative treatment. *Chirurg*, 70(11): 1202-8.

**Bostrom, M. P., Lane, J. M., Berberian, W. S., Missri, A. A., Tomin, E., Weiland, A., Doty, S. B., Glaser, D., Rosen, V. M.** (1995): Immunolocalization and expression of bone morphogenetic proteins 2 and 4 in fracture healing. *J Orthop Res*, 13(3): 357-67.

**Boyan, B. D., Caplan, A. I., Heckman, J. D., Lennon, D. P., Ehler, W., Schwartz, Z.** (1999): Osteochondral progenitor cells in acute and chronic canine nonunions. *J Orthop Res*, 17(2): 246-55.

**Braun, W., Ruter, A.** (1996): Fracture healing. Morphologic and physiologic aspects. *Unfallchirurg*, 99(1): 59-67.

**Breitbart, A. S., Grande, D. A., Kessler, R., Ryaby, J. T., Fitzsimmons, R. J., Grant, R. T.** (1998): Tissue engineered bone repair of calvarial defects using cultured periosteal cells. *Plast Reconstr Surg*, 101(3): 567-74.

**Brighton, C. T.** (1984): The biology of fracture repair. *Instr Course Lect.*, 33: 60-82.

**Brighton, C. T., Hunt, R. M.** (1997): Early histologic and ultrastructural changes in microvessels of periosteal callus. *J Orthop Trauma*, 11(4): 244-53.

**Bruder, S. P., Jaiswal, N., Haynesworth, S. E.** (1997): Growth kinetics, self-renewal, and the osteogenic potential of purified human mesenchymal stem cells during extensive subcultivation and following cryopreservation. *J Cell Biochem*, 64(2): 278-94.

**Bruder, S. P., Kurth, A. A., Shea, M., Hayes, W. C., Jaiswal, N., Kadiyala, S.** (1998): Bone regeneration by implantation of purified, culture-expanded human mesenchymal stem cells. *J Orthop Res*, 16(2): 155-62.

**Bucher, O., Wartenberg, H.** Cytologie, Histologie und mikroskopische Anatomie des Menschen. 183-200, Bern, 1997

**Cancedda, R., Bianchi, G., Derubeis, A., Quarto, R.** (2003): Cell therapy for bone disease: a review of current status. *Stem Cells*, 21(5): 610-9.

**Caplan, A. I.** (1991): Mesenchymal stem cells. *J Orthop Res*, 9(5): 641-50.

**Caplan, A. I.** (1994): The mesengenic process. *Clin Plast Surg*, 21(3): 429-35.

**Claes, L., Heitemeyer, U., Krischak, G., Braun, H., Hierholzer, G.** (1999): Fixation technique influences osteogenesis of comminuted fractures. *Clin Orthop Relat Res*, (365): 221-9.

**Connolly, J. F., Guse, R., Tiedeman, J., Dehne, R.** (1989): Autologous marrow injection for delayed unions of the tibia: a preliminary report. *J Orthop Trauma*, 3(4): 276-82.

**Cruess, R. L., Dumont, J.** (1975): Fracture healing. *Can J Surg.*, 18(5): 403-13.

**Cullinane, D. M., Fredrick, A., Eisenberg, S. R., Pacicca, D., Elman, M. V., Lee, C., Salisbury, K., Gerstenfeld, L. C., Einhorn, T. A.** (2002): Induction of a neoarthrosis by precisely controlled motion in an experimental mid-femoral defect. *J Orthop Res*, 20(3): 579-86.

**Devine, M. J., Mierisch, C. M., Jang, E., Anderson, P. C., Balian, G.** (2002): Transplanted bone marrow cells localize to fracture callus in a mouse model. *J Orthop Res*, 20(6): 1232-9.

- Einhorn, T. A.** (1995): Enhancement of fracture-healing. *J Bone Joint Surg Am*, 77(6): 940-56.
- Einhorn, T. A.** (1998a): Breakout session. 1: Definitions of fracture repair. *Clin Orthop Relat Res*, (355 Suppl): 353.
- Einhorn, T. A.** (1998b): The cell and molecular biology of fracture healing. *Clin Orthop Relat Res.*, (355 Suppl): 7-21.
- Einhorn, T. A., Lane, J. M., Burstein, A. H., Kopman, C. R., Vigorita, V. J.** (1984): The healing of segmental bone defects induced by demineralized bone matrix. A radiographic and biomechanical study. *J Bone Joint Surg Am*, 66(2): 274-9.
- Eitel, F., Schweiberer, L.** (1982): Knochentransplantation bei Pseudarthrosen. *Z Orthop*, 129: 540-541.
- Eitel, F., Seiler, H., Schweiberer, L.** (1981): Morphological examination of animal-experiment results: comparison with regeneration of the human bone-structure. II. Research results. *Unfallheilkunde*, 84(6): 255-64.
- Ellender, G., Feik, S. A., Carach, B. J.** (1988): Periosteal structure and development in a rat caudal vertebra. *J Anat*, 158: 173-87.
- Engler, A. J., Sen, S., Sweeney, H. L., Discher, D. E.** (2006): Matrix elasticity directs stem cell lineage specification. *Cell*, 126(4): 677-89.
- Erben, R. G.** (1996): Trabecular and endocortical bone surfaces in the rat: modeling or remodeling? *Anat Rec*, 246(1): 39-46.
- Flygare, L., Hosoki, H., Rohlin, M., Petersson, A.** (1997): Bone histomorphometry using interactive image analysis. A methodological study with application on the human temporomandibular joint. *Eur J Oral Sci*, 105(1): 67-73.
- Friedenstein, A. J., Chailakhjan, R. K., Lalykina, K. S.** (1970): The development of fibroblast colonies in monolayer cultures of guinea-pig bone marrow and spleen cells. *Cell Tissue Kinet*, 3(4): 393-403.

**Friedenstein, A. J., Gorskaja, J. F., Kulagina, N. N.** (1976): Fibroblast precursors in normal and irradiated mouse hematopoietic organs. *Exp Hematol*, 4(5): 267-74.

**Frost, H. M.** (1989): The biology of fracture healing. An overview for clinicians. Part I. *Clin Orthop Relat Res*, (248): 283-93.

**Frost, H. M.** (1989): The biology of fracture healing. An overview for clinicians. Part II. *Clin Orthop Relat Res*, (248): 294-309.

**Gazdag, A. R., Lane, J. M., Glaser, D., Forster, R. A.** (1995): Alternatives to Autogenous Bone Graft: Efficacy and Indications. *J Am Acad Orthop Surg*, 3(1): 1-8.

**Gebauer, D., Mayr, E., Orthner, E., Ryaby, J. P.** (2005): Low-intensity pulsed ultrasound: effects on nonunions. *Ultrasound Med Biol*, 31(10): 1391-402.

**Glowacki, J.** (1998): Angiogenesis in fracture repair. *Clin Orthop Relat Res.*, (355 Suppl): 82-9.

**Goodship, A. E., Kenwright, J.** (1985): The influence of induced micromovement upon the healing of experimental tibial fractures. *J Bone Joint Surg Br*, 67(4): 650-5.

**Grundnes, O., Reikeras, O.** (1993): The importance of the hematoma for fracture healing in rats. *Acta Orthop Scand.*, 64(3): 340-2.

**Haas, N. P.** (2000): Callus modulation-fiction or reality? *Chirurg*, 71(9): 987-8.

**Harrison, L. J., Cunningham, J. L., Stromberg, L., Goodship, A. E.** (2003): Controlled induction of a pseudarthrosis: a study using a rodent model. *J Orthop Trauma*, 17(1): 11-21.

**Heckman, J. D., Ryaby, J. P., McCabe, J., Frey, J. J., Kilcoyne, R. F.** (1994): Acceleration of tibial fracture-healing by non-invasive, low-intensity pulsed ultrasound. *J Bone Joint Surg Am*, 76(1): 26-34.

**Hernigou, P., Beaujean, F.** (1997): Abnormalities in the bone marrow of the iliac crest in patients who have osteonecrosis secondary to corticosteroid therapy or alcohol abuse. *J Bone Joint Surg Am*, 79(7): 1047-53.

- Hernigou, P., Poignard, A., Manicom, O., Mathieu, G., Rouard, H.** (2005): The use of percutaneous autologous bone marrow transplantation in nonunion and avascular necrosis of bone. *J Bone Joint Surg Br*, 87(7): 896-902.
- Hietaniemi, K., Peltonen, J., Paavolainen, P.** (1995): An experimental model for non-union in rats. *Injury*, 26(10): 681-6.
- Hulth, A.** (1989): Current concepts of fracture healing. *Clin Orthop Relat Res*, (249): 265-84.
- Hunt, J. A., Williams, D. F.** (1995): Quantifying the soft tissue response to implanted materials. *Biomaterials*, 16(3): 167-70.
- Ito, Y., Fitzsimmons, J. S., Sanyal, A., Mello, M. A., Mukherjee, N., O'Driscoll, S. W.** (2001): Localization of chondrocyte precursors in periosteum. *Osteoarthritis Cartilage*, 9(3): 215-23.
- Jäger, M., Westhoff, B., Wild, A., Krauspe, R.** (2005a): Bone harvesting from the iliac crest. *Orthopade*, 34(10): 976-82, 984, 986-90, 992-4.
- Jäger, M., Sager, M., Lensing-Hohn, S., Krauspe, R.** (2005b): The critical size bony defect in a small animal for bone healing studies (I): Comparative anatomical study on rats' femur. *Biomed Tech (Berl)*, 50(4): 107-10.
- Jäger, M., Sager, M., Lensing-Hohn, S., Krauspe, R.** (2005c): The critical size bony defect in a small animal for bone healing studies (II): implant evolution and surgical technique on a rat's femur. *Biomed Tech (Berl)*, 50(5): 137-42.
- Jaiswal, N., Haynesworth, S. E., Caplan, A. I., Bruder, S. P.** (1997): Osteogenic differentiation of purified, culture-expanded human mesenchymal stem cells in vitro. *J Cell Biochem*, 64(2): 295-312.
- Jaquier, C., Schaeren, S., Farhadi, J., Mainil-Varlet, P., Kunz, C., Zeilhofer, H. F., Heberer, M., Martin, I.** (2005): In vitro osteogenic differentiation and in vivo bone-forming capacity of human isogenic jaw periosteal cells and bone marrow stromal cells. *Ann Surg*, 242(6): 859-67.

**Jones, C. B., Mayo, K. A.** (2005): Nonunion treatment: iliac crest bone graft techniques. *J Orthop Trauma*, 19(10 Suppl): 11-3.

**Josten, C., Kremer, M., Muhr, G.** (1996): Ilizarov procedure in pseudarthrosis. *Orthopade*, 25(5): 405-15.

**Junqueira, L. C., Carneiro, J., Schiebler, T. H.** Histologie: Zytologie, Histologie und mikroskopische Anatomie des Menschen unter besonderer Berücksichtigung der Histophysiologie. 190-212, Berlin-Heidelberg, Springer Verlag Berlin Heidelberg, 1996

**Kadiyala, S., Young, R. G., Thiede, M. A., Bruder, S. P.** (1997): Culture expanded canine mesenchymal stem cells possess osteochondrogenic potential in vivo and in vitro. *Cell Transplant*, 6(2): 125-34.

**Kahn, A., Gibbons, R., Perkins, S., Gazit, D.** (1995): Age-related bone loss. A hypothesis and initial assessment in mice. *Clin Orthop Relat Res*, (313): 69-75.

**Kanou, M., Ueno, T., Kagawa, T., Fujii, T., Sakata, Y., Ishida, N., Fukunaga, J., Sugahara, T.** (2005): Osteogenic potential of primed periosteum graft in the rat calvarial model. *Ann Plast Surg*, 54(1): 71-8.

**Klaushofer, K., Peterlik, M.** (1994): Pathophysiology of fracture healing. *Radiologe*, 34(12): 709-14.

**Kokubu, T., Hak, D. J., Hazelwood, S. J., Reddi, A. H.** (2003): Development of an atrophic nonunion model and comparison to a closed healing fracture in rat femur. *J Orthop Res*, 21(3): 503-10.

**Kon, E., Muraglia, A., Corsi, A., Bianco, P., Marcacci, M., Martin, I., Boyde, A., Ruspantini, I., Chistolini, P., Rocca, M., et al.** (2000): Autologous bone marrow stromal cells loaded onto porous hydroxyapatite ceramic accelerate bone repair in critical-size defects of sheep long bones. *J Biomed Mater Res*, 49(3): 328-37.

**König, H. E., Liebig, H. G.** Anatomie der Haussäugetiere. 8-17, Stuttgart-New York, 2005

- Kopen, G. C., Prockop, D. J., Phinney, D. G.** (1999): Marrow stromal cells migrate throughout forebrain and cerebellum, and they differentiate into astrocytes after injection into neonatal mouse brains. *Proc Natl Acad Sci U S A*, 96(19): 10711-6.
- Kotton, D. N., Ma, B. Y., Cardoso, W. V., Sanderson, E. A., Summer, R. S., Williams, M. C., Fine, A.** (2001): Bone marrow-derived cells as progenitors of lung alveolar epithelium. *Development*, 128(24): 5181-8.
- Kowalski, M. J., Schemitsch, E. H., Kregor, P. J., Senft, D., Swiontkowski, M. F.** (1996): Effect of periosteal stripping on cortical bone perfusion: a laser doppler study in sheep. *Calcif Tissue Int*, 59(1): 24-6.
- Kuner, E. H., Berwarth, H., Lucke, S. V.** (1996): Treatment principles in aseptic pseudarthrosis. *Orthopade*, 25(5): 394-404.
- Kutsch-Lissberg, F., Hopf, K. F.** (2003): Why do osteosyntheses fail? The problem with biomechanics and biology. *Unfallchirurg*, 106(9): 708-21.
- Lane, J. M., Tomin, E., Bostrom, M. P.** (1999): Biosynthetic bone grafting. *Clin Orthop Relat Res*, (367 Suppl): 107-17.
- Lazarus, H. M., Haynesworth, S. E., Gerson, S. L., Rosenthal, N. S., Caplan, A. I.** (1995): Ex vivo expansion and subsequent infusion of human bone marrow-derived stromal progenitor cells (mesenchymal progenitor cells): implications for therapeutic use. *Bone Marrow Transplant*, 16(4): 557-64.
- Le Blanc, K., Ringden, O.** (2006): Mesenchymal stem cells: properties and role in clinical bone marrow transplantation. *Curr Opin Immunol*, 18(5): 586-91.
- Lee, J. W., Kim, Y. H., Kim, S. H., Han, S. H., Hahn, S. B.** (2004): Chondrogenic differentiation of mesenchymal stem cells and its clinical applications. *Yonsei Med J*, 45 Suppl: 41-7.
- Li, G., White, G., Connolly, C., Marsh, D.** (2002): Cell proliferation and apoptosis during fracture healing. *J Bone Miner Res.*, 17(5): 791-9.

**Lieberman, J. R., Daluiski, A., Stevenson, S., Wu, L., McAllister, P., Lee, Y. P., Kabo, J. M., Finerman, G. A., Berk, A. J., Witte, O. N.** (1999): The effect of regional gene therapy with bone morphogenetic protein-2-producing bone-marrow cells on the repair of segmental femoral defects in rats. *J Bone Joint Surg Am*, 81(7): 905-17.

**Liebig, H. G.** Funktionelle Histologie der Haussäugetiere. 72-84, Stuttgart-New York, 1999

**Lienau, J., Schell, H., Duda, G. N., Seebeck, P., Muchow, S., Bail, H. J.** (2005): Initial vascularization and tissue differentiation are influenced by fixation stability. *J Orthop Res*, 23(3): 639-45.

**Lovell-Badge, R.** (2001): The future for stem cell research. *Nature*, 414(6859): 88-91.

**Macnab, I., De Haas, W. G.** (1974): The role of periosteal blood supply in the healing of fractures of the tibia. *Clin Orthop Relat Res*, (105): 27-33.

**Majumdar, M. K., Thiede, M. A., Mosca, J. D., Moorman, M., Gerson, S. L.** (1998): Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *J Cell Physiol*, 176(1): 57-66.

**Makino, T., Hak, D. J., Hazelwood, S. J., Curtiss, S., Reddi, A. H.** (2005): Prevention of atrophic nonunion development by recombinant human bone morphogenetic protein-7. *J Orthop Res*, 23(3): 632-8.

**Malizos, K. N., Papatheodorou, L. K.** (2005): The healing potential of the periosteum molecular aspects. *Injury*, 36 Suppl 3: 13-9.

**Marsh, D.** (1998): Concepts of fracture union, delayed union, and nonunion. *Clin Orthop Relat Res*, (355 Suppl): 22-30.

**Marsh, D. R., Li, G.** (1999): The biology of fracture healing: optimising outcome. *Br Med Bull*, 55(4): 856-69.

**Mathon, D. H., Autefage, A.** (1998): Development of a segmental long-bone defect model in sheep. *Vet Comp Orthop Traumatol*, 11: 1-7.

**McKibbin, B.** (1978): The biology of fracture healing in long bones. *J Bone Joint Surg Br.*, 60-B(2): 150-62.

**Muraglia, A., Cancedda, R., Quarto, R.** (2000): Clonal mesenchymal progenitors from human bone marrow differentiate in vitro according to a hierarchical model. *J Cell Sci*, 113: 1161-6.

**Muschler, G. F., Boehm, C., Easley, K.** (1997): Aspiration to obtain osteoblast progenitor cells from human bone marrow: the influence of aspiration volume. *J Bone Joint Surg Am*, 79(11): 1699-709.

**Muschler, G. F., Nitto, H., Boehm, C. A., Easley, K. A.** (2001): Age- and gender-related changes in the cellularity of human bone marrow and the prevalence of osteoblastic progenitors. *J Orthop Res*, 19(1): 117-25.

**Nakahara, H., Bruder, S. P., Haynesworth, S. E., Holecek, J. J., Baber, M. A., Goldberg, V. M., Caplan, A. I.** (1990): Bone and cartilage formation in diffusion chambers by subcultured cells derived from the periosteum. *Bone*, 11(3): 181-8.

**Ng, K. W., Romas, E., Donnan, L., Findlay, D. M.** (1997): Bone biology. *Baillieres Clin Endocrinol Metab.*, 11(1): 1-22.

**Noble, B. S., Reeve, J.** (2000): Osteocyte function, osteocyte death and bone fracture resistance. *Mol Cell Endocrinol.*, 159(1-2): 7-13.

**Nunamaker, D. M.** (1998): Experimental models of fracture repair. *Clin Orthop Relat Res*, (355 Suppl): 56-65.

**Ohgushi, H., Goldberg, V. M., Caplan, A. I.** (1989): Repair of bone defects with marrow cells and porous ceramic. Experiments in rats. *Acta Orthop Scand*, 60(3): 334-9.

**Okumura, M., Ohgushi, H., Dohi, Y., Katuda, T., Tamai, S., Koerten, H. K., Tabata, S.** (1997): Osteoblastic phenotype expression on the surface of hydroxyapatite ceramics. *J Biomed Mater Res*, 37(1): 122-9.

**Olah, A. J., Simon, A., Gaudy, M., Herrmann, W., Schenk, R. K.** (1977): Differential staining of calcified tissues in plastic embedded microtome sections by a modification of Movat's pentachrome stain. *Stain Technol.*, 52(6): 331-7.

**Owen, M.** (1970): The origin of bone cells. *Int Rev Cytol.*, 28: 213-38.

**Ozaki, A., Tsunoda, M., Kinoshita, S., Saura, R.** (2000): Role of fracture hematoma and periosteum during fracture healing in rats: interaction of fracture hematoma and the periosteum in the initial step of the healing process. *J Orthop Sci.*, 5(1): 64-70.

**Penning, D.** (1990): The biology of bones and of bone fracture healing. *Unfallchirurg.*, 93(11): 488-91.

**Pereira, R. F., Halford, K. W., O'Hara, M. D., Leeper, D. B., Sokolov, B. P., Pollard, M. D., Bagasra, O., Prockop, D. J.** (1995): Cultured adherent cells from marrow can serve as long-lasting precursor cells for bone, cartilage, and lung in irradiated mice. *Proc Natl Acad Sci U S A*, 92(11): 4857-61.

**Petite, H., Viateau, V., Bensaid, W., Meunier, A., de Pollak, C., Bourguignon, M., Oudina, K., Sedel, L., Guillemin, G.** (2000): Tissue-engineered bone regeneration. *Nat Biotechnol.*, 18(9): 959-63.

**Phinney, D. G., Kopen, G., Isaacson, R. L., Prockop, D. J.** (1999): Plastic adherent stromal cells from the bone marrow of commonly used strains of inbred mice: variations in yield, growth, and differentiation. *J Cell Biochem.*, 72(4): 570-85.

**Pittenger, M. F., Mackay, A. M., Beck, S. C., Jaiswal, R. K., Douglas, R., Mosca, J. D., Moorman, M. A., Simonetti, D. W., Craig, S., Marshak, D. R.** (1999): Multilineage potential of adult human mesenchymal stem cells. *Science*, 284(5411): 143-7.

**Probst, A., Spiegel, H. U.** (1997): Cellular mechanisms of bone repair. *J Invest Surg.*, 10(3): 77-86.

**Prockop, D. J.** (1997): Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science*, 276(5309): 71-4.

- Puelacher, W. C., Vacanti, J. P., Ferraro, N. F., Schloo, B., Vacanti, C. A.** (1996): Femoral shaft reconstruction using tissue-engineered growth of bone. *Int J Oral Maxillofac Surg*, 25(3): 223-8.
- Radasch, R. M.** (1999): Biomechanics of bone and fractures. *Vet Clin North Am Small Anim Pract*, 29(5): 1045-82.
- Reddi, A. H.** (1992): Regulation of cartilage and bone differentiation by bone morphogenetic proteins. *Curr Opin Cell Biol*, 4(5): 850-5.
- Redlich, A., Perka, C., Schultz, O., Spitzer, R., Haupl, T., Burmester, G. R., Sittinger, M.** (1999): Bone engineering on the basis of periosteal cells cultured in polymer fleeces. *J Mater Sci Mater Med*, 10(12): 767-72.
- Remedios, A.** (1999): Bone and bone healing. *Vet Clin North Am Small Anim Pract.*, 29(5): 1029-44.
- Rhinelander, F. W.** (1974): Tibial blood supply in relation to fracture healing. *Clin Orthop Relat Res*, (105): 34-81.
- Roach, H. I., Shearer, J. R., Archer, C.** (1989): The choice of an experimental model. A guide for research workers. *J Bone Joint Surg Br*, 71(4): 549-53.
- Runkel, M., Rommens, P. M.** (2000): Pseudoarthrosis. *Unfallchirurg*, 103(1): 51-63.
- Rüter, A., Mayr, E.** (1999): Pseudarthrosis. *Chirurg*, 70(11): 1239-45.
- Sakou, T.** (1998): Bone morphogenetic proteins: from basic studies to clinical approaches. *Bone*, 22(6): 591-603.
- Sanchez-Ramos, J., Song, S., Cardozo-Pelaez, F., Hazzi, C., Stedeford, T., Willing, A., Freeman, T. B., Saporta, S., Janssen, W., Patel, N., et al.** (2000): Adult bone marrow stromal cells differentiate into neural cells in vitro. *Exp Neurol*, 164(2): 247-56.

**Sarmiento, A., Gersten, L. M., Sobol, P. A., Shankwiler, J. A., Vangsness, C. T.** (1989): Tibial shaft fractures treated with functional braces. Experience with 780 fractures. *J Bone Joint Surg Br*, 71(4): 602-9.

**Sarugaser, R., Lickorish, D., Baksh, D., Hosseini, M. M., Davies, J. E.** (2005): Human umbilical cord perivascular (HUCPV) cells: a source of mesenchymal progenitors. *Stem Cells*, 23(2): 220-9.

**Schebitz, H., Brass, W., Wintzer, H.** Allgemeine Chirurgie für Tierärzte und Studiere. 165-168, Verlag Paul Parey, berlin, Hamburg, 1993

**Schell, H., Lienau, J., Epari, D. R., Seebeck, P., Exner, C., Muchow, S., Bragulla, H., Haas, N. P., Duda, G. N.** (2006): Osteoclastic activity begins early and increases over the course of bone healing. *Bone*, 38(4): 547-54.

**Schenk, R. K.** (1975): Histology of primary bone healing. *Fortschr Kiefer Gesichtschir.*, 19: 8-12.

**Schmitz, J. P., Hollinger, J. O.** (1986): The critical size defect as an experimental model for craniomandibulofacial nonunions. *Clin Orthop Relat Res*, (205): 299-308.

**Schmitz, M. A., Finnegan, M., Natarajan, R., Champine, J.** (1999): Effect of smoking on tibial shaft fracture healing. *Clin Orthop Relat Res*, (365): 184-200.

**Schoellner, C., Rompe, J. D., Decking, J., Heine, J.** (2002): High energy extracorporeal shockwave therapy (ESWT) in pseudarthrosis. *Orthopade*, 31(7): 658-62.

**Schweiberer, L., Baumgart, R., Deiler, S.** (1999): The biological reaction in atrophic and hypertrophic pseudarthrosis of diaphysis of long bone. Causes and forms of appearance. *Chirurg*, 70(11): 1193-201.

**Shao, Z., Liu, B., Peng, Q., Liu, W., Liu, Y., Liu, R., Xu, Y., Liu, L.** (2007): Transplantation of osteoblast-like cells to the distracted callus in the rabbit mandible. *Plast Reconstr Surg*, 119(2): 500-7.

- Shirley, D., Marsh, D., Jordan, G., McQuaid, S., Li, G.** (2005): Systemic recruitment of osteoblastic cells in fracture healing. *J Orthop Res*, 23(5): 1013-21.
- Simmons, D. J.** (1985): Fracture healing perspectives. *Clin Orthop Relat Res.*, (200): 100-13.
- Simon, T. M., Van Sickle, D. C., Kunishima, D. H., Jackson, D. W.** (2003): Cambium cell stimulation from surgical release of the periosteum. *J Orthop Res.*, 21(3): 470-80.
- Squier, C. A., Ghoneim, S., Kremenak, C. R.** (1990): Ultrastructure of the periosteum from membrane bone. *J Anat*, 171: 233-9.
- Stürmer, K. M.** (1996): Pathophysiology of disrupted bone healing. *Orthopade.*, 25(5): 386-93.
- Takushima, A., Kitano, Y., Harii, K.** (1998): Osteogenic potential of cultured periosteal cells in a distracted bone gap in rabbits. *J Surg Res*, 78(1): 68-77.
- Tonna, A. E., Cronkite, E. P.** (1963): The periosteum: autoradiographic studies on cellular proliferation and transformation utilizing tritiated thymidine. 218-32.
- Tonna, E. A., Cronkite, E. P.** (1961): Use of tritiated thymidine for the study of the origin of the osteoclast. *Nature*, 190: 459-60.
- Trostle, S. S., Markel, M. D.** (1996): Fracture biology, biomechanics, and internal fixation. *Vet Clin North Am Food Anim Pract.*, 12(1): 19-46.
- Tuan, R. S., Boland, G., Tuli, R.** (2003): Adult mesenchymal stem cells and cell-based tissue engineering. *Arthritis Res Ther*, 5(1): 32-45.
- Vögelin, E., Brekke, J. H., Jones, N. F.** (2000): Heterotopic and orthotopic bone formation with a vascularized periosteal flap, a matrix and rh-BMP-2 (bone morphogenetic protein) in the rat model. *Mund Kiefer Gesichtschir*, 4 Suppl 2: 454-8.
- Volpon, J. B.** (1994): Nonunion using a canine model. *Arch Orthop Trauma Surg*, 113(6): 312-7.

**Wakitani, S., Saito, T., Caplan, A. I.** (1995): Myogenic cells derived from rat bone marrow mesenchymal stem cells exposed to 5-azacytidine. *Muscle Nerve*, 18(12): 1417-26.

**Webb, J. C. J., Tricker, J.** (2000): Bone biology- a review of fracture healing. *Current Orthopaedics*, 12: 457-63.

**Weber, B.** (1982): Klassifizierung der Pseudarthrosen nach morphologischen und klinischen Gesichtspunkten. *Z Orthop*, 120: 528-531.

**Weinand, C., Pomerantseva, I., Neville, C. M., Gupta, R., Weinberg, E., Madisch, I., Shapiro, F., Abukawa, H., Troulis, M. J., Vacanti, J. P.** (2006): Hydrogel-beta-TCP scaffolds and stem cells for tissue engineering bone. *Bone*, 38(4): 555-63.

**Weiner, S., Traub, W., Wagner, H. D.** (1999): Lamellar bone: structure-function relations. *J Struct Biol.*, 126(3): 241-55.

**Weiner, S., Wagner, H. D.** (1998): The material bone: structure-mechanical function relations. *Annu Rev Mater Sci*, 28: 271-98.

**Willenegger, H., Perren, S. M., Schenk, R.** (1971): Primary and secondary healing of bone fractures. *Chirurg*, 42(6): 241-52.

**Wright, C. D., Vedi, S., Garrahan, N. J., Stanton, M., Duffy, S. W., Compston, J. E.** (1992): Combined inter-observer and inter-method variation in bone histomorphometry. *Bone*, 13(3): 205-8.

**Würzler, K. K., Kubler, N. R., Reuther, J. F.** (2000): Effect of periosteum on induced bone formation by autolyzed, antigen-extracted, allogeneic bone. Determination of the extent of bone formation using quantitative computerized tomography. *Mund Kiefer Gesichtschir*, 4 Suppl 2: 459-64.

**Yoo, J. U., Johnstone, B.** (1998): The role of osteochondral progenitor cells in fracture repair. *Clin Orthop Relat Res*, (355 Suppl): 73-81.

**Yoshikawa, T., Ohgushi, H., Tamai, S.** (1996): Immediate bone forming capability of prefabricated osteogenic hydroxyapatite. *J Biomed Mater Res*, 32(3): 481-92.

**Zvaipler, N. J., Marinova-Mutafchieva, L., Adams, G., Edwards, C. J., Moss, J., Burger, J. A., Maini, R. N.** (2000): Mesenchymal precursor cells in the blood of normal individuals. *Arthritis Res*, 2(6): 477-88.