

8 Literaturverzeichnis

- Adachi H, Konishi K, Horikoshi I.** The effects of 1,2,3,4,6-penta-O-galloyl-beta-D-glucose on rat liver mitochondrial respiration. *Chem Pharm Bull (Tokyo)*, 1989, **37**(5): 1341-4
- Affar EB, Duriez PJ, Shah RG, Sallmann FR, Bourassa S, Kupper JH, Burkle A, Poirier GG.** Immunodot blot method for the detection of poly(ADP-ribose) synthesized in vitro and in vivo. *Anal Biochem*, 1998, **259**(2), 280-3
- Affar EB, Germain M, Winstall E, Vodenicharov M, Shah RG, Salvesen GS, Poirier GG.** Caspase-3-mediated processing of poly(ADP-ribose) glycohydrolase during apoptosis. *J Biol Chem*, 2001, **276**(4), 2935-42
- Althaus FR, Richter C.** ADP-Ribosylation of proteins. *Mol. Biol. Biochem. Biophys.*, 1987, **37**, 1-125
- Althaus FR.** Poly ADP-ribosylation: a histone shuttle mechanism in DNA excision repair. *J Cell Sci*, 1992, **102**(4), 663-70
- Alvarez-Gonzalez R, Pacheco-Rodriguez G, Mendoza-Alvarez H.** Enzymology of ADP-ribose polymer synthesis. *Mol Cell Biochem*, 1994, **138**(1-2), 33-7
- Ame JC, Jacobson EL, Jacobson MK.** Molecular heterogeneity and regulation of poly(ADP-ribose) glycohydrolase. *Mol Cell Biochem*, 1999, **193**(1-2), 75-81
- Ame JC, Spenlehauer C, de Murcia G.** The PARP superfamily. *Bioessays*, 2004, **26**(8), 882-93
- Atorino L, Di Meglio S, Farina B, Jones R, Quesada P.** Rat germinal cells require PARP for repair of DNA damage induced by gamma-irradiation and H₂O₂ treatment. *Eur J Cell Biol*, 2001, **80**(3), 222-29
- Aoki K, Nishimura K, Abe H, Maruta H, Sakagami H, Hatano T, Okuda T, Yoshida T, Tsai YJ, Uchiumi F, et al.** Novel inhibitors of poly(ADP-ribose) glycohydrolase. *Biochim Biophys Acta*, 1993, **1158**(3), 251-6
- Aoki K, Maruta H, Uchiumi F, Hatano T, Yoshida T, Tanuma S.** A macrocircular ellagitannin, oenothein B, suppresses mouse mammary tumor gene expression via inhibition of poly(ADP-ribose) glycohydrolase. *Biochem Biophys Res Commun*, 1995, **210**(2), 329-37
- Ariumi Y, Masutani M, Copeland TD, Mimori T, Sugimura T, Shimotohno K, Ueda K, Hatanaka M, Noda M.** Suppression of the poly(ADP-ribose) polymerase activity by DNA-dependent protein kinase in vitro. *Oncogene*, 1999, **18**(32), 4616-25
- Au WW, Salama SA, Sierra-Torres CH.** Functional characterization of polymorphisms in DNA repair genes using cytogenetic challenge assays. *Environ Health Perspect*, 2003, **111**(15), 1843-50
- Audebert M, Salles B, Calsou P.** Involvement of poly(ADP-ribose) polymerase-1 and XRCC1/DNA ligase III in an alternative route for DNA double-strand breaks rejoining. *J Biol Chem*, 2004, **279**(53), 55117-26

- Augustin A, Spenlehauer C, Dumond H, Menissier-De Murcia J, Piel M, Schmit AC, Apiou F, Vonesch JL, Kock M, Bornens M, De Murcia G.** PARP-3 localizes preferentially to the daughter centriole and interferes with the G1/S cell cycle progression. *J Cell Sci.*, 2003, **116(8)**, 1551-62
- Austin CJ, Mizdrak J, Matin A, Sirijovski N, Kosim-Satyaputra P, Willows RD, Roberts TH, Truscott RJ, Polekhina G, Parker MW, Jamie JF.** Optimised expression and purification of recombinant human indoleamine 2,3-dioxygenase. *Protein Expr Purif.*, 2004, **37(2)**, 392-8
- Bakondi E, Bai P, Erdelyi K, Szabo C, Gergely P, Virág L.** Cytoprotective effect of gallotannin in oxidatively stressed HaCaT keratinocytes: the role of poly(ADP-ribose) metabolism. *Exp Dermatol.*, 2004, **13(3)**, 170-8
- Bauer PI, Buki KG, Hakam A, Kun E.** Macromolecular association of ADP-ribosyltransferase and its correlation with enzymic activity. *Biochem J.*, 1990, **270(1)**, 17-26
- Beardsley DI, Kim WJ, Brown KD.** N-methyl-N'-nitro-N-nitrosoguanidine activates cell-cycle arrest through distinct mechanisms activated in a dose-dependent manner. *Mol Pharmacol.*, 2005, **68(4)**, 1049-60
- Beausoleil SA, Jedrychowski M, Schwartz D, Elias JE, Villen J, Li J, Cohn MA, Cantley LC, Gygi SP.** Large-scale characterization of HeLa cell nuclear phosphoproteins. *Proc Natl Acad Sci USA.*, 2004, **101(33)**, 12130-5
- Beernink PT, Hwang M, Ramirez M, Murphy MB, Doyle SA, Thelen MP.** Specificity of protein interactions mediated by BRCT domains of the XRCC1 DNA repair protein. *J Biol Chem.*, 2005, **280(34)**, 30206-13
- Bellocchi D, Costantino G, Pellicciari R, Re N, Marrone A, Coletti C.** Poly(ADP-Ribose)-Polymerase-Catalyzed Hydrolysis of NAD(+): QM/MM Simulation of the Enzyme Reaction. *ChemMedChem.*, 2006, **1(5)**, 533-539
- Benjamin RC, Gill DM.** ADP-ribosylation in mammalian cell ghosts. Dependence of poly(ADP-ribose) synthesis on strand breakage in DNA. *J Biol Chem.*, 1980, **255(21)**, 10493-501
- Bennick A.** Interaction of plant polyphenols with salivary proteins. *Crit Rev Oral Biol Med.*, 2002, **13(2)**, 184-96
- Berger NA.** Poly(ADP-ribose) in the cellular response to DNA damage. *Radiat Res.*, 1985, **101(1)**, 4-15
- Berger F, Ramirez-Hernandez MH, Ziegler M.** The new life of a centenarian: signalling functions of NAD(P). *Trends Biochem Sci.*, 2004, **29(3)**, 111-8
- Berger F, Lau C, Dahlmann M, Ziegler M.** Subcellular compartmentation and differential catalytic properties of the three human nicotinamide mononucleotide adenylyltransferase isoforms. *J Biol Chem.*, 2005, **280(43)**, 36334-41
- Berghammer H, Auer B.** "Easypreps": fast and easy plasmid minipreparation for analysis of recombinant clones in *E. coli*. *Biotechniques.*, 1993, **14(4)**, 524
- Bernstein C, Bernstein H, Payne CM, Garewal H.** DNA repair/pro-apoptotic dual-role proteins in five major DNA repair pathways: fail-safe protection against carcinogenesis. *Mutat Res.*, 2002, **511(2)**, 145-78

- Bertram B, Bollow U, Rajaei-Behbahani N, Burkle A, Schmezer P.** Induction of poly(ADP-ribosylation) and DNA damage in human peripheral lymphocytes after treatment with (-)-epigallocatechin-gallate. *Mutat Res.*, 2003, **534(1-2)**, 77-84
- Blackwell JR, Horgan R.** A novel strategy for production of a highly expressed recombinant protein in an active form. *FEBS Lett.*, **295(1-3)**, 10-2
- Blenn C, Althaus FR, Malanga M.** Poly(ADP-ribose) glycohydrolase silencing protects against H₂O₂-induced cell death. *Biochem J.*, 2006, **396(3)**, 419-29
- Bonicalzi ME, Vodenicharov M, Coulombe M, Gagne JP, Poirier GG.** Alteration of poly(ADP-ribose) glycohydrolase nucleocytoplasmic shuttling characteristics upon cleavage by apoptotic proteases. *Biol Cell.*, 2003, **95(9)**, 635-44
- Bonicalzi ME, Haince JF, Droit A, Poirier GG.** Regulation of poly(ADP-ribose) metabolism by poly(ADP-ribose) glycohydrolase: where and when? *Cell Mol Life Sci.*, 2005, **62(7-8)**, 739-50
- Bouchard VJ, Rouleau M, Poirier GG.** PARP-1, a determinant of cell survival in response to DNA damage. *Exp Hematol.*, 2003, **31(6)**: 446-54
- Boulikas T.** Poly(ADP-ribose) synthesis and degradation in mammalian nuclei. *Anal Biochem.*, 1992, **203(2)**, 252-8
- Bradford MM.** A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.*, 1976, **72**, 248-54
- Bratton SB, Cohen GM.** Apoptotic death sensor: an organelle's alter ego? *Trends Pharmacol Sci.*, 2001, **22(6)**, 306-15
- Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E, Kyle S, Meuth M, Curtin NJ, Helleday T.** Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature.*, 2005, **434(7035)**, 913-7
- Bujard H, Gentz R, Lanzer M, Stueber D, Mueller M, Ibrahimi I, Haeuptle MT, Dobberstein B.** A T5 promoter-based transcription-translation system for the analysis of proteins in vitro and in vivo. *Methods Enzymol.*, 1987, **155**, 416-33
- Buki KG, Bauer PI, Hakam A, Kun E.** Identification of domains of poly(ADP-ribose) polymerase for protein binding and self-association. *J Biol Chem.*, 1995, **270(7)**, 3370-7
- Burzio LO, Riquelme PT, Ohtsuka E, Koide SS.** Evidence for two variants of poly(adenosine diphosphate ribose) glycohydrolase in rat testis. *Arch Biochem Biophys.*, 1976, **173(1)**, 306-19
- Cahill D, Connor B, Carney JP.** Mechanisms of eukaryotic DNA double strand break repair. *Front Biosci.*, 2006, **11**, 1958-76
- Caldecott KW, McKeown CK, Tucker JD, Ljungquist S, Thompson LH.** An interaction between the mammalian DNA repair protein XRCC1 and DNA ligase III. *Mol Cell Biol.*, 1994, **14(1)**, 8-76
- Caldecott KW, Tucker JD, Stanker LH, Thompson LH.** Characterization of the XRCC1-DNA ligase III complex in vitro and its absence from mutant hamster cells. *Nucleic Acids Res.*, 1995, **23(23)**, 4836-43

- Caldecott KW, Aoufouchi S, Johnson P, Shall S.** XRCC1 polypeptide interacts with DNA polymerase beta and possibly poly (ADP-ribose) polymerase, and DNA ligase III is a novel molecular 'nick-sensor' in vitro. *Nucleic Acids Res.*, 1996, **24(22)**, 4387-94
- Caldecott KW.** Mammalian DNA single-strand break repair: an X-ra(y)ted affair. *Bioessays.*, 2001, **23(5)**, 447-55
- Caldecott KW.** XRCC1 and DNA strand break repair. *DNA Repair (Amst.)*, 2003, **2(9)**, 955-69
- Cande C, Cohen I, Daugas E, Ravagnan L, Larochette N, Zamzami N, Kroemer G.** Apoptosis-inducing factor (AIF): a novel caspase-independent death effector released from mitochondria. *Biochimie.*, 2002, **84(2-3)**, 215-22
- Cappelli E, Taylor R, Cevasco M, Abbondandolo A, Caldecott K, Frosina G.** Involvement of XRCC1 and DNA ligase III gene products in DNA base excision repair. *J Biol Chem.*, 1997, **272(38)**, 3970-5
- Chambon P, Weill JD, Mandel P.** Nicotinamide mononucleotide activation of new DNA-dependent polyadenylic acid synthesizing nuclear enzyme. *Biochem Biophys Res Commun.*, 1963, **11**, 39-43
- Chambon P, Weill. JD, Doly J, Strosser MT, Mandel P.** On the formation of a novel adenylic compound by enzymatic extracts of liver nuclei. *Biochem Biophys Res Commun.*, 1966, **6**, 638-643
- Chang P, Coughlin M, Mitchison TJ.** Tankyrase-1 polymerization of poly(ADP-ribose) is required for spindle structure and function. *Nat Cell Biol.*, 2005, **7(11)**, 1133-9
- Chang W, Dynek JN, Smith S.** NuMA is a major acceptor of poly(ADP-ribosylation) by tankyrase 1 in mitosis. *Biochem J.*, 2005, **391(2)**, 177-84
- Chi NW, Lodish HF.** Tankyrase is a golgi-associated mitogen-activated protein kinase substrate that interacts with IRAP in GLUT4 vesicles. *J Biol Chem.*, 2000, **275(49)**, 38437-44
- Chiarugi A.** Poly(ADP-ribose) polymerase: killer or conspirator? The 'suicide hypothesis' revisited. *Trends Pharmacol Sci.*, 2002, **23(3)**, 122-9
- Chiarugi A.** "Simple but not simpler": toward a unified picture of energy requirements in cell death. *FASEB J.*, 2005, **19(13)**, 1783-8
- Christmann M, Tomicic MT, Roos WP, Kaina B.** Mechanisms of human DNA repair: an update. *Toxicology.*, 2003, **193(1-2)**, 3-34
- Cistulli C, Lavrik OI, Prasad R, Hou E, Wilson SH.** AP endonuclease and poly(ADP-ribose) polymerase-1 interact with the same base excision repair intermediate. *DNA Repair (Amst.)*, 2004, **3(6)**, 581-91
- Clements PM, Breslin C, Deeks ED, Byrd PJ, Ju L, Bieganski P, Brenner C, Moreira MC, Taylor AM, Caldecott KW.** The ataxia-oculomotor apraxia 1 gene product has a role distinct from ATM and interacts with the DNA strand break repair proteins XRCC1 and XRCC4. *DNA Repair (Amst.)*, 2004, **3(11)**, 1493-502
- Cook BD, Dynek JN, Chang W, Shostak G, Smith S.** Role for the related poly(ADP-Ribose) polymerases tankyrase 1 and 2 at human telomeres. *Mol Cell Biol.*, 2002, **22(1)**, 332-42

- Cooke MS, Evans MD, Dizdaroglu M, Lunec J.** Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J.*, 2003, **17(10)**, 1195-214
- Cortes U, Tong WM, Coyle DL, Meyer-Ficca ML, Meyer RG, Petrilli V, Herceg Z, Jacobson EL, Jacobson MK, Wang ZQ.** Depletion of the 110-kilodalton isoform of poly(ADP-ribose) glycohydrolase increases sensitivity to genotoxic and endotoxic stress in mice. *Mol Cell Biol.*, 2004, **24(16)**, 7163-78
- Cottet F, Blanche H, Verasdonck P, Le Gall I, Schachter F, Burkle A, Muiras ML.** New polymorphisms in the human poly(ADP-ribose) polymerase-1 coding sequence: lack of association with longevity or with increased cellular poly(ADP-ribosyl)ation capacity. *J Mol Med.*, 2000, **78(8)**, 431-40
- Cregan SP, Dawson VL, Slack RS.** Role of AIF in caspase-dependent and caspase-independent cell death. *Oncogene.*, 2004, **23(16)**, 2785-96
- Curtin NJ.** PARP inhibitors for cancer therapy. *Expert Rev Mol Med.*, 2005, **7(4)**, 1-20
- Cuzzocrea S, Wang ZQ.** Role of poly(ADP-ribose) glycohydrolase (PARG) in shock, ischemia and reperfusion. *Pharmacol Res.*, 2005, **52(1)**, 100-8
- d'Adda di Fagagna F, Hande MP, Tong WM, Lansdorp PM, Wang ZQ, Jackson SP.** Functions of poly(ADP-ribose) polymerase in controlling telomere length and chromosomal stability. *Nat Genet.*, 1999, **23(1)**, 76-80
- D'Amours D, Desnoyers S, D'Silva I, Poirier GG.** Poly(ADP-ribosyl)ation reactions in the regulation of nuclear functions. *Biochem J.*, 1999, **342 (2)**, 249-68
- D'Amours D, Sallmann FR, Dixit VM, Poirier GG.** Gain-of-function of poly(ADP-ribose) polymerase-1 upon cleavage by apoptotic proteases: implications for apoptosis. *J Cell Sci.*, 2001, **114(20)**, 3771-8
- Dantzer F, Nasheuer HP, Vonesch JL, de Murcia G, Menissier-de Murcia J.** Functional association of poly(ADP-ribose) polymerase with DNA polymerase alpha-primase complex: a link between DNA strand break detection and DNA replication. *Nucleic Acids Res.*, 1998, **26(8)**, 1891-8
- Dantzer F, de La Rubia G, Menissier-De Murcia J, Hostomsky Z, de Murcia G, Schreiber V.** Base excision repair is impaired in mammalian cells lacking Poly(ADP-ribose) polymerase-1. *Biochemistry.*, 2000, **39(25)**, 7559-69
- Dantzer F, Giraud-Panis MJ, Jaco I, Ame JC, Schultz I, Blasco M, Koering CE, Gilson E, Menissier-de Murcia J, de Murcia G, Schreiber V.** Functional interaction between poly(ADP-Ribose) polymerase 2 (PARP-2) and TRF2: PARP activity negatively regulates TRF2. *Mol Cell Biol.*, 2004, **24(4)**, 1595-607
- Daugas E, Susin SA, Zamzami N, Ferri KF, Irinopoulou T, Larochette N, Prevost MC, Leber B, Andrews D, Penninger J, Kroemer G.** Mitochondrio-nuclear translocation of AIF in apoptosis and necrosis. *FASEB J.*, 2000, **14(5)**, 729-39
- Davidovic L, Vodenicharov M, Affar EB, Poirier GG.** Importance of poly(ADP-ribose) glycohydrolase in the control of poly(ADP-ribose) metabolism. *Exp Cell Res.*, 2001, **268(1)** 7-13

- Desmarais Y, Menard L, Lagueux J, Poirier GG.** Enzymological properties of poly(ADP-ribose)polymerase: characterization of automodification sites and NADase activity. *Biochim Biophys Acta.*, 1991, **1078**(2), 179-86
- de Murcia G, Schreiber V, Molinete M, Saulier B, Poch O, Masson M, Niedergang C, Menissier de Murcia J.** Structure and function of poly(ADP-ribose) polymerase. *Mol Cell Biochem.*, 1994, **138**(1-2), 15-24
- Desnoyers S, Shah GM, Brochu G, Hoflack JC, Verreault A, Poirier GG.** Biochemical properties and function of poly(ADP-ribose) glycohydrolase. *Biochimie.*, 1995, **77**(6), 433-8
- Desnoyers S, Kaufmann SH, Poirier GG.** Alteration of the nucleolar localization of poly(ADP-ribose) polymerase upon treatment with transcription inhibitors. *Exp Cell Res.*, 1996, **227**(1), 146-53
- Desnoyers S., St-Laurent JF., Dequen F., Gagnon S.** Poly (ADP-ribose) glycohydrolases are needed for survival after DNA damage in the nematode *C. elegans*. 2005, *Med Sci Monit.*, **11**(1), 42
- Diefenbach J, Burkle A.** Introduction to poly(ADP-ribose) metabolism. *Cell Mol Life Sci.*, 2005, **62**(7-8), 721-30
- Drablos F, Feyzi E, Aas PA, Vaagbo CB, Kavli B, Bratlie MS, Pena-Diaz J, Otterlei M, Slupphaug G, Krokan HE.** Alkylation damage in DNA and RNA--repair mechanisms and medical significance. *DNA Repair (Amst.)*, 2004, **3**(11), 1389-407
- Dignam JD, Lebovitz RM, Roeder RG.** Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.* 1983, **11**(5), 1475-89
- Du L, Zhang X, Han YY, Burke NA, Kochanek PM, Watkins SC, Graham SH, Carcillo JA, Szabo C, Clark RS.** Intra-mitochondrial poly(ADP-ribosylation) contributes to NAD⁺ depletion and cell death induced by oxidative stress. *J Biol Chem.*, 2003, **278**(20), 18426-33
- Dumitriu IE, Voll RE, Kolowos W, Gaippl US, Heyder P, Kalden JR, Herrmann M.** UV irradiation inhibits ABC transporters via generation of ADP-ribose by concerted action of poly(ADP-ribose) polymerase-1 and glycohydrolase. *Cell Death Differ.*, 2004, **11**(3), 314-20
- Durkacz BW, Omidiji O, Gray DA, Shall S.** (ADP-ribose)n participates in DNA excision repair. *Nature.*, 1980, **283**(5747), 593-6
- Dyne JN, Smith S.** Resolution of sister telomere association is required for progression through mitosis. *Science.*, 2004, **304**(5667), 97-100
- Earnshaw WC, Martins LM, Kaufmann SH.** Mammalian caspases: structure, activation, substrates, and functions during apoptosis. *Annu Rev Biochem.*, 1999, **68**, 383-424
- Edinger AL, Thompson CB.** Death by design: apoptosis, necrosis and autophagy. *Curr Opin Cell Biol.*, 2004, **16**(6), 663-9
- Eguchi Y, Srinivasan A, Tomaselli KJ, Shimizu S, Tsujimoto Y.** ATP-dependent steps in apoptotic signal transduction. *Cancer Res.*, 1999, **59**(9), 2174-81
- El-Khamisy SF, Masutani M, Suzuki H, Caldecott KW.** A requirement for PARP-1 for the assembly or stability of XRCC1 nuclear foci at sites of oxidative DNA damage. *Nucleic Acids Res.*, 2003, **31**(19), 5526-33

- Erdelyi K, Kiss A, Bakondi E, Bai P, Szabo C, Gergely P, Erdodi F, Virag L.** Gallotannin inhibits the expression of chemokines and inflammatory cytokines in A549 cells. *Mol Pharmacol.*, 2005, **68(3)**, 895-904
- Falsig J, Christiansen SH, Feuerhahn S, Burkle A, Oei SL, Keil C, Leist M.** Poly(ADP-ribose) glycohydrolase as a target for neuroprotective intervention: assessment of currently available pharmacological tools. *Eur J Pharmacol.*, 2004, **497(1)**, 7-16
- Fan J, Otterlei M, Wong HK, Tomkinson AE, Wilson DM 3rd.** XRCC1 co-localizes and physically interacts with PCNA. *Nucleic Acids Res.*, 2004, **32(7)**, 2193-201
- Fan J, Wilson DM 3rd.** Protein-protein interactions and posttranslational modifications in mammalian base excision repair. *Free Radic Biol Med.*, 2005, **38(9)**, 1121-38
- Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB, Santarosa M, Dillon KJ, Hickson I, Knights C, Martin NM, Jackson SP, Smith GC, Ashworth A.** Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature.*, 2005, **434(7035)**, 917-21
- Ferri KF, Kroemer G.** Mitochondria--the suicide organelles. *Bioessays.*, 2001, **23(2)**, 111-5
- Finlay TH, Troll V, Levy M, Johnson AJ, Hodgins LT.** New methods for the preparation of biospecific adsorbents and immobilized enzymes utilizing trichloro-s-triazine. *Anal Biochem.*, 1978, **87(1)**, 77-90
- Frangioni JV, Neel BG.** Solubilization and purification of enzymatically active glutathione S-transferase (pGEX) fusion proteins. *Anal Biochem.*, 1993, **210(1)**, 179-87
- Frosina G.** Commentary: DNA base excision repair defects in human pathologies. *Free Radic Res.*, 2004, **38(10)**, 1037-54
- Gagne JP, Hunter JM, Labrecque B, Chabot B, Poirier GG.** A proteomic approach to the identification of heterogeneous nuclear ribonucleoproteins as a new family of poly(ADP-ribose)-binding proteins. *Biochem J.*, 2003, **371(2)**, 331-40
- Gagne JP, Bonicalzi ME, Gagne P, Ouellet ME, Hendzel MJ, Poirier GG.** Poly(ADP-ribose) glycohydrolase is a component of the FMRP-associated messenger ribonucleoparticles. *Biochem J.*, 2005, **392(3)**, 499-509
- Gagne JP, Hendzel MJ, Droit A, Poirier GG.** The expanding role of poly(ADP-ribose) metabolism: current challenges and new perspectives. *Curr Opin Cell Biol.*, 2006, **18(2)**, 145-51
- Genovese T, Di Paola R, Catalano P, Li JH, Xu W, Massuda E, Caputi AP, Zhang J, Cuzzocrea S.** Treatment with a novel poly(ADP-ribose) glycohydrolase inhibitor reduces development of septic shock-like syndrome induced by zymosan in mice. *Crit Care Med.*, 2004, **32(6)**, 1365-74
- Germain M, Affar EB, D'Amours D, Dixit VM, Salvesen GS, Poirier GG.** Cleavage of automodified poly(ADP-ribose) polymerase during apoptosis. Evidence for involvement of caspase-7. *J Biol Chem.*, 1999, **274(40)**, 28379-84
- Geyer R, Bennett A, Rohr A.** Fatty acids of the triglycerides and phospholipids of HeLa cells and strain L fibroblasts. *J Lipid Res.*, 1962, **3(1)**, 80-83

- Giuliano M, Bellavia G, Lauricella M, D'Anneo A, Vassallo B, Vento R, Tesoriere G.** Staurosporine-induced apoptosis in Chang liver cells is associated with down-regulation of Bcl-2 and Bcl-XL. *Int J Mol Med.*, 2004, **13(4)**, 565-71
- Gomez M, Wu J, Schreiber V, Dunlap J, Dantzer F, Wang Y, Liu Y.** PARP1 Is a TRF2-associated poly(ADP-ribose)polymerase and protects eroded telomeres. *Mol Biol Cell.*, 2006, **17(4)**, 1686-96
- Goth-Goldstein R, Hughes M.** Cell killing by various monofunctional alkylating agents in Chinese hamster ovary cells. *Mutat Res.*, 1987, **177(2)**, 267-76
- Griesenbeck J, Oei SL, Mayer-Kuckuk P, Ziegler M, Buchlow G, Schweiger M.** Protein-protein interaction of the human poly(ADP-ribosyl)transferase depends on the functional state of the enzyme. *Biochemistry*, 1997, **36(24)**, 7297-304
- Gueven N, Becherel OJ, Kijas AW, Chen P, Howe O, Rudolph JH, Gatti R, Date H, Onodera O, Taucher-Scholz G, Lavin MF.** Aprataxin, a novel protein that protects against genotoxic stress. *Hum Mol Genet.*, 2004, **13(10)**, 1081-93
- Ha HC, Snyder SH.** Poly(ADP-ribose) polymerase is a mediator of necrotic cell death by ATP depletion. *Proc Natl Acad Sci USA.*, 1999, **96(24)**, 13978-82
- Hagerman AE, Klucher KM.** Tannin-protein interactions. *Prog Clin Biol Res.*, 1986, **213**, 67-76
- Haince JF, Ouellet ME, McDonald D, Hendzel MJ, Poirier GG.** Dynamic relocation of poly(ADP-ribose) glycohydrolase isoforms during radiation-induced DNA damage. *Biochim Biophys Acta.*, 2006, **1763(2)**, 226-37
- Hayashi K, Tanaka M, Shimada T, Miwa M, Sugimura T.** Size and shape of poly(ADP-ribose): examination by gel filtration, gel electrophoresis and electron microscopy. *Biochem Biophys Res Commun.*, 1983, **112(1)**, 102-7
- Hanai S, Kanai M, Ohashi S, Okamoto K, Yamada M, Takahashi H, Miwa M.** Loss of poly(ADP-ribose) glycohydrolase causes progressive neurodegeneration in *Drosophila melanogaster*. *Proc Natl Acad Sci USA.*, 2004, **101(1)**, 82-6
- Haslam E.** Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J Nat Prod.*, 1996, **59(2)**, 205-15
- Hassa PO, Buerki C, Lombardi C, Imhof R, Hottiger MO.** Transcriptional coactivation of nuclear factor-kappaB-dependent gene expression by p300 is regulated by poly(ADP)-ribose polymerase-1. *J Biol Chem.*, 2003, **278(46)**, 45145-53
- Hatakeyama K, Nemoto Y, Ueda K, Hayaishi O.** Purification and characterization of poly(ADP-ribose) glycohydrolase. Different modes of action on large and small poly(ADP-ribose). *J Biol Chem.*, 1986, **261(32)**, 14902-11
- Hayaishi O, Ueda K.** Poly(ADP-ribose) and ADP-ribosylation of proteins. *Annu Rev Biochem.*, 1977, **46**, 95-116
- Hengartner MO.** The biochemistry of apoptosis. *Nature.*, 2000, **407(6805)**, 770-6

- Herceg Z, Wang ZQ.** Functions of poly(ADP-ribose) polymerase (PARP) in DNA repair, genomic integrity and cell death. *Mutat Res.*, 2001, **477(1-2)**, 97-110
- Hoeijmakers JH.** Genome maintenance mechanisms for preventing cancer. *Nature.*, 2001, **411(6835)**, 366-74
- Hong SJ, Dawson TM, Dawson VL.** Nuclear and mitochondrial conversations in cell death: PARP-1 and AIF signaling. *Trends Pharmacol Sci.*, 2004, **25(5)**, 259-64
- Huh NW, Porter NA, McIntosh TJ, Simon SA.** The interaction of polyphenols with bilayers: conditions for increasing bilayer adhesion. *Biophys J.*, 1996, **71(6)**, 3261-77
- Iftner T, Elbel M, Schopp B, Hiller T, Loizou JI, Caldecott KW, Stubenrauch F.** Interference of papillomavirus E6 protein with single-strand break repair by interaction with XRCC1. *EMBO J.*, 2002, **21(17)**,: 4741-8
- Ikejima M, Bohannon D, Gill DM, Thompson LH.** Poly(ADP-ribose) metabolism appears normal in EM9, a mutagen-sensitive mutant of CHO cells. *Mutat Res.*, 1984, **128(2)**, 213-20
- Ikejima M, Gill DM.** Poly(ADP-ribose) degradation by glycohydrolase starts with an endonucleolytic incision. *J Biol Chem.*, 1988, **263(23)**, 11037-40
- Jacobson EL, Jacobson MK.** Tissue NAD as a biochemical measure of niacin status in humans. *Methods Enzymol.*, 1997, **280**, 221-30
- Jacobson EL, Cervantes-Laurean D, Jacobson MK.** ADP-ribose in glycation and glycoxidation reactions. *Adv Exp Med Biol.*, 1997, **419**, 371-9
- Jain AK, Shimoji K, Nakamura Y, Kada T, Hara Y, Tomita I.** Crude tea extracts decrease the mutagenic activity of N-methyl-N'-nitro-N-nitrosoguanidine in vitro and in intragastric tract of rats. *Mutat Res.*, 1989, **210(1)**, 1-8
- James P, Halladay J, Craig EA.** Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast. *Genetics*, 1996, **144(4)**, 1425-36
- Jonsson GG, Menard L, Jacobson EL, Poirier GG, Jacobson MK.** Effect of hyperthermia on poly(adenosine diphosphate-ribose) glycohydrolase. *Cancer Res.*, 1988, **48(15)**, 4240-3
- Kameshita I, Matsuda Z, Taniguchi T, Shizuta Y.** Poly (ADP-Ribose) synthetase. Separation and identification of three proteolytic fragments as the substrate-binding domain, the DNA-binding domain, and the automodification domain. *J Biol Chem.*, 1984, **259(8)**, 4770-6
- Kanai M, Miwa M, Kuchino Y, Sugimura T.** Presence of branched portion in poly(adenosine diphosphate ribose) in vivo. *J Biol Chem.*, 1982, **257(11)**, 6217-23
- Kanai M, Tong WM, Sugihara E, Wang ZQ, Fukasawa K, Miwa M.** Involvement of poly(ADP-Ribose) polymerase 1 and poly(ADP-Ribosylation) in regulation of centrosome function. *Mol Cell Biol.*, 2003, **23(7)**, 2451-62
- Kandra L, Gyemant G, Zajacz A, Batta G.** Inhibitory effects of tannin on human salivary alpha-amylase. *Biochem Biophys Res Commun.*, 2004, **319(4)**, 1265-71
- Kashiwada Y, Nonaka G, Nishioka I, Lee KJ, Bori I, Fukushima Y, Bastow KF, Lee KH.** Tannins as potent inhibitors of DNA topoisomerase II in vitro. *J Pharm Sci.*, 1993, **82(5)**, 487-92

- Kaufmann SH, Desnoyers S, Ottaviano Y, Davidson NE, Poirier GG.** Specific proteolytic cleavage of poly(ADP-ribose) polymerase: an early marker of chemotherapy-induced apoptosis. *Cancer Res.*, 1993, **53(17)**, 3976-85
- Kawaichi M, Ueda K, Hayaishi O.** Initiation of poly(ADP-ribosyl) histone synthesis by poly(ADP-ribose) synthetase. *J Biol Chem.*, 1980, **255(3)**, 816-9
- Kawamitsu H, Hoshino H, Okada H, Miwa M, Momoi H, Sugimura T.** Monoclonal antibodies to poly(adenosine diphosphate ribose) recognize different structures. *Biochemistry.*, 1984, **23(16)**, 3771-77
- Kerr JF, Wyllie AH, Currie AR.** Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer.*, 1972, **26(4)**, 239-57
- Kim JW, Kim K, Kang K, Joe CO.** Inhibition of homodimerization of poly(ADP-ribose) polymerase by its C-terminal cleavage products produced during apoptosis. *J Biol Chem.*, 2000, **275(11)**, 8121-5
- Kim MY, Mauro S, Gevry N, Lis JT, Kraus WL.** NAD+-dependent modulation of chromatin structure and transcription by nucleosome binding properties of PARP-1. *Cell.*, 2004, **119(6)**, 803-14
- Kim MY, Zhang T, Kraus WL.** Poly(ADP-ribosylation) by PARP-1: 'PAR-laying' NAD+ into a nuclear signal. *Genes Dev.*, 2005, **19(17)**, 1951-67
- Koh DW., Ame C., Coyle D., Ramsinghani S., Slama JT., Jacobson MK.** Expression, purification and characterization of recombinant poly (ADP-ribose) glycohydrolase. 1997, *Dev Mol Cell Biochem.*, Kluwer Academic Publisher
- Koh DW, Lawler AM, Poitras MF, Sasaki M, Wattler S, Nehls MC, Stoger T, Poirier GG, Dawson VL, Dawson TM.** Failure to degrade poly(ADP-ribose) causes increased sensitivity to cytotoxicity and early embryonic lethality. *Proc Natl Acad Sci U S A.*, 2004, **101(51)**, 17699-704
- Krammer PH.** CD95's deadly mission in the immune system. *Nature.*, 2000, **407(6805)**, 789-95
- Kraus WL, Lis JT.** PARP goes transcription. *Cell.*, 2003, **113(6)**, 677-83
- Kroemer G, Dallaporta B, Resche-Rigon M.** The mitochondrial death/life regulator in apoptosis and necrosis. *Annu Rev Physiol.*, 1998, **60**, 619-42
- Kroemer G, Martin SJ.** Caspase-independent cell death. *Nat Med.*, 2005, **11(7)**, 725-30
- Kubota Y, Nash RA, Klungland A, Schar P, Barnes DE, Lindahl T.** Reconstitution of DNA base excision-repair with purified human proteins: interaction between DNA polymerase beta and the XRCC1 protein. *EMBO J.*, 1996, **15(23)**, 6662-70
- Kun E, Kirsten E, Mendeleyev J, Ordahl CP.** Regulation of the enzymatic catalysis of poly(ADP-ribose) polymerase by dsDNA, polyamines, Mg²⁺, Ca²⁺, histones H1 and H3, and ATP. *Biochemistry.*, 2004, **43(1)**, 210-6
- Labieniec M, Gabryelak T, Falcioni G.** Antioxidant and pro-oxidant effects of tannins in digestive cells of the freshwater mussel *Unio tumidus*. *Mutat Res.*, 2003, **539(1-2)**, 19-28
- Labieniec M, Gabryelak T.** Effects of tannins on Chinese hamster cell line B14. *Mutat Res.*, 2003, **539(1-2)**, 127-35

- Ladiges WC.** Mouse models of XRCC1 DNA repair polymorphisms and cancer. *Oncogene.*, 2006, **25(11)**, 1612-9
- Laemmli UK.** Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.*, 1970, **227(5259)**, 680-5
- Lan L, Nakajima S, Oohata Y, Takao M, Okano S, Masutani M, Wilson SH, Yasui A.** In situ analysis of repair processes for oxidative DNA damage in mammalian cells. *Proc Natl Acad Sci U S A.*, 2004, **101(38)**, 13738-43
- Lazebnik YA, Kaufmann SH, Desnoyers S, Poirier GG, Earnshaw WC.** Cleavage of poly(ADP-ribose) polymerase by a proteinase with properties like ICE. *Nature.*, 1994, **371(6495)**, 346-7
- Leist M, Single B, Castoldi AF, Kuhnle S, Nicotera P.** Intracellular adenosine triphosphate (ATP) concentration: a switch in the decision between apoptosis and necrosis. *J Exp Med.*, 1997, **185(8)**, 1481-6
- Leist M, Single B, Kunstle G, Volbracht C, Hentze H, Nicotera P.** Apoptosis in the absence of poly-(ADP-ribose) polymerase. *Biochem Biophys Res Commun.*, 1997, **233(2)**, 518-22
- Leppard JB, Dong Z, Mackey ZB, Tomkinson AE.** Physical and functional interaction between DNA ligase IIIalpha and poly(ADP-Ribose) polymerase 1 in DNA single-strand break repair. *Mol Cell Biol.*, 2003, **23(16)**, 5919-27
- Levy N, Martz A, Bresson A, Spenlehauer C, de Murcia G, Menissier-de Murcia J.** XRCC1 is phosphorylated by DNA-dependent protein kinase in response to DNA damage. *Nucleic Acids Res.*, 2006, **34(1)**, 32-41
- Li Y, Kim J, Li J, Liu F, Liu X, Himmeldirk K, Ren Y, Wagner TE, Chen X.** Natural anti-diabetic compound 1,2,3,4,6-penta-O-galloyl-D-glucopyranose binds to insulin receptor and activates insulin-mediated glucose transport signaling pathway. *Biochem Biophys Res Commun.*, 2005, **336(2)**, 430-7
- Lin W, Ame JC, Aboul-Ela N, Jacobson EL, Jacobson MK.** Isolation and characterization of the cDNA encoding bovine poly(ADP-ribose) glycohydrolase. *J Biol Chem.*, 1997, **272(18)**, 11895-901
- Loizou JI, El-Khamisy SF, Zlatanou A, Moore DJ, Chan DW, Qin J, Sarno S, Meggio F, Pinna LA, Caldecott KW.** The protein kinase CK2 facilitates repair of chromosomal DNA single-strand breaks. *Cell.*, 2004, **117(1)**, 17-28
- Lu XC, Massuda E, Lin Q, Li W, Li JH, Zhang J.** Post-treatment with a novel PARG inhibitor reduces infarct in cerebral ischemia in the rat. *Brain Res.*, 2003, **978(1-2)**, 99-103
- Luo H, Chan DW, Yang T, Rodriguez M, Chen BP, Leng M, Mu JJ, Chen D, Songyang Z, Wang Y, Qin J.** A new XRCC1-containing complex and its role in cellular survival of methyl ethanesulfonate treatment. *Mol Cell Biol.*, 2004, **24(19)**, 8356-65
- Lyons RJ, Deane R, Lynch DK, Ye ZS, Sanderson GM, Eyre HJ, Sutherland GR, Daly RJ.** Identification of a novel human tankyrase through its interaction with the adaptor protein Grb14. *J Biol Chem.*, 2001, **276(20)**, 17172-80

- Mackey ZB, Niedergang C, Murcia JM, Leppard J, Au K, Chen J, de Murcia G, Tomkinson AE.** DNA ligase III is recruited to DNA strand breaks by a zinc finger motif homologous to that of poly(ADP-ribose) polymerase. Identification of two functionally distinct DNA binding regions within DNA ligase III. *J Biol Chem.*, 1999, **274**(31), 21679-87
- Makrides SC.** Strategies for achieving high-level expression of genes in *Escherichia coli*. *Microbiol Rev.*, 1996, **60**(3), 512-38
- Malanga M, Althaus FR.** Poly(ADP-ribose) molecules formed during DNA repair in vivo. *J Biol Chem.*, 1994, **269**(26), 17691-6
- Malanga M, Althaus FR.** The role of poly(ADP-ribose) in the DNA damage signaling network. *Biochem Cell Biol.*, 2005, **83**(3), 354-64
- Marintchev A, Mullen MA, Maciejewski MW, Pan B, Gryk MR, Mullen GP.** Solution structure of the single-strand break repair protein XRCC1 N-terminal domain. *Nat Struct Biol.*, 1999, **6**(9), 884-93
- Marsin S, Vidal AE, Sossou M, Menissier-de Murcia J, Le Page F, Boiteux S, de Murcia G, Radicella JP.** Role of XRCC1 in the coordination and stimulation of oxidative DNA damage repair initiated by the DNA glycosylase hOGG1. *J Biol Chem.*, 2003, **278**(45), 44068-74
- Maruta H, Inageda K, Aoki T, Nishina H, Tanuma S.** Characterization of two forms of poly(ADP-ribose) glycohydrolase in guinea pig liver. *Biochemistry.*, 1991, **30**(24), 5907-12
- Maruta H, Matsumura N, Tanuma S.** Role of (ADP-ribose)n catabolism in DNA repair. *Biochem Biophys Res Commun.*, 1997, **236**(2), 265-9
- Masson M, Niedergang C, Schreiber V, Muller S, Menissier-de Murcia J, de Murcia G.** XRCC1 is specifically associated with poly(ADP-ribose) polymerase and negatively regulates its activity following DNA damage. *Mol Cell Biol.*, 1998, **18**(6), 3563-71
- Masutani M, Nozaki T, Nishiyama E, Shimokawa T, Tachi Y, Suzuki H, Nakagama H, Wakabayashi K, Sugimura T.** Function of poly(ADP-ribose) polymerase in response to DNA damage: gene-disruption study in mice. *Mol Cell Biochem.*, 1999, **193**(1-2), 149-52
- Masutani, M., Shimikawa, T., Igarashi, M., Hamada, M., Shibata, A., Oami, S., Nozaki, T., Nakagama, H., Sugimura, T., Takeuchi, T., Hori, M.** Inhibition of poly (ADP-ribose) glycohydrolase activity by cyclic peptide antibiotics containing piperazic acid residues. *Proc. Japan Acad.*, 2002, **78**, 15-17
- McKelvey-Martin VJ, Green MH, Schmezer P, Pool-Zobel BL, De Meo MP, Collins A.** The single cell gel electrophoresis assay (comet assay): a European review. *Mutat Res.*, 1993, **288**(1), 47-63
- McLennan AG.** The Nudix hydrolase superfamily. *Cell Mol Life Sci.*, 2006, **63**(2), 123-43
- Menard L, Thibault L, Poirier GG.** Reconstitution of an in vitro poly(ADP-ribose) turnover system. *Biochim Biophys Acta.*, 1990, **1049**(1), 45-58
- Mendoza-Alvarez H, Alvarez-Gonzalez R.** Poly(ADP-ribose) polymerase is a catalytic dimer and the automodification reaction is intermolecular. *J Biol Chem.*, 1993, **268**(30), 22575-80
- Mendoza-Alvarez H, Chavez-Bueno S, Alvarez-Gonzalez R.** Chain length analysis of ADP-ribose polymers generated by poly(ADP-ribose) polymerase (PARP) as a function of beta-NAD⁺ and enzyme concentrations. *IUBMB Life.*, 2000, **50**(2), 145-9

- Mendoza-Alvarez H, Alvarez-Gonzalez R.** Regulation of p53 sequence-specific DNA-binding by covalent poly(ADP-ribosyl)ation. *J Biol Chem.*, 2001, **276**(39), 36425-30
- Menissier de Murcia M, Niedergang C, Trucco C, Ricoul M, Dutrillaux B, Mark M, Oliver FJ, Masson M, Dierich A, LeMeur M, Walztinger C, Chambon P, de Murcia G.** Requirement of poly(ADP-ribose) polymerase in recovery from DNA damage in mice and in cells. *Proc Natl Acad Sci USA.*, 1997, **94**(14), 7303-7
- Menissier de Murcia M, Ricoul M, Tartier L, Niedergang C, Huber A, Dantzer F, Schreiber V, Ame JC, Dierich A, LeMeur M, Sabatier L, Chambon P, de Murcia G.** Functional interaction between PARP-1 and PARP-2 in chromosome stability and embryonic development in mouse. *EMBO J.*, 2003, **22**(9): 2255-63
- Meyer RG, Meyer-Ficca ML, Jacobson EL, Jacobson MK.** Human poly(ADP-ribose) glycohydrolase (PARG) gene and the common promoter sequence it shares with inner mitochondrial membrane translocase 23 (TIM23). *Gene.*, 2003, **314**, 181-90
- Meyer-Ficca ML, Meyer RG, Coyle DL, Jacobson EL, Jacobson MK.** Human poly(ADP-ribose) glycohydrolase is expressed in alternative splice variants yielding isoforms that localize to different cell compartments. *Exp Cell Res.*, 2004, **297**(2), 521-32
- Minaga T, Kun E.** Probable helical conformation of poly(ADP-ribose). The effect of cations on spectral properties. *J Biol Chem.*, 1983, **258**(9), 5726-30
- Miramar MD, Costantini P, Ravagnan L, Saraiva LM, Haouzi D, Brothers G, Penninger JM, Peleato ML, Kroemer G, Susin SA.** NADH oxidase activity of mitochondrial apoptosis-inducing factor. *J Biol Chem.*, 2001, **276**(19), 16391-8
- Mitic SS, Miletic GZ, Kostic DA.** Kinetic determination of traces of iodide by its catalytic effect on oxidation of sodium pyrogallol-5-sulfonate by hydrogen peroxide. *Anal Sci.*, 2003, **19**(6), 913-6
- Miwa M, Sugimura T.** Splitting of the ribose-ribose linkage of poly(adenosine diphosphate-ribose) by a calf thymus extract. *J Biol Chem.*, 1971, **246**(20), 6362-4
- Miwa M, Tanaka M, Matsushima T, Sugimura T.** Purification and properties of glycohydrolase from calf thymus splitting ribose-ribose linkages of poly(adenosine diphosphate ribose). *J Biol Chem.*, 1974, **249**(11), 3475-82
- Miwa M, Saito H, Sakura H, Saikawa N, Watanabe F, Matsushima T, Sugimura T.** A ¹³C NMR study of poly(adenosine diphosphate ribose) and its monomers: evidence of alpha-(1' leads to 2') ribofuranosyl ribofuranoside residue. *Nucleic Acids Res.*, 1977, **4**(11), 3997-4005
- Miwa M, Saikawa N, Yamaizumi Z, Nishimura S, Sugimura T.** Structure of poly(adenosine diphosphate ribose): identification of 2'-[1"-ribosyl-2"-(or 3"-)(1""-ribosyl)]adenosine-5',5",5'''-tris(phosphate) as a branch linkage. *Proc Natl Acad Sci USA.*, 1979, **76**(2), 595-9
- Miwa M, Sugimura T.** Structure of poly(ADP-ribose). *Methods Enzymol.*, 1984, **106**, 441-50
- Morinaga N, Iwamaru Y, Yahiro K, Tagashira M, Moss J, Noda M.** Differential activities of plant polyphenols on the binding and internalization of cholera toxin in vero cells. *J Biol Chem.*, 2005, **280**(24): 23303-9

- Modjtahedi N, Giordanetto F, Madeo F, Kroemer G.** Apoptosis-inducing factor: vital and lethal. *Trends Cell Biol.*, 2006, **16(5)**, 264-72
- Muftuoglu M, Wong HK, Imam SZ, Wilson DM 3rd, Bohr VA, Opresko PL.** Telomere repeat binding factor 2 interacts with base excision repair proteins and stimulates DNA synthesis by DNA polymerase beta. *Cancer Res.*, 2006, **66(1)**, 113-24
- Murakami S, Muramatsu M, Otomo S.** Inhibitory effect of tannic acid on gastric H⁺,K(+)-ATPase. *J Nat Prod.*, 1992, **55(4)**, 513-6
- Nakagawa T, Yokozawa T.** Direct scavenging of nitric oxide and superoxide by green tea. *Food Chem Toxicol.*, 2002, **40(12)** 1745-50
- Naegeli H, Althaus FR.** Regulation of poly(ADP-ribose) polymerase. Histone-specific adaptations of reaction products. *J Biol Chem.*, 1991, **266(16)**, 10596-601
- Nakamura J, Asakura S, Hester SD, de Murcia G, Caldecott KW, Swenberg JA.** Quantitation of intracellular NAD(P)H can monitor an imbalance of DNA single strand break repair in base excision repair deficient cells in real time. *Nucleic Acids Res.*, 2003, **31(17)**, e104
- Nash RA, Caldecott KW, Barnes DE, Lindahl T.** XRCC1 protein interacts with one of two distinct forms of DNA ligase III. *Biochemistry.*, 1997, **36(17)**, 5207-11
- Newmeyer DD, Ferguson-Miller S.** Mitochondria: releasing power for life and unleashing the machineries of death. *Cell.*, 2003, **112(4)**, 481-90
- Nicotera P, Leist M, Ferrando-May E.** Intracellular ATP, a switch in the decision between apoptosis and necrosis. *Toxicol Lett.*, 1998, **102-103**, 139-42
- Nicotera P, Melino G.** Regulation of the apoptosis-necrosis switch. *Oncogene.*, 2004, **23(16)**, 2757-65
- Norton WT, Pduslo SE.** Neuronal perikarya and astroglia of rat brain: chemical composition during myelation. *J Lipid Res.*, 1971, **12**, 84-90
- Nunez-Duran H.** Tannic acid as an electron microscope tracer for permeable cell membranes. *Stain Technol.*, 1980, **55(6)**, 361-5
- Okano S, Lan L, Caldecott KW, Mori T, Yasui A.** Spatial and temporal cellular responses to single-strand breaks in human cells. *Mol Cell Biol.*, 2003, **23(11)**, 3974-81
- Oei SL, Griesenbeck J, Buchlow G, Jorcke D, Mayer-Kuckuk P, Wons T, Ziegler M.** NAD⁺ analogs substituted in the purine base as substrates for poly(ADP-ribosyl) transferase. *FEBS Lett.*, 1996, **397(1)**, 17-21
- Oei SL, Griesenbeck J, Schweiger M, Babich V, Kropotov A, Tomilin N.** Interaction of the transcription factor YY1 with human poly(ADP-ribosyl) transferase. *Biochem Biophys Res Commun.*, 1997, **240(1)**, 108-11
- Oei SL, Griesenbeck J, Schweiger M, Ziegler M.** Regulation of RNA polymerase II-dependent transcription by poly(ADP-ribosylation) of transcription factors. *J Biol Chem.*, 1998, **273(48)**, 31644-7
- Oei SL, Ziegler M.** ATP for the DNA ligation step in base excision repair is generated from poly(ADP-ribose). *J Biol Chem.*, 2000, **275(30)**, 23234-9

- Oei SL, Keil C, Ziegler M.** Poly(ADP-ribosylation) and genomic stability. *Biochem Cell Biol.*, 2005, **83(3)**, 263-9
- Oliveira MA, Koh D, Patel CN, Jacobson MK.** Structure-based characterization of a novel anticancer target poly (ADP-ribose) glycohydrolase (PARG): evidence for the presence of an ADP-ribosyltransferase (ADPRT) fold. *Proc. Assoc. Cancer Res.*, 2001, **42**, 832
- Oka J, Ueda K, Hayaishi O, Komura H, Nakanishi K.** ADP-ribosyl protein lyase. Purification, properties, and identification of the product. *J Biol Chem.*, 1984, **259(2)**, 986-95
- Oka S, Kato J, Moss J.** Identification and characterization of a mammalian 39-kDa poly(ADP-ribose) glycohydrolase. *J Biol Chem.*, 2006, **281(2)**, 705-13
- Okano S, Lan L, Caldecott KW, Mori T, Yasui A.** Spatial and temporal cellular responses to single-strand breaks in human cells. *Mol Cell Biol.*, 2003, **23(11)**, 3974-81
- Okano S, Lan L, Tomkinson AE, Yasui A.** Translocation of XRCC1 and DNA ligase IIIalpha from centrosomes to chromosomes in response to DNA damage in mitotic human cells. *Nucleic Acids Res.*, 2005, **33(1)**, 422-9
- Otera H, Ohsakaya S, Nagaura Z, Ishihara N, Mihara K.** Export of mitochondrial AIF in response to proapoptotic stimuli depends on processing at the intermembrane space. *EMBO J.*, 2005, **24(7)**, 1375-86
- Panzeter PL, Althaus FR.** High resolution size analysis of ADP-ribose polymers using modified DNA sequencing gels. *Nucleic Acids Res.*, 1990, **18(8)**, 2194
- Panzeter PL, Zweifel B, Althaus FR.** Synthesis of poly(ADP-ribose)-agarose beads: an affinity resin for studying (ADP-ribose)n-protein interactions. *Anal Biochem.*, 1992, **207(1)**, 157-62
- Park SY, Cheng YC.** Poly(ADP-ribose) polymerase-1 could facilitate the religation of topoisomerase I-linked DNA inhibited by camptothecin. *Cancer Res.*, 2005, **65(9)**, 3894-902
- Park CJ, Choi BS.** The protein shuffle. Sequential interactions among components of the human nucleotide excision repair pathway. *FEBS J.*, 2006, **273(8)**, 1600-8
- Parson JL, Dianov GL.** Monitoring base excision repair proteins on damaged DNA using human cell extracts. *Biochem. Soc. Trans.*, 2004, **32**, 962-963
- Peddi SR, Chattopadhyay R, Naidu CV, Izumi T.** The human apurinic/apyrimidinic endonuclease-1 suppresses activation of poly(adp-ribose) polymerase-1 induced by DNA single strand breaks. *Toxicology.*, 2006, **224(1-2)**, 44-55
- Peter ME, Krammer PH.** The CD95(APO-1/Fas) DISC and beyond. *Cell Death Differ.*, 2003, **10(1)**, 26-35
- Petermann E, Ziegler M, Oei SL.** ATP-dependent selection between single nucleotide and long patch base excision repair. *DNA Repair (Amst)*, 2003, **2(10)**, 1101-14
- Petermann E, Keil C, Oei SL.** Importance of poly(ADP-ribose) polymerases in the regulation of DNA-dependent processes. *Cell Mol Life Sci.*, 2005, **62(7-8)**, 731-8
- Petrucco S.** Sensing DNA damage by PARP-like fingers. *Nucleic Acids Res.*, 2003, **31(23)**, 6689-99
- Pion E, Ullmann GM, Ame JC, Gerard D, de Murcia G, Bombarda E.** DNA-induced dimerization of poly(ADP-ribose) polymerase-1 triggers its activation. *Biochemistry.*, 2005, **44(44)**, 14670-81

- Pleschke JM, Kleczkowska HE, Strohm M, Althaus FR.** Poly(ADP-ribose) binds to specific domains in DNA damage checkpoint proteins. *J Biol Chem.*, 2000, **275**(52), 40974-80
- Plo I, Liao ZY, Barcelo JM, Kohlhagen G, Caldecott KW, Weinfeld M, Pommier Y.** Association of XRCC1 and tyrosyl DNA phosphodiesterase (Tdp1) for the repair of topoisomerase I-mediated DNA lesions. *DNA Repair (Amst.)*, 2003, **2**(10), 1087-100
- Prasad R, Lavrik OI, Kim SJ, Kedar P, Yang XP, Vande Berg BJ, Wilson SH.** DNA polymerase beta – mediated long patch base excision repair. Poly(ADP-ribose)polymerase-1 stimulates strand displacement DNA synthesis. *J Biol Chem.*, 2001, **276**(35), 32411-4
- Reed JC.** Bcl-2 family proteins. *Oncogene.*, 1998, **17**(25), 3225-36
- Riedl SJ, Shi Y.** Molecular mechanisms of caspase regulation during apoptosis. *Nat Rev Mol Cell Biol.*, 2004, **5**(11), 897-907
- Rongvaux A, Andris F, Van Gool F, Leo O.** Reconstructing eukaryotic NAD metabolism. *Bioessays.*, 2003, **25**(7), 683-90
- Rosenberg AH, Goldman E, Dunn JJ, Studier FW, Zubay G.** Effects of consecutive AGG codons on translation in Escherichia coli, demonstrated with a versatile codon test system. *J Bacteriol.*, 1993, **175**(3), 716-22
- Rossi L, Denegri M, Torti M, Poirier GG, Ivana Scovassi A.** Poly(ADP-ribose) degradation by post-nuclear extracts from human cells. *Biochimie.*, 2002, **84**(12), 1229-35
- Rouleau M, Aubin RA, Poirier GG.** Poly(ADP-ribosyl)ated chromatin domains: access granted. *J Cell Sci.*, 2004, **117**(6), 815-25
- Rudolph, R., Böhm, G., Lilie, H., Jaenicke, R.** Folding proteins. *Protein Function: A Practical Approach*, 1997, pp 57-99
- Rodriguez-Vico F, Martinez-Cayuela M, Garcia-Peregrin E, Ramirez H.** A procedure for eliminating interferences in the lowry method of protein determination. *Anal Biochem.*, 1989, **183**(2), 275-8
- Rouleau M, Aubin RA, Poirier GG.** Poly(ADP-ribosyl)ated chromatin domains: access granted. *J Cell Sci.*, 2004, **117**(6), 815-25
- Rouleau M, McDonald D, Gagne P, Ouellet ME, Droit A, Hunter JM, Dutertre S, Prigent C, Hendzel MJ, Poirier GG.** PARP-3 associates with polycomb group bodies and with components of the DNA damage repair machinery. *J Cell Biochem.*, 2006, in press
- Rossi L, Denegri M, Torti M, Poirier GG, Ivana Scovassi A.** Poly(ADP-ribose) degradation by post-nuclear extracts from human cells. *Biochimie.*, 2002, **84**(12), 1229-35
- Ruf A, Mennissier de Murcia J, de Murcia G, Schulz GE.** Structure of the catalytic fragment of poly(AD-ribose) polymerase from chicken. *Proc Natl Acad Sci U S A.*, 1996, **93**(15), 7481-5
- Ruscetti T, Lehnert BE, Halbrook J, Le Trong H, Hoekstra MF, Chen DJ, Peterson SR.** Stimulation of the DNA-dependent protein kinase by poly(ADP-ribose) polymerase. *J Biol Chem.*, 1998, **273**(23), 14461-7
- Saelens X, Festjens N, Vande Walle L, van Gurp M, van Loo G, Vandenebeele P.** Toxic proteins released from mitochondria in cell death. *Oncogene.*, 2004, **23**(16), 2861-74

- Sambrook J, Fritsch E. F., Maniatis T.** Molecular Cloning : A Laboratory Manual, 2nd ed., 1989, Cold Spring Harbor University Press, Cold Spring Harbour, NY
- Samejima K, Earnshaw WC.** Trashing the genome: the role of nucleases during apoptosis. *Nat Rev Mol Cell Biol.*, 2005, **6(9)**, 677-88
- Satoh K, Nagai F, Ushiyama K, Yasuda I, Seto T, Kano I.** Inhibition of Na⁺,K(+)-ATPase by 1,2,3,4,6-penta-O-galloyl-beta-D-glucose, a major constituent of both moutan cortex and Paeoniae radix. *Biochem Pharmacol.*, 1997, **53(4)**, 611-4
- Sawada H, Hamatake M, Hara A, Nakagawa M, Nakayama T.** Inhibition of human placenta aldose reductase by tannic acid. *Chem Pharm Bull (Tokyo).*, 1989, **37(6)**, 1662-4
- Saxena A, Saffery R, Wong LH, Kalitsis P, Choo KH.** Centromere proteins Cenpa, Cenpb, and Bub3 interact with poly(ADP-ribose) polymerase-1 protein and are poly(ADP-ribosyl)ated. *J Biol Chem.*, 02, **277(30)**, 26921-6
- Schafer ZT, Kornbluth S.** The apoptosome: physiological, developmental, and pathological modes of regulation. *Dev Cell.*, 2006, **10(5)**, 549-61
- Schein CH.** Optimizing protein folding to the native state in bacteria. *Curr Opin Biotechnol.*, 1991, **2(5)**, 746-50
- Schreiber E, Matthias P, Muller MM, Schaffner W.** Rapid detection of octamer binding proteins with 'mini-extracts', prepared from a small number of cells. *Nucleic Acids Res.*, 1989, **17(15)**, 6419
- Schreiber V, Molinete M, Boeuf H, de Murcia G, Menissier-de Murcia J.** The human poly(ADP-ribose) polymerase nuclear localization signal is a bipartite element functionally separate from DNA binding and catalytic activity. *EMBO J.*, 1992, **11(9)**, 3263-9
- Schreiber V, Ame JC, Dolle P, Schultz I, Rinaldi B, Fraulob V, Menissier-de Murcia J, de Murcia G.** Poly(ADP-ribose) polymerase-2 (PARP-2) is required for efficient base excision DNA repair in association with PARP-1 and XRCC1. *J Biol Chem.*, 2002, **277(25)**, 23028-36
- Schreiber V, Dantzer F, Ame JC, de Murcia G.** Poly(ADP-ribose): novel functions for an old molecule. *Nat Rev Mol Cell Biol.*, 2006, **7(7)**, 517-28
- Scovassi AI, Torriglia A.** Activation of DNA-degrading enzymes during apoptosis. *Eur J Histochem.*, 2003, **47(3)**, 185-94
- Shah GM, Poirier D, Duchaine C, Brochu G, Desnoyers S, Lagueux J, Verreault A, Hoflack JC, Kirkland JB, Poirier GG.** Methods for biochemical study of poly(ADP-ribose) metabolism in vitro and in vivo. *Anal Biochem.*, 1995, **227(1)**, 1-13
- Shall S, de Murcia G.** Poly(ADP-ribose) polymerase-1: what have we learned from the deficient mouse model? *Mutat Res.*, 2000, **460(1)**, 1-15
- Shall S.** Poly (ADP-ribosylation)--a common control process? *Bioessays.*, 2002, **24(2)**, 197-201
- Shen MR, Zdienicka MZ, Mohrenweiser H, Thompson LH, Thelen MP.** Mutations in hamster single-strand break repair gene XRCC1 causing defective DNA repair. *Nucleic Acids Res.*, 1998, **26(4)**, 1032-7

- Shieh WM, Ame JC, Wilson MV, Wang ZQ, Koh DW, Jacobson MK, Jacobson EL.** Poly(ADP-ribose) polymerase null mouse cells synthesize ADP-ribose polymers. *J Biol Chem.*, 1998, **273(46)**, 30069-72
- Shimokawa T, Masutani M, Nagasawa S, Nozaki T, Ikota N, Aoki Y, Nakagama H, Sugimura T.** Isolation and cloning of rat poly(ADP-ribose) glycohydrolase: presence of a potential nuclear export signal conserved in mammalian orthologs. *J Biochem (Tokyo).*, 1999, **126(4)**, 748-55
- Simbulan-Rosenthal CM, Rosenthal DS, Hilz H, Hickey R, Malkas L, Applegren N, Wu Y, Bers G, Smulson ME.** The expression of poly(ADP-ribose) polymerase during differentiation-linked DNA replication reveals that it is a component of the multiprotein DNA replication complex. *Biochemistry.*, 1996, **35(36)**, 11622-33
- Simbulan-Rosenthal CM, Rosenthal DS, Iyer S, Boulares AH, Smulson ME.** Transient poly(ADP-ribosylation) of nuclear proteins and role of poly(ADP-ribose) polymerase in the early stages of apoptosis. *J Biol Chem.* 1998 May 29;273(22):13703-12
- Simbulan-Rosenthal CM, Rosenthal DS, Iyer S, Boulares H, Smulson ME.** Involvement of PARP and poly(ADP-ribosylation) in the early stages of apoptosis and DNA replication. *Mol Cell Biochem.*, 1999, **193(1-2)**, 137-48
- Simbulan-Rosenthal CM, Ly DH, Rosenthal DS, Konopka G, Luo R, Wang ZQ, Schultz PG, Smulson ME.** Misregulation of gene expression in primary fibroblasts lacking poly(ADP-ribose) polymerase. *Proc Natl Acad Sci USA.*, 2000, **97(21)**, 11274-9
- Simbulan-Rosenthal CM, Rosenthal DS, Luo R, Samara R, Espinoza LA, Hassa PO, Hottiger MO, Smulson ME.** PARP-1 binds E2F-1 independently of its DNA binding and catalytic domains, and acts as a novel coactivator of E2F-1-mediated transcription during re-entry of quiescent cells into S phase. *Oncogene.*, 2003, **22(52)**, 8460-71
- Simon SA, Disalvo EA, Gawrisch K, Borovyagin V, Toone E, Schiffman SS, Needham D, McIntosh TJ.** Increased adhesion between neutral lipid bilayers: interbilayer bridges formed by tannic acid. *Biophys J.*, 1994, **66(6)**, 1943-58
- Simonin F, Menissier-de Murcia J, Poch O, Muller S, Gradwohl G, Molinete M, Penning C, Keith G, de Murcia G.** Expression and site-directed mutagenesis of the catalytic domain of human poly(ADP-ribose)polymerase in Escherichia coli. Lysine 893 is critical for activity. *J Biol Chem.*, 1990, **265(31)**, 19249-56
- Simonin F, Hofferer L, Panzeter PL, Muller S, de Murcia G, Althaus FR.** The carboxyl-terminal domain of human poly(ADP-ribose) polymerase. Overproduction in Escherichia coli, large scale purification, and characterization. *J Biol Chem.*, 1993, **268(18)**, 13454-61
- Skerra A.** Use of the tetracycline promoter for the tightly regulated production of a murine antibody fragment in Escherichia coli. *Gene.*, 1994, **151(1-2)**, 131-5
- Skulachev VP.** Bioenergetic aspects of apoptosis, necrosis and mitoptosis. *Apoptosis.*, 2006, **11(4)**, 473-85

- Slama JT, Aboul-Ela N, Goli DM, Cheesman BV, Simmons AM, Jacobson MK.** Specific inhibition of poly(ADP-ribose) glycohydrolase by adenosine diphosphate (hydroxymethyl)pyrrolidinediol. *J Med Chem.*, 1995, **38**(2), 389-93
- Slama JT, Aboul-Ela N, Jacobson MK.** Mechanism of inhibition of poly(ADP-ribose) glycohydrolase by adenosine diphosphate (hydroxymethyl)pyrrolidinediol. *J Med Chem.*, 1995, **38**(21), 4332-6
- Smith S.** The world according to PARP. *Trends Biochem Sci.*, 2001, **26**(3), 174-9
- Smith AH, Zoetendal E, Mackie RI.** Bacterial mechanisms to overcome inhibitory effects of dietary tannins. *Microb Ecol.*, 2005, **50**(2), 197-205
- Soldani C, Scovassi AI.** Poly(ADP-ribose) polymerase-1 cleavage during apoptosis: an update. *Apoptosis.*, 2002, **7**(4), 321-8
- Song D, Sakamoto S, Taniguchi T.** Inhibition of poly(ADP-ribose) polymerase activity by Bcl-2 in association with the ribosomal protein S3a. *Biochemistry.*, 2002, **41**(3), 929-34
- Strandberg L, Enfors SO.** Factors influencing inclusion body formation in the production of a fused protein in Escherichia coli. *Appl Environ Microbiol.*, 1991, **57**(6), 1669-74
- Sugimura T, Fujimura S, Hasegawa S, Kawamura Y.** Polymerization of the adenosine 5'-diphosphate ribose moiety of NAD by rat liver nuclear enzyme. *Biochim Biophys Acta.*, 1967, **138**(2), 438-41
- Sundberg L, Porath J.** Preparation of adsorbents for biospecific affinity chromatography. Attachment of group-containing ligands to insoluble polymers by means of bifunctional oxiranes. *J Chromatogr.*, 1974, **90**(1), 87-98
- Susin SA, Lorenzo HK, Zamzami N, Marzo I, Snow BE, Brothers GM, Mangion J, Jacotot E, Costantini P, Loeffler M, Larochette N, Goodlett DR, Aebersold R, Siderovski DP, Penninger JM, Kroemer G.** Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature.*, 1999, **397**(6718), 441-6
- Susin SA, Daugas E, Ravagnan L, Samejima K, Zamzami N, Loeffler M, Costantini P, Ferri KF, Irinopoulou T, Prevost MC, Brothers G, Mak TW, Penninger J, Earnshaw WC, Kroemer G.** Two distinct pathways leading to nuclear apoptosis. *J Exp Med.*, 2000, **192**(4), 571-80
- Takanami T, Nakamura J, Kubota Y, Horiuchi S.** The Arg280His polymorphism in X-ray repair cross-complementing gene 1 impairs DNA repair ability. *Mutat Res.*, 2005, **582**(1-2), 135-45
- Tan GT, Lee S, Lee IS, Chen J, Leitner P, Besterman JM, Kinghorn AD, Pezzuto JM.** Natural-product inhibitors of human DNA ligase I. *Biochem J.*, 1996, **314**(3), 993-1000
- Tanaka S, Takehashi M, Iida S, Kitajima T, Kamanaka Y, Stedeford T, Banasik M, Ueda K.** Mitochondrial impairment induced by poly(ADP-ribose) polymerase-1 activation in cortical neurons after oxygen and glucose deprivation. *J Neurochem.*, 2005, **95**(1), 179-90
- Tanuma S, Kawashima K, Endo H.** Occurrence of (ADP-ribose)_n glycohydrolase in human erythrocytes. *Biochem Biophys Res Commun.*, 1986, **136**(3), 1110-5
- Tanuma S, Kawashima K, Endo H.** Purification and properties of an (ADP-ribose)_n glycohydrolase from guinea pig liver nuclei. *J Biol Chem.*, 1986, **261**(2), 965-9

- Tanuma S, Sakagami H, Endo H.** Inhibitory effect of tannin on poly(ADP-ribose) glycohydrolase from human placenta. *Biochem Int.*, 1989, **18(4)**, 701-8
- Tanuma S, Tsai YJ, Sakagami H, Konno K, Endo H.** Lignin inhibits (ADP-ribose)n glycohydrolase activity. *Biochem Int.*, 1989, **19(6)**: 1395-402
- Tanuma Y., Okano T., Nakajima T.** Lignin glycoside and its use. Patentanmeldung, 1990, **Application number 03205402 A**
- Tanuma S, Tsai YJ, Sakagami H, Konno K, Endo H.** Lignin inhibits (ADP-ribose)n glycohydrolase activity. *Biochem Int.*, 1989, **19(6)**, 1395-402
- Tartier L, Spenlehauer C, Newman HC, Folkard M, Prise KM, Michael BD, Menissier-de Murcia J, de Murcia G.** Local DNA damage by proton microbeam irradiation induces poly(ADP-ribose) synthesis in mammalian cells. *Mutagenesis.*, 2003, **18(5)**, 411-6
- Tavassoli M, Tavassoli MH, Shall S.** Isolation and purification of poly(ADP-ribose) glycohydrolase from pig thymus. *Eur J Biochem.*, 1983, **135(3)**, 449-55
- Tavassoli M, Tavassoli MH, Shall S.** Effect of DNA intercalators on poly(ADP-ribose) glycohydrolase activity. *Biochim Biophys Acta.*, 1985, **827(3)**, 28-34
- Taylor RM, Thistlethwaite A, Caldecott KW.** Central role for the XRCC1 BRCT I domain in mammalian DNA single-strand break repair. *Mol Cell Biol.*, 2002, **22(8)**, 2556-63
- Tebbs RS, Flannery ML, Meneses JJ, Hartmann A, Tucker JD, Thompson LH, Cleaver JE, Pedersen RA.** Requirement for the Xrcc1 DNA base excision repair gene during early mouse development. *Dev Biol.*, 1999, **208(2)**, 513-29
- Tebbs RS, Thompson LH, Cleaver JE.** Rescue of Xrcc1 knockout mouse embryo lethality by transgene-complementation. *DNA Repair (Amst)*, 2003, **2(12)**, 1405-17
- Tebib K, Rouanet JM, Besancon P.** Effect of grape seed tannins on the activity of some rat intestinal enzyme activities. *Enzyme Protein.*, 1994-1995, **48(1)**, 51-60
- Tentori L, Leonetti C, Scarsella M, Muzi A, Vergati M, Forini O, Lacal PM, Ruffini F, Gold B, Li W, Zhang J, Graziani G.** Poly(ADP-ribose) glycohydrolase inhibitor as chemosensitiser of malignant melanoma for temozolomide. *Eur J Cancer.*, 2005, **41(18)**, 2948-57
- Tewari M, Quan LT, O'Rourke K, Desnoyers S, Zeng Z, Beidler DR, Poirier GG, Salvesen GS, Dixit VM.** Yama/CPP32 beta, a mammalian homolog of CED-3, is a CrmA-inhibitable protease that cleaves the death substrate poly(ADP-ribose) polymerase. *Cell.*, 1995, **81(5)**, 801-9
- Thomassin H, Jacobson MK, Guay J, Verreault A, Aboul-ela N, Menard L, Poirier GG.** An affinity matrix for the purification of poly(ADP-ribose) glycohydrolase. *Nucleic Acids Res.*, 1990, **18(16)**, 4691-4
- Thompson LH, Brookman KW, Jones NJ, Allen SA, Carrano AV.** Molecular cloning of the human XRCC1 gene, which corrects defective DNA strand break repair and sister chromatid exchange. *Mol Biol.*, 1990, **10(12)**, 6160-71

- Tong WM, Hande MP, Lansdorp PM, Wang ZQ.** DNA strand break-sensing molecule poly(ADP-Ribose) polymerase cooperates with p53 in telomere function, chromosome stability, and tumor suppression. *Mol Cell Biol.*, 2001, **21(12)**, 4046-54
- Trucco C, Oliver FJ, de Murcia G, Menissier-de Murcia J.** DNA repair defect in poly(ADP-ribose) polymerase-deficient cell lines. *Nucleic Acids Res.*, 1998, **26(11)**, 2644-9
- Tsai YJ, Abe H, Maruta H, Hatano T, Nishina H, Sakagami H, Okuda T, Tanuma S.** Effects of chemically defined tannins on poly (ADP-ribose) glycohydrolase activity. *Biochem Int.*, 1991, **24(5)**, 889-97
- Tsai YJ, Aoki T, Maruta H, Abe H, Sakagami H, Hatano T, Okuda T, Tanuma S.** Mouse mammary tumor virus gene expression is suppressed by oligomeric ellagitannins, novel inhibitors of poly(ADP-ribose) glycohydrolase. *J Biol Chem.*, 1992, **267(20)**, 14436-42
- Tsuchiya H.** Effects of green tea catechins on membrane fluidity. *Pharmacology.*, 1999, **59(1)**, 34-44
- Tulin A, Spradling A.** Chromatin loosening by poly(ADP)-ribose polymerase (PARP) at Drosophila puff loci. *Science.*, 2003, **299(5606)**, 560-2
- Tulin A, Naumova NM, Menon AK, Spradling AC.** Drosophila poly(ADP-ribose) glycohydrolase mediates chromatin structure and SIR2-dependent silencing. *Genetics.*, 2006, **172(1)**, 363-71
- Uchida K, Suzuki H, Maruta H, Abe H, Aoki K, Miwa M, Tanuma S.** Preferential degradation of protein-bound (ADP-ribose)n by nuclear poly(ADP-ribose) glycohydrolase from human placenta. *J Biol Chem.*, 1993, **268(5)** 3194-200
- Uchiumi F, Maruta H, Inoue J, Yamamoto T, Tanuma S.** Inhibitory effect of tannic acid on human immunodeficiency virus promoter activity induced by 12-O-tetra decanoylephorbol-13-acetate in Jurkat T-cells. *Biochem Biophys Res Commun.*, 1996, **220(2)**, 411-7
- Uchiumi F, Ikeda D, Tanuma S.** Changes in the activities and gene expressions of poly(ADP-ribose) glycohydrolases during the differentiation of human promyelocytic leukemia cell line HL-60. *Biochim Biophys Acta.*, 2004, **1676(1)**, 1-11
- Ueda K, Hayaishi O.** ADP-ribosylation. *Annu Rev Biochem.*, 1985, **54**, 73-100
- Vahsen N, Cande C, Briere JJ, Benit P, Joza N, Larochette N, Mastroberardino PG, Pequignot MO, Casares N, Lazar V, Feraud O, Debili N, Wissing S, Engelhardt S, Madeo F, Piacentini M, Penninger JM, Schagger H, Rustin P, Kroemer G.** AIF deficiency compromises oxidative phosphorylation. *EMBO J.*, 2004, **23(23)**, 4679-89
- Vahsen N, Cande C, Dupaigne P, Giordanetto F, Kroemer RT, Herker E, Scholz S, Modjtahedi N, Madeo F, Le Cam E, Kroemer G.** Physical interaction of apoptosis-inducing factor with DNA and RNA. *Oncogene.*, 2006, **25(12)**, 1763-74
- Verstraeten SV, Oteiza PI, Fraga CG.** Membrane effects of cocoa procyanidins in liposomes and Jurkat T cells. *Biol Res.*, 2004, **37(2)**, 293-300
- Virag L, Salzman AL, Szabo C.** Poly(ADP-ribose) synthetase activation mediates mitochondrial injury during oxidant-induced cell death. *J Immunol.*, 1998, **161(7)**, 3753-9

- Virag L, Szabo C.** The therapeutic potential of poly(ADP-ribose) polymerase inhibitors. *Pharmacol Rev.*, 2002, **54**(3), 375-429
- Virág L.** PARP-1 and the shape of cell death. *Poly (ADP-Ribosyl)ation*, 2005, Landes Biosciences/Eurekah, Georgetown
- Vidal AE, Boiteux S, Hickson ID, Radicella JP.** XRCC1 coordinates the initial and late stages of DNA abasic site repair through protein-protein interactions. *EMBO J.*, 2001, **20**(22), 6530-9
- Vodenicharov MD, Sallmann FR, Satoh MS, Poirier GG.** Base excision repair is efficient in cells lacking poly(ADP-ribose) polymerase 1. *Nucleic Acids Res.*, 2000, **28**(20), 3887-96
- Voss S, Skerra A.** Mutagenesis of a flexible loop in streptavidin leads to higher affinity for the Strep-tag II peptide and improved performance in recombinant protein purification. *Protein Eng.*, 1997, **10**(8), 975-82
- Walker JW, Jijon HB, Madsen KL.** AMP-activated protein kinase is a positive regulator of poly(ADP-ribose) polymerase. *Biochem Biophys Res Commun.*, 2006, **342**(1), 336-41
- Wang ZQ, Auer B, Stingl L, Berghammer H, Haidacher D, Schweiger M, Wagner EF.** Mice lacking ADPRT and poly(ADP-ribosylation) develop normally but are susceptible to skin disease. *Genes Dev.*, 1995, **9**(5), 509-20
- Wang BH, Foo LY, Polya GM.** Differential inhibition of eukaryote protein kinases by condensed tannins. *Phytochemistry.*, 1996, **43**(2), 359-65
- Wang ZQ, Stingl L, Morrison C, Jantsch M, Los M, Schulze-Osthoff K, Wagner EF.** PARP is important for genomic stability but dispensable in apoptosis. *Genes Dev.*, 1997, **11**(18), 2347-58
- Weickert MJ, Doherty DH, Best EA, Olins PO.** Optimization of heterologous protein production in *Escherichia coli*. *Curr Opin Biotechnol.*, 1996, **7**(5), 494-9
- Wesierska-Gadek J, Wojciechowski J, Schmid G.** Central and carboxy-terminal regions of human p53 protein are essential for interaction and complex formation with PARP-1. *J Cell Biochem.*, 2003, **89**(2), 220-32
- West JD, Ji C, Marnett LJ.** Modulation of DNA fragmentation factor 40 nuclease activity by poly(ADP-ribose) polymerase-1. *J Biol Chem.*, 2005, **280**(15), 15141-7
- Whitehouse CJ, Taylor RM, Thistlethwaite A, Zhang H, Karimi-Busheri F, Lasko DD, Weinfeld M, Caldecott KW.** XRCC1 stimulates human polynucleotide kinase activity at damaged DNA termini and accelerates DNA single-strand break repair. *Cell.*, 2001, **104**(1), 107-17
- Widlak P, Garrard WT.** Discovery, regulation, and action of the major apoptotic nucleases DFF40/CAD and endonuclease G. *J Cell Biochem.*, 2005, **94**(6), 1078-87
- Wieler S, Gagne JP, Vaziri H, Poirier GG, Benchimol S.** Poly(ADP-ribose) polymerase-1 is a positive regulator of the p53-mediated G1 arrest response following ionizing radiation. *J Biol Chem.*, 2003, **278**(21), 18914-21
- Williams JC, Chambers JP, Liehr JG.** Glutamyl ribose 5-phosphate storage disease. A hereditary defect in the degradation of poly(ADP-ribosylated) proteins. *J Biol Chem.*, 1984, **259**(2), 1037-42

- Winstall E, Affar EB, Shah R, Bourassa S, Scovassi AI, Poirier GG.** Poly(ADP-ribose) glycohydrolase is present and active in mammalian cells as a 110-kDa protein. *Exp Cell Res.*, 1999, **246**(2), 395-8
- Winstall E, Affar EB, Shah R, Bourassa S, Scovassi IA, Poirier GG.** Preferential perinuclear localization of poly(ADP-ribose) glycohydrolase. *Exp Cell Res.*, 1999, **251**(2), 372-8
- Wong HK, Kim D, Hogue BA, McNeill DR, Wilson DM 3rd.** DNA damage levels and biochemical repair capacities associated with XRCC1 deficiency. *Biochemistry.*, 2005, **44**(43), 14335-43
- Wu LT, Chu CC, Chung JG, Chen CH, Hsu LS, Liu JK, Chen SC.** Effects of tannic acid and its related compounds on food mutagens or hydrogen peroxide-induced DNA strands breaks in human lymphocytes. *Mutat Res.*, 2004, **556**(1-2), 75-82
- Ye H, Cande C, Stephanou NC, Jiang S, Gurbuxani S, Larochette N, Daugas E, Garrido C, Kroemer G, Wu H.** DNA binding is required for the apoptogenic action of apoptosis inducing factor. *Nat Struct Biol.*, 2002, **9**(9), 680-4
- Yeh TY, Meyer TN, Schwesinger C, Tsun ZY, Lee RM, Chi NW.** Tankyrase recruitment to the lateral membrane in polarized epithelial cells. Regulation by cell-cell contact and protein PARsylation. *Biochem J.* 2006, in press.
- Ying W, Swanson RA.** The poly(ADP-ribose) glycohydrolase inhibitor gallotannin blocks oxidative astrocyte death. *Neuroreport.*, 2000, **11**(7), 1385-8
- Ying W, Sevigny MB, Chen Y, Swanson RA.** Poly(ADP-ribose) glycohydrolase mediates oxidative and excitotoxic neuronal death. *Proc Natl Acad Sci USA.*, 2001, **98**(21), 12227-32
- Yu SW, Wang H, Poitras MF, Coombs C, Bowers WJ, Federoff HJ, Poirier GG, Dawson TM, Dawson VL.** Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing actor. *Science.*, 2002, **297**(5579), 259-63
- Yu M, Schreek S, Cerni C, Schamberger C, Lesniewicz K, Poreba E, Vervoorts J, Walsemann G, Grotzinger J, Kremmer E, Mehraein Y, Mertsching J, Kraft R, Austen M, Luscher-Firzlaff J, Luscher B.** PARP-10, a novel Myc-interacting protein with poly(ADP-ribose) polymerase activity, inhibits transformation. *Oncogene.*, 2005, **24**(12), 1982-93
- Yung TM, Sato S, Satoh MS.** Poly(ADP-ribosyl)ation as a DNA damage-induced post-translational modification regulating poly(ADP-ribose) polymerase-1-topoisomerase I interaction. *J Biol Chem.*, 2004, **279**(38), 39686-96
- Zhang XD, Gillespie SK, Hersey P.** Staurosporine induces apoptosis of melanoma by both caspase-dependent and -independent apoptotic pathways. *Mol Cancer Ther.*, 2004, **3**(2), 187-97
- Zhang J, Dawson VL, Dawson TM, Snyder SH.** Nitric oxide activation of poly(ADP-ribose) synthetase in neurotoxicity. *Science.*, 1994, **263**(5147), 687-9
- Ziegler M, Oei SL.** A cellular survival switch: poly(ADP-ribosyl)ation stimulates DNA repair and silences transcription. *Bioessays.*, 2001, **23**(6), 543-8
- Zong WX, Ditsworth D, Bauer DE, Wang ZQ, Thompson CB.** Alkylating DNA damage stimulates a regulated form of necrotic cell death. *Genes Dev.*, 2004, **18**(11), 1272-82

9 Abkürzungsverzeichnis

3-ABA	3-Aminobenzamid
AD	Aktivatordomäne
ADP	Adenosindiphosphat
ADP-Ribose	Adenosindiphosphat-Ribose
AK	Antikörper
Agarose LM-MP	niedrigschmelzende Agarose
AMD	Automodifikationsdomäne
Amp	Ampizillin
AMP	Adenosinmonophosphat
AP	Alkalische Phosphatase
ATP	Adenosintriphosphat
BCIP	5-Brom-4-Chlor-3-Indolylphosphat-p-tolidinsalz
BER	<i>Base Excision Repair</i>
bp	Basenpaare
BrCN	Bromcyan
BRCT	<i>BRCA-1 carboxyterminus</i>
BSA	Rinderserumalbumin
Chl	Chloramphenicol
CHO	<i>chinese hamster ovary</i>
C-Terminus	Carboxyterminus
Da	Dalton
DABCO	1,4-Diazabicyclo-Octan
DAPI	Diaminophenylindol
DBD	DNA-bindende Domäne
DC	Dünnschichtchromatographie
DMEM	<i>Dulbeccos Modified Eagle Medium</i>
DMSO	Dimethylsulfoxid
DNA	Desoxyribonukleinsäure
DNase	Desoxyribonuklease
dNTP	Desoxyribonukleotid
DTT	Dithiothreitol
EDTA	Ethyldiamintetraacetat
EGTA	Ethylenglykoltetraacetat
EMS	Ethylmethansulfat
FCS	Fötales Kälberserum

<i>fwd</i>	vorwärts
Gdm-HCl	Guanidiniumhydrochlorid
GFP	<i>Green Fluorescent Protein</i>
GSH	Glutathion
GST	Glutathion-S-Transferase
HRP	Meerrettich-Peroxidase
IF	Immunfluoreszenz
IgG	Immunglobulin
IPTG	Isopropyl- β -D-Thiogalaktosid
Kan	Kanamycin
kbp	Kilobasenpaare
KE	Kernextrakt
λ	Wellenlänge
LB	Luria-Bertani-Medium
Lig III	DNA-Ligase III
MEM	<i>Modified Eagle Medium</i>
MMS	Methylmethanosulfat
MNNG	N-Methyl-N'-Nitro-N-Nitrosoguanin
mRNA	Boten-Ribonukleinsäure
MTT	Methylthiazoltetrazoliumsalz
NAAD ⁺	Nikotinsäure-Adenin-Dinukleotid
NAD ⁺	Nikotinamid-Adenin-Dinukleotid
NBT	Nitrotetrazoliumblauchlorid
NER	<i>Nukleotide Excision Repair</i>
NES	Kernexportsequenz
NLS	Kernlokalisationssequenz
bp NLS	bipartite Kernlokalisationssequenz
NP-40	Nonidet P-40
N-Terminus	Aminoterminus
OD ₆₀₀	optische Dichte bei $\lambda= 600$ nm
P	Phosphat
³² P	Phosphor-32
PAR	Poly (ADP-Ribose)
PARG	Poly (ADP-Ribose) Glycohydrolase
PARP	Poly (ADP-Ribose) Polymerase
PCR	Polymerasekettenreaktion
PFA	Paraformaldehyd
PMS	Phenylmethansulfonylfluorid

PNKP	Polynukleotidkinase/Phosphatase
Polβ	DNA-Polymerase β
PR-AMP	Phosphoribose-AMP
(PR) ₂ -AMP	Di-Phosphoribose-AMP
PRPs	<i>prolin rich proteins</i>
RNA	Ribonukleinsäure
RNase	Ribonuklease
<i>rev</i>	rückwärts
rpm	Umdrehung pro Minute
RPMI	<i>Roswell Park Memorial Institute developed Media</i>
SD-Medium	synthetisches Hefeminimalmedium
SDS	Natriumdodecylsulfat
SDS-PAG	SDS-Polyacrylamidgel
SDS-PAGE	SDS-Polyacrylamidgelektrophorese
SSBR	<i>Single Strand Break Repair System</i>
TCA	Trichloressigsäure
Temp	Temperatur
Tris	Tris-(hydroxymethyl)-Aminomethan
UV	ultraviolett
v/v	Volumenverhältnis
WB	Western-Blot
WT	Wildtyp
w/v	Massevolumenverhältnis
XRCC1	<i>X-ray repair cross-complementing 1</i>
YPD	Hefevollmedium
YY1	Transkriptionsfaktor Ying Yang 1
ZF	Zinkfinger