

8 Literaturverzeichnis

- Abramjuk C. (2001)
Magnetresonanztomographische und histologische Untersuchungen zum biologischen und physikalischen Verhalten eines Monomer- stabilisierten superparamagnetischen Kontrastmittels für die Magnetresonanztomographie am Modell der Ratte
Vet. Med. Diss. FU Berlin
- Aguas A.P., Grande N.R., Carvalho E. (1991)
Inflammatory macrophages in the dog contain high amounts of intravesicular ferritin and are associated with pouches of connective tissue fibers
Am J Anat 1991 Jan; 190(1):89-96
- Ahluwalia N., Gordon M., Handte G., Mahlon M., Li N.Q., Beard J., Weinstock D., Ross C. (2000)
Iron status and stores decline with age in Lewis rats
J.Nutr.2000, 130: 2378-2383
- Andrew E. R.(1982)
In: P. N. T. Wells
Scientific Basis of med. Imaging
Edinburg, Churchill Livingstone ,1982
- Anzai Y., Prince M. R. (1997)
Iron oxide-enhanced MR lymphography: the evaluation of cervical lymph node metastases in head and neck cancer
J. Magn. Reson. Imaging 7(1): 75-81
- Baker L.L., Parker B.R., Donaldson S.S., Castellino R.A. (1990)
Staging of Hogkin disease in children: Comparison of CT and lymphography with laparotomy
Am J Roentgenol.1990 Jun 154(6):1251-5
- Baudisch E. (1988)
Grundlagen der medizinischen Radiologie: ein Leitfaden für Studierende
Verl. Volk und Gesundheit, Berlin, 1988
- Belisle C., Sainte-Marie G. (1981)
Topography of the deep cortex of the lymph nodes of various mammalian species
Anat Rec 1981 Nov 201(3): 553-61
- Bengele H.H., Palmacci S., Rogers J., Jung C.W., Crenshaw J., Josephson L. (1994)
Biodistribution of an USPIO colloid BMS 180549 by different routes of administration
Magn Reson Imaging 1994, 12(3), 433-42
- Berg R. (1993)
Lehrbuch der Veterinär- Anatomie, Band 3
Die großen Versorgungs- und Steuerungssysteme
Gustav Fischer, Jena, Stuttgart
- Bilder G.E. (1975)
Studies on immune competence in the rat: changes with age, sex and strain
J Gerontol 1975 Nov, 30(6): 641-6
- Boudghène F.P., Bach-Gansmo T., Grange J.-D., Lame S., Nantois C., Wallays C., Bigot J.-M. (1993)
Contribution of oral magnetic particles in MRI of the Abdomen with spin-echo and gradient-echo sequences
JMRI 3 (1993) 107-112

- Brouwer A., Horan M.A., Barelds R.J., Knook D.L. (1986)
Cellular aging of the reticuloendothelial system
Arch Gerontol Geriatr 1986 Dec, 5(4):317-24
- Castellino R.A., Bergiron C., Markovits P. (1977)
Experiment with 658 consecutive lymphograms in children
Cancer 1977 Sep, 40(3):1097-101
- Castenholz A. (1998)
Functional microanatomy of initial lymphatics with special consideration of the extracellular matrix
Lymphology 1998 Sep, 31(3):101-18
- Cihal S., Maas R., Delling G., Klich S., Nielsen P., Lodemann K.P., Bucheler E. (1999)
USPIO particles: the signal behavior and relaxation times of the bone marrow after repeated iv. application in the pig model at 1.5 tesla
RoFo Fortschr Geb Rontgenstr Neue Bildgeb Verfahr 1999 Aug 171(2):149-54
- Chai J.W., Lin Y.C., Chen J.H., Wu C.C., Lee C.P. Chu W.C., Lee S.K. (2001)
In vivo MR study of fatty liver: importance of intracellular ultrastructural alteration for MR tissue parameters change
J Magn Reson Imaging 2001, 14:35-41
- Chouly C., Bordanave L., Bareille R., Guerin V., Baquey V., Pouliquen D., Baquey C., Jallet P. (1994)
In vitro study of the hemocompatibility of superparamagnetic contrast agent for MRI
Clin. Mater. 15/4 (293-301)
- De Leeuw A.M., Brouwer A., Barelds R.J., Knook D.L. (1983)
Maintenance cultures of Kupffer cells isolated from rats of various ages: ultrastructure, enzyme cytochemistry and endocytosis
Hepatology 1983 Jul-Aug, 3(4):497-506
- Elizondo G., Weissleder R. (1990)
Hepatic cirrhosis and hepatitis: MR imaging enhanced with superparamagnetic iron oxide
Radiology 174: 797-801
- Elste V., (1996a)
Magentresonanztomographische Lymphographie mit superparamagnetischen Eisenoxidpartikeln:
Untersuchungen zum Verteilungsverhalten nach intravenöser Injektion an der Ratte
Vet. Med. Diss. FU Berlin
- Elste V., Wagner S., Taupitz M., Pfefferer D., Kresse M., Hamm B., Berg R., Wolf K.J., Semmel W. (1996b)
MR Lymphography in rats: effects of muscular activity and hyperthermia on the lymphnode uptake of intravenous injected superparamagnetic iron oxide particles
Acad-Radiologica 3(8): 660-666
- Faich G., Strobos J. (1999)
Sodium ferric gluconate complex in sucrose: safer intravenous iron therapy than iron dextrans
Am J Kidney Dis 1999 Mar; 33(3): 464-70
- Feldman K.S., Sahasrabudhe K., Smith R.S., Scheuchenzuber W.J. (1999)
Immunostimulation by plant polyphenols: a relationship between tumor necrosis factor-alpha production and tannin structure
Bioorg Med Chem Lett 1999 Apr 5, 9(7):985-90

- Flesch M., Schwarz A., Bohm M. (1998)
Effects of red and white wine on endothelium- dependent vasorelaxation of rat aorta and human coronary arteries
Am J Physiol 1998 Oct, 275 (4 Pt 2): H1183-90
- Fuchs M. (1998)
Histologische Untersuchung zur Aufnahme und Verteilung von superparamagnetische Eisendextrancontrastmitteln in Lymphknoten nach subkutaner Applikation beim Meerschweinchen
Vet. med. Diss., FU Berlin
- Gushchin I.S., Prozorovsky N.S., Yamshchikova O.A., Kiselev A.V. (1991)
Mast cell regulatory effect on lymphoid cell proliferation
Agents Actions 1991 May, 33(1-2):185-8
- Hahn P.F., Stark D.D., Lewis J.M., Saini S., Elizondo G., Weissleder R., Fretz C.J., Ferrucci J.T. (1990)
First clinical trial of a new superparamagnetic Iron Oxide for use as an oral gastrointestinal contrast agent in MRI
Radiology 175 (1990) 695-700
- Hamm B., Vogl T.J., Branding G., Schnell B., Taupitz M, Wolf K.-J., Lissner J. (1992)
Focal liver lesions: MR Imaging with Mn- DPDP. Initial clinical results in 40 patients
Radiology 182 (1992): 167-174
- Harika L., Weissleder R., Poss K., Zimmer C., Papisov M.I., Brady T.J. (1995)
MR Lymphography with a lymphotropic T1- Type MR contrast agent: GD-DTPA-PGM
Magn-Reson-Med. 33: 88-92
- Harika L., Weissleder R., Poss K., Papisov M.I. (1996)
Macromolecular intravenous contrast agent for MR Lymphography: Characterization and efficacy studies
Radiology 198: 365-370
- Harisinghani M.G., Weissleder R., Saini S., Hahn P.F., Yantiss R.K., Tempany C., Wood B.J., Mueller P.R. (1999)
MR lymphangiography using USPIO in patients with primary abdominal and pelvic Malignancies: Radiographic- pathologic correlation
AJR 172/5: 1347-1351
- Harisinghani M.G., Weissleder R., Saini S., Hahn P.F., Rubin D., DeLange E., Harms S., Weinreb J., Small W., Sukerkar A., Brown J.J. Zelch J., Lucas M., Morris M. (2001)
Splenic imaging with USPIO ferumoxtran-10 (AMI-7227): preliminary observations
J Comp Assist Tomogr 2001 Sep-Oct 25(5):770-6
- Hässig A., Liang W.X., Schwabl H., Stampfli K. (1999):
Flavonoids and Tannins: plant-based antioxidants with vitamin character
Medical Hypotheses (1999) 52 (5), 479-481
- Heidemann P.H., Stubbe P., Schurnbrand P., Prindull G. (1982)
Iodine induced Hypothyreoidism and goiter following lipiodol TM lymphography
Eur J Pediatr 1982 Feb 138 (1), 82-3
- Hünigen H (2002)
Persönliche Mitteilung
- Ikeda T., Masamura S, Fujii H., Hiramatsu H., Mukai M., Matsui A. (2000)
Sentinel lymph node biopsy using tin colloid RI and blue dye method
Breast cancer 2000, 7(4):284-6

- Ikomi F., Hunt J., Hanna G., Schmid-Schönbein G.W. (1996)
Interstitial fluid, plasma protein, colloid and leucocyte uptake into initial lymphatics
J Appl Physiol 1996, 81(5):2060-2067
- Kaplan M., Hayek T., Raz A., Coleman R., Dornfeld L., Vaya J., Aviram M. (2001)
Pomegranate juice supplementation to atherosclerotic mice reduces macrophage lipid peroxidation, cellular cholesterol accumulation and development of atherosclerosis
J Nutr 2001 Aug, 131(8): 2082-9
- Kehagias D.T., Goulimos A.D., Smyniotis V., Vlahos L.J., (2001)
Diagnostic efficacy and safety of MRI of the liver with superparamagnetic iron oxide particles (SH U 555A)
J Magn Reson Imaging 2001 Nov, 14:595-601
- Klinke R., Silbernagl S. (2001)
Lehrbuch der Physiologie
Thieme, Stuttgart, 3. Auflage
- Köchli V. D., Marincek B. (1998)
Wie funktioniert MRI?
Springer, Berlin
- Komárek V. (2000)
Gross Anatomy of the laboratory rat
Aus: The laboratory Rat von G.J. Krinke
Academic Press, London
- Krebsregister (1998)
Krebsregister des Robert-Koch-Instituts
www.RKI.de/gbe/krebs
- Kreosfky T.J., Russel J.A., Rohrbach M.S. (1990)
Inhibition of alveolar macrophage spreading and phagocytosis by cotton bract Tannin. A potential mechanism in the pathogenesis of byssinosis.
Am J Pathol 1990 Aug, 137 (2): 263-74
- Kühnel W. (1992)
Taschenatlas der Zytologie, Histologie und mikroskopischen Anatomie
Thieme Verlag, Stuttgart, 11. Auflage
- Lauterbur P.C. (1973)
Image formation by induced local interactions: Examples employing nuclear magnetic resonance
Nature 242, 190-191
- Lawaczek R., Bauer H. Frenzel T., Hasegawa M., Ito Y., Kito K., Miwa N., Tsutsui H., Vogler H., Weinmann H.J. (1997)
Magnetic iron oxide particles coated with carboxydextran for parenteral administration and liver contrasting. Pre-clinical Profile of SHU 555A
Acta Radiol 1997 Jul, 38 (4Pt 1):584-97
- Lee A.S., Weissleder R., Brady T.J., Wittenberg J. (1991)
Lymph nodes: microstructural anatomy at MR imaging
Radiology 178: 519-522
- Liebich H.-G. (1993)
Funktionelle Histologie:
Farbatlas und Kurzlehrbuch der mikroskopischen Anatomie der Haussäugetiere
Schattauer, Stuttgart, New York, 2. Auflage

- Listinsky C.M. (1988)
Common reactive erythrophagocytosis in axillary lymph nodes
Am J Clin Pathol 1988 Aug, 90(2):189-92
- Macdougall I.C. (1999)
Strategies for iron supplementation: oral versus intravenous
Kidney Int Suppl 1999 Mar, 69:S61-6
- Majeed S.K. (1994)
Mast cell distribution in rats
Arzneimittelforschung 1994 Mar, 44 (3):370-4
- Marglin S.I., Castellin R.A. (1979)
Severe pulmonary hemorrhage following lymphography
Cancer 1979 Feb, 43(2):482-3
- Mariani G., Morsco L., Viale G., Villa G., Bagnasco M., Canavese G., Buscombe J., Strauss H.W., Paganelli G. (2001)
Radioguided sentinel lymph node biopsy in breast cancer surgery
J Nucl Med 2001, 42:1198-1215
- Matsuno T., Mori M., Awai M. (1985)
Distribution of ferritin and hemosiderin in the liver, spleen and bone marrow of normal, phlebotomized and iron overloaded rats
Acta Med Okayama 1985 Oct, 39(5):347-60
- Maurer J., Willam C., Steinkamp H.J., Knollmann F.D., Felix R. (1996)
Keratinization and necrosis. Morphologic aspects of lymphatic metastases in ultrasound
Invest Radiol 1996 Sep, 31(9):545-9
- McLachlan S.J., Morris M.R., Lucas M.a., Fisco R.A., Eakins M.N., Fowler D.R., Scheetz R.B., Olukotun A.Y. (1994)
Phase I clinical evaluation of a new iron oxide MR contrast agent
J Magn Reson Imaging 1994 May-Jun, 4 (3): 301-7
- Miotti R. (1965)
Die Lymphknoten und Lymphgefäße der weißen Ratte
Acta Anatomica, Vol 62, Nr 4
- Misselwitz B., Sachse A. (1997)
Interstitial MR Lymphography using Gd-carrying liposomes
Acta. Radiol. Suppl. 412, 51-55
- Misselwitz B., Platzek J., Raduchel B., Oellinger J. J., Weinmann H.J. (1999)
Gadofluorine 8: Initial experience with a new contrast medium for interstitial MR Lymphography
Magn. Resonan. Materials in Physics, Biology and Medicine 8/3 (190-195)
- Miyata K., Takaya K. (1985)
Acid-Phosphatase activity of reticular cells and macrophages in the lymph node of the rat after ingestion of mast-cell granules
Histochemistry 1985, 83(3):201-5
- Moore A., Weissleder R, Bogdanov A. (1997)
Uptake of dextran coated monocrySTALLINE iron oxides in tumor cells and macrophages
J Magn Reson Imaging 1997 Nov-Dec, 7(6):1140-5
- Mühler A., Zang X., Wang H. Lawaczeck R., Weinmann H.- J. (1995)
Investigation of mechanisms influencing the accumulation of USPIO in lymph nodes
Invest. Radiology 1995-30, 98-103

- Niendorf H. P., Balzer T. (1997)
 In: Reiser, Semmel
 Magnetresonanztomographie
 Springer, Berlin, 2. Auflage
- Oehmichen M., Wietholter H., Wolburg H. (1982)
 Enhanced phagocytic activity of lymph node macrophages after intranodular injection of
 autologous red blood cells
 Z Rechtsmed 1982, 88(4):285-96
- Okazaki S., Hatayama K., Tamura K. Saito M., Watanabe S. (1994)
 One-month intravenous toxicity study of AMI 25 with a recovery period of one month
 in rats
 Jpn- Pharmacol.- Ther. 22: 57-91
- Okon E., Pouliquen D., Okon P., Kovaleva Z.V., Stepanova T.P. Lavit S.G., Kudryavtsev
 B.N., Jallet P. (1994)
 Biodegradation of magnetite dextran nanoparticles in the rat: A histologic and biophysical
 study
 Lab Invest 1994 Dec, 71(6):895-903
- Penzes L. (1977)
 The rodent as an experimental tool in gerontological research
 Aktuelle Gerontol 1977 Oct, 7(10):513-20
- Percy H., Barthold S. (1993)
 Pathology of laboratory rodents and rabbits
 Iowa State Univ. Press 1993
- Petzinger E. (1996)
 In: Frey H.-H., Loescher W
 Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin
 Enke, Stuttgart
- Pfefferer D., Wagner S., Taupitz M., Kresse M., Elste V., Ebert W. (1993)
 Intravenous MR Lymphography with ultrasmall iron oxide particles: Dose and time response
 to evaluate the interlymphnodal distribution.
 Book of Abstracts: Society of Magnetic Resonance in Medicine
 New York
- Pond G.D., Castellino R.A., Horning S., Hoppe R.T. (1989)
 Non-Hodkin Lymphoma: influence of Lymphography, CT and bone marrow biopsy on staging
 and management
 Radiology 1989 Jan, 170 :159-64
- Pouliquen D., Le Jeune J.J., Perdrisot R., Ermias A., Jallet P. (1991)
 Iron oxide nanoparticles for use as an MRI contrast agent: pharmacokinetics and metabolism
 Magn Reson Imaging 1991, 9(3): 275-83
- Predescu D., Palade G.E., (1993)
 Plasmalemmal vesicles represent the large pore system of continuous microvascular
 endothelium
 Am J Physiol 1993 Aug, 265 (2Pt 2):H725-33
- Rankin S.C., Cook G.J.R. (1999)
 Lymphoma: monitoring response to therapy
 Imaging 1999, 11, 269-281
- Reimer P., Weissleder R., Lee A.S., Wittenberg J. Brady T.J. (1990)
 Receptor imaging: Application to MR imaging of liver cancer
 Radiology 177 (1990) 729-734

- Reimer P., Weissleder R., Lee A.S., Buettner S., Wittenberg J. Brady T.J. (1991)
Asialoglycoprotein Receptor function in benign Liver disease: Evaluation with MRI
Radiology 178 (1991)769-774
- Reimer P., Weissleder R., Shen T., Knoefel W.T., Brady T.J. (1994)
Pancreatic Receptors: Initial feasibility studies with a targeted contrast agent for MRI
Radiology 193: 527-531
- Reinhardt M., Ehrhrit-Braun C., Vogelsang D., Ihling C., Högerle S., Mix M., Moser E.,
Krause T. (2001)
Metastatic lymph nodes in patients with cervical cancer: Detection with MR imaging and
FDG PET
Radiology 2001, 218: 776-782
- Réty F., Clément O., Siauve N., Cuénod C-A., Carnot F., Sich M., Buisine A., Frija G.
(2000)
MR Lymphography using iron oxide nanoparticles in rats: pharmacokinetics in the lymphatic
system after intravenous injection
J Magn Res Imaging 12:734-739 (2000)
- Richard D.M., Mumaw V.R., Schoenbert M.D. (1961)
The transport and distribution of colloidal iron and its relation to the ultrastructure of the cell
J. Ultrastructure Research 5: 244-256
- Rink T.; Heuser T., Fitz H., Schroth H.J., Weller E., Zippel H.H. (2001)
Results of a standardized protocol for sentinel node imaging in breast cancer with Tc-99m
labeled nanocolloidal albumin
Nuklearmedizin 2001 Jun, 40(3):80-5
- Rohrbach M.S., Kreofsky T.J., Vuk-Pavlovic Z., Lauque D. (1992)
Cotton condensed Tannin: a potent modulator of alveolar macrophage host-defense
function
Basic Life Sci 1992, 59:803-24
- Ruehm S., Schroeder T., Debatin J. (2001)
Interstitial MR Lymphography with Gadoterate Meglumine: Initial experience in humans
Radiology 2001: 220:816-21
- Rutledge J.C., Curry F.E., Blanche P., Krauss R.M. (1995)
Solvent drag of LDL across mammalian endothelial barriers with increased permeability
Am J Physiol 1995 May, 268 (5Pt2):H1982-91
- Saini S., Stark D.D., Hahn P.F., Wittenberg J., Brady T.J. Ferrucci J.T. (1987)
Ferrite particles: A superparamagnetic MR contrast agent for the Reticuloendothelial System
Radiology 162 (1987) 211-216
- Sainte-Marie G., Peng F.S., Belisle C. (1982)
Overall Architecture and Pattern of Lymph Flow in the Rat Lymph Node
The American Journal of Anatomy 164:275-309
- Scheunert A., Trautmann A. (1987)
Lehrbuch der Veterinär Physiologie
Parey, Berlin 1987
- Schlittgen, R. (1996):
Einführung in die Statistik. Analyse und Modellierung von Daten
6. Aufl., Oldenbourg München, Wien

- Schmidt H. (1987)
 In: Lissner, Seiderer
 Klin. Kernspintomographie
 Enke, Stuttgart
- Schnorr, B. (1999)
 Makroskopisch- anatomische, kernspintomographische und histologische Untersuchung zum
 Zeitverlauf der Aufnahme sowie zur Verteilung interstitiell verabreichter eisenhaltiger
 Kontrastmittel in den Lymphknoten der Ratte
 Diss Vet. Med., FU Berlin
- Schnorr J., Taupitz M., Wagner S., Pilgrimm H., Hansel J., Hamm B. (2000)
 Age-related blood half-life of particulate contrast material: experimental results with a USPIO
 in rats
 J Magn Res Imaging 12:740-744 (2000)
- Schnorr, J. (2002)
 Persönliche Mitteilung
- Schuhmann-Giampieri G., Schmitt-Willich H., Press W.-R., Negishi C., Weinmann H.-J.,
 Speck U. (1992)
 Preclinical evaluation of Gd-EOB- DTPA as a contrast agent in MR Imaging of the
 hepatobiliary system
 Radiology 183 (1992) 59-64
- Schulze E., Ferrucci J. T., Poss K., Lapointe L., Bogdanova A., Weissleder R. (1995)
 Cellular uptake and trafficking of a prototypical magnetic iron oxide label in vitro
 Invest Radiol 1995 Oct, 30(10): 604-10
- Seiderer M. (1987)
 In: Lissner, Seiderer
 Klin. Kernspintomographie
 Enke, Stuttgart
- Semmler W. (1997)
 In: Reiser, Semmel
 Magnetresonanztomographie
 Springer, Berlin
- Staatz G., Nolte-Ernsting C., Adam G., Grosskortenhaus S., Miselwitz B., Bücken A., Günther
 R. (2001)
 Interstitial T1-weighted MR Lymphography: lipophilic perfluorinated Gadolinium chelates in
 pigs
 Radiology 2001, 220:129-134
- Steinkamp H. J., Zwicker C., Langer M., Mathe M., Ehrlich C., Neumann K., Felix R. (1992)
 Reactive enlargement of cervical lymph nodes and cervical lymph node metastases: sonography
 (M/Q quotient) and computed tomography
 Aktuelle Radiol 1992 Jul, 2 (4):188-95
- Steinkamp H.J., Cornehl M., Hosten N., Pegios W., Vogl T., Felix R. (1995)
 Cervical lymphadenopathy: ratio of long-to-short-axis diameter as a predictor of malignancy
 Br J Radiol 1995 Mar, 68 (807):266-70
- Steinkamp H.J., Teichgraber U.K., Mueffelman M., Hosten N., Kenzel P., Felix R. (1999)
 Differential diagnosis of lymph node lesions. A semiquantitative approach with power
 Doppler sonography
 Invest Radiol 1999 Aug; 34(8):509-15

- Swift S., Husband J.E.S. (1999)
The identification of lymph node disease in lymphoma
Imaging 1999, 11, 230-239
- Tanimoto A., Oshio K., Suematsu M., Pouliquen D., Stark D. (2001)
Relaxation effects of clustered particles
J Magn Reson Imaging 2001, 14:72-77
- Taupitz M., Wagner S., Hamm B. (1996)
Kontrastmittel für die Magnetresonanztomographische Lymphknotendiagnostik (MR Lymphographie)
Der Radiologe 1996, 36(2), 134-140
- Theil E.C., Sayers D.E., Brown M.A., (1979)
Similarity of the structure of ferritin and iron-dextran (imferon) determined by extended X-ray absorption fine structure analysis
J Biol Chem 1979 Sep 10, 254 (17): 8132-4
- Tiedjen K.U., Kosberg R., Kruger G. (1986)
Pulmonary microlithiasis following iodine lipiodol Lymphography
Radiologe 1986 Okt., 26 (19):460-3
- Torchia M., Nason R., Danzinger R., Lewis J., Thliveris J. (2001)
Interstitial MR Lymphangiography for the detection of sentinel lymph nodes
J Surg Onkol 2001, 78:151-157
- Van Beers B.E. Sempoux C., Materne R., Delos M. Smith A.M. (2001)
Biodistribution of ultrasmall iron oxide particles in the rat liver
J Magn Reson Imaging 2001 Apr, 13 (4):594-9
- Vogl T.J., Pegios W., Waitzinger J., Pirovano G., Balzer J., Lissner J. (1992)
Kernspintomographie der Leber mit dem neuen Kontrastmittel Gd- BOPTA: Ergebnisse einer In- Vivo- Phase-I- Prüfung
Fortschr. Röntgenstr. 156, 5 (1992) 465-470
- Vogl T.J., Hammerstingl R., Pegios W. (1994)
Wertigkeit des leberspezifischen superparamagnetischen Kontrastmittels AMI-25 für die Detektion und Differenzierung lebereigener Tumoren versus Metastasen.
Fortschr. Röntgenstr. 160 (1994) 319-328
- Vogl T.J., Schwarz W., Hammerstingl R., Kummel S., Balzer J.O., Mack M.G., Diebold T., Balzer T., Shamsi K., Bechstein W.O., Schrem H., Neuhaus P., Felix R. (1997)
Dynamic and static MRI using superparamagnetic MRI contrast medium RESOVIST for visualizing primary and secondary liver tumors
Fortschr. Röntgenstr. 167 (1997) 254-273
- Wagner S., Pfefferer D., Ebert W., Kresse M., Taupitz M., Hamm B., Lawczeck R., Wolf K.-J. (1995)
Intravenous MR lymphography with superparamagnetic iron oxide particles: experimental studies in rats and rabbits.
Eur. Radiol. 5
- Wagner S., Schnorr S., Pilgrimm H., Hamm., Taupitz M. (2002a)
Monomer-coated Very Small Superparamagnetic Iron Oxide Particles (VSOP) as contrast medium for MRI: Preclinical in vivo Characterization
Invest Radiology im Druck
- Wagner S. (2002b)
Persönliche Mitteilung

- Wang Y., Hussain S., Krestin G. (2001)
 Superparamagnetic iron oxide contrast agents: physicochemical characteristics and applications in MR imaging
 Eur. Radiol. (2001) 11: 2319-2331
- Weir M.P., Gibson J.F., Peters T.J. (1984)
 Haemosiderosis and tissue damage
 Cell Biochem Funct 2: 186-194
- Weiss L., (1983)
 Histology
 Urban & Schwarzenberg, München
- Weissleder R., Hahn P.F., Stark D.D., Rummeny E., Saini S., Wittenberg J., Ferrucci J.T. (1987)
 Mr Imaging of splenic metastases: ferrite enhanced detection in rats
 Radiology 149 (1987) 723-726
- Weissleder R., Stark D.D., Rummeny E., Compton C.C., Ferrucci J.T. (1988)
 Splenic Lymphoma: Ferrite-enhanced MR Imaging in rats
 Radiology 166 (1988) 423-430
- Weissleder R., Stark D. D., Engelstad B.L., Bacon B.R., Compton C.C., White D.L., Jacobs P., Lewis J. (1989a)
 Superparamagnetic Iron Oxide: Pharmacokinetics and toxicity
 AJR 152: 167-173
- Weissleder R., Elizondo G., Stark D.D., Hahn P.F. Marfil J., Gonzales J.F., Saini S., Todd L.E. (1989b)
 The diagnosis of splenic lymphoma by MR imaging: value of superparamagnetic iron oxide
 AJR 152: 175-180
- Weissleder R., Elizondo G., Josephson L., Compton C.C., Fretz C.J., Stark D.D., Ferruci J.T. (1989c)
 Experimental lymph node metastases: enhanced detection with MR lymphography
 Radiology 171: 835-839
- Weissleder R., Elizondo G., Wittenberg J., Rabito C. A., Bengel H.H., Josephson L. (1990a)
 Ultrasmall superparamagnetic iron oxide: characterization of a new class of contrast agents for MR imaging
 Radiology 1990 May, 175 (2):489-93
- Weissleder R., Elizondo G., Wittenberg J., Lee A.S. Josephson L., Brady T.J. (1990b)
 USPIO: an intravenous contrast agent for assessing lymph nodes with MR imaging
 Radiology 1990 May, 175(2): 494-8
- Weissleder R., Lee A.S., Fischman A.J., Reimer P., Shen T., Wilkinson R. Callahan R.J., Brady T.J. (1991)
 Polyclonal human Immunoglobulin G labeled with polymeric Iron Oxide: Antibody MRI
 Radiology 181 (1991) 245-249
- Weissleder R., Heautot J. F., Schaffer B. K., Nossiff N., Papisov M. I., Bogdanov A., Brady T.J. (1994)
 Mr Lymphography: study of a high- efficiency lymphotropic agent.
 Radiology 191:225-230
- Wunderbaldinger P., Josephson L., Bremer C., Moore A., Weissleder R. (2002)
 Detection of lymph node metastases by contrast enhanced MRI in an experimental model
 Magn Reson in Medicine 2002, 47:292-297

- Yagi T, Sato A., Hayakawa H., Ide K., (1997)
Failure of aged rats to accumulate eosinophils in allergic inflammation of the airway
J Allergy Clin Immunol 1997 Jan, 99 (1 Pt 1): 38-47
- Young L.R. (1999)
In: Dawson P., Cosgrove D., Grainger R.
Textbook of contrast media
Isis Medical Media Ltd, Oxford, 1999
- Yuan X.M., Anders W.L., Olsson A.G., Brunk U.T. (1996)
Iron in human atheroma and LDL oxidation by macrophages following erythrophagocytosis
Atherosclerosis 1996 Jul; 124 (1):61-73