

## 5 Literatur

- Abdou M, Storring JM, Riad M, Paquette Y, Albert PR, Drobetsky E und Kouassi E (2001). "Transcriptional mechanisms for induction of 5-HT1A receptor mRNA and protein in activated B and T lymphocytes." *J Biol Chem* **276**(6): 4382-8.
- Adayev T, El-Sherif Y, Barua M, Penington NJ und Banerjee P (1999). "Agonist stimulation of the serotonin1A receptor causes suppression of anoxia-induced apoptosis via mitogen-activated protein kinase in neuronal HN2-5 cells." *J Neurochem* **72**(4): 1489-96.
- Adham N, Zgombick JM, Bard J und Branchek TA (1998). "Functional characterization of the recombinant human 5-hydroxytryptamine7(a) receptor isoform coupled to adenylate cyclase stimulation." *J Pharmacol Exp Ther* **287**(2): 508-14.
- Aghajanian GK (1995). Electrophysiology of serotonin receptor subtypes and signal transduction pathways. In: Bloom FR und Kupfer DJ (Eds.), *Psychopharmacology: The Fourth Generation of Progress*. New York, Raven: pp. 1451-1459.
- Aghajanian GK, Andrade R, Azmitia EC, Baumgarten HG und al. e (1997). Handbook of Experimental Pharmacology, edn.; Baumgarten HG und Göthert M, Serotonergic Neurons and 5-HT Receptors in the CNS. Berlin Heidelberg New York, Springer-Verlag.
- Aghajanian GK und Gallager DW (1975). "Raphe origin of serotonergic nerves terminating in the cerebral ventricles." *Brain Res* **88**(2): 221-31.
- Aghajanian GK und Wang RY (1977). "Habenular and other midbrain raphe afferents demonstrated by a modified retrograde tracing technique." *Brain Res* **122**(2): 229-42.
- Aguzzi A und Theuring F (1994). "Improved in situ beta-galactosidase staining for histological analysis of transgenic mice." *Histochemistry* **102**(6): 477-81.
- Albert PR, Lembo P, Storring JM, Charest A und Saucier C (1996). "The 5-HT1A receptor: signaling, desensitization, and gene transcription." *Neuropsychopharmacology* **14**(1): 19-25.
- Albert PR und Tiberi M (2001). "Receptor signaling and structure: insights from serotonin-1 receptors." *Trends Endocrinol Metab* **12**(10): 453-60.
- Albert PR, Zhou QY, Van Tol HH, Bunzow JR und Civelli O (1990). "Cloning, functional expression, and mRNA tissue distribution of the rat 5-hydroxytryptamine1A receptor gene." *J Biol Chem* **265**(10): 5825-32.
- Alberts B, Bray D, Lewis J, Raff M, Roberts K und Watson JD (1995). Molekularbiologie der Zelle, 3 edn.; Jaenicke L, Weinheim, Wiley-VCH Verlag.
- Alberts GL, Chio CL und Im WB (2001). "Allosteric modulation of the human 5-HT(7A) receptor by lipidic amphipathic compounds." *Mol Pharmacol* **60**(6): 1349-55.
- Alvarez FJ, Pearson JC, Harrington D, Dewey D, Torbeck L und Fyffe RE (1998). "Distribution of 5-hydroxytryptamine-immunoreactive boutons on alpha-motoneurons in the lumbar spinal cord of adult cats." *J Comp Neurol* **393**(1): 69-83.

Arriza JL, Weinberger C, Cerelli G, Glaser TM, Handelin BL, Housman DE und Evans RM (1987). "Cloning of human mineralocorticoid receptor complementary DNA: structural and functional kinship with the glucocorticoid receptor." Science **237**(4812): 268-75.

Ayoubi TA, Creemers JW, Roebroek AJ und Van de Ven WJ (1994). "Expression of the dibasic proprotein processing enzyme furin is directed by multiple promoters." J Biol Chem **269**(12): 9298-303.

Azizkhan JC, Jensen DE, Pierce AJ und Wade M (1993). "Transcription from TATA-less promoters: dihydrofolate reductase as a model." Crit Rev Eukaryot Gene Expr **3**(4): 229-54.

Azmitia EC (1978). The serotonin-producing neurons of the midbrain median and dorsal raphe nuclei, edn.; In Iversen LL, Iversen SD und Snyder SH, Handbook of Psychopharmacology, vol. 9. New York, Plenum Press pp 233-314.

Azmitia EC und Segal M (1978). "An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat." J Comp Neurol **179**(3): 641-67.

Azmitia EC und Whitaker-Azmitia PM (1997). Handbook of experimental pharmacology: Serotonergic neurons and 5-HT receptors in the CNS, edn.; In: Baumgarten HG und Goertner M, Development and Neuroplasticity of Central Serotonergic Neurons., pp 1-39.

Bacon SJ, Zagon A und Smith AD (1990). "Electron microscopic evidence of a monosynaptic pathway between cells in the caudal raphe nuclei and sympathetic preganglionic neurons in the rat spinal cord." Exp Brain Res **79**(3): 589-602.

Bacon WL und Beck SG (2000). "5-Hydroxytryptamine(7) receptor activation decreases slow afterhyperpolarization amplitude in CA3 hippocampal pyramidal cells." J Pharmacol Exp Ther **294**(2): 672-9.

Baker LP, Nielsen MD, Impey S, Metcalf MA, Poser SW, Chan G, Obrietan K, Hamblin MW und Storm DR (1998). "Stimulation of type 1 and type 8 Ca<sup>2+</sup>/calmodulin-sensitive adenylyl cyclases by the Gs-coupled 5-hydroxytryptamine subtype 5-HT<sup>7A</sup> receptor." J Biol Chem **273**(28): 17469-76.

Bard JA, Zgombick J, Adham N, Vaysse P, Branchek TA und Weinshank RL (1993). "Cloning of a novel human serotonin receptor (5-HT<sup>7</sup>) positively linked to adenylate cyclase." J Biol Chem **268**(31): 23422-6.

Barnes NM und Sharp T (1999). "A review of central 5-HT receptors and their function." Neuropharmacology **38**(8): 1083-152.

Behzadi G, Kalen P, Parvopassu F und Wiklund L (1990). "Afferents to the median raphe nucleus of the rat: retrograde cholera toxin and wheat germ conjugated horseradish peroxidase tracing, and selective D-[<sup>3</sup>H]aspartate labelling of possible excitatory amino acid inputs." Neuroscience **37**(1): 77-100.

Bezuglov VV, Bobrov M und Archakov AV (1998). "Bioactive amides of fatty acids." Biochemistry (Mosc) **63**(1): 22-30.

Bianchi C, Siniscalchi A und Beani L (1990). "5-HT1A agonists increase and 5-HT3 agonists decrease acetylcholine efflux from the cerebral cortex of freely-moving guinea-pigs." *Br J Pharmacol* **101**(2): 448-52.

Blake MC, Jambou RC, Swick AG, Kahn JW und Azizkhan JC (1990). "Transcriptional initiation is controlled by upstream GC-box interactions in a TATAA-less promoter." *Mol Cell Biol* **10**(12): 6632-41.

Boger DL, Henriksen SJ und Cravatt BF (1998). "Oleamide: an endogenous sleep-inducing lipid and prototypical member of a new class of biological signaling molecules." *Curr Pharm Des* **4**(4): 303-14.

Bossone SA, Asselin C, Patel AJ und Marcu KB (1992). "MAZ, a zinc finger protein, binds to c-MYC and C2 gene sequences regulating transcriptional initiation and termination." *Proc Natl Acad Sci U S A* **89**(16): 7452-6.

Bourson A, Kapps V, Zwingelstein C, Rudler A, Boess FG und Sleight AJ (1997). "Correlation between 5-HT7 receptor affinity and protection against sound-induced seizures in DBA/2J mice." *Naunyn Schmiedebergs Arch Pharmacol* **356**(6): 820-6.

Bunin MA und Wightman RM (1999). "Paracrine neurotransmission in the CNS: involvement of 5-HT." *Trends Neurosci* **22**(9): 377-82.

Buratowski S (1997). "Multiple TATA-binding factors come back into style." *Cell* **91**(1): 13-5.

Burge CB, Tuschl T und Sharp PA (1999). Splicing of precursors to mRNAs by splicosomes. In: Gesteland RF, Cech TR und Atkins JF (Eds.), *The RNA World*. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press: pp. 525-560.

Burnet PW, Mefford IN, Smith CC, Gold PW und Sternberg EM (1992). "Hippocampal 8-[<sup>3</sup>H]hydroxy-2-(di-n-propylamino) tetralin binding site densities, serotonin receptor (5-HT1A) messenger ribonucleic acid abundance, and serotonin levels parallel the activity of the hypothalamopituitary-adrenal axis in rat." *J Neurochem* **59**(3): 1062-70.

Bussemaker HJ, Li H und Siggia ED (2000). "Regulatory element detection using a probabilistic segmentation model." *Proc Int Conf Intell Syst Mol Biol* **8**: 67-74.

Canton H, Emeson RB, Barker EL, Backstrom JR, Lu JT, Chang MS und Sanders-Bush E (1996). "Identification, molecular cloning, and distribution of a short variant of the 5-hydroxytryptamine2C receptor produced by alternative splicing." *Mol Pharmacol* **50**(4): 799-807.

Cato AC und Wade E (1996). "Molecular mechanisms of anti-inflammatory action of glucocorticoids." *Bioessays* **18**(5): 371-8.

Chalmers DT, Kwak SP, Mansour A, Akil H und Watson SJ (1993). "Corticosteroids regulate brain hippocampal 5-HT1A receptor mRNA expression." *J Neurosci* **13**(3): 914-23.

Chalmers DT, Lopez JF, Vazquez DM, Akil H und Watson SJ (1994). "Regulation of hippocampal 5-HT1A receptor gene expression by dexamethasone." *Neuropsychopharmacology* **10**(3): 215-22.

- Chaouloff F, Berton O und Mormede P (1999). "Serotonin and stress." *Neuropsychopharmacology* **21**(2 Suppl): 28S-32S.
- Chapin EM und Andrade R (2001). "A 5-HT(7) receptor-mediated depolarization in the anterodorsal thalamus. I. Pharmacological characterization." *J Pharmacol Exp Ther* **297**(1): 395-402.
- Chapin EM und Andrade R (2001). "A 5-HT(7) receptor-mediated depolarization in the anterodorsal thalamus. II. Involvement of the hyperpolarization-activated current I(h)." *J Pharmacol Exp Ther* **297**(1): 403-9.
- Charney DS, Krystal JH, Delgado PL und Heninger GR (1990). "Serotonin-specific drugs for anxiety and depressive disorders." *Annu Rev Med* **41**: 437-46.
- Chatelin S, Wehrle R, Mercier P, Morello D, Sotelo C und Weber MJ (2001). "Neuronal promoter of human aromatic L-amino acid decarboxylase gene directs transgene expression to the adult floor plate and aminergic nuclei induced by the isthmus." *Brain Res Mol Brain Res* **97**(2): 149-60.
- Chen NH und Reith ME (1995). "Monoamine interactions measured by microdialysis in the ventral tegmental area of rats treated systemically with (+/-)-8-hydroxy-2-(di-n-propylamino)tetralin." *J Neurochem* **64**(4): 1585-97.
- Chong JA, Tapia-Ramirez J, Kim S, Toledo-Aral JJ, Zheng Y, Boutros MC, Altshuller YM, Frohman MA, Kraner SD, et al. (1995). "REST: a mammalian silencer protein that restricts sodium channel gene expression to neurons." *Cell* **80**(6): 949-57.
- Clark AJ, Bissinger P, Bullock DW, Damak S, Wallace R, Whitelaw CB und Yull F (1994). "Chromosomal position effects and the modulation of transgene expression." *Reprod Fertil Dev* **6**(5): 589-98.
- Connaughton M, Priestley JV, Sofroniew MV, Eckenstein F und Cuello AC (1986). "Inputs to motoneurones in the hypoglossal nucleus of the rat: light and electron microscopic immunocytochemistry for choline acetyltransferase, substance P and enkephalins using monoclonal antibodies." *Neuroscience* **17**(1): 205-24.
- Consolo S, Arnaboldi S, Ramponi S, Nannini L, Ladinsky H und Baldi G (1996). "Endogenous serotonin facilitates in vivo acetylcholine release in rat frontal cortex through 5-HT 1B receptors." *J Pharmacol Exp Ther* **277**(2): 823-30.
- Contesse V, Lenglet S, Grumolato L, Anouar Y, Lihrmann I, Lefebvre H, Delarue C und Vaudry H (1999). "Pharmacological and molecular characterization of 5-hydroxytryptamine(7) receptors in the rat adrenal gland." *Mol Pharmacol* **56**(3): 552-61.
- Corradetti R, Laaris N, Hanoun N, Laporte AM, Le Poul E, Hamon M und Lanfumey L (1998). "Antagonist properties of (-)-pindolol and WAY 100635 at somatodendritic and postsynaptic 5-HT1A receptors in the rat brain." *Br J Pharmacol* **123**(3): 449-62.
- Cummings JP und Felten DL (1979). "A raphe dendrite bundle in the rabbit medulla." *J Comp Neurol* **183**(1): 1-23.

- Darnell J, Lodish H und Baltimore D (1990). Molekulare Zellbiologie, edn.; Berlin, Walter de Gruyter & Co 305-366.
- Davidson JR, DuPont RL, Hedges D und Haskins JT (1999). "Efficacy, safety, and tolerability of venlafaxine extended release and buspirone in outpatients with generalized anxiety disorder." *J Clin Psychiatry* **60**(8): 528-35.
- De Kloet ER (1991). "Brain corticosteroid receptor balance and homeostatic control." *Front Neuroendocrinol* **12**: 95-164.
- Della Rocca GJ, Mukhin YV, Garnovskaya MN, Daaka Y, Clark GJ, Luttrell LM, Lefkowitz RJ und Raymond JR (1999). "Serotonin 5-HT1A receptor-mediated Erk activation requires calcium/calmodulin-dependent receptor endocytosis." *J Biol Chem* **274**(8): 4749-53.
- Diamond MI, Miner JN, Yoshinaga SK und Yamamoto KR (1990). "Transcription factor interactions: selectors of positive or negative regulation from a single DNA element." *Science* **249**(4974): 1266-72.
- Done CJ und Sharp T (1994). "Biochemical evidence for the regulation of central noradrenergic activity by 5-HT1A and 5-HT2 receptors: microdialysis studies in the awake and anaesthetized rat." *Neuropharmacology* **33**(3-4): 411-21.
- Doupnik CA, Dessauer CW, Slepak VZ, Gilman AG, Davidson N und Lester HA (1996). "Time resolved kinetics of direct G beta 1 gamma 2 interactions with the carboxyl terminus of Kir3.4 inward rectifier K<sup>+</sup> channel subunits." *Neuropharmacology* **35**(7): 923-31.
- Drouin J, Sun YL, Chamberland M, Gauthier Y, De Lean A, Nemer M und Schmidt TJ (1993). "Novel glucocorticoid receptor complex with DNA element of the hormone-repressed POMC gene." *Embo J* **12**(1): 145-56.
- Dumont A, Hehner SP, Schmitz ML, Gustafsson JA, Liden J, Okret S, van der Saag PT, Wissink S, van der Burg B, et al. (1998). "Cross-talk between steroids and NF-kappa B: what language?" *Trends Biochem Sci* **23**(7): 233-5.
- Duncan MJ, Short J und Wheeler DL (1999). "Comparison of the effects of aging on 5-HT7 and 5-HT1A receptors in discrete regions of the circadian timing system in hamsters." *Brain Res* **829**(1-2): 39-45.
- Edery I (2000). "Circadian rhythms in a nutshell." *Physiol Genomics* **3**(2): 59-74.
- Edgar DM, Miller JD, Prosser RA, Dean RR und Dement WC (1993). "Serotonin and the mammalian circadian system: II. Phase-shifting rat behavioral rhythms with serotonergic agonists." *J Biol Rhythms* **8**(1): 17-31.
- Ehlen JC, Grossman GH und Glass JD (2001). "In vivo resetting of the hamster circadian clock by 5-HT7 receptors in the suprachiasmatic nucleus." *J Neurosci* **21**(14): 5351-7.
- Erdmann J, Nothen MM, Shimron-Abarbanell D, Rietschel M, Albus M, Borrmann M, Maier W, Franzek E, Korner J, et al. (1996). "The human serotonin 7 (5-HT7) receptor gene: genomic organization and systematic mutation screening in schizophrenia and bipolar affective disorder." *Mol Psychiatry* **1**(5): 392-7.

- Evans RM und Arriza JL (1989). "A molecular framework for the actions of glucocorticoid hormones in the nervous system." *Neuron* **2**(2): 1105-12.
- Fargin A, Raymond JR, Regan JW, Cotecchia S, Lefkowitz RJ und Caron MG (1989). "Effector coupling mechanisms of the cloned 5-HT1A receptor." *J Biol Chem* **264**(25): 14848-52.
- Fischer H (1971). Vergleichende Pharmakologie von Überträgersubstanzen in tiersystematischer Darstellung., edn.; Berlin, Heidelberg, New York, Springer Verlag.
- Fishburn CS, Belleli D, David C, Carmon S und Fuchs S (1993). "A novel short isoform of the D3 dopamine receptor generated by alternative splicing in the third cytoplasmic loop." *J Biol Chem* **268**(8): 5872-8.
- Forbes IT, Dabbs S, Duckworth DM, Jennings AJ, King FD, Lovell PJ, Brown AM, Collin L, Hagan JJ, et al. (1998). "(R)-3,N-dimethyl-N-[1-methyl-3-(4-methyl-piperidin-1-yl)propyl]benzenesulfonamide: the first selective 5-HT<sub>7</sub> receptor antagonist." *J Med Chem* **41**(5): 655-7.
- Freund TF (1992). "GABAergic septal and serotonergic median raphe afferents preferentially innervate inhibitory interneurons in the hippocampus and dentate gyrus." *Epilepsy Res Suppl* **7**: 79-91.
- Galter D und Unsicker K (2000). "Sequential activation of the 5-HT1(A) serotonin receptor and TrkB induces the serotonergic neuronal phenotype." *Mol Cell Neurosci* **15**(5): 446-55.
- Garattini S und Valzelli L (1965). Serotonin, edn.; Amsterdam TN, Elsevier.
- Garrick D, Fiering S, Martin DI und Whitelaw E (1998). "Repeat-induced gene silencing in mammals." *Nat Genet* **18**(1): 56-9.
- Gelernter J, Rao PA, Pauls DL, Hamblin MW, Sibley DR und Kidd KK (1995). "Assignment of the 5HT<sub>7</sub> receptor gene (HTR7) to chromosome 10q and exclusion of genetic linkage with Tourette syndrome." *Genomics* **26**(2): 207-9.
- Gershon MD (1991). "Serotonin: its role and receptors in enteric neurotransmission." *Adv Exp Med Biol* **294**: 221-30.
- Gill CH, Soffin EM, Hagan JJ und Davies CH (2002). "5-HT<sub>7</sub> receptors modulate synchronized network activity in rat hippocampus." *Neuropharmacology* **42**(1): 82-92.
- Gingrich JA und Hen R (2001). "Dissecting the role of the serotonin system in neuropsychiatric disorders using knockout mice." *Psychopharmacology (Berl)* **155**(1): 1-10.
- Giros B, Martres MP, Pilon C, Sokoloff P und Schwartz JC (1991). "Shorter variants of the D3 dopamine receptor produced through various patterns of alternative splicing." *Biochem Biophys Res Commun* **176**(3): 1584-92.
- Glennon RA und Lucki I (1988). Behavioural models of serotonin receptor activation. In: Sanders-Bush E (Eds.), *The Serotonin Receptors*. Clifton, NJ, The Humana: pp. 253-293.
- Gottlicher M, Heck S und Herrlich P (1998). "Transcriptional cross-talk, the second mode of steroid hormone receptor action." *J Mol Med* **76**(7): 480-9.

- Gozlan H, El Mestikawy S, Pichat L, Glowinski J und Hamon M (1983). "Identification of presynaptic serotonin autoreceptors using a new ligand: 3H-PAT." Nature **305**(5930): 140-2.
- Gozlan H, Thibault S, Laporte AM, Lima L und Hamon M (1995). "The selective 5-HT1A antagonist radioligand [3H]WAY 100635 labels both G-protein-coupled and free 5-HT1A receptors in rat brain membranes." Eur J Pharmacol **288**(2): 173-86.
- Graveleau C, Paust HJ, Schmidt-Grimminger D und Mukhopadhyay AK (2000). "Presence of a 5-HT7 receptor positively coupled to adenylate cyclase activation in human granulosa-lutein cells." J Clin Endocrinol Metab **85**(3): 1277-86.
- Green AR und Grahame-Smith DG (1976). "Effects of drugs on the processes regulating the functional activity of brain 5-hydroxytryptamine." Nature **260**(5551): 487-91.
- Green AR und Heal DJ (1985). The effects of drugson serotonin-mediated behavioural models. In: Green AR (Eds.), Neuropharmacology of Serotonin. Oxford, Oxford University: pp. 326-365.
- Gronostajski RM (2000). "Roles of the NFI/CTF gene family in transcription and development." Gene **249**(1-2): 31-45.
- Gross C, Santarelli L, Brunner D, Zhuang X und Hen R (2000). "Altered fear circuits in 5-HT(1A) receptor KO mice." Biol Psychiatry **48**(12): 1157-63.
- Gross C, Zhuang X, Stark K, Ramboz S, Oosting R, Kirby L, Santarelli L, Beck S und Hen R (2002). "Serotonin1A receptor acts during development to establish normal anxiety-like behaviour in the adult." Nature **416**(6879): 396-400.
- Gudermann T, Kalkbrenner F und Schultz G (1996). "Diversity and selectivity of receptor-G protein interaction." Annu Rev Pharmacol Toxicol **36**: 429-59.
- Gustafson EL, Durkin MM, Bard JA, Zgombick J und Branchek TA (1996). "A receptor autoradiographic and in situ hybridization analysis of the distribution of the 5-ht7 receptor in rat brain." Br J Pharmacol **117**(4): 657-66.
- Guy J, Bosler O, Dusticier G, Pelletier G und Calas A (1987). "Morphological correlates of serotonin-neuropeptide Y interactions in the rat suprachiasmatic nucleus: combined radioautographic and immunocytochemical data." Cell Tissue Res **250**(3): 657-62.
- Haj-Dahmane S, Hamon M und Lanfumey L (1991). "K<sup>+</sup> channel and 5-hydroxytryptamine1A autoreceptor interactions in the rat dorsal raphe nucleus: an in vitro electrophysiological study." Neuroscience **41**(2-3): 495-505.
- Hamblin MW, Guthrie CR, Kohen R und Heidmann DE (1998). "Gs protein-coupled serotonin receptors: receptor isoforms and functional differences." Ann N Y Acad Sci **861**: 31-7.
- Hamon M, Gozlan H, el Mestikawy S, Emerit MB, Bolanos F und Schechter L (1990). "The central 5-HT1A receptors: pharmacological, biochemical, functional, and regulatory properties." Ann N Y Acad Sci **600**: 114-29; discussion 129-31.

- Hamprecht B (1977). "Structural, electrophysiological, biochemical, and pharmacological properties of neuroblastoma-glioma cell hybrids in cell culture." *Int Rev Cytol* **49**: 99-170.
- Handley SL (1995). "5-Hydroxytryptamine pathways in anxiety and its treatment." *Pharmacol Ther* **66**(1): 103-48.
- Hapgood JP, Riedemann J und Scherer SD (2001). "Regulation of gene expression by GC-rich DNA cis-elements." *Cell Biol Int* **25**(1): 17-31.
- Heck S, Kullmann M, Gast A, Ponta H, Rahmsdorf HJ, Herrlich P und Cato AC (1994). "A distinct modulating domain in glucocorticoid receptor monomers in the repression of activity of the transcription factor AP-1." *Embo J* **13**(17): 4087-95.
- Hedlund PB, Carson MJ, Sutcliffe JG und Thomas EA (1999). "Allosteric regulation by oleamide of the binding properties of 5-hydroxytryptamine<sub>7</sub> receptors." *Biochem Pharmacol* **58**(11): 1807-13.
- Heidmann DE, Metcalf MA, Kohen R und Hamblin MW (1997). "Four 5-hydroxytryptamine<sub>7</sub> (5-HT<sub>7</sub>) receptor isoforms in human and rat produced by alternative splicing: species differences due to altered intron-exon organization." *J Neurochem* **68**(4): 1372-81.
- Heidmann DE, Szot P, Kohen R und Hamblin MW (1998). "Function and distribution of three rat 5-hydroxytryptamine<sub>7</sub> (5-HT<sub>7</sub>) receptor isoforms produced by alternative splicing." *Neuropharmacology* **37**(12): 1621-32.
- Heisler LK, Chu HM, Brennan TJ, Danao JA, Bajwa P, Parsons LH und Tecott LH (1998). "Elevated anxiety and antidepressant-like responses in serotonin 5-HT1A receptor mutant mice." *Proc Natl Acad Sci U S A* **95**(25): 15049-54.
- Hellendall RP, Schambra U, Liu J, Breese GR, Millhorn DE und Lauder JM (1992). "Detection of serotonin receptor transcripts in the developing nervous system." *J Chem Neuroanat* **5**(4): 299-310.
- Hendricks T, Francis N, Fyodorov D und Deneris ES (1999). "The ETS domain factor Pet-1 is an early and precise marker of central serotonin neurons and interacts with a conserved element in serotonergic genes." *J Neurosci* **19**(23): 10348-56.
- Henikoff S (1998). "Conspiracy of silence among repeated transgenes." *Bioessays* **20**(7): 532-5.
- Hermann DM, Luppi PH, Peyron C, Hinckel P und Jouvet M (1997). "Afferent projections to the rat nuclei raphe magnus, raphe pallidus and reticularis gigantocellularis pars alpha demonstrated by iontophoretic application of choleratoxin (subunit b)." *J Chem Neuroanat* **13**(1): 1-21.
- Hery F, Boulenguez P, Semont A, Hery M, Becquet D, Faudon M, Deprez P und Fache MP (1999). "Identification and role of serotonin 5-HT1A and 5-HT1B receptors in primary cultures of rat embryonic rostral raphe nucleus neurons." *J Neurochem* **72**(5): 1791-801.
- Hery M, Francois-Bellan AM, Hery F, Deprez P und Becquet D (1997). "Serotonin directly stimulates luteinizing hormone-releasing hormone release from GT1 cells via 5-HT<sub>7</sub> receptors." *Endocrine* **7**(2): 261-5.

Hillion J, Milne-Edwards JB, Catelon J, de Vitry F, Gros F und Hamon M (1993). "Prenatal developmental expression of rat brain 5-HT1A receptor gene followed by PCR." Biochem Biophys Res Commun **191**(3): 991-7.

Hirst WD, Price GW, Rattray M und Wilkin GP (1997). "Identification of 5-hydroxytryptamine receptors positively coupled to adenylyl cyclase in rat cultured astrocytes." Br J Pharmacol **120**(3): 509-15.

Hjorth S, Carlsson A, Lindberg P, Sanchez D, Wikstrom H, Arvidsspm H, Hacksell U und Nilsson JLG (1982). "8-Hydroxy-2-(dim-propylamino)tetralin, 8-OH-DPAT, a potent and selective simplifies ergot congener with central 5-HT receptor stimulating activity." J Neural Transm **55**: 169-188.

Hogan B, Beddington R, Constantini F und Lacy E (1994). Manipulating the Mouse Embryo, 2nd edn.; Cold Spring Harbor, Cold Spring Harbor Press.

Holsboer F (1995). Neuroendocrinology of mood disorders. In: Bloom FR und Kupfer DJ (Eds.), Psychopharmacology: The Fourth Generation of Progress. New York, Raven: pp. 957-969.

Hornung JP und Celio MR (1992). "The selective innervation by serotoninergic axons of calbindin-containing interneurons in the neocortex and hippocampus of the marmoset." J Comp Neurol **320**(4): 457-67.

Hoyer D, Clarke DE, Fozard JR, Hartig PR, Martin GR, Mylecharane EJ, Saxena PR und Humphrey PP (1994). "International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (Serotonin)." Pharmacol Rev **46**(2): 157-203.

Huidobro-Toro JP und Harris RA (1996). "Brain lipids that induce sleep are novel modulators of 5-hydroxytryptamine receptors." Proc Natl Acad Sci U S A **93**(15): 8078-82.

Hwang CK, D'Souza UM, Eisch AJ, Yajima S, Lammers CH, Yang Y, Lee SH, Kim YM, Nestler EJ, et al. (2001). "Dopamine receptor regulating factor, DRRF: a zinc finger transcription factor." Proc Natl Acad Sci U S A **98**(13): 7558-63.

Iken K, Chheng S, Fargin A, Goulet AC und Kouassi E (1995). "Serotonin upregulates mitogen-stimulated B lymphocyte proliferation through 5-HT1A receptors." Cell Immunol **163**(1): 1-9.

Iyer VR, Horak CE, Scafe CS, Botstein D, Snyder M und Brown PO (2001). "Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF." Nature **409**(6819): 533-8.

Jasper JR, Kosaka A, To ZP, Chang DJ und Eglen RM (1997). "Cloning, expression and pharmacology of a truncated splice variant of the human 5-HT7 receptor (h5-HT7b)." Br J Pharmacol **122**(1): 126-32.

Kahle W und Frotscher M (2001). Taschenatlas der Anatomie, Band 3: Nervensystem und Sinnesorgane, 7 edn.; Stuttgart, New York, Georg Thieme Verlag, Deutscher Taschenbuchverlag.

Kia HK, Brisorgueil MJ, Daval G, Langlois X, Hamon M und Verge D (1996). "Serotonin1A receptors are expressed by a subpopulation of cholinergic neurons in the rat medial septum and diagonal band of Broca--a double immunocytochemical study." Neuroscience **74**(1): 143-54.

Kia HK, Brisorgueil MJ, Hamon M, Calas A und Verge D (1996). "Ultrastructural localization of 5-hydroxytryptamine1A receptors in the rat brain." *J Neurosci Res* **46**(6): 697-708.

Kikuchi C, Nagaso H, Hiranuma T und Koyama M (1999). "Tetrahydrobenzindoles: selective antagonists of the 5-HT7 receptor." *J Med Chem* **42**(4): 533-5.

Kilpatrick GJ, Dautzenberg FM, Martin GR und Eglen RM (1999). "7TM receptors: the splicing on the cake." *Trends Pharmacol Sci* **20**(7): 294-301.

Kirchgessner AL, Liu MT, Raymond JR und Gershon MD (1996). "Identification of cells that express 5-hydroxytryptamine1A receptors in the nervous systems of the bowel and pancreas." *J Comp Neurol* **364**(3): 439-455.

Kobayashi H, Magnoni MS, Govoni S, Izumi F, Wada A und Trabucchi M (1985). "Neuronal control of brain microvessel function." *Experientia* **41**(4): 427-34.

Konings PN, Shahid M, van Alebeek C, Makkink WK, Stam NJ, Ruigt GS und Vanderheyden PM (1995). "Combined in situ hybridisation, northern blot analysis, and receptor binding studies in clones expressing different levels of the human 5-HT1A receptor." *J Recept Signal Transduct Res* **15**(1-4): 443-55.

Krobert KA, Bach T, Syversveen T, Kværingdal AM und Levy FO (2001). "The cloned human 5-HT7 receptor splice variants: a comparative characterization of their pharmacology, function and distribution." *Naunyn Schmiedebergs Arch Pharmacol* **363**(6): 620-32.

Krobert KA und Levy FO (2002). "The human 5-HT(7) serotonin receptor splice variants: constitutive activity and inverse agonist effects." *Br J Pharmacol* **135**(6): 1563-1571.

Laaris N, Le Poul E, Hamon M und Lanfumey L (1997). "Stress-induced alterations of somatodendritic 5-HT1A autoreceptor sensitivity in the rat dorsal raphe nucleus--in vitro electrophysiological evidence." *Fundam Clin Pharmacol* **11**(3): 206-14.

Lanfumey L und Hamon M (2000). "Central 5-HT(1A) receptors: regional distribution and functional characteristics." *Nucl Med Biol* **27**(5): 429-35.

Lassig JP, Vachirasomtoon K, Hartzell K, Leventhal M, Courchesne E, Courchesne R, Lord C, Leventhal BL und Cook EH, Jr. (1999). "Physical mapping of the serotonin 5-HT(7) receptor gene (HTR7) to chromosome 10 and pseudogene (HTR7P) to chromosome 12, and testing of linkage disequilibrium between HTR7 and autistic disorder." *Am J Med Genet* **88**(5): 472-5.

Lauder JM, Moiseiwitsch JR, Liu J und Wilkie MB (1994). *Alfred Benzon Symp*. **37**: 60-72.

Le Corre S, Sharp T, Young AH und Harrison PJ (1997). "Increase of 5-HT7 (serotonin-7) and 5-HT1A (serotonin-1A) receptor mRNA expression in rat hippocampus after adrenalectomy." *Psychopharmacology (Berl)* **130**(4): 368-74.

Le Poul E, Boni C, Hanoun N, Laporte AM, Laaris N, Chauveau J, Hamon M und Lanfumey L (2000). "Differential adaptation of brain 5-HT1A and 5-HT1B receptors and 5-HT transporter in rats treated chronically with fluoxetine." *Neuropharmacology* **39**(1): 110-22.

Lefebvre H, Contesse V, Delarue C, Vaudry H und Kuhn JM (1998). "Serotonergic regulation of adrenocortical function." *Horm Metab Res* **30**(6-7): 398-403.

Lemonde S, Morris SJ, Bakish D, Du L, Hrdina PD, Kushwahal N und Albert PR (2000). "Abrogated protein-DNA interaction at a C(-1019)G polymorphism in the 5-HT1A receptor gene associated with major depression." Soc. Neurosci. Abstr.: 690.5.

Lenglet S, Delarue C, Lefebvre H, Vaudry H und Contesse V (2000). "Rat glomerulosa cells express functional 5-HT7 receptors." Endocr Res **26**(4): 597-602.

Loots GG, Locksley RM, Blankenspoor CM, Wang ZE, Miller W, Rubin EM und Frazer KA (2000). "Identification of a coordinate regulator of interleukins 4, 13, and 5 by cross-species sequence comparisons." Science **288**(5463): 136-40.

López-Revilla R, Chávez-Dueñas L und Azamar Y (1999). "Improved transgene detection in mouse ear digests." Focus(21): 14.

Lorez HP und Richards JG (1982). "Supra-ependymal serotonergic nerves in mammalian brain: morphological, pharmacological and functional studies." Brain Res Bull **9**(1-6): 727-41.

Lovenberg TW, Baron BM, de Lecea L, Miller JD, Prosser RA, Rea MA, Foye PE, Racke M, Slone AL, et al. (1993). "A novel adenylyl cyclase-activating serotonin receptor (5-HT<sub>7</sub>) implicated in the regulation of mammalian circadian rhythms." Neuron **11**(3): 449-58.

Lucki I (1992). "5-HT1 receptors and behavior." Neurosci Biobehav Rev **16**(1): 83-93.

Lüllmann H, Mohr K und Wehling M (1999). Pharmakologie und Toxikologie. Arzneimittelwirkungen verstehen - Medikamente gezielt einsetzen, 14 edn.; Stuttgart, Thieme Verlag.

Madison DV und Nicoll JA (1986). "Cyclic adenosine 3',5'-monophosphate mediates beta-receptor actions of noradrenaline in rat hippocampal pyramidal cells." J Physiol **372**: 245-59.

Malkoski SP, Handanos CM und Dorin RI (1997). "Localization of a negative glucocorticoid response element of the human corticotropin releasing hormone gene." Mol Cell Endocrinol **127**(2): 189-99.

Marino LR (1993). "Characterization of the mouse carbonic anhydrase II gene promoter." J Biol Chem **268**(10): 7081-9.

McQuade R und Sharp T (1997). "Functional mapping of dorsal and median raphe 5-hydroxytryptamine pathways in forebrain of the rat using microdialysis." J Neurochem **69**(2): 791-6.

Medanic M und Gillette MU (1992). "Serotonin regulates the phase of the rat suprachiasmatic circadian pacemaker in vitro only during the subjective day." J Physiol **450**: 629-42.

Meijer OC und de Kloet ER (1994). "Corticosterone suppresses the expression of 5-HT1A receptor mRNA in rat dentate gyrus." Eur J Pharmacol **266**(3): 255-61.

Meijer OC und de Kloet ER (1995). "A role for the mineralocorticoid receptor in a rapid and transient suppression of hippocampal 5-HT1A receptor mRNA by corticosterone." J Neuroendocrinol **7**(8): 653-7.

Meijer OC, Williamson A, Dallman MF und Pearce D (2000). "Transcriptional repression of the 5-HT1A receptor promoter by corticosterone via mineralocorticoid receptors depends on the cellular context." *J Neuroendocrinol* **12**(3): 245-54.

Menard J und Treit D (1999). "Effects of centrally administered anxiolytic compounds in animal models of anxiety." *Neurosci Biobehav Rev* **23**(4): 591-613.

Meyerhof W, Obermuller F, Fehr S und Richter D (1993). "A novel rat serotonin receptor: primary structure, pharmacology, and expression pattern in distinct brain regions." *DNA Cell Biol* **12**(5): 401-9.

Mezey E, Leranth C, Brownstein MJ, Friedman E, Krieger DT und Palkovits M (1984). "On the origin of the serotonergic input to the intermediate lobe of the rat pituitary." *Brain Res* **294**(2): 231-7.

Millan MJ, Bervoets K und Colpaert FC (1991). "5-hydroxytryptamine (5-HT)1A receptors and the tail-flick response. I. 8-hydroxy-2-(di-n-propylamino) tetralin HBr-induced spontaneous tail-flicks in the rat as an in vivo model of 5-HT1A receptor-mediated activity." *J Pharmacol Exp Ther* **256**(3): 973-82.

Miner JN und Yamamoto KR (1992). "The basic region of AP-1 specifies glucocorticoid receptor activity at a composite response element." *Genes Dev* **6**(12B): 2491-501.

Minowa T, Minowa MT und Mouradian MM (1994). "Negative modulator of the rat D2 dopamine receptor gene." *J Biol Chem* **269**(15): 11656-62.

Moiseiwitsch JR und Lauder JM (1995). "Serotonin regulates mouse cranial neural crest migration." *Proc Natl Acad Sci U S A* **92**(16): 7182-6.

Moiseiwitsch JR, Raymond JR, Tamir H und Lauder JM (1998). "Regulation by serotonin of tooth-germ morphogenesis and gene expression in mouse mandibular explant cultures." *Arch Oral Biol* **43**(10): 789-800.

Moller M (1976). "The ultrastructure of the human fetal pineal gland. II. Innervation and cell junctions." *Cell Tissue Res* **169**(1): 7-21.

Mollgard K und Wiklund L (1979). "Serotonergic synapses on ependymal and hypendymal cells of the rat subcommissural organ." *J Neurocytol* **8**(4): 445-67.

Molliver ME (1987). "Serotonergic neuronal systems: what their anatomic organization tells us about function." *J Clin Psychopharmacol* **7**(6 Suppl): 3S-23S.

Moukhles H, Bosler O, Bolam JP, Vallee A, Umbriaco D, Geffard M und Doucet G (1997). "Quantitative and morphometric data indicate precise cellular interactions between serotonin terminals and postsynaptic targets in rat substantia nigra." *Neuroscience* **76**(4): 1159-71.

Mulcahy RT und Gipp JJ (1995). "Identification of a putative antioxidant response element in the 5'-flanking region of the human gamma-glutamylcysteine synthetase heavy subunit gene." *Biochem Biophys Res Commun* **209**(1): 227-33.

Mullins UL, Gianutsos G und Eison AS (1999). "Effects of antidepressants on 5-HT7 receptor regulation in the rat hypothalamus." *Neuropsychopharmacology* **21**(3): 352-67.

Nam D, Qian IH, Kusumi I, Ulpian C, Tallerico T, Liu IS und Seeman P (1998). "The human serotonin-7 receptor pseudogene: variation and chromosome location." J Psychiatry Neurosci **23**(4): 214-6.

Nelson CS, Cone RD, Robbins LS, Allen CN und Adelman JP (1995). "Cloning and expression of a 5HT7 receptor from *Xenopus laevis*." Receptors Channels **3**(1): 61-70.

Neumaier JF, Sexton TJ, Yracheta J, Diaz AM und Brownfield M (2001). "Localization of 5-HT(7) receptors in rat brain by immunocytochemistry, in situ hybridization, and agonist stimulated cFos expression." J Chem Neuroanat **21**(1): 63-73.

Nirenberg M (1978). "Studies on synapse formation and opiate dependence." Natl Cancer Inst Monogr(48): 339-42.

Novina CD und Roy AL (1996). "Core promoters and transcriptional control." Trends Genet **12**(9): 351-5.

Oleskevich S (1995). "G alpha o1 decapeptide modulates the hippocampal 5-HT1A potassium current." J Neurophysiol **74**(5): 2189-93.

Olsen MA, Nawoschik SP, Schurman BR, Schmitt HL, Burno M, Smith DL und Schechter LE (1999). "Identification of a human 5-HT6 receptor variant produced by alternative splicing." Brain Res Mol Brain Res **64**(2): 255-63.

Olsen MA und Schechter LE (1999). "Cloning, mRNA localization and evolutionary conservation of a human 5-HT7 receptor pseudogene." Gene **227**(1): 63-9.

Osterlund MK, Halldin C und Hurd YL (2000). "Effects of chronic 17beta-estradiol treatment on the serotonin 5-HT(1A) receptor mRNA and binding levels in the rat brain." Synapse **35**(1): 39-44.

Osterlund MK und Hurd YL (1998). "Acute 17 beta-estradiol treatment down-regulates serotonin 5HT1A receptor mRNA expression in the limbic system of female rats." Brain Res Mol Brain Res **55**(1): 169-72.

Ou XM, Jafar-Nejad H, Storring JM, Meng JH, Lemonde S und Albert PR (2000). "Novel dual repressor elements for neuronal cell-specific transcription of the rat 5-HT1A receptor gene." J Biol Chem **275**(11): 8161-8.

Ou XM, Storring JM, Kushwaha N und Albert PR (2001). "Heterodimerization of mineralocorticoid and glucocorticoid receptors at a novel negative response element of the 5-HT1A receptor gene." J Biol Chem **276**(17): 14299-307.

Parks CL, Robinson PS, Sibille E, Shenk T und Toth M (1998). "Increased anxiety of mice lacking the serotonin1A receptor." Proc Natl Acad Sci U S A **95**(18): 10734-9.

Parks CL und Shenk T (1996). "The serotonin 1a receptor gene contains a TATA-less promoter that responds to MAZ and Sp1." J Biol Chem **271**(8): 4417-30.

- Pearce D, Matsui W, Miner JN und Yamamoto KR (1998). "Glucocorticoid receptor transcriptional activity determined by spacing of receptor and nonreceptor DNA sites." *J Biol Chem* **273**(46): 30081-5.
- Pearce D und Yamamoto KR (1993). "Mineralocorticoid and glucocorticoid receptor activities distinguished by nonreceptor factors at a composite response element." *Science* **259**(5098): 1161-5.
- Pecins-Thompson M und Bethea CL (1999). "Ovarian steroid regulation of serotonin-1A autoreceptor messenger RNA expression in the dorsal raphe of rhesus macaques." *Neuroscience* **89**(1): 267-77.
- Pedarzani P, Krause M, Haug T, Storm JF und Stuhmer W (1998). "Modulation of the Ca<sup>2+</sup>-activated K<sup>+</sup> current sIAHP by a phosphatase-kinase balance under basal conditions in rat CA1 pyramidal neurons." *J Neurophysiol* **79**(6): 3252-6.
- Peroutka SJ und Howell TA (1994). "The molecular evolution of G protein-coupled receptors: focus on 5-hydroxytryptamine receptors." *Neuropharmacology* **33**(3-4): 319-24.
- Peroutka SJ und Snyder SH (1979). "Multiple serotonin receptors: differential binding of [<sup>3</sup>H]5-hydroxytryptamine, [<sup>3</sup>H]lysergic acid diethylamide and [<sup>3</sup>H]spiroperidol." *Mol Pharmacol* **16**(3): 687-99.
- Perrais M, Pigny P, Ducourouble MP, Petitprez D, Porchet N, Aubert JP und Van Seuningen I (2001). "Characterization of human mucin gene MUC4 promoter: importance of growth factors and proinflammatory cytokines for its regulation in pancreatic cancer cells." *J Biol Chem* **276**(33): 30923-33.
- Pickel VM, Joh TH und Reis DJ (1978). Immunocytochemical evidence for serotonergic innervation of noradrenergic neurons in nucleus locus ceruleus. In Garattini S et al. (eds), *Interactions Between Putative Neurotransmitters in the Brain. Monographs of the Mario Negri Institute for Pharmacological Research*. Milan, NY, Mario Negri Institute, pp. 369–382
- Pietrantonio PV, Jagge C und McDowell C (2001). "Cloning and expression analysis of a 5HT7-like serotonin receptor cDNA from mosquito Aedes aegypti female excretory and respiratory systems." *Insect Mol Biol* **10**(4): 357-69.
- Plassat JL, Amlaiky N und Hen R (1993). "Molecular cloning of a mammalian serotonin receptor that activates adenylate cyclase." *Mol Pharmacol* **44**(2): 229-36.
- Plotsky PM und Sawchenko PE (1987). "Hypophysial-portal plasma levels, median eminence content, and immunohistochemical staining of corticotropin-releasing factor, arginine vasopressin, and oxytocin after pharmacological adrenalectomy." *Endocrinology* **120**(4): 1361-9.
- Prosser RA, Dean RR, Edgar DM, Heller HC und Miller JD (1993). "Serotonin and the mammalian circadian system: I. In vitro phase shifts by serotonergic agonists and antagonists." *J Biol Rhythms* **8**(1): 1-16.
- Qian IH, Kusumi I, Ulpian C, Tallerico T, Nam D, Liu IS, Seeman MV und Seeman P (1998). "A human serotonin-7 receptor pseudogene." *Brain Res Mol Brain Res* **53**(1-2): 339-43.

Ramboz S, Oosting R, Amara DA, Kung HF, Blier P, Mendelsohn M, Mann JJ, Brunner D und Hen R (1998). "Serotonin receptor 1A knockout: an animal model of anxiety-related disorder." *Proc Natl Acad Sci U S A* **95**(24): 14476-81.

Rapport MM, Green AA und Page IH (1948). "Crystalline Serotonin." *Science* **108**: 329-330.

Rapport MM, Green AA und Page IH (1948). "Partial purification of the vasoconstrictor in beef serum." *J Biol Chem* **174**: 735-738.

Raymond JR, Mukhin YV, Gettys TW und Garnovskaya MN (1999). "The recombinant 5-HT1A receptor: G protein coupling and signalling pathways." *Br J Pharmacol* **127**(8): 1751-64.

Raymond JR, Olsen CL und Gettys TW (1993). "Cell-specific physical and functional coupling of human 5-HT1A receptors to inhibitory G protein alpha-subunits and lack of coupling to Gs alpha." *Biochemistry* **32**(41): 11064-73.

Razani H, Hedlund PB, Hansson AC, Fuxe K und Ogren SO (1997). "5-HT1A receptor activation: short-term effects on the mRNA expression of the 5-HT1A receptor and galanin in the raphe nuclei." *Neuroreport* **8**(16): 3565-70.

Ren B, Robert F, Wyrick JJ, Aparicio O, Jennings EG, Simon I, Zeitlinger J, Schreiber J, Hannett N, et al. (2000). "Genome-wide location and function of DNA binding proteins." *Science* **290**(5500): 2306-9.

Reul JM und de Kloet ER (1985). "Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation." *Endocrinology* **117**(6): 2505-11.

Riad M, Garcia S, Watkins KC, Jodoin N, Doucet E, Langlois X, el Mestikawy S, Hamon M und Descarries L (2000). "Somatodendritic localization of 5-HT1A and preterminal axonal localization of 5-HT1B serotonin receptors in adult rat brain." *J Comp Neurol* **417**(2): 181-94.

Ridet I und Privat A (2000). "Volume transmission." *Trends Neurosci* **23**(2): 58-9.

Robinson DS, Rickels K, Feighner J, Fabre LF, Jr., Gammans RE, Shrotriya RC, Alms DR, Andary JJ und Messina ME (1990). "Clinical effects of the 5-HT1A partial agonists in depression: a composite analysis of buspirone in the treatment of depression." *J Clin Psychopharmacol* **10**(3 Suppl): 67S-76S.

Roth BL, Craig SC, Choudhary MS, Uluer A, Monsma FJ, Jr., Shen Y, Meltzer HY und Sibley DR (1994). "Binding of typical and atypical antipsychotic agents to 5-hydroxytryptamine-6 and 5-hydroxytryptamine-7 receptors." *J Pharmacol Exp Ther* **268**(3): 1403-10.

Ruat M, Traiffort E, Leurs R, Tardivel-Lacombe J, Diaz J, Arrang JM und Schwartz JC (1993). "Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT7) activating cAMP formation." *Proc Natl Acad Sci U S A* **90**(18): 8547-51.

Rudge JS, Eaton MJ, Mather P, Lindsay RM und Whittemore SR (1996). "CNTF induces raphe neuronal precursors to switch from a serotonergic to a cholinergic phenotype in vitro." *Mol Cell Neurosci* **7**(3): 204-21.

- Rupprecht R, Arriza JL, Spengler D, Reul JM, Evans RM, Holsboer F und Damm K (1993). "Transactivation and synergistic properties of the mineralocorticoid receptor: relationship to the glucocorticoid receptor." *Mol Endocrinol* 7(4): 597-603.
- Saffer JD, Jackson SP und Annarella MB (1991). "Developmental expression of Sp1 in the mouse." *Mol Cell Biol* 11(4): 2189-99.
- Sah P (1996). "Ca(2+)-activated K<sup>+</sup> currents in neurones: types, physiological roles and modulation." *Trends Neurosci* 19(4): 150-4.
- Sambrook J, Fritsch E und Maniatis T (1989). Molecular Cloning, A Laboratory Manual, edn.; Cold Spring Harbour, New York, Cold Spring Harbour Laboratory Press.
- Scheibel ME, Tomiyasu U und Scheibel AB (1975). "Do raphe nuclei of the reticular formation have a neurosecretory or vascular sensor function?" *Exp Neurol* 47(2): 316-29.
- Schoenherr CJ und Anderson DJ (1995). "The neuron-restrictive silencer factor (NRSF): a coordinate repressor of multiple neuron-specific genes." *Science* 267(5202): 1360-3.
- Schoenherr CJ und Anderson DJ (1995). "Silencing is golden: negative regulation in the control of neuronal gene transcription." *Curr Opin Neurobiol* 5(5): 566-71.
- Schwarz DA, Barry G, Eliasof SD, Petroski RE, Conlon PJ und Maki RA (2000). "Characterization of gamma-aminobutyric acid receptor GABAB(1e), a GABAB(1) splice variant encoding a truncated receptor." *J Biol Chem* 275(41): 32174-81.
- Sharp T und Hjorth S (1990). "Application of brain microdialysis to study the pharmacology of the 5-HT1A autoreceptor." *J Neurosci Methods* 34(1-3): 83-90.
- Shen Y, Monsma FJ, Jr., Metcalf MA, Jose PA, Hamblin MW und Sibley DR (1993). "Molecular cloning and expression of a 5-hydroxytryptamine7 serotonin receptor subtype." *J Biol Chem* 268(24): 18200-4.
- Shimizu M, Nishida A, Zensho H, Miyata M und Yamawaki S (1997). "Down-regulation of 5-hydroxytryptamine7 receptors by dexamethasone in rat frontocortical astrocytes." *J Neurochem* 68(6): 2604-9.
- Shimizu M, Nishida A, Zensho H, Miyata M und Yamawaki S (1998). "Agonist-induced desensitization of adenylyl cyclase activity mediated by 5-hydroxytryptamine7 receptors in rat frontocortical astrocytes." *Brain Res* 784(1-2): 57-62.
- Shimizu M, Nishida A, Zensho H und Yamawaki S (1996). "Chronic antidepressant exposure enhances 5-hydroxytryptamine7 receptor-mediated cyclic adenosine monophosphate accumulation in rat frontocortical astrocytes." *J Pharmacol Exp Ther* 279(3): 1551-8.
- Sibug RM, Compaan JC, Meijer OC, Van der Gugten J, Olivier B und De Kloet ER (1998). "Flesinoxan treatment reduces 5-HT1A receptor mRNA in the dentate gyrus independently of high plasma corticosterone levels." *Eur J Pharmacol* 353(2-3): 207-14.
- Sleight AJ, Carolo C, Petit N, Zwingelstein C und Bourson A (1995). "Identification of 5-hydroxytryptamine7 receptor binding sites in rat hypothalamus: sensitivity to chronic antidepressant treatment." *Mol Pharmacol* 47(1): 99-103.

Smith TA (1971). "The occurrence, metabolism and functions of amines in plants." Biol Rev Camb Philos Soc **46**(2): 201-41.

Snyder LA, Roberts JL und Sealfon SC (1991). "Alternative transcripts of the rat and human dopamine D3 receptor." Biochem Biophys Res Commun **180**(2): 1031-5.

Stam NJ, Roesink C, Dijcks F, Garritsen A, van Herpen A und Olijve W (1997). "Human serotonin 5-HT7 receptor: cloning and pharmacological characterisation of two receptor variants." FEBS Lett **413**(3): 489-94.

Stamford JA, Davidson C, McLaughlin DP und Hopwood SE (2000). "Control of dorsal raphe 5-HT function by multiple 5-HT(1) autoreceptors: parallel purposes or pointless plurality?" Trends Neurosci **23**(10): 459-65.

Stargell LA und Struhl K (1996). "Mechanisms of transcriptional activation in vivo: two steps forward." Trends Genet **12**(8): 311-5.

Stokes PE und Sikes CR (1987). Hypothalamic-Pituitary-Adrenal Axis in Affective Disorders. In: Meltzer HY (Eds.), Psychopharmacology: The Third Generation of Progress. New York, Raven: pp. 957-969.

Storring JM, Charest A, Cheng P und Albert PR (1999). "TATA-driven transcriptional initiation and regulation of the rat serotonin 5-HT1A receptor gene." J Neurochem **72**(6): 2238-47.

Stowe RL und Barnes NM (1998). "Selective labelling of 5-HT7 receptor recognition sites in rat brain using [<sup>3</sup>H]5-carboxamidotryptamine." Neuropharmacology **37**(12): 1611-9.

Stryer L (1990). Biochemie, 3 edn.; Heidelberg, Spektrum der Wissenschaft Verlagsgesellschaft mbH 735-764.

Suzuki M, Matsuda T, Asano S, Somboonthum P, Takuma K und Baba A (1995). "Increase of noradrenaline release in the hypothalamus of freely moving rat by postsynaptic 5-hydroxytryptamine1A receptor activation." Br J Pharmacol **115**(4): 703-11.

Suzuki Y, Tsunoda T, Sese J, Taira H, Mizushima-Sugano J, Hata H, Ota T, Isogai T, Tanaka T, et al. (2001). "Identification and characterization of the potential promoter regions of 1031 kinds of human genes." Genome Res **11**(5): 677-84.

Tejani-Butt SM und Labow DM (1994). "Time course of the effects of adrenalectomy and corticosterone replacement on 5-HT1A receptors and 5-HT uptake sites in the hippocampus and dorsal raphe nucleus of the rat brain: an autoradiographic analysis." Psychopharmacology (Berl) **113**(3-4): 481-6.

Theuring F, Aguzzi A, Kropp C, Wohn KD, Hoffmann S und Schleuning WD (1995). "Analysis of the human tissue type plasminogen activator gene promoter activity during embryogenesis of transgenic mice and rats." Fibrinol Proteol **9**: 277-287.

Thomas DR, Atkinson PJ, Ho M, Bromidge SM, Lovell PJ, Villani AJ, Hagan JJ, Middlemiss DN und Price GW (2000). "[<sup>3</sup>H]-SB-269970--A selective antagonist radioligand for 5-HT(7) receptors." Br J Pharmacol **130**(2): 409-17.

Thomas EA, Carson MJ, Neal MJ und Sutcliffe JG (1997). "Unique allosteric regulation of 5-hydroxytryptamine receptor-mediated signal transduction by oleamide." *Proc Natl Acad Sci U S A* **94**(25): 14115-9.

Thomas EA, Cravatt BF und Sutcliffe JG (1999). "The endogenous lipid oleamide activates serotonin 5-HT<sub>7</sub> neurons in mouse thalamus and hypothalamus." *J Neurochem* **72**(6): 2370-8.

Tjian R (1996). "The biochemistry of transcription in eukaryotes: a paradigm for multisubunit regulatory complexes." *Philos Trans R Soc Lond B Biol Sci* **351**(1339): 491-9.

To ZP, Bonhaus DW, Eglen RM und Jakeman LB (1995). "Characterization and distribution of putative 5-HT<sub>7</sub> receptors in guinea-pig brain." *Br J Pharmacol* **115**(1): 107-16.

Traber J und Glaser T (1987). "5-HT<sub>1A</sub>-related anxiolytics." *Trends Pharmacol Sci* **8**: 432-437.

Tsou AP, Kosaka A, Bach C, Zuppan P, Yee C, Tom L, Alvarez R, Ramsey S, Bonhaus DW, et al. (1994). "Cloning and expression of a 5-hydroxytryptamine<sub>7</sub> receptor positively coupled to adenylyl cyclase." *J Neurochem* **63**(2): 456-64.

Turek FW (1985). "Circadian neural rhythms in mammals." *Annu Rev Physiol* **47**: 49-64.

Twarog BM (1954). "Responses of a molluscan smooth muscle to acetylcholine and 5-hydroxytryptamine." *J Cell Comp Physiol* **44**(141-163).

Twarog BM und Page IH (1953). "Serotonin content of some mammalian tissues and urine and a method for its determination." *Am J Physiol* **175**(157-161).

Van Hartesveldt C, Moore B und Hartman BK (1986). "Transient midline raphe glial structure in the developing rat." *J Comp Neurol* **253**(2): 174-84.

Vanhoenacker P, Haegeman G und Leysen JE (2000). "5-HT<sub>7</sub> receptors: current knowledge and future prospects." *Trends Pharmacol Sci* **21**(2): 70-7.

Verhofstad AA, Steinbusch HW, Penke B, Varga J und Joosten HW (1981). "Serotonin-immunoreactive cells in the superior cervical ganglion of the rat. Evidence for the existence of separate serotonin- and catecholamine-containing small ganglionic cells." *Brain Res* **212**(1): 39-49.

Waeber C und Moskowitz MA (1995). "Autoradiographic visualisation of [<sup>3</sup>H]5-carboxamidotryptamine binding sites in the guinea pig and rat brain." *Eur J Pharmacol* **283**(1-3): 31-46.

Whitaker-Azmitia PM und Azmitia EC (1994). "Astroglial 5-HT<sub>1A</sub> receptors and S-100 beta in development and plasticity." *Perspect Dev Neurobiol* **2**(3): 233-8.

White LA, Eaton MJ, Castro MC, Klose KJ, Globus MY, Shaw G und Whittemore SR (1994). "Distinct regulatory pathways control neurofilament expression and neurotransmitter synthesis in immortalized serotonergic neurons." *J Neurosci* **14**(11 Pt 1): 6744-53.

Wilkinson LO, Middlemiss DN und Hutson PH (1994). "5-HT<sub>1A</sub> receptor activation increases hippocampal acetylcholine efflux and motor activity in the guinea pig: agonist efficacy influences functional activity in vivo." *J Pharmacol Exp Ther* **270**(2): 656-64.

Wirz-Justice A und Van den Hoofdakker RH (1999). "Sleep deprivation in depression: what do we know, where do we go?" Biol Psychiatry **46**(4): 445-53.

Wissink S, Meijer O, Pearce D, van Der Burg B und van Der Saag PT (2000). "Regulation of the rat serotonin-1A receptor gene by corticosteroids." J Biol Chem **275**(2): 1321-6.

Wissink S, van der Burg B, Katzenellenbogen BS und van der Saag PT (2001). "Synergistic activation of the serotonin-1A receptor by nuclear factor-kappa B and estrogen." Mol Endocrinol **15**(4): 543-52.

Wu S und Comings DE (1999). "A common C-1018G polymorphism in the human 5-HT1A receptor gene." Psychiatr Genet **9**(2): 105-6.

Wurch T, Chastagnier C, Palmier C, Colpaert FC und Pauwels PJ (1996). "A 413 bp Region Upstream the Human 5-HT1A Receptor Gene is Sufficient for its *in vitro* Expression." Neurosci Res Commun **19**(2): 75-82.

Yajima S, Lee SH, Minowa T und Mouradian MM (1998). "Sp family transcription factors regulate expression of rat D2 dopamine receptor gene." DNA Cell Biol **17**(5): 471-9.

Yau JL, Noble J und Seckl JR (2001). "Acute restraint stress increases 5-HT7 receptor mRNA expression in the rat hippocampus." Neurosci Lett **309**(3): 141-4.

Yau JL, Noble J, Widdowson J und Seckl JR (1997). "Impact of adrenalectomy on 5-HT6 and 5-HT7 receptor gene expression in the rat hippocampus." Brain Res Mol Brain Res **45**(1): 182-6.

Yau JL, Olsson T, Noble J und Seckl JR (1999). "Serotonin receptor subtype gene expression in the hippocampus of aged rats following chronic amitriptyline treatment." Brain Res Mol Brain Res **70**(2): 282-7.

Ying SW und Rusak B (1997). "5-HT7 receptors mediate serotonergic effects on light-sensitive suprachiasmatic nucleus neurons." Brain Res **755**(2): 246-54.

Yu GD, Liu YL, Jiang XH, Guo SY, Zhang HQ, Yin QZ und Hisamitsu T (2001). "The inhibitory effect of serotonin on the spontaneous discharge of suprachiasmatic neurons in hypothalamic slice is mediated by 5-HT(7) receptor." Brain Res Bull **54**(4): 395-8.

Zhang L, Barker JL, Xing G, Giorgi O, Ma W, Chang YH, Hu Q, Choi N und Rubinow DR (1997). "5-HT1A receptor mRNA expressions differ in the embryonic spinal cord of male and female rats." Neurosci Lett **237**(1): 41-4.

Zhong P und Ciaranello RD (1995). "Transcriptional regulation of hippocampal 5-HT1a receptors by corticosteroid hormones." Brain Res Mol Brain Res **29**(1): 23-34.

Zhukovskaya NL und Neumaier JF (2000). "Clozapine downregulates 5-hydroxytryptamine6 (5-HT6) and upregulates 5-HT7 receptors in HeLa cells." Neurosci Lett **288**(3): 236-40.