

Bibliography

- AG Bodenkunde (ed.) (2005). *Bodenkundliche Kartieranleitung*. 5 ed. Hannover.
- Ahnert, F. (1996). *Einführung in die Geomorphologie*. Ulmer, Stuttgart.
- Aitchison, J. (1986). *The statistical analysis of compositional data*. Monographs on statistics and applied probability. Chapman and Hall, London.
- Aitchison, J. (1999). Logratios and natural laws in compositional data analysis. *Mathematical Geology*, **31**, 563–580.
- Aitchison, J., Egozcue, J. J. (2005). Compositional data analysis: where are we and where should we be heading. *Mathematical Geology*, **37**, 829–850.
- Aitken, M. J. (1998). *Introduction to optical dating*. Oxford University Press, Oxford.
- Akademiya Nauk (1990). *Mongolskaya Narodua Respublika Nacionalny Atlas*. Adademiya Nauk.
- Alley, R. B., Agustsdottir, A. M. (2005). The 8k event: cause and consequences of a major Holocene abrupt climate change. *Quaternary Science Reviews*, **24**, 10-11, 1123–1149.
- Allsen, T. T. (1997). Ever closer encounters: the appropriation of culture and the apportionment of peoples in the Mongol empire. *Journal of Early Modern History*, **1**, 2–23.
- An, C.-B., Feng, Z.-D., Barton, L. (2006). Dry or humid? Mid-Holocene humidity changes in arid and semi-arid China. *Quaternary Science Reviews*, **25**, 3-4, 351–361.
- An, C.-B., Tang, L., Barton, L., Chen, F.-H. (2005). Climate change and cultural response around 4000 cal yr B.P. in the western part of Chinese Loess Plateau. *Quaternary Research*, **63**, 347–352.
- An, Z. (2000). The history and variability of the East Asian paleomonsoon climate. *Quaternary Science Reviews*, **19**, 171–187.
- An, Z., Kutzbach, J. E., Prell, W. L., Porter, S. C. (2001). Evolution of Asian monsoons and phased uplift of the Himalaya-Tibetan plateau since Late Miocene times. *Nature*, **411**, 62–66.
- An, Z., Porter, S. P., Kutzbach, J. E., Xihao, W., Suming, W., Xiaodong, L., L. Xiaoqiang, Weijian, Z. (2000). Asynchronous Holocene optimum of the East Asian monsoon. *Quaternary Science Reviews*, **19**, 743–762.
- Aspinall, A., Gaffney, C. F. (2001). The Schlumberger array – potential and pitfalls in archaeological prospection. *Archaeological Prospection*, **8**, 199–209.
- Baines, D., Smith, D. G., Froese, D. G., Bauman, P., Nimeck, G. (2002). Electrical resistivity ground imaging (ERGI): a new tool for mapping the lithology and geometry of channel-belts and valley-fills. *Sedimentology*, **49**, 441–449.
- Bao, C. (ed.) (1987). *Synoptic meteorology in China*. China Ocean Press, Beijing.

Bibliography

- Barthel, H. (1983). Die regionale und jahreszeitliche Differenzierung des Klimas in der Mongolischen Volksrepublik. In: Barthel, H., Brunner, H., Haase, G. (eds.), *Physisch-Geographische Studien in Asien, Studia Geographica*, vol. 34. BRNO, pp. 3–91.
- Barthel, H., Haase, G., Richter, H. (1962). Die mongolische Volksrepublik. *Zeitschrift für den Erdkundeunterricht*, **14**, 288–318.
- Batkhisig, O., Lehmkuhl, F. (2003). Degradation und Desertifikation in der Mongolei. *Petermanns Geographische Mitteilungen*, **147**, 5, 48–49.
- Batmunkh, D., Tosun, S., Gündongdu, B., Candansayar, M. E., Ulugergerli, E. U. (2004). HLEM and magnetic surveys: Examples from the Orkhun Valley, Mongolia. *Archaeological Prospection*, **11**, 133–144.
- Batsaikhan, Z., Munkhdelger, J., Zorigt, K., Munkhbold, Z. (2006). Mongolian-American joint archaeological survey and excavation in Tamiriin Ulaan khoshuum Ugiinuur Sum, Arkhangai aimag. *Mongolian Journal of Anthropology, Archaeology and Ethnology*, **2**, 1, 206–208.
- Bemmann, J., Pohl, E., Schütt, B., Schwanghart, W. (2008). Archeological findings in the Upper and Middle Orchon Valley and their geographical setup. In: Bemmann, J., Hüttel, H. G., Pohl, E. (eds.), *Mongolian-German Qara Qorum-Expedition*, FAAK (Forschungen zur Archäologie Außereuropäischer Kulturen). Bonn, p. in press.
- Berger, A., Loutre, M. F. (1991). Insolation values for the climate of the last 10 million years. *Quaternary Science Reviews*, **10**, 297–317.
- Berkey, C. P., Morris, K. K. (1927). *Geology of Mongolia*. American Museum of Natural History, New York.
- Besler, H. (1992). *Geomorphologie der ariden Gebiete, Erträge der Forschung*, vol. 280. Wissenschaftliche Buchgesellschaft, Darmstadt.
- Böhner, J. (2006). General climatic controls and topoclimatic variations in Central and High Asia. *Boreas*, **35**, 279–295.
- Birkeland, P. W. (1999). *Soils and geomorphology*. 3 ed. Oxford University Press, New York, Oxford.
- Blümel, W. D. (1981). *Pedologische und geomorphologische Aspekte der Kalkkrustenbildung in Südwestafrika und Südostspanien, Karlsruher Geographische Hefte*, vol. 10. Karlsruhe.
- Blyakharchuk, T., Wright, H., Borodavko, P., van der Knaap, W., Ammann, B. (2004). Late Glacial and Holocene vegetational changes on the Ulagan high-mountain plateau, Altai Mountains, southern Siberia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **209**, 259–279.
- Boës, X., Piotrowska, N., Fagel, N. (2005). High-resolution diatom/clay record in Lake Baikal from grey scale, and magnetic susceptibility over Holocene and Termination I. *Global and Planetary Change*, **46**, 299–313.
- Bohannon, J. (2008). The big thaw reaches Mongolia's pristine north. *Science*, **319**, 567–568.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., Bonani, G. (2001). Persistent solar influence on North Atlantic climate during the Holocene. *Science*, **294**, 2130–2136.
- Boyle, J. F. (2001). Inorganic geochemical methods in paleolimnology. In: Last, W. M., Smol, J. P. (eds.), *Tracking environmental change using lake sediments. Volume 2: Physical and geochemical methods*. Kluwer, Dordrecht, Boston, London, pp. 83–142.

Bibliography

- Bradley, R. S. (1999). *Paleoclimatology, International Geophysics Series*, vol. 64. 2 ed. Harcourt Academic Press, San Diego, London, Boston.
- Broccoli, A. J., Manabe, S. (1992). The effects of orography on midlatitude northern hemisphere dry climates. *Journal of Climate*, **5**, 1181–1201.
- Broughton, P. L. (1972). Monohydrocalcite in speleothems: an alternative interpretation. *Contributions to Mineralogy and Petrology*, **36**, 171–174.
- Brunsdon, D. (1996). Geomorphological events and landform change. *Zeitschrift für Geomorphologie*, **40**, 273–288.
- Bullard, J. E. (2004). Arid geomorphology. *Progress in Physical Geography*, **28**, 130–144.
- Bush, A. B. (2005). CO₂/H₂O and orbitally driven climate variability over central Asia through the Holocene. *Quaternary International*, **136**, 1, 15–23.
- Bush, A. B. G. (2004). Modelling of late Quaternary climate over Asia: a synthesis. *Boreas*, **33**, 155–163.
- Bush, A. B. G., Little, E. C., Rokosh, D., White, D., Rutter, N. W. (2004). Investigation of the spatio-temporal variability in Eurasian Late Quaternary loess-paleosol sequences using a coupled atmosphere-ocean general circulation model. *Quaternary Science Reviews*, **23**, 481–498.
- Chen, F., Yu, Z., Yang, M., Ito, E., Wang, S., Madsen, D. B., Huang, X., Zhao, Y., Sato, T., Birks, H. J. B., Boomer, I., Chen, J., An, C., Wünnemann, B. (2008). Holocene moisture evolution in arid central Asia and its out-of-phase relationship with Asian monsoon history. *Quaternary Science Reviews*, **27**, 351–364.
- Chen, F.-H., Cheng, B., Zhao, Y., Zhu, Y., Madsen, D. B. (2006). Holocene environmental change inferred from a high-resolution pollen record, Lake Zhuyeze, arid China. *The Holocene*, **16**, 5, 675–684.
- Chlachula, J. (2003). The Siberian loess record and its significance for reconstruction of Pleistocene climate change in north-central Asia. *Quaternary Science Reviews*, **22**, 18-19, 1879–1906.
- Claussen, M., Kubatzki, C., Brovkin, V., Ganopolski, A., Hoelzmann, P., Pachur, H. J. (1999). Simulation of an abrupt change in Saharan vegetation in the mid-Holocene. *Geophysical Research Letters*, **26**, 2037–2040.
- Cronin, T. M. (1999). *Principles of Paleoclimatology*. Columbia University Press, New York.
- Dahl, K., Buchardt, B. (2006). Monohydrocalcite in the Arctic Ikka Fjord, SW Greenland: First Reported Marine Occurrence. *Journal of Sedimentary Research*, **76**, 460–471.
- Dahlin, T. (2001). The development of DC resistivity imaging techniques. *Computers and Geosciences*, **27**, 1019–1029.
- Danzeglocke, U., Jöris, O., Weninger, B. (2008). CalPal-2007 online. <http://www.calpal-online.de/>, accessed 2008-02-29.
- D'Arrigo, R., Jacoby, G., Frank, D., Pederson, N., Cook, E., Buckley, B., Nachin, B., Mijiddorj, R., Dugarjav, C. (2001). 1738 years of Mongolian temperature variability inferred from a tree-ring width chronology of Siberian pine. *Geophysical Research Letters*, **28**, 543–546.
- D'Arrigo, R., Jacoby, G., Wilson, R., Panagiotopoulos, F. (2005). A reconstructed Siberian High index since A.D. 1599 from Eurasian and North American tree rings. *Geophysical Research Letters*, **32**, L05705: 1–4.

Bibliography

- Davis, J. C. (1986). *Statistics and Data Analysis in Geology*. 2 ed. John Wiley & Sons, New York, Chichester, Brisbane.
- Davis, M. E., Thompson, L. G., Yao, T., Wang, N. (2005). Forcing of the Asian monsoon on the Tibetan Plateau: Evidence from high-resolution ice core and tropical coral records. *Journal of Geophysical Research*, **110**, D04101.
- Dean, W. E. (1999). The carbon cycle and biogeochemical dynamics in lake sediments. *Journal of Paleolimnology*, **21**, 375–393.
- Demske, D., Heumann, G., Granoszewski, W., Nita, M., Mamakowa, K., Tarasov, P. E., Oberhansli, H. (2005). Late glacial and Holocene vegetation and regional climate variability evidenced in high-resolution pollen records from Lake Baikal. *Global and Planetary Change*, **46**, 1-4, 255–279.
- Demske, D., Mohr, B., Oberhansli, H. (2002). Late Pliocene vegetation and climate of the Lake Baikal region, southern East Siberia, reconstructed from palynological data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **184**, 1-2, 107–129.
- Derbyshire, E., Meng, X., Kemp, R. A. (1998). Provenance, transport and characteristics of modern aeolian dust in western Gansu Province, China, and interpretation of the Quaternary loess record. *Journal of Arid Environments*, **39**, 497–516.
- Dikau, R. (2006). Komplexe Systeme in der Geomorphologie. *Mitteilungen der Österreichischen Geographischen Gesellschaft*, **148**, 125–150.
- Dill, H. G., Khishigsuren, S., Majigsuren, Y., Myagmarsuren, S., Bulgamaa, J. (2006). Geomorphological studies along a transect from the taiga to the desert in Central Mongolia—evolution of landforms in the mid-latitude continental interior as a function of climate and vegetation. *Journal of Asian Earth Sciences*, **27**, 2, 241–264.
- Domrös, M., Peng, G. (1988). *The climate of China*. Springer, Berlin, Heidelberg, New York.
- Dong, Z., Xunming, W., Liu, L. (2000). Wind erosion in arid and semiarid China: an overview. *Journal of Soil and Water Conservation*, **55**, 439–444.
- Dorofeyuk, N. I., Tarasov, P. E. (1998). Vegetation and lake levels in Northern Mongolia in the last 12,500 years as indicated by data of pollen and diatom analyses. *Stratigraphy and Geological Correlation*, **6**, 1, 70–83.
- Drompp, M. R. (1999). Breaking the Orkhon Tradition: Kirghiz adherence to the Yenisei Region after A. D. 840. *Journal of the American Oriental Society*, **119**, 3, 390–403.
- Drompp, M. R. (2005). *Tang China and the collapse of the Uighur Empire: A documentary history*. BRILL.
- Duan, A. M., Wu, G. X. (2005). Role of the Tibetan Plateau thermal forcing in the summer climate patterns over subtropical Asia. *Climate Dynamics*, **24**, 793–807.
- Dulam, J. (2005). Discriminate analysis for dust storm prediction in the gobi and steppe regions in Mongolia. *Water, Air, and Soil Pollution*, **5**, 37–49.
- Dulamsuren, C., Hauck, M., Mühlenberg, M. (2005a). Ground vegetation in the Mongolian taiga forest-steppe ecotone does not offer evidence for the human origin of grasslands. *Applied Vegetation Science*, **8**, 149–154.
- Dulamsuren, C., Hauck, M., Mühlenberg, M. (2005b). Vegetation at the taiga forest-steppe borderline in the western Khentey Mountains, northern Mongolia. *Annales Botanici Fennici*, **42**, 411–426.

Bibliography

- Dunne, T., Zhang, W., Aubry, B. (1991). Effects of rainfall, vegetation and microtopography on infiltration and runoff. *Water Resources Research*, **27**, 2271–2285.
- Dykoski, C. A., Edwards, R. L., Cheng, H., Yuan, D., Cai, Y., Zhang, M., Lin, Y., Qing, J., An, Z., Revenaugh, J. (2005). A high resolution, absolute-dated Holocene and deglacial Asian monsoon record from Dongge Cave, China. *Earth and Planetary Science Letters*, **233**, 71–86.
- Erdenebat, U., Pohl, E. (2005). Aus der Mitte der Hauptstadt – Die Ausgrabungen der Universität Bonn im Zentrum von Qara Qorum. In: *Dschingis Khan und seine Erben*. Hirmer, München, pp. 168–175.
- Fagan, B. (2008). *The great warming: climate change and the rise and fall of civilizations*. Bloomsbury Press, New York.
- Fang, X. M., Ono, Y., Fukusawa, H., Bao-Tian, P., Li, J. J., Dong-Hong, G., Oi, K., Tsukamoto, S., Torii, M., Mishima, T. (1999). Asian summer monsoon instability during the past 60,000 years: magnetic susceptibility and pedogenic evidence from the western Chinese Loess Plateau. *Earth and Planetary Science Letters*, **168**, 219–232.
- Fedotov, A. P., Chebykin, E. P., Yu, S. M., Vorobyova, S. S., Yu, O. E., Golobokova, L. P., Pogodaeva, T. V., Zheleznyakova, T. O., Grachev, M. A., Tomurhuu, D. (2004). Changes in the volume and salinity of Lake Khubsugul (Mongolia) in response to global climate changes in the upper Pleistocene and the Holocene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **209**, 1–4, 245–257.
- Fernandez-Gimenez, M., Allen-Diaz, B. (2001). Vegetation change along gradients from water sources in three grazed Mongolian ecosystems. *Plant Ecology*, **157**, 101–118.
- Fischbeck, R., Müller, G. (1971). Monohydrocalcite, hydromagnesite, nesquehonite, dolomite, aragonite and calcite in speleothems of the Fränkische Schweiz, Western Germany. *Contributions to Mineralogy and Petrology*, **33**, 87–92.
- Foley, J. A., Kutzbach, J. E., Coe, M. T., Levis, S. (1994). Feedbacks between climate and boreal forests during the Holocene epoch. *Nature*, **371**, 52–53.
- Fowell, S. J., Hansen, B. C., Peck, J. A., Khosbayar, P., Ganbold, E. (2003). Mid to late Holocene climate evolution of the Lake Telmen basin, north central Mongolia, based on palynological data. *Quaternary Research*, **59**, 3, 353–363.
- Franken, C. (2005). Die Brennöfen im Palastbezirk von Karakorum. In: *Dschingis Khan und seine Erben*. Hirmer, München, pp. 147–149.
- Frechen, M. (1999). Luminescence dating of loessic sediments from the Loess Plateau, China. *Geologische Rundschau*, **87**, 675–684.
- Frechen, M., Dodonov, A. E. (1998). Loess chronology of the Middle and Upper Pleistocene in Tadjikistan. *Geologische Rundschau*, **87**, 2–20.
- Frechen, M., Oches, E. A., Kohfeld, K. E. (2003). Loess in Europe – mass accumulation rates during the Last Glacial Period. *Quaternary Science Reviews*, **22**, 1835–1857.
- Frechen, M., Schweitzer, U., Zander, A. (1996). Improvements in sample preparation for the fine grain technique. *Ancient TL*, **14**, 15–17.
- Fu, C. (2003). Potential impacts of human-induced land cover change on East Asia monsoon. *Global and Planetary Change*, **37**, 219–229.
- Fu, C., de Vries, F. P. (2006). *Initial science plan of the Monsoon Asia Integrated Study, MAIRS working paper series*, vol. 1. International Program Office, MAIRS.

Bibliography

- Fuhlendorf, S. D., Smeins, F. E. (1996). Spatial scale influence on longterm temperoral patterns of a semi-arid grassland. *Landscape Ecology*, **11**, 107–113.
- Gao, T., Su, L., Ma, Q., Li, H., Li, X., Yu, X. (2003). Climatic analyses on increasing dust storm frequency in the springs of 2000 and 2001 in Inner Mongolia. *International Journal of Climatology*, **23**, 1743–1755.
- Geerts, B. (2003). Empirical estimation of the monthly-mean daily temperature range. *Theoretical and Applied Climatology*, **74**, 145–165.
- Gelegdorj, E., Chunag, A., Gordon, R. B., Park, J.-S. (2007). Transitions in cast iron technology of the nomads in Mongolia. *Journal of Archaeological Science*, **34**, 1187–1196.
- Goroncy, F. (2006). Räumliche und zeitliche Analyse von Klimaelementen in Zentralasien für die Erklärung der Genese von Zud-Ereignissen in der Mongolei. Diploma thesis at the Freie Universität Berlin.
- Goudie, A. S., Middleton, N. J. (1992). The Changing Frequency of Dust Storms Through Time. *Climatic Change*, **20**, 197–225.
- Grunert, J., Dasch, D. (2004). Dynamics and evolution of dune fields on the northern rim of the Gobi Desert (Mongolia). *Zeitschrift für Geomorphologie/Supplement*, **133**, 81–106.
- Grunert, J., Klein, M. (1998). Binnendünen im nördlichen Zentralasien (Uvs Nuur, westliche Mongolei). In: Böse, M., Hofmann, J. (eds.), *Forschungsergebnisse zur Klimageschichte und Reliefentwicklung Nordafrikas und Asiens, Berliner Geographische Abhandlungen*, vol. 63. Selbstverlag des Geomorphologischen Labors der Freien Universität Berlin, Berlin, pp. 45–66.
- Grunert, J., Klein, M., Stumböck, M., Dasch, M. (1999). Bodenentwicklung auf Altdünen im Uvs Nuur Becken. *Die Erde*, **130**, 97–116.
- Grunert, J., Lehmkuhl, F. (2004). Aeolian sedimentation in arid and semi-arid environments of Western Mongolia. In: Smykatz-Kloss, W., Felix-Henningsen, P. (eds.), *Paleoecology of Quaternary Drylands, Lecture Notes in Earth Sciences*, vol. 102. Springer, Berlin, New York, Toronto, pp. 195–218.
- Grunert, J., Lehmkuhl, F., Walther, M. (2000). Paleoclimatic evolution of the Uvs Nuur basin and adjacent areas (Western Mongolia). *Quaternary International*, **65-66**, 171–192.
- Gunin, P. D., Vostokova, E. A., Dorofeyuk, N. I., Tarasov, P. E., Black, C. C. (1999). *Vegetation dynamics of Mongolia, Geobotany*, vol. 26. Kluwer, Dordrecht, Boston, London.
- Guo, Z. T., Ruddiman, W. F., Hao, Q. Z., Wu, H. B., Qiao, Y. S., Zhu, R. X., Peng, S. Z., Wei, J. J., Y., Y. B., Liu, T. S. (2002). Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China. *Nature*, **416**, 159–163.
- Haase, G. (1983). Beiträge zur Bodengeographie der Mongolischen Volksrepublik. In: Barthel, H., Brunner, H., Haase, G. (eds.), *Physisch-Geographische Studien in Asien, Studia Geographica*, vol. 34. BRNO, pp. 231–367.
- Harmon, R. S. (1983). The mineralogy of Castleguard Cave, Columbia Icefields, Alberta, Canada. *Arctic and Alpine Research*, **15**, 503–516.
- Harrison, S. P., Yu, G., Tarasov, P. E. (1996). Late Quaternary lake-level record from northern Eurasia. *Quaternary Research*, **45**, 138–159.
- Hartmann, K., Wünnemann, B. (2007). Hydrological changes and Holocene climate variations in NW China, inferred from lake sediments of Juyanze palaeolake by factor analyses. *Quaternary International*, **in press**.

Bibliography

- Hauck, C., Vonder Mühl, D. (2003). Evaluation of geophysical techniques for application in mountain permafrost studies. *Zeitschrift für Geomorphologie, Suppl.-Bd.* **132**, 161–190.
- Heikinen, K. (1990). Seasonal changes in iron transport and nature of dissolved organic matter in a humic river in Northern Finland. *Earth Surface Processes and Landforms*, **15**, 583–596.
- Herzschuh, U. (2006). Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years. *Quaternary Science Reviews*, **25**, 1-2, 163–178.
- Herzschuh, U., Tarasov, P., Wünnemann, B., Hartmann, K. (2004). Holocene vegetation and climate of the Alashan Plateau, NW China, reconstructed from pollen data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **211**, 1–17.
- Herzschuh, U., Winter, K., Wünnemann, B., Li, S. (2006). A general cooling trend on the central Tibetan Plateau throughout the Holocene recorded by the Lake Zigetang pollen spectra. *Quaternary International*, **154-155**, 113–121.
- Hilbig, W. (1995). *The vegetation of Mongolia*. SPB Academic Publishing.
- Hilbig, W. (2000). Forest distribution and retreat in the forest steppe ecotone of Mongolia. *Marburger Geographische Schriften*, **135**, 171–187.
- Hill, J., Schütt, B. (2000). The use of remote sensing satellites for mapping complex patterns of erosion and stability in arid mediterranean ecosystems. *Remote Sensing of Environment*, **74**, 557–569.
- Håkanson, L., Jansson, M. (1983). *Principles of lake sedimentology*. Springer, Berlin, Heidelberg, New York.
- Huntley, D. J., Lamothe, M. (2001). Ubiquity of anomalous fading in K-feldspars and the measurement and correction for it in optical dating. *Canadian Journal of Earth Science*, **38**, 1093–1106.
- Hövsgöl Drilling Project Group (2007). Structure of bottom sediments in Lake Hövsgöl: geological and climate controls. *Russian Geology and Geophysics*, **48**, 863–885.
- Ilyashuk, B. P., Ilyashuk, E. A. (2007). Chironomid record of Late Quaternary climatic and environmental changes from two sites in Central Asia (Tuva Republic, Russia) - local, regional or global causes? *Quaternary Science Reviews*, **26**, 705–731.
- Jacoby, G. C., D'Arrigo, R. D., Davaaajamts, T. (1996). Mongolian tree rings and 20th-century warming. *Science*, **273**, 771–773.
- Jadambaa, N., Grimmelmann, W., Kampe, A. (2003). Hydrogeological map of Mongolia 1:1,000,000. Explanatory Notes. *Geologisches Jahrbuch*, **C69**, 3–50.
- Jiang, W., Guo, Z., Sun, X., Wu, H., Chu, G., Yuan, B., Hatte, C., Guiot, J. (2006). Reconstruction of climate and vegetation changes of Lake Bayanchagan (Inner Mongolia): Holocene variability of the East Asian monsoon. *Quaternary Research*, **65**, 411–420.
- Jäkel, D. (2004). Beobachtungen zur Dynamik und zu den Ursachen von Sand- und Staubstürmen in ariden Räumen, angestellt in Nordafrika und China. *Die Erde*, **3-4**, 341–367.
- Jäkel, D., Grunert, J. (2003). Die Wüsten Zentralasiens. *Petermanns Geographische Mitteilungen*, **147**, 5, 26–33.
- Jobbágy, E. G., Jackson, R. B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications*, **10**, 423–436.

Bibliography

- Karabanov, E. B., Prokopenko, A. A., Williams, D. F., Khursevich, G. K. (2000). A new record of Holocene climate change from the bottom sediments of Lake Baikal. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **156**, 3-4, 211–224.
- Karabanov, E. B., Williams, D., Kuzmin, M., Sideleva, V., Khursevich, G., Prokopenko, A., Solotchina, E., Tkachenko, L., Fedenya, S., Kerber, E. (2004). Ecological collapse of Lake Baikal and Lake Hovsgol ecosystems during the Last Glacial and consequences for aquatic species diversity. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **209**, 1-4, 227–243.
- Klinge, M. (2001). Glazialgeomorphologische Untersuchungen im Mongolischen Altai als Beitrag zur jungquartären Landschafts- und Klimageschichte der Westmongolei. *Aachener Geographische Arbeiten*, vol. 35. Geographisches Institut der RWTH Aachen im Selbstverlag, Aachen.
- Kneisel, C. (2003). Electrical resistivity tomography as a tool for geomorphological investigations – some case studies. *Zeitschrift für Geomorphologie, Suppl.-Bd.* **132**, 37–49.
- Kneisel, C. (2004). New insights into mountain permafrost occurrence and characteristics in glacier forefields at high altitude through the application of 2D resistivity imaging. *Permafrost and Periglacial Processes*, **15**, 221–227.
- Kneisel, C. (2005). Detecting mountain permafrost and hazard assessment for debris flows through the application of electrical resistivity tomography - an example from the Swiss Alps. *Zeitschrift für Geomorphologie, Suppl.-Bd.* **138**, 1–10.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F. (2006). World map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, **15**, 259–263.
- Kovalenko, D. V., Yarmolyuk, V. V., Solov'ev, A. V. (1997). Migration of volcanic centers of the south Khangai hot spot: paleomagnetic evidence. *Geotectonics*, **31**, 228–235.
- Kowalkowski, A. (2001). Hauptzüge der quartären Entwicklung der Bodenlandschaften im mongolischen Hochgebirge. In: Bussemer, S. (ed.), *Das Erbe der Eiszeit - Festschrift zum 70. Geburtstag von Joachim Marcinek*. Beier und Beran, Langenweißbach, pp. 17–32.
- Krauskopf, K. B. (1967). *Introduction to Geochemistry*. McGraw-Hill, New York.
- Krautblatter, M., Hauck, C. (2007). Electrical resistivity tomography monitoring of permafrost in solid rock walls. *Journal of Geophysical Research*, **112**, F02S20.
- Küster, Y., Hetzel, R., Krbetschek, M., Tao, M. (2006). Holocene loess sedimentation along the Qilian Shan (China): significance for understanding the processes and timing of loess deposition. *Quaternary Science Reviews*, **25**, 1-2, 114–125.
- Kucera, M., Malmgren, B. A. (1998). Logratio transformation of compositional data – a resolution of the constant sum constraint. *Marine Micropaleontology*, **34**, 117–120.
- Kuhle, M. (2004). The High Glacial (Last Ice Age and LGM) ice cover in High and Central Asia. In: Ehlers, J., Gibbard, P. L. (eds.), *Quaternary glaciations extent and chronology. Part III: South America, Asia, Africa, Australasia, Antarctica, Developments in Quaternary Science*, vol. 2. Elsevier, Amsterdam, Boston, Heidelberg, pp. 175–200.
- Kutzbach, J., Guetter, P. (1986). The influence of changing orbital parameters and surface boundary conditions on climate simulations for the past 180,000 years. *Journal of Atmospheric Sciences*, **43**, 1726–1759.
- Lal, R. (2004). Carbon sequestration in dryland ecosystems. *Environmental Management*, **33**, 528–544.

Bibliography

- Langbein, W. B. (1961). Salinity and hydrology of closed lakes. *U.S. Geology Survey Professional Paper*, **421**, 1–20.
- Lehmkuhl, F. (1997). The spatial distribution of loess and loess-like sediments in the mountain areas of Central and High Asia. *Zeitschrift für Geomorphologie N.F.*, **Suppl.-Bd. 111**, 97–116.
- Lehmkuhl, F., Haselein, F. (2000). Quaternary palaeoenvironmental change on the Tibetan Plateau and adjacent areas (Western China and Mongolia). *Quaternary International*, **65/66**, 121–145.
- Lehmkuhl, F., Klinge, M., Stauch, G. (2004). The extent of Late Pleistocene glaciations in the Altai and Khangai Mountains. In: Ehlers, J., Gibbard, P. L. (eds.), *Quaternary glaciations extent and chronology. Part III: South America, Asia, Africa, Australasia, Antarctica, Developments in Quaternary Science*, vol. 2. Elsevier, Amsterdam, Boston, Heidelberg, pp. 243–254.
- Lehmkuhl, F., Lang, A. (2001). Geomorphological investigations and luminescence dating in the southern part of the Khangai and the Valley of the Gobi Lakes (Central Mongolia). *Journal of Quaternary Science*, **16**, 1, 69–87.
- Lehmkuhl, F., Owen, L. A. (2005). Late Quaternary glaciation of Tibet and the bordering mountains: a review. *Boreas*, **34**, 87–100.
- Lehmkuhl, F., Zander, A., Frechen, M. (2007). Luminescence chronology of fluvial and aeolian deposits in the Russian Altai (Southern Siberia). *Quaternary Geochronology*, **2**, 195–201.
- Lerman, A. (1979). *Geochemical processes, water and sediment environment*. John Wiley and Sons.
- Li, F.-R., Zhao, L.-Y., Zhang, H., Zhang, T.-H., Shirato, Y. (2004). Wind erosion and airborne dust deposition in farmland during spring in the Horqin Sandy Land of eastern Inner Mongolia, China. *Soil and Tillage Research*, **75**, 2, 121 – 130.
- Li, J., Dong, Z., Wang, X., Qian, G., Luo, W. (2008). Seasonal variations in Dustfall and its iron content over North China. *Advances in Atmospheric Sciences*, **25**, 467–473.
- Lippmann, F. (1973). *Sedimentary carbonate minerals, Minerals, rocks and inorganic materials. Monograph series of theoretical and experimental studies*, vol. 6. Springer, Berlin, Heidelberg, New York.
- Liu, J., Diamond, J. (2005). China's environment in a globalizing world. *Nature*, **453**, 1179–1186.
- Liu, Q., Banerjee, S. K., Jackson, M. J., Deng, C., Pan, Y., Zhu, R. (2005). Inter-profile correlation of the Chinese loess/paleosol sequences during Marine Oxygen Isotope Stage 5 and indications of pedogenesis. *Quaternary Science Reviews*, **24**, 1-2, 195–210.
- Liu, T., Ding, Z. (1998). Chinese loess and the paleomonsoon. *Annual Review of Earth and Planetary Sciences*, **26**, 111–145.
- Löwner, M.-O., Preston, N. J., Dikau, R. (2005). Reconstruction of a colluvial body using geoelectrical resistivity. *Zeitschrift für Geomorphologie*, **49**, 225–238.
- Mackereth, F. J. H. (1966). Some chemical observations on post-glacial lake sediments. *Phil. Trans. R. Soc. B.*, **250**, 165–213.
- Madsen, D. B., Jingzen, L., Elston, R. G., Cheng, X., Bettinger, R. L., Kan, G., Brantingham, P. J., Kan, Z. (1998). The loess/paleosol record and the nature of the younger dryas climate in Central China. *Geoarchaeology: An International Journal*, **13**, 8, 847–869.
- Maher, B. A. (2008). Holocene variability of the East Asian summer monsoon from Chinese cave records: a re-assessment. *The Holocene*, **18**, 861–866.

Bibliography

- Maher, B. A., Hu, M. (2006). A high-resolution record of Holocene rainfall variations from the western Chinese Loess Plateau: antiphase behaviour of the African/Indian and East Asian summer monsoons. *The Holocene*, **16**, 309–319.
- Maher, B. A., Thomson, R. (1991). Mineral magnetic record of the Chinese loess and paleosols. *Geology*, **19**, 3–6.
- Mainguet, M., Da Silva, G. G. (1998). Desertification and drylands development: what can be done? *Land Degradation and Development*, **9**, 375–382.
- Maki, A., Kenji, T., Kiyokazu, K., Teruo, H. (2007). Morphological and physico-chemical characteristics of soils in a steppe region of the Kherlen River basin, Mongolia. *Journal of Hydrology*, **333**, 100–108.
- Marschner, H. (1969). Hydrocalcite ($\text{CaCO}_3 \cdot \text{H}_2\text{O}$) and nesquehonite ($\text{MgCO}_3 \cdot 3\text{H}_2\text{O}$) in carbonate scales. *Science*, **165**, 1119–1121.
- Mathess, A., Pekdeger, A. (1980). Chemisch-biochemische Umsetzungen bei der Grundwasserneubildung. *Gas- und Wasserfach*, **121**, 214–219.
- Meyers, P. A. (1994). Preservation of elemental and isotopic source identification of sedimentary organic matter. *Chemical Geology*, **114**, 289–302.
- Meyers, P. A., Lallier-Verges, E. (1999). Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates. *Journal of Paleolimnology*, **21**, 345–372.
- Middleton, N. (1991). Dust storms in the Mongolian People's Republic. *Journal of Arid Environments*, **20**, 287–297.
- Miehe, G., Schlütz, F., Miehe, S., Opgenoorth, L., Cermak, J., Samiya, R., Jäger, E. J., Wesche, K. (2007). Mountain forest islands and Holocene environmental changes in Central Asia: A case study from the southern Gobi Altay, Mongolia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **250**, 150–166.
- Minorsky, V. (1948). Tamim ibn Bahr's journey to the Uyghurs. *Bulletin of the School of Oriental and African Studies, University of London*, **12**, 2, 275–305.
- Mischke, S., Demske, D., Schudack, M. E. (2003). Hydrologic and climatic implications of a multi-disciplinary study of the mid to late Holocene Lake Easter Juyanze. *Chinese Science Bulletin*, **48**, 1411–1417.
- Molnar, P., England, P., Martinod, J. (1993). Mantle dynamics, uplift of the Tibetan Plateau, and the Indian monsoon. *Reviews of Geophysics*, **31**, 357–396.
- Moore, T. R. (1973). The distribution of iron, manganese and aluminium in some soils from north-east Scotland. *Journal of Soil Sciences*, **24**, 162–171.
- Morley, D. W., Leng, M. J., Mackay, A. W., Sloane, H. J. (2005). Late glacial and Holocene environmental change in the Lake Baikal region documented by oxygen isotopes from diatom silica. *Global and Planetary Change*, **46**, 221–233.
- Mussett, A. E., Khan, M. A. (2000). *Looking into the earth. An introduction to geological geophysics*. Cambridge University Press, Cambridge.
- Nandintsetseg, B., Green, J. S., Goulden, C. E. (2007). Trends in extreme daily precipitation and temperature near Lake Hövsgöl, Mongolia. *International Journal of Climatology*, **27**, 341–347.
- Natsagdorj, L., Jugder, D., Chung, Y. S. (2003). Analysis of dust storms observed in Mongolia during 1937–1999. *Atmospheric Environment*, **37**, 1401–1411.

Bibliography

- Naumann, S. (1999). Spät- und postglaziale Landschaftsentwicklung im Bajan Nuur Seebecken (Nordwestmongolei). *Die Erde*, **130**, 117–130.
- Opp, C., Hilbig, W. (2003). Verbreitungsregeln von Böden und Pflanzengesellschaften im nördlichen Zentralasien unter besonderer Berücksichtigung des Uvs-Nuur-Beckens. *Petermanns Geographische Mitteilungen*, **147**, 5, 16–23.
- Overpeck, J., Anderson, D., Trumbore, S., Prell, W. (1996). The southwest Indian Monsoon over the last 18,000 years. *Climate Dynamics*, **12**, 213–225.
- Overpeck, J. T. (1995). Paleoclimatology and climate system dynamics. *Reviews of Geophysics*, **33**, S1, 863–872.
- Owen, L. A., Richards, B., Rhodes, E. J., Cunningham, W. D., Windley, B. F., Badamgarav, J., Dorjnamjaa, D. (1998). Relic permafrost structures in the Gobi of Mongolia: age and significance. *Journal of Quaternary Science*, **13**, 539–547.
- Owen, L. A., Windley, B. F., Cunningham, W. D., Badamgarav, J., Dorjnamjaa, D. (1997). Quaternary alluvial fans in the Gobi of southern Mongolia: evidence for neotectonics and climate change. *Journal of Quaternary Science*, **12**, 239–252.
- Pachur, H.-J., Wünnemann, B., Zhang, H. (1995). Lake Evolution in the Tengger Desert, Northwestern China, during the Last 40,000 Years. *Quaternary Research*, **44**, 2, 171–180.
- Peck, J. A., Khosbayar, P., Fowell, S. J., Pearce, R. B., Ariunbileg, S., Hansen, B. C., Soninkhishig, N. (2002). Mid to Late Holocene climate change in north central Mongolia as recorded in the sediments of Lake Telmen. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **183**, 1-2, 135–153.
- Pederson, N., Jacoby, G. C., D'Arrigo, R. D., Cook, E. R., Buckley, B. M. (2001). Hydrometeorological reconstructions for Northeastern Mongolia derived from tree rings: 1651–1995. *Journal of Climate*, **14**, 872–881.
- Pekala, K., Repelewska-Pekalowa, J. (1993). Solifluction and related slope processes in the mountains of Central Asia (Mongolia). *Palaeoclimate Research, Special Issue: ESF Project 'European Palaeoclimate and Man'*, **6**, 11, 87–101.
- Pessenda, L. C. R., Gouveia, S. E. M., Aravena, R. (2001). Radiocarbon dating of total soil organic matter and humin fraction and its comparison with ^{14}C ages of fossil charcoal. *Radiocarbon*, **43**, 2B, 595–601.
- Peters, D. P. C., Pielke, R. A. S., Bestelmeyer, B. T., Allen, C. D., Munson-McGee, S., Havstad, K. M. (2004). Cross-scale interactions, nonlinearities, and forecasting catastrophic events. *Proceedings of the National Academy of Sciences of the USA*, **101**, 15130–15135.
- Phillips, J. D. (2003). Sources of nonlinearity and complexity in geomorphic systems. *Progress in Physical Geography*, **27**, 1–23.
- Pickup, G., Chewings, V. H. (1988). Forecasting patterns of erosion in arid land forms from Landsat MSS data. *International Journal of Remote Sensing*, **9**, 69–84.
- Piro, S., Mauriello, P., Cammarano, F. (2000). Quantitative integration of geophysical methods for archaeological prospection. *Archaeological Prospection*, **7**, 203–213.
- Poberezhnaya, A. E., Fedotov, A. P., Sitnikova, T. Y., Semenov, M. Y., Ziborova, G. A., Otinova, E. L., Khabuev, A. V. (2006). Paleocological and paleoenvironmental record of the Late Pleistocene record of Lake Khubsugul (Mongolia) based on ostracod remains. *Journal of Paleolimnology*, **36**, 133–149.

Bibliography

- Porter, S. C. (2001). Chinese loess record of monsoon climate during the last glacial-interglacial cycle. *Earth-Science Reviews*, **54**, 115–128.
- Porter, S. C., An, Z. (1995). Correlation between climate events in the North Atlantic and China during the last glaciation. *Nature*, **375**, 305–308.
- Prescott, J. R., Hutton, J. T. (1994). Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. *Radiation measurements*, **23**, 497–500.
- Prokopenko, A. A., Khursevich, G. K., Bezrukova, E. V., Kuzmin, M. I., Boes, X., Williams, D. F., Fedenya, S. A., Kulagina, N. V., Letunova, P. P., Abzaeva, A. A. (2007). Paleoenvironmental proxy records from Lake Hovsgol, Mongolia, and a synthesis of Holocene climate change in the Lake Baikal watershed. *Quaternary Research*, **68**, 2–17.
- Prokopenko, A. A., Kuzmin, M. I., Williams, D. F., Gelety, V. F., Kalmychkov, G. V., Gvozdkov, A. N., Solotchin, P. A. (2005). Basin-wide sedimentation changes and deglacial lake-level rise in the Hovsgol basin, NW Mongolia. *Quaternary International*, **136**, 1, 59–69.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., Gill, T. E. (2002). Environmental characterization of global sources of atmospheric soil dust derived from NIMBUS 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Reviews of Geophysics*, **40**, 1, 2.1–2.31.
- Pye, K. (1995). The nature, origin and accumulation of loess. *Quaternary Science Reviews*, **14**, 653–667.
- Ren, G. (2000). Decline of the mid- to late Holocene forests in China: climatic change or human impact. *Journal of Quaternary Science*, **15**, 273–281.
- Richter, H., Haase, G., Barthel, H. (1963). Die Bildung von Gebirgsfußflächen im Gobi-Altai. In: Harig, G., Steinmets, M. (eds.), *Lehre – Forschung – Praxis. Die Karl-Marx-Universität zum zehnten Jahrestag ihrer Namensgebung am 5. Mai*. pp. 198–213.
- Richthofen, F. v. (1877). *China. Ergebnisse eigener Reisen. 1. Band*. Berlin.
- Roberts, H. M., Wintle, A. G., Maher, B. A., Hu, M. (2001). Holocene sediment-accumulation rates in the western Loess Plateau, China, and a 2500-year record of agricultural activity revealed by OSL dating. *The Holocene*, **11**, 477–483.
- Rost, K. T. (2001). Late Holocene loess deposits and dust accumulation in the alpine meadow belt of the Wutai Shan, China. *Quaternary International*, **76/77**, 85–92.
- Rost, K. T., Böhner, J., Pörtge, K.-H. (2003). Landscape degradation and desertification in the Mu Us Shamo, Inner Mongolia – an ecological and climatic problem since historical times? *Erdkunde*, **57**, 110–125.
- Rösch, M., Fischer, E., Märkle, T. (2005). Human diet and land use in the time of the Khans - Archaeobotanical research in the capital of the Mongolian Empire, Qara Qorum, Mongolia. *Vegetation History and Archaeobotany*, **14**, 485–492.
- Ruddiman, W. (2003). The anthropogenic greenhouse era began thousands of years ago. *Climatic Change*, **61**, 261–293.
- Running, S. W., Loveland, T. R., Merchant, J. W., Ohlen, D. O. (1995). A remote sensing based vegetation classification logic for global land cover analysis. *Remote Sensing of Environment*, **51**, 3948–3952.
- Sack, D. (2001). Shoreline and basin configuration techniques in paleolimnology. In: Last, W. M., Smol, J. P. (eds.), *Tracking environmental change using lake sediments. Volume 2: Physical and geochemical methods*. Kluwer, Dordrecht, Boston, London, pp. 49–71.

Bibliography

- Salzmann, U., Hoelzmann, P. (2005). The Dahomey Gap: an abrupt climatically induced rain forest fragmentation in West Africa during the late Holocene. *The Holocene*, **15**, 190–199.
- Sapozhnikov, D. G., Tsvetkov, A. I. (1959). Precipitation of hydrous calcium carbonate on the bottom of Lake Issyk-Kul. *Dokl. Akad. Nauk SSSR*, **124**, 402–405.
- Sato, T., Kimura, F., Kitoh, A. (2007). Projection of global warming onto regional precipitation over Mongolia using a regional climate model. *Journal of Hydrology*, **333**, 144–154.
- Scheffer, F. (2002). *Lehrbuch der Bodenkunde / Scheffer/Schachtschabel*. 15 ed. Spektrum, Heidelberg, Berlin.
- Scheffer, M. (2004). *Ecology of shallow lakes, Population and community biology series*, vol. 22. Kluwer Academic Publishers, Dordrecht, Boston, London.
- Schlichting, E., Schweikle, V. (1980). Interpedon translocations and soil classification. *Soil Science*, **130**, 200–204.
- Schlütz, F., Dulamsuren, C., Wieckowska, M., Mühlenberg, M., Hauck, M. (2008). Late Holocene vegetation history suggests natural origin of steppes in the northern Mongolian mountain taiga. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **261**, 203–217.
- Schlütz, F., Lehmkuhl, F. (2007). Climatic change in the Russian Altai, southern Siberia, based on palynological and geomorphological results, with implications for climatic teleconnections and human history since the middle Holocene. *Vegetation History and Archaeobotany*, **16**, 101–118.
- Schotterer, U., Fröhlich, K., Gäggeler, H. W., Sandjordj, S., Stichler, W. (1997). Isotope records from Mongolian and alpine ice cores as climate indicators. *Climatic Change*, **36**, 519–530.
- Schütt, B. (1998). Reconstruction of Holocene paleoenvironments in the endorheic basin of Laguna de Gallocanta, Central Spain by investigation of mineralogical and geochemical characters from lacustrine sediments. *Journal of Paleolimnology*, **20**, 217–234.
- Schütt, B. (2004a). The chemistry of playa-lake-sediments as a tool for the reconstruction of Holocene environmental conditions – a case study from the central Ebro basin. In: Smykatz-Kloss, W., Felix-Henningsen, P. (eds.), *Paleoecology of Quaternary Drylands, Lecture Notes in Earth Sciences*, vol. 102. Springer, Berlin, New York, Toronto, pp. 5–30.
- Schütt, B. (2004b). *Zum holozänen Klimawandel der zentralen Iberischen Halbinsel, Relief Boden Paläoklima*, vol. 20. Bornträger, Berlin, Stuttgart.
- Schwanghart, W., Schütt, B. (2008). Holocene morphodynamics in the Ugii Nuur basin, Mongolia – insights from a sediment profile and 1D electrical resistivity tomography. *Zeitschrift für Geomorphologie*, **52**, Suppl. 2, 35–55.
- Schwanghart, W., Schütt, B., Walther, M. (2008). Holocene climate evolution of the Ugii Nuur basin, Mongolia. *Advances in Atmospheric Sciences, MAIRS special issue*, in press.
- Shi, Y., Zhaozheng, K., Sumin, W., Lingyu, T., Wang, F., Tandong, Y., Xitao, Z., Zhang, P., Shi, S. (1993). Mid-Holocene climates and environments in China. *Global and Planetary Change*, **7**, 219–233.
- Shiraishi, N. (2004). Seasonal migrations of the Mongol emperors and the peri-urban area of Kharakhorum. *The International Journal of Asian Studies*, **1**, 105–119.
- Shukla, J., Mintz, Y. (1982). The influence of land-surface evapotranspiration on Earth's climate. *Science*, **215**, 1496–1501.

Bibliography

- Sirocko, F., Sarntheim, M., Erlenkeuser, H., Lange, H., Arnold, M., Duplessy, J. C. (1993). Century-scale events in monsoonal climate over the past 24,000 years. *Nature*, **364**, 322–324.
- Skinner, H. C. W., Osbaldiston, G. W., Wilner, A. N. (1977). Monohydrocalcite in a guinea pig bladder stone, a novel occurrence. *American Mineralogist*, **62**, 273–277.
- Sladen, C., Traynor, J. J. (2000). Lakes during the evolution of Mongolia. In: Gierlowski-Kordesch, E. H., Kelts, K. R. (eds.), *Lake basins through space and time*, vol. 46. American Association of Petroleum Geologists Studies in Geology, pp. 35–57.
- Sommer, M., Treter, U. (1999). Die Lärchenwälder der Gebirgswaldsteppe in den Randgebirgen des Uvs Nuur-Beckens. *Die Erde*, **130**, 173–188.
- Sonnenfeld, P. (1984). *Brines and evaporites*. Academic Press, Orlando.
- Stadelbauer, J. (2003). Mittelasien – Zentralasien: Raumbegriffe zwischen wissenschaftlicher Strukturierung und politischer Konstruktion. *Petermanns Geographische Mitteilungen*, **147**, 5, 58–63.
- Stallins, J. A. (2006). Geomorphology and ecology: unifying theses for complex systems in biogeomorphology. *Geomorphology*, **77**, 207–216.
- Street-Perrot, F. A. (1980). The relative importance of climate and hydrogeological factors in influencing lake level fluctuations. *Paleoecology of Africa*, **12**, 137–158.
- Street-Perrot, F. A., Harrison, S. P. (1985). Lake levels and climate reconstructions. In: Hecht, A. D. (ed.), *Palaeoclimate analysis and modelling*. New York, pp. 291–340.
- Talbot, M. R., Johannessen, T. (1992). A high resolution palaeoclimatic record for the last 27,500 years in tropical West Africa from carbon and nitrogen isotopic composition of lacustrine organic matter. *Earth and Planetary Science Letters*, **110**, 23–37.
- Tarasov, P., Dorofeyuk, N., Metel’Tseva, E. (2000). Holocene vegetation and climate changes in Hoton-Nur basin, northwest Mongolia. *Boreas*, **29**, 117–126.
- Tarasov, P. E., Bezrukova, E., Karabanov, E., Nakagawa, T., Wagner, M., Kulagina, N., Letunova, P., Abzaeva, A., Granoszewski, W., Riedel, F. (2007a). Vegetation and climate dynamics during the Holocene and Eemian interglacials derived from Lake Baikal pollen records. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **252**, 440–457.
- Tarasov, P. E., Guiot, J., Cheddadi, R., Andreev, A. A., Bezusko, L. G., Blyakharchuk, T. A., Dorofeyuk, N. I., Filimonova, L. V., Volkova, V. S., Zernitskaya, V. P. (1999a). Climate in northern Eurasia 6000 years ago reconstructed from pollen data. *Earth and Planetary Science Letters*, **171**, 635–645.
- Tarasov, P. E., Peyron, O., Guiot, J., Brewer, S., Volkova, V. S., Bezusko, L. G., Dorofeyuk, N. I., Kvavadze, E. V., Osipova, I. M., Panova, N. K. (1999b). Last Glacial Maximum climate of the former Soviet Union and Mongolia reconstructed from pollen and plant macrofossil data. *Climate Dynamics*, **15**, 227–240.
- Tarasov, P. E., Pushenko, M. Y., Harrison, S. P., Saarse, L., Andreev, A. A., Aleshinskaya, Z. V., Davydova, N. N., Dorofeyuk, N. I., Efremov, Y. V., Elina, G. A., Elovicheva, Y. K., Filimonova, L. V., Gunova, V. S., Khomutova, V. I., Kvavadze, E. V., Neustrueva, I. Y., Pisareva, V. V., Sevastyanov, D. V., Shelekhova, T. S., Subetto, D. A., Uspenskaya, O. N., Zernitskaya, V. P. (1996). *Lake status records from the Former Soviet Union and Mongolia: Documentation of the second version of the data base*. No. 5 in NOAA Paleoclimatology Publications Series Report. NOAA, Boulder.
- Tarasov, P. E., Wagner, M., Jin, G. (2007b). Paläobotanische Archive: Jungquartäre und holozäne Klimaentwicklung in chinesischen Trockengebieten. *Geographische Rundschau*, **59**, 4, 12–20.

Bibliography

- Tarasov, P. E., Williams, J. W., Andreev, A., Nakagawa, T., Bezrukova, E., Herzschuh, U., Igarashi, Y., Müller, S., Werner, K., Zheng, Z. (2007c). Satellite- and pollen-based quantitative woody cover reconstructions for northern Asia: Verification and application to late-Quaternary pollen data. *Earth and Planetary Science Letters*, **264**, 284–298.
- Taylor, G. F. (1975). The occurrence of monohydrocalcite in two small lakes in the south-east of South Australia. *American Mineralogist*, **60**, 690–697.
- Thompson, L. G., Yao, T., Davis, M. E., Henderson, K. A., Thompson, E. M., Lin, P.-N., Beer, J., Synal, H.-A., Cole-Dai, J., Bolzan, J. F. (1997). Tropical climate instability: the last glacial cycle from a Quighai-Tibetan ice core. *Science*, **276**, 1821–1825.
- Thompson, L. G., Yao, T., Mosley-Thompson, E., Davis, M. E., Henderson, K. A., Lin, P.-N. (2000). A high-resolution millennial record of the South Asian Monsoon from Himalayan ice cores. *Science*, **289**, 1916–1999.
- Traynor, J. J., Sladen, C. (1995). Tectonic and stratigraphic evolution of the Mongolian People's Republic and its influence on hydrocarbon geology and potential. *Marine and Petroleum Geology*, **12**, 35–52.
- Treter, U. (1996). Gebirgs-Waldsteppe in der Mongolei. *Geographische Rundschau*, **48**, 11, 655–661.
- Vallentyne, J. R. (1962). Solubility and the decomposition of organic matter in nature. *Archives of Hydrobiology*, **58**, 423–434.
- Vipper, P. B., Dorofeyuk, N. I., Meteltzeva, E. I., Sokolovskaya, V. P. (1989). Landscape and climate changes in central Mongolia during the Holocene. In: Khotinsky, N. A. (ed.), *Paeoklimaty pozdnelednikovya i golotsena (Paleoclimates of Late Glacial and the Holocene)*. Nauka, Moscow, pp. 160–167. In Russian.
- Völker, J. (2005). Hydrographie und aktueller Gewässerzustand. In: Walther, M., Gegeensuvd, T. (eds.), *Ugi Nuur (Central Mongolia) paleoenvironmental studies of lake level fluctuations and Holocene climate change*, *Geographica-Oekologica*, vol. 2. MOLARE Research Centre, Ulaanbaatar, pp. 2–6.
- von den Driesch, A., Peters, J., Delgermaa, L. (2008). Animal economy in the ancient Mongolian Town of Qara Qorum (Kharkhorin). Preliminary report on the faunal remains. In: Bemann, J., Hüttel, H. G., Pohl, E. (eds.), *Mongolian-German Qara Qorum-Expedition*, FAAK (Forschungen zur Archäologie Außereuropäischer Kulturen). Bonn, p. in press.
- Walker, R. T., Molor, E., Fox, M., Bayasgalan, A. (2008). Active tectonics of an apparently aseismic region: distributed active strike-slip faulting in the Hangay Mountains of central Mongolia. *Geophysical Journal International*, **174**, 1121–1137.
- Wallinga, J., Murray, A. S., Wintle, A. G. (2000). The single-aliquot regenerative-dose (SAR) protocol applied to coarse-grain feldspar. *Radiation measurements*, **32**, 529–533.
- Walther, M. (1998). Paläoklimatische Untersuchungen zur jungpleistozänen Landschaftsentwicklung im Changai-Bergland und in der nördlichen Gobi (Mongolei). *Petermanns Geographische Mitteilungen*, **142**, 3/4, 207–217.
- Walther, M. (1999). Befunde zur jungquartären Klimaentwicklung rekonstruiert am Beispiel der Seespiegelstände des Uvs Nuur-Beckens (NW-Mongolei). *Die Erde*, **130**, 131–150.
- Walther, M. (2005). Ein idealer Ort für ein festes Lager. Zur Geographie des Orchontals und der Umgebung von Charchorin (Karakorum). In: *Dschingis Khan und seine Erben. Das Weltreich der Mongolen*. Hirmer, pp. 128–132.

Bibliography

- Walther, M., Gegeensuvd, T. (2005). *Ugii Nuur (Central Mongolia) paleo environmental studies of lake level fluctuations and Holocene climate change*, *Geographica-Oekologica*, vol. 2. MOLARE Research Centre, Ulaanbaatar.
- Walther, M., Naumann, S. (1997). Beobachtungen zur Fu"ssl"achenbildung im ariden bis semiariden Bereich der West- und S"udmongolei (N"ordliches Zentralasien). *Stuttgarter Geographische Studien*, **126**, 154–171.
- Walther, M., W"unnemann, B., Tschimeksaichan, A. (2003). Seen und Pal"oeseen in der Mongolei und Nordwestchina. *Petermanns Geographische Mitteilungen*, **147**, 5, 40–47.
- Wang, L., Sarnthein, M., Erlenkeuser, H., Grimalt, J., Grootes, P., Heilig, S., Ivanova, E., Kienast, M., Pelejero, C., Pflaumann, U. (1999). East Asian monsoon climate during the Late Pleistocene: high-resolution sediment records from the South China Sea. *Marine Geology*, **156**, 245–284.
- Wang, P. (1999). Response of western pacific marginal seas to glacial cycles: paleoceanographic and sedimentological features. *Marine Geology*, **156**, 5–39.
- Wang, P., Clemens, S., Beaufort, L., Braconnot, P., Ganssen, G., Zhimin, J., Kershaw, P., Sarnthein, M. (2005a). Evolution and variability of the Asian monsoon system: state of the art and outstanding issues. *Quaternary Science Reviews*, **34**, 595–629.
- Wang, X., Xia, D., Wang, T., Xue, X., Li, J. (2008a). Dust sources in arid and semiarid China and southern Mongolia: Impacts of geomorphological setting and surface materials. *Geomorphology*, **97**, 583–600.
- Wang, Y., Amundson, R., Trumbore, S. (1996). Radiocarbon dating of soil organic matter. *Quaternary Research*, **45**, 282–288.
- Wang, Y., Cheng, H., Edwards, L., He, Y., Kong, X., An, Z., Wu, J., Kelly, M. J., Dykoski, C. A., Li, X. (2005b). The Holocene Asian Monsoon: Links to solar changes and North Atlantic climate. *Science*, **308**, 854–857.
- Wang, Y., Cheng, H., Edwards, R. L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X., An, Z. (2008b). Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years. *Nature*, **451**, 1090–1093.
- Wang, Y. J., Cheng, H., Edwards, R. L., An, Z. S., Wu, Y., J, Shen, C.-C., Dorale, J. A. (2001). A high-resolution absolute-dated Late Pleistocene monsoon record from Hulu Cave, China. *Science*, **294**, 2345–2348.
- Weiers, M. (2005). Steppe und Steppenreiche bis Cinggis Khan. In: *Dschingis Khan und seine Erben*. Hirmer, M"unchen, pp. 36–38.
- Weischet, W., Endlicher, W. (2000). *Regionale Klimatologie, Teil 2, Die Alte Welt (Europa, Afrika, Asien)*. Teubner, Stuttgart, Leipzig.
- Weninger, B., J"oris, O. (2007). Towards an absolute chronology at the Middle to Upper Palaeolithic transition in western Eurasia: A new GreenlandHulu time-scale based on U/Th ages. *Journal of Human Evolution*, **in press**.
- Westover, K. S., Fritz, S. C., Blyakharchuk, T. A., Wright, H. E. (2006). Diatom paleolimnological record of Holocene climatic and environmental change in the Altai Mountains, Siberia. *Journal of Paleolimnology*, **35**, 519–541.

Bibliography

- Wünnemann, B., Chen, F., Riedel, F., Zhang, C., Mischke, S., Chen, G., Demske, D., Ming, J. (2003). Holocene lake deposits of Bosten Lake, southern Xinjiang, China. *Chinese Science Bulletin*, **48**, 14, 1429–1432.
- Wünnemann, B., Hartmann, K. (2002). Morphodynamics and Paleohydrography of the Gaxun Nur Basin, Inner Mongolia, China. *Zeitschrift für Geomorphologie/Supplement*, **126**, 147–168.
- Wünnemann, B., Hartmann, K., Altmann, N., Hambach, U., Pachur, H. J. (2005). Interglacial and glacial fingerprints from lake deposits in the Gobi Desert, NW China. In: Sirocko, F., Claussen, M., Litt, T., Goñi, M. F. (eds.), *The climate of the past interglacials. Developments in Quaternary Science*. Elsevier, Amsterdam, p. in press.
- Wünnemann, B., Pachur, H., Li, J. (1998). Chronologie der pleistozänen und holozänen Seespiegelschwankungen des Gaxun Nur / Sogo Nur und Baijian Hu, Innere Mongolei, Nordwestchina. *Petermanns Geographische Mitteilungen*, **142**, 3/4, 191–206.
- Wright, J. (2007). Organizational principles of Khirigsuur monuments in the lower Egiin Gol valley, Mongolia. *Journal of Anthropological Archaeology*, **26**, 350–365.
- Wright, J. S. (2001). Making loess-sized quartz silt: data from laboratory simulations and implications for sediment transport pathways and the formation of desert loess deposits associated with the Sahara. *Quaternary International*, **76/77**, 7–19.
- Wright, V. P., Tucker, M. E. (1991). Calcretes. An introduction. In: Wright, V. P., Tucker, M. E. (eds.), *Calcretes*, Reprint series vol. 2 of the International Association of Sedimentologists. Blackwell, Oxford, London, pp. 1–22.
- Wu, W., Liu, T. (2004). Possible role of the Holocene Event 3 on the collapse of Neolithic cultures around the Central Plain of China. *Quaternary International*, **117**, 153–166.
- Xiao, J., Nakamura, T., Lu, H., Zhang, G. (2002). Holocene climate changes over the desert/loess transition of north-central China. *Earth and Planetary Science Letters*, **197**, 11–18.
- Xiao, J., Wu, J., Si, B., Liang, W., Nakamura, T., Liu, B., Inouchi, Y. (2006). Holocene climate changes in the monsoon/arid transition reflected by carbon concentration in Dahai Lake of Inner Mongolia. *The Holocene*, **16**, 551–560.
- Xiao, J., Xu, Q., Nakamura, T., Yang, X., Liang, W., Inouchi, Y. (2004). Holocene vegetation variation in the Daihai Lake region of north-central China: a direct indication of the Asian monsoon climatic history. *Quaternary Science Reviews*, **23**, 14–15, 1669–1679.
- Xuan, J., Sokolik, I. N., Hao, J., Guo, F., Mao, H., Yang, G. (2004). Identification and characterization of sources of atmospheric mineral dust in East Asia. *Atmospheric Environment*, **38**, 36, 6239–6252.
- Xue, Y., Juang, H.-M. H., Li, W.-P., Prince, S., DeFries, R., Jiao, Y., Vasic, R. (2004). Role of land surface processes in monsoon development - East Asia and West Africa. *Journal of Geophysical Research*, **109**, D3.
- Yair, A., Bryan, R. B. (2000). Hydrological response of desert margins to climatic change: the effect of changing surface properties. In: McLaren, S. J., Kniveton, D. R. (eds.), *Linking climate change to land surface changes*. Kluwer Academic Publishers, London, pp. 49–64.
- Yancheva, G., Nowaczyk, N. R., Mingram, J., Dulski, P., Schettler, G., Negendank, J. F. W., Liu, J., Sigman, D. M., Peterson, L. C., Haug, G. H. (2007). Influence of the intertropical convergence zone on the East Asia Monsoon. *Nature*, **445**, 74–77.

Bibliography

- Yang, S., K.-M., L. (1998). Influences of sea surface temperature and ground wetness on Asian summer monsoon. *Journal of Climate*, **11**, 3230–3246.
- Yang, X., Rost, K. T., Lehmkuhl, F., Zhenda, Z., Dodson, J. (2004). The evolution of dry lands in northern China and in the Republic of Mongolia since the Last Glacial Maximum. *Quaternary International*, **118-119**, 69–85.
- Yao, T., Thompson, L. G., Shi, Y., Qin, D., Jiao, K., Yang, Z., Tian, L., Thompson, E. M. (1997). Climate variation since the last interglaciation recorded in the Guliya ice core. *Science in China Series D (Earth Sciences)*, **40**, 662–668.
- Yuan, D., Cheng, H., Edwards, R. L., Dykoski, C. A., Kelly, M. J., Zhang, M., Qing, J., Lin, Y., Wang, Y., Wu, J., Dorale, J. A., An, Z., Cai, Y. (2004). Timing, duration, and transitions of the last interglacial Asian Monsoon. *Science*, **304**, 575–578.
- Zech, W., Bäumlner, R., Sovoskul, O., Sauer, G. (1996). Zur Problematik der pleistozänen und holozänen Vergletscherung des westlichen Tienshan. *Eiszeitalter und Gegenwart*, **46**, 144–151.
- Zender, C. S., Bian, H., Newman, D. (2003). Mineral dust entrainment and deposition (DEAD) model: Description and 1990s dust climatology. *Journal of Geophysical Research*, **108**, AAC 8.1–19.
- Zhang, X. Y., Arimoto, R., Zhu, G. H., Chen, T., Zhang, G. Y. (1998). Concentration, size-distribution and deposition of mineral aerosol over Chinese desert regions. *Tellus B*, **50**, 317–330.
- Zhang, Y., Ohata, T., Yang, D., Davaa, G. (2004). Bias correction of daily precipitation measurements for Mongolia. *Hydrological Processes*, **18**, 2991–3005.
- Zhang, Y., X., Arimoto, R., An, Z. S. (1997). Dust emission from Chinese desert sources linked to variations in atmospheric circulation. *Journal of Geophysical Research*, **102**, D23, 28041–28047.
- Zhao, T. L., Gong, S. L., Zhang, X. Y., McKendry, I. G. (2003). Modeled size-segregated wet and dry deposition budgets of soil dust aerosol during ACE-Asia 2001: Implications for trans-Pacific transport. *Journal of Geophysical Research*, **108**, 108(D23), 8665.
- Zhu, R., Liu, Q., Jackson, M. J. (2004). Paleoenvironmental significance of the magnetic fabrics in Chinese loess-paleosols since the last interglacial (< 130 ka). *Earth and Planetary Science Letters*, **221**, 55–69.