

## 7 Literaturverzeichnis

- Aberle, H., Bauer, A., Stappert, J., Kispert, A. und Kemler, R. (1997).  $\beta$ -catenin is a target of the ubiquitin-proteasome pathway. *EMBO J* 16, 3797-3804.
- Ahmed, K., Gerber, D. A. und Cochet, C. (2002). Joining the cell survival squad: an emerging role for protein kinase CK2. *Trends Cell Biol* 12, 226-230.
- Ahmed, K. und Tawfic, S. (1994). Mechanism of intracellular regulation of protein kinase CK2: role of stimulus-mediated subnuclear association. *Cell Mol Biol Res* 40, 539-545.
- Amit, S., Hatzubai, A., Birman, Y., Andersen, J. S., Ben-Shushan, E., Mann, M., Ben-Neriah, Y. und Alkalay, I. (2002). Axin-mediated CK1 phosphorylation of  $\beta$ -catenin at Ser 45: a molecular switch for the Wnt-pathway. *Genes Dev* 16, 1066-1076.
- Arnold, S. J., Stappert, J., Bauer, A., Kispert, A., Hermann, B. G. und Kemler, R. (2000). Brachyury is a target gene of the Wnt/ $\beta$ -catenin signaling pathway. *Mech Dev* 91, 249-258.
- Barth, A. I., Nathke, I. S. und Nelson, W. J. (1997). Cadherins, catenins and APC protein: interplay between cytoskeletal complexes and signaling pathways. *Curr Opin Cell Biol* 9, 683-690.
- Bauer, A., Chauvet, S., Huber, O., Usseglio, F., Rothbächer, U., Aragnol, D., Kemler, R. und Pradel, J. (2000). Pontin52 and Reptin52 function as antagonistic regulators of  $\beta$ -catenin signalling activity. *EMBO J* 19, 6121-6130.
- Bauer, A., Huber, O. und Kemler, R. (1998). Pontin52, an interaction partner of  $\beta$ -catenin, binds to the TATA box binding protein. *Proc Natl Acad Sci USA* 95, 14787-14792.
- Behrens, J., von Kries, J. P., Kühl, M., Bruhn, L., Wedlich, D., Grosschedl, R. und Birchmeier, W. (1996). Functional interaction of  $\beta$ -catenin with the transcription factor LEF-1. *Nature* 382, 638-642.
- Bek, S. und Kemler, R. (2002). Protein kinase CKII regulates the interaction of  $\beta$ -catenin with  $\alpha$ -catenin and its protein stability. *J Cell Sci* 115, 4743-4753.
- Bienz, M. (2001). Spindles cotton on to junctions, APC and EB1. *Nat Cell Biol* 3, E67-E68.
- Bienz, M. und Clevers, H. (2000). Linking colorectal cancer to Wnt signaling. *Cell* 103, 311-320.
- Birnboim, H. C. und Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 7, 1513-1523.
- Boutros, M., Paricio, N., Strutt, D. I. und Mlodzik, M. (1998). Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and wingless signaling. *Cell* 94, 109-118.
- Brannon, M., Brown, J. D., Bates, R., Kimelman, D. und Moon, R. T. (1999). XCtBP is a XTcf-3 co-repressor with roles throughout Xenopus development. *Development* 126, 3159-3170.
- Brannon, M., Gomperts, M., Sumoy, L., Moon, R. T. und Kimelman, D. (1997). A  $\beta$ -catenin/XTcf-3 complex binds to the siamois promotor to regulate dorsal axis specification in *Xenopus*. *Genes Dev* 11, 2359-2370.

- Brantjes, H., Roose, J., van de Wetering, M. und Clevers, H. (2001). All Tcf HMG box transcription factors interact with Groucho-related co-repressors. *Nucleic Acids Res* 29, 1410-1419.
- Bruhn, L., Munnerlyn, A. und Grosschedl, R. (1997). ALY, a context-dependent coactivator of LEF-1 and AML-1, is required for TCR $\alpha$  enhancer function. *Genes Dev* 11, 640-653.
- Brunner, E., Peter, O., Schweizer, L. und Basler, K. (1997). Pangolin encodes a Lef-1 homologue that acts downstream of Armadillo to transduce the Wingless signal in Drosophila. *Nature* 385, 829-833.
- Carlsson, P., Waterman, M. und Jones, K. A. (1993). The hLEF/TCF-1  $\alpha$  HMG protein contains a context-dependent transcriptional activation domain that induces the TCR  $\alpha$  enhancer in T cells. *Genes Dev* 7, 2418-2430.
- Casnellie, J. F. (1991). Assay of protein kinases using peptides with basic residues for phosphocellulose binding. *Methods Enzymol* 200, 115-120.
- Cavallo, R. A., Cox, R. T., Moline, M. M., Roose, J., Polevoy, G. A., Clevers, H., Peifer, M. und Bejsovec, A. (1998). Drosophila Tcf and Groucho interact to repress Wingless signalling activity. *Nature* 395, 604-608.
- Chen, G., Fernandez, J., Mische, S. und Courey, A. J. (1999). A functional interaction between the histone deacetylase Rpd3 and the corepressor groucho in Drosophila development. *Genes Dev* 13, 2218-2230.
- Chijiwa, T., Hagiwara, M. und Hidaka, H. (1989). A newly synthesized Casein kinase I inhibitor, N-(2-Aminoethyl)-5-chloroisoquinoline-8-sulfonamide, and affinity purification of casein kinase I from bovine testis. *J Biol Chem* 264, 4924-4927.
- Critchfield, J., Coligan, J., Folks, T. und Butera, S. (1997). Casein kinase II is a selective target of HIV-1 transcriptional inhibitors. *Proc Natl Acad Sci USA* 94, 6110-6115.
- Daniels, D. L. und Weis, W. I. (2002). ICAT inhibits  $\beta$ -Catenin binding to Tcf/Lef-family transcription factors and the general coactivator p300 using independent structural modules. *Mol Cell* 10, 573-584.
- Djiane, A., Riou, J. F., Umbhauer, M., Boucaut, J. C. und Shi, D. L. (2000). Role of frizzled-7 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* 127, 3091-3100.
- Eastman, Q. und Grosschedl, R. (1999). Regulation of LEF-1/TCF transcription factors by Wnt and other signals. *Curr Opin Cell Biol* 11, 233-240.
- Easwaran, V., Pishvaian, M., Salimuddin und Byers, S. (1999). Cross-regulation of  $\beta$ -catenin-LEF/TCF and retinoid signaling pathways. *Curr Biol* 9, 1415-1418.
- Eide, E. J. und Virshup, D. M. (2001). Casein kinase I: another cog in the circadian clockworks. *Chronobiol Int* 18, 389-398.
- Fagotto, F., Jho, E., Zeng, L., Kurth, T., Joos, T., Kaufmann, C. und Costantini, F. (1999). Domains of axin involved in protein-protein interactions, Wnt pathway inhibition, and intracellular localization. *J Cell Biol* 145, 741-756.

- Fiol, C. J., Mahrenholz, A. M., Wang, Y., Roeske, R. W. und Roach, P. J. (1987). Formation of protein kinase recognition sites by covalent modification of the substrate. Molecular mechanism for the synergistic action of casein kinase II and glycogen synthase kinase 3. *J Biol Chem* 262, 14042-14048.
- Fish, K. J., Cegielska, A., Getman, M. E., Landes, G. M. und Virshup, D. M. (1995). Isolation and characterization of human casein kinase I $\varepsilon$  (CKI), a novel member of the CKI gene family. *J Biol Chem* 270, 14875-14883.
- Gao, Z.-H., Seeling, J. M., Hill, V., Yochum, A. und Virshup, D. M. (2002). Casein Kinase I phosphorylates and destabilizes the  $\beta$ -catenin degradation complex. *Proc Natl Acad Sci USA* 99, 1182-1187.
- Giese, K., Amsterdam, A. und Grosschedl, R. (1991). DNA-binding properties of the HMG domain of the lymphoid-specific transcriptional regulator LEF-1. *Genes Dev* 5, 2567-2578.
- Giese, K. und Grosschedl, R. (1993). LEF-1 contains an activation domain that stimulates transcription only in a specific context of factor-binding sites. *EMBO J* 12, 4667-4676.
- Giese, K., Kingsley, C., Kirshner, J. R. und Grosschedl, R. (1995). Assembly and function of a TCR $\alpha$  enhancer complex is dependent on LEF-1-induced DNA bending and multiple protein-protein interactions. *Genes Dev* 9, 995-1008.
- Gradl, D., König, A. und Wedlich, D. (2002). Functional diversity of Xenopus lymphoid enhancer factor/T-cell factor transcription factors relies on combinations of activating and repressing elements. *J Biol Chem* 277, 14159-14171.
- Graham, T. A., Clements, W. K., Kimelman, D. und Xu, W. (2002). The crystal structure of the  $\beta$ -catenin/ICAT complex reveals the inhibitory mechanism of ICAT. *Mol Cell* 10, 563-571.
- Graham, T. A., Weaver, C., Mao, F. und Kimelman, D. (2000). Crystal structure of a  $\beta$ -catenin/Tcf complex. *Cell* 103, 885-896.
- Grammer, T., Liu, K., Mariani, F. und Harland, R. (2000). Use of large-scale expression cloning screens in the *Xenopus laevis* tadpole to identify gene function. *Dev Biol* 228, 197-210.
- Gross, S. D. und Anderson, R. A. (1998). Casein kinase I: spatial organization and positioning of a multifunctional protein kinase family. *Cell Signal* 10, 699-711.
- Gross, S. D., Simerly, C., Schatten, G. und Anderson, R. A. (1997). A casein kinase I isoform is required for proper cell cycle progression in fertilized mouse oocyte. *J Cell Sci* 110, 3083-3090.
- Grosschedl, R., Giese, K. und Pagel, J. (1994). HMG domain proteins: architectural elements in the assembly of nucleoprotein structures. *Trends Genet* 10, 94-100.
- Guerra, B., Boldyreff, B., Sarno, S., Cesaro, L., Issinger, O.-G. und Pinna, L. A. (1999). CK2: a protein kinase in need of control. *Pharmacol Ther* 82, 303-313.
- Hamaguchi, M., Matsuyoshi, N., Ohnishi, Y., Gotoh, B., Takeichi, M. und Nagai, Y. (1993). p60v-src causes tyrosine phosphorylation and inactivation of the N-cadherin-catenin cell adhesion system. *EMBO J* 12, 307-314.

- Hannahan, D. (1983). Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 166, 557-580.
- Hart, M. J., Concorde, J. P., Lassot, I., Albert, I., de los Santos, R., Durand, H., Perret, C., Rubinfeld, B., Margottin, F., Benarous, R. und Polakis, P. (1999). The F-box protein  $\beta$ -TrCP associates with phosphorylated  $\beta$ -catenin and regulates its activity in the cell. *Curr Biol* 9, 207-210.
- Hart, M. J., de los Santos, R., Albert, I. N., Rubinfeld, B. und Polakis, P. (1998). Downregulation of  $\beta$ -catenin by human Axin and its association with the APC tumor suppressor,  $\beta$ -catenin and GSK3 $\beta$ . *Curr Biol* 8, 573-581.
- Hathaway, G., Lubben, T. und Traugh, J. (1980). Inhibition of casein kinase II by heparin. *J Biol Chem* 255, 8038-8041.
- He, X., Sparks, A. B., Rago, C., Hermeking, H., Zawel, L., da Costa, L. T., Morin, P. J., Vogelstein, B. und Kinzler, K. W. (1998). Identification of c-MYC as a target of the APC pathway. *Science* 281, 1509-1512.
- Hecht, A., Litterst, C. M., Huber, O. und Kemler, R. (1999). Functional characterization of multiple transactivating elements in  $\beta$ -catenin, some of which interact with the TATA-binding protein *in vitro*. *J Biol Chem* 274, 18017-18025.
- Hecht, A., Vleminckx, K., Stemmler, M. P., van Roy, M. und Kemler, R. (2000). The p300/CBP acetyltransferases function as transcriptional coactivators of  $\beta$ -catenin in vertebrates. *EMBO J* 19, 1839-1850.
- Heisenberg, C. P., Tada, M., Rauch, G. J., Saude, L., Concha, M. L., Geisler, R., Stemple, D. L., Smith, D. L., Smith, J. C. und Wilson, S. W. (2000). Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* 405, 76-81.
- Homma, M. K., Li, D., Krebs, E. G., Yuasa, Y. und Homma, Y. (2002). Association and regulation of casein kinase 2 activity by adenomatous polyposis coli protein. *Proc Natl Acad Sci USA* 99, 5959-5964.
- Hsu, S. C., Galceran, J. und Grosschedl, R. (1998). Modulation of transcriptional regulation by LEF-1 in response to Wnt-1 signaling and association with  $\beta$ -catenin. *Mol Cell Biol* 18, 4807-4818.
- Huber, A. H., Nelson, W. J. und Weis, W. I. (1997). Three-dimensional structure of the armadillo repeat region of  $\beta$ -catenin. *Cell* 90, 871-882.
- Huber, A. H. und Weis, W. I. (2001). The structure of the  $\beta$ -catenin/E-cadherin complex and the molecular basis of the diverse ligand recognition by  $\beta$ -catenin. *Cell* 105, 391-402.
- Huber, O., Korn, R., McLaughlin, J., Ohsugi, M., Hermann, B. G. und Kemler, R. (1996). Nuclear localization of  $\beta$ -catenin by interaction with transcription factor LEF-1. *Mech Dev* 59, 3-10.
- Hülsken, J. und Birchmeier, W. (2001). New aspects of Wnt signaling pathways in higher vertebrates. *Curr Opin Gen Dev* 11, 547-553.
- Hülsken, J., Birchmeier, W. und Behrens, J. (1994). E-cadherin and APC compete for the interaction with  $\beta$ -catenin and the cytoskeleton. *J Cell Biol* 127, 2061-2069.

- Ikeda, S., Kishida, M., Matsuura, Y., Usui, H. und Kikuchi, A. (2000). GSK-3 $\beta$ -dependent phosphorylation of adenomatous polyposis coli gene product can be modulated by  $\beta$ -catenin and protein phosphatase 2A complexed with Axin. *Oncogene* 19, 537-545.
- Jiang, J. und Struhl, G. (1998). Regulation of the Hedgehog and Wingless signaling pathways by the F-box/WD40-repeat protein Slimb. *Nature* 391, 493-496.
- Kinzler, K. W. und Vogelstein, B. (1996). Lessons from hereditary colorectal cancer. *Cell* 87, 159-170.
- Kishida, M., Hino, S., Michiue, T., Yamamoto, H., Kishida, S., Fukui, A., Asashima, M. und Kikuchi, A. (2001). Synergistic activation of the Wnt signaling pathway by Dvl and casein kinase I $\epsilon$ . *J Biol Chem* 276, 33147-33155.
- Kishida, S., Yamamoto, H., Hino, S., Ikeda, S., Kishida, M. und Kikuchi, A. (1999). DIX domains of Dvl and axin are necessary for protein interactions and their ability to regulate  $\beta$ -catenin stability. *Mol Cell Biol* 19, 4414-4422.
- Kitagawa, M., Hatakeyama, S., Shirane, M., Matsumoto, K., Ishida, N., Hattori, K., Nakamichi, I., Kikuchi, A. und Nakayama, K. (1999). An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of  $\beta$ -catenin. *EMBO J* 18, 2401-2410.
- Korinek, V., Barker, N., Morin, P. J., van Wichen, D., de Weger, R., Kinzler, K. W., Vogelstein, B. und Clevers, H. (1997). Constitutive transcriptional activation by a  $\beta$ -catenin-Tcf complex in APC -/- colon carcinoma. *Science* 275, 1784-1787.
- Korinek, V., Barker, N., Willert, K., Molenaar, M., Roose, J., Wagenaar, G., Markman, M., Lamers, W., Destree, O. und Clevers, H. (1998). Two members of the Tcf family implicated in Wnt/ $\beta$ -catenin signaling during embryogenesis in the mouse. *Mol Cell Biol* 18, 1248-1256.
- Kramps, T., Peter, O., Brunner, E., Nellen, D., Froesch, B., Chatterjee, S., Murone, M. und Zullig, S. (2002). Wnt/Wingless signaling requires BCL9/legless-mediated recruitment of pygopus to the nuclear  $\beta$ -catenin-TCF complex. *Cell* 109, 47-60.
- Kühl, M., Sheldahl, L. C., Park, M., Miller, J. R. und Moon, R. T. (2000). The Wnt/Ca<sup>2+</sup> pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet* 16, 279-283.
- Labbe, E., A., L. und Attisano, L. (2000). Association of Smads with lymphoid enhancer binding factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor- $\beta$  and wnt pathways. *Proc Natl Acad Sci USA* 97, 8358-8363.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227, 660-685.
- Landesman-Bollag, E., Song, D. H., Romieu-Mourez, R., Sussman, D. J., Cardiff, R. D., Soneshein, G. E. und Seldin, D. C. (2001). Protein kinase CK2: signaling and tumorigenesis in the mammary gland. *Mol Cell Biochem* 277, 153-165.
- Laudet, V., Stehelin, D. und Clevers, H. (1993). Ancestry and diversity of the HMG box superfamily. *Nucleic Acids Res* 21, 2493-2501.
- Lee, E., Salic, A. und Kirschner, M. W. (2001). Physiological regulation of  $\beta$ -catenin stability by Tcf3 and CK1 $\epsilon$ . *J Cell Biol* 154, 983-993.

- Li, L., Yuan, H., Weaver, C. D., Mao, J., Farr, G. H., Sussman, D. J., Jonkers, J., Kimelman, D. und Wu, D. (1999). Axin and Frat1 interact with Dvl and GSK, bridging Dvl to GSK in Wnt-mediated regulation of LEF-1. *EMBO J* 18, 4233-4240.
- Li, X., Yost, H. J., Virshup, D. M. und Seeling, J. M. (2001). Protein phosphatase 2A and its B56 regulatory subunit inhibit Wnt signaling in *Xenopus*. *EMBO J* 20, 4122-4131.
- Lickert, H., Bauer, A., Kemler, R. und Stappert, J. (2000). Casein kinase II phosphorylation of E-cadherin increases E-cadherin/β-catenin interaction and strengthens cell-cell adhesion. *J Biol Chem* 275, 5090-5095.
- Litchfield, D. W. (2003). Protein kinase CK2: structure, regulation and role in cellular decisions of life and death. *Biochem J* 369, 1-15.
- Liu, C., Kato, Y., Zhang, Y., Do, V. M., Yanker, B. A. und He, X. (1999). β-TrCP couples β-catenin phosphorylation-degradation and regulates *Xenopus* axis formation. *Proc Natl Acad Sci USA* 96, 6273-6278.
- Liu, C., Li, Y., Semenov, M., Han, C., Baeg, G. H., Tan, Y., Zhang, Z., Lin, X. und He, X. (2002). Control of β-catenin phosphorylation/degradation by a dual kinase mechanism. *Cell* 108, 837-847.
- Liu, W., Dong, X., Mai, M., Seelan, R. S., Taniguchi, K., Krishnadath, K. K., Halling, K. C., Cunningham, J. M., Qian, C. und Christensen, E. (2000). Mutations in AXIN2 cause colorectal cancer with defective mismatch repair by activating β-catenin-Tcf signaling. *Nat Genet* 26, 146-147.
- Love, J. J., Li, X., Case, D. A., Giese, K. und Grosschedl, R. (1995). Structural basis for DNA bending by the architectural transcription factor LEF-1. *Nature* 376, 791-795.
- Lu, B., Roegiers, F., Jan, L. Y. und Jan, Y. N. (2001). Adherens junctions inhibit asymmetric division in the *Drosophila* epithelium. *Nature* 409, 522-525.
- Mann, B., Gelos, M., Siedow, A., Hanski, M. L., Gratchev, A., Ilyas, M., Bodmer, W. F., Moyer, M. P., Riecken, E. O., Buhr, H. J. und Hanski, C. (1999). Target genes of β-catenin-T cell-factor/lymphoid-enhancer-factor signaling in human colorectal carcinomas. *Proc Natl Acad Sci USA* 96, 1603-1608.
- Mao, J., Wang, J., Liu, B., Pan, W., Farr, G. H., Flynn, C., Yuan, H., Takada, S., Kimelman, D., Li, L. und Wu, D. (2001). Low-density lipoprotein receptor-related protein-5 binds to axin and regulates the canonical Wnt signaling pathway. *Mol Cell* 7, 801-809.
- Marikawa, Y. und Elinson, R. P. (1998). β-TrCP is a negative regulator of Wnt/β-catenin signaling pathway and dorsal axis formation in *Xenopus* embryos. *Mech Dev* 77, 75-80.
- McCrea, P. D. und Gumbiner, B. M. (1991). Purification of a 92 kDa cytoplasmic protein tightly associated with the cell-cell adhesion molecule E-cadherin (uvomorulin). Characterization and extractability of protein complex from the cell cytostructure. *J Biol Chem* 266, 4514-4520.
- McGrew, L. L., Takemaru, K. I., Bates, R. und Moon, R. T. (1999). Direct regulation of the *Xenopus engrailed-2* promotor by the Wnt signaling pathway, and a molecular screen for Wnt-responsive genes, confirm a role for Wnt signaling during neural patterning in *Xenopus*. *Mech Dev* 87, 21-32.
- McKay, R. M., Peters, J. M. und Graff, J. M. (2001). The casein kinase I family in Wnt signaling. *Dev Biol* 235, 388-396.

- Miravet, S., Piedra, J., Miro, F., Itarte, E., Garcia de Herreros, A. und Dunach, M. (2002). The transcriptional factor Tcf-4 contains different binding sites for  $\beta$ -catenin and plakoglobin. *J Biol Chem* 277, 1884-1891.
- Mlodzik, M. (2000). Spiny legs and prickled bodies: new insights and complexities in planar polarity establishment. *Bioessays* 22, 311-315.
- Molenaar, M., van de Wetering, M., Oosterwegel, M., Petersonmaduro, J., Godsake, S., Korinek, V., Roose, J., Destree, O. und Clevers, H. (1996). XTCF-3 transcription factor mediates  $\beta$ -catenin-induced axis formation in Xenopus embryos. *Cell* 86, 391-399.
- Morin, P. J. (1999).  $\beta$ -catenin signaling in cancer. *Bioessays* 21, 1021-1030.
- Morin, P. J., Sparks, A. B., Korinek, V., Barker, N., Clevers, H., Vogelstein, B. und Kinzler, K. W. (1997). Activation of  $\beta$ -catenin-Tcf signaling in colon cancer by mutations in  $\beta$ -catenin or APC. *Science* 275, 1787-1790.
- Munemitsu, S., Albert, I., Souza, B., Rubinfeld, B. und Polakis, P. (1995). Regulation of intracellular  $\beta$ -catenin levels by the adenomatous polyposis coli (APC) tumor-suppressor protein. *Proc Natl Acad Sci USA* 92, 3046-3050.
- Münstermann, U., Fritz, G., Seitz, G., Yiping, L., Schneider, H. und Issinger, O.-G. (1990). Casein kinase II is elevated in solid tumours and rapidly proliferating tissue. *Eur J Biochem* 189, 251-257.
- Noordermeer, J., Klingensmith, J., Perrimon, N. und Nusse, R. (1994). Dishevelled and armadillo act in the wingless signalling pathway in Drosophila. *Nature* 367, 80-83.
- Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. und Nakatani, Y. (1996). The transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell* 87, 953-959.
- Oosterwegel, M., van de Wetering, M., Dooijes, D., Klomp, L., Winoto, A., Georgopoulos, K., Meijlink, F. und Clevers, H. (1991). Cloning a murine TCF-1, a T cell-specific transcription factor interacting with functional motifs in the CD3 $\epsilon$  and T cell receptor  $\alpha$  enhancers. *J Exp Med* 173, 1133-1142.
- Orsulic, S., Huber, O., Aberle, H., Arnold, S. J. und Kemler, R. (1999). E-cadherin binding prevents  $\beta$ -catenin nuclear localization and  $\beta$ -catenin/LEF-1-mediated transactivation. *J Cell Sci* 112, 1237-1245.
- Oyama, T., Kanai, Y., Ochiai, A., Akimoto, S., Oda, T., Yanagihara, K., Nagafuchi, A., Tsukita, S., Shibamoto, S. und Ito, F. (1994). A truncated  $\beta$ -catenin disrupts the interaction between E-cadherin and  $\alpha$ -catenin: a cause of loss of intracellular adhesiveness in human cancer cell lines. *Cancer Res* 54, 6282-6287.
- Ozawa, M., Baribault, H. und Kemler, R. (1989). The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species. *EMBO J* 8, 1711-1717.
- Ozawa, M. und Kemler, R. (1992). Molecular organization of the uvomorulin-catenin complex. *J Cell Biol* 116, 989-996.

- Pawlowski, J. E., Ertel, J. R., Allen, M. P., Xu, M., Butler, C., Wilson, E. M. und Wierman, M. E. (2002). Liganded androgen receptor interaction with  $\beta$ -catenin: nuclear co-localization and modulation of transcriptional activity in neuronal cells. *J Biol Chem* 277, 20702-20710.
- Peifer, M. (1997). Cancer -  $\beta$ -catenin as oncogene - the smoking gun. *Science* 275, 1752-1753.
- Penner, C. G., Wang, Z. und Litchfield, D. W. (1997). Expression and localization of epitope-tagged protein kinase CK2. *J Cell Biol* 64, 525-537.
- Peters, J. M., McKay, R. M., McKay, J. P. und Graff, J. M. (1999). Casein kinase I transduces Wnt signals. *Nature* 401, 345-350.
- Pinna, L. A. (1990). Casein kinase 2: An 'eminence grise' in cellular regulation? *Biochim Biophys Acta* 1054, 267-284.
- Pinson, K. I., Brennan, J., Monkley, S., Avery, B. J. und Skarnes, W. C. (2000). An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* 407, 535-538.
- Pistorius, K., Seitz, G., Remberger, K. und Issinger, O.-G. (1991). Differential CKII activities in human colorectal mucosa, adenomas and carcinomas. *Onkologie* 14, 256-260.
- Polakis, P. (2000). Wnt signaling and cancer. *Genes Dev* 14, 1837-1851.
- Pukrop, T., Gradl, D., Henningfeld, K. A., Knöchel, W., Wedlich, D. und Kühl, M. (2001). Identification of two regulatory elements within the high mobility group box transcription factor XTCF-4. *J Biol Chem* 276, 8968-8978.
- Ratcliff, M. J., Itoh, K. und Sokol, S. Y. (2000). A positive role for the PP2A catalytic subunit in Wnt signal transduction. *J Biol Chem* 275, 35680-35683.
- Riese, J., Yu, X., Munnerlyn, A., Eresh, S., Hsu, S. C., Grosschedl, R. und Bienz, M. (1997). LEF-1, a nuclear factor coordinating signaling inputs from wingless and decapentaplegic. *Cell* 88, 777-787.
- Riggleman, B., Schedl, P. und Wieschaus, E. (1990). Spatial organization of the Drosophila segment polarity gene armadillo is posttranscriptionally regulated by wingless. *Cell* 63, 549-560.
- Roose, J., Molenaar, M., Peterson, J., Hurenkamp, J., Brantjes, H., Moerer, P., van de Wetering, M., Destree, O. und Clevers, H. (1998). The Xenopus Wnt XTcf-3 interacts with Groucho-related transcriptional repressors. *Nature* 395, 608-612.
- Rubinfeld, B., Albert, I., Porfiri, E., Fiol, C., Munemitsu, S. und Polakis, P. (1996). Binding of GSK3 $\beta$  to the APC- $\beta$ -catenin complex and regulation of complex assembly. *Science* 272, 1023-1026.
- Rubinfeld, B., Albert, I., Porfiri, E., Munemitsu, S. und Polakis, P. (1997a). Loss of  $\beta$ -catenin regulation by the APC tumor suppressor protein correlates with loss of structure due to common somatic mutations of the gene. *Cancer Res* 57, 4624-4630.
- Rubinfeld, B., Robbins, P., El-Gamil, M., Albert, I., Porfiri, E. und Polakis, P. (1997b). Stabilization of  $\beta$ -catenin by genetic defects in melanoma cell lines. *Science* 275, 1790-1792.

- Rubinfeld, B., Tice, D. A. und Polakis, P. (2001). Axin-dependent phosphorylation of the adenomatous polyposis coli protein mediated by casein kinase 1 $\epsilon$ . *J Biol Chem* 276, 39037-39045.
- Sakanaka, C., Leong, P., Harrison, S. D. und Williams, L. T. (1999). Casein kinase 1 $\epsilon$  in the Wnt pathway: regulation of  $\beta$ -catenin function. *Proc Natl Acad Sci USA* 96, 12548-12552.
- Salic, A., Lee, E., Mayer, L. und Kirschner, M. W. (2000). Control of  $\beta$ -catenin stability: Reconstitution of the cytoplasmic steps of the wnt pathway in Xenopus egg extracts. *Mol Cell* 5, 523-532.
- Sanger, F., Nicklen, S. und Coulson, A. R. (1992). DNA sequencing with chain-terminating inhibitors. 1977. *Biotechnology* 24, 104-108.
- Santos, J., Logarinho, E., Tapia, C., Allende, C., Allende, J. und Sunkel, C. (1996). The casein kinase I $\alpha$  gene of *Drosophila melanogaster* is developmentally regulated and the kinase activity of the protein induced by DNA damage. *J Cell Sci* 109, 1847-1856.
- Satoh, S., Daigo, Y., Furukawa, Y., Kato, T., Miwa, N., Nishiwaki, T., Kawasoe, T., Ishiguro, H., Fujita, M. und Tokino, T. (2000). Axin1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1. *Nat Genet* 24, 245-250.
- Schwarz-Romond, T., Asbrand, C., Bakkers, J., Kühl, M., Schaeffer, H.-J., Hülsken, J., Behrens, J., Hammerschmidt, M. und Birchmeier, W. (2002). The ankyrin repeat protein Diversin recruits casein kinase I $\epsilon$  to the  $\beta$ -catenin degradation complex and acts in both canonical Wnt and Wnt/JNK signaling. *Genes Dev* 16, 2073-2084.
- Seeling, J. M., Miller, J. R., Gil, R., Moon, R. T., White, R. und Virshup, D. M. (1999). Regulation of  $\beta$ -catenin signaling by the B56 subunit of protein phosphatase 2A. *Science* 283, 2089-2091.
- Seidenstricker, M. J. und Behrens, J. (2000). Biochemical interactions in the Wnt pathway. *Biophys Acta* 1495, 168-182.
- Seldin, D. C. und Leder, P. (1995). Casein kinase II $\alpha$  transgene-induced murine lymphoma: relation to theileriosis in cattle. *Science* 267, 894-897.
- Semplici, F., Meggio, F., Pinna, L. A. und Oliviero, S. (2002). CK2-dependent phosphorylation of the E2 ubiquitin conjugating enzyme UBC3B induces its interaction with  $\beta$ -TrCP and enhances  $\beta$ -catenin degradation. *Oncogene* 21, 3978-3987.
- Shtutman, M., Zhurinsky, J., Simcha, I., Albanese, C., D'Amico, M., Pestell, R. und Ben-Ze'ev, A. (1999). The cyclin D1 gene is a target of the  $\beta$ -catenin/LEF-1 pathway. *Proc Natl Acad Sci USA* 96, 5522-5527.
- Siegfried, E., Wilder, E. und Perrimon, N. (1994a). Components of wingless signalling in *Drosophila*. *Nature* 367, 76-80.
- Smalley, M. J., Sara, E., Paterson, H., Naylor, S., Cook, D., Jayatilake, H., Fryer, L. G., Hutchinson, L., Fry, M. J. und Dale, T. C. (1999). Interaction of axin and Dvl-2 proteins regulates Dvl-2-stimulated TCF-dependent transcription. *EMBO J* 18, 2823-2835.
- Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., Fujimoto, E. K., Goeke, N. M., Olson, B. J. und Klenk, D. C. (1985). Measurement of protein using bicinchoninic acid. *Anal Biochem* 150, 76-85.

- Smits, R., Kielman, M. F., Breukel, C., Zurcher, C., Neufeld, K., Jagmohan-Changur, S., Hofland, N., van Dijk, J., White, R. und Edelmann, W. (1999). Apc<sup>1638T</sup>: a mouse model delineating critical domains of the adenomatous polyposis coli protein involved in tumorigenesis and development. *Genes Dev* 13, 1309-1321.
- Song, D. H., Sussman, D. J. und Seldin, D. C. (2000). Endogenous protein kinases CK2 participates in Wnt signaling in mammary epithelial cells. *J Biol Chem* 275, 23790-23797.
- Stalter, G., Siemer, S., Becht, E., Ziegler, M., Remberger, K. und Issinger, O.-G. (1994). Asymmetric expression of protein kinase CK2 subunits in human kidney tumors. *Biochem Biophys Res Commun* 202, 141-147.
- Strovel, E. T. und Sussman, D. J. (1999). Transient overexpression of murine dishevelled genes results in apoptotic cell death. *Exp Cell Res* 253, 637-648.
- Strovel, E. T., Wu, D. und Sussman, D. J. (2000). Protein phosphatase 2C $\alpha$  dephosphorylates axin and activates LEF-1-dependent transcription. *J Biol Chem* 275, 2399-2403.
- Su, L. K., Vogelstein, B. und Kinzler, K. W. (1993). Association of the APC tumor suppressor protein with catenins. *Science* 262, 1734-1737.
- Sun, T.-Q., Lu, B., Reinhard, C., Jan, Y. N., Fant, W. J. und Williams, L. T. (2001). PAR-1 is a dishevelled-associated kinase and a positive regulator of Wnt signalling. *Nature Cell Biol* 3, 628-636.
- Tago, K., Nakamura, T., Nishita, M., Hyodo, J., Nagai, S., Murata, Y., Adachi, S., Ohwada, S., Morishita, Y., Shibuya, H. und Akiyama, T. (2000). Inhibition of Wnt signaling by ICAT, a novel  $\beta$ -catenin-interacting protein. *Genes Dev* 14, 1741-1749.
- Takemaru, K. I. und Moon, R. T. (2000). The transcriptional coactivator CBP interacts with  $\beta$ -catenin to activate gene expression. *J Cell Biol* 149, 249-254.
- Tamai, K., Semenov, M., Kato, Y., Spokony, R., Liu, C., Katsuyama, Y., Hess, F., Saint-Jeannet, J. P. und He, X. (2000). LDL-receptor-related proteins in Wnt Signal transduction. *Nature* 407, 530-535.
- Tetsu, O. und McCormick, F. (1999).  $\beta$ -Catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* 398, 422-426.
- Torres, M., Stoykova, A., Huber, O., Chowdhury, K., Bonaldo, P., Mansouri, A., Butz, S., Kemler, R. und Gruss, P. (1997). An  $\alpha$ -E-catenin gene trap mutation defines its function in preimplantation development. *Proc Natl Acad Sci USA* 94, 901-906.
- Towbin, H., Stähelin, T. und Gordon, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* 76, 4350-4354.
- Travis, A., Amsterdam, A., Belanger, C. und Grosschedl, R. (1991). LEF-1, a gene encoding a lymphoid-specific protein with an HMG domain, regulates T-cell receptor  $\alpha$  enhancer function. *Genes Dev* 5, 880-894.
- Tuazon, P. T. und Traugh, J. A. (1991). Casein kinase I and II-multipotential protein kinases: structure, function and regulation. *Adv Second Messenger Phosphoprotein Res* 23, 123-163.

- van Beest, M., Dooijes, D., van de Wetering, M., Kjaerulff, S., Bonvin, A., Nielsen, O. und Clevers, H. (2000). Sequence-specific high mobility group box factors recognize 10-12 base pair minor groove motifs. *J Biol Chem* 275, 27266-27273.
- van de Wetering, M., Cavallo, R. A., Dooijes, D., van Beest, M., van Es, J., Loureiro, J., Ypma, A., Hursh, D., Jones, T., Bejsovec, A., et al. (1997). Armadillo coactivates transcription driven by the product of the *Drosophila* segment polarity gene dTCF. *Cell* 88, 789-799.
- van de Wetering, M., Oosterwegel, M., Dooijes, D. und Clevers, H. (1991). Identification and cloning of TCF-1, a T lymphocyte-specific transcription factor containing a sequence-specific HMG box. *EMBO J* 10, 123-132.
- van der Geer, P., Luo, K., Sefton, B. M. und Hunter, T. (1995). Phosphopeptide mapping and phosphoamino acid analysis on cellulose thin-layer plates. In *Protein Phosphorylation - A Practical Approach*, D. G. Hardie, ed. (Oxford University Press), pp. 31-58.
- Vilk, G., Saulnier, R. B., St Pierre, R. und Litchfield, D. W. (1999). Inducible expression of protein kinase CK2 in mammalian cells. Evidence for functional specialization of CK2 isoforms. *J Biol Chem* 274, 14406-14414.
- Vleminckx, K., Kemler, R. und Hecht, A. (1999). The C-terminal transactivation domain of  $\beta$ -catenin is necessary and sufficient for signaling by the LEF-1/ $\beta$ -catenin complex in *Xenopus laevis*. *Mech Dev* 81, 65-74.
- von Kries, J. P., Winbeck, G., Asbrand, C., Schwarz-Romond, T., Sochnikova, N., Dell'Oro, A. und Birchmeier, W. (2000). Hot spots in  $\beta$ -catenin for interactions with LEF-1, conductin and APC. *Nat Struct Biol* 7, 800-807.
- Wagner, U., Brownlees, J., Irving, N. G., Lucas, F. R., Salinas, P. C. und Miller, C. C. (1997). Overexpression of the mouse dishevelled-1 protein inhibits GSK3 $\beta$  mediated phosphorylation of tau in transfected mammalian cells. *FEBS Lett* 411, 369-372.
- Waltzer, L. und Bienz, M. (1998). Drosophila CBP represses the transcription factor TCF to antagonize Wingless signalling. *Nature* 395, 521-525.
- Waterman, M., Fischer, W. H. und Jones, K. A. (1991). A thymus-specific member of the HMG protein family regulates the human T cell receptor C $\alpha$  enhancer. *Genes Dev* 5, 656-669.
- Wehrli, M., Dougan, S. T., Caldwell, K., O'Keefe, L., Schwartz, S., Vaizel-Ohayon, D., Schejter, E., Tomlinson, A. und DiNardo, S. (2000). Arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* 407, 527-530.
- Willert, K., Brink, M., Wodarz, A., Varmus, H. und und Nusse, R. (1997). Casein kinase 2 associates with and phosphorylates Dishevelled. *EMBO J* 16, 3089-3096.
- Willert, K., Shibamoto, S. und Nusse, R. (1999). Wnt-induced dephosphorylation of axin releases  $\beta$ -catenin from the axin complex. *Genes Dev* 13, 1768-1773.
- Wodarz, A. und Nusse, R. (1998). Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 14, 59-88.

- Yamamoto, H., Kishida, S., Kishida, M., Ikeda, S., Takada, S. und Kikuchi, A. (1999). Phosphorylation of axin, a Wnt signal negative regulator, by glycogen synthase-3 $\beta$  regulates its stability. *J Biol Chem* 274, 10681-10684.
- Yanagawa, S., Matsuda, Y., Lee, Y., Matsubayashi, H., Sese, S., Kadokawa, T. und Ishimoto, A. (2002). Casein kinase I phosphorylates the Armadillo protein and induces its degradation in Drosophila. *EMBO J* 21, 1733-1742.
- Yang, F., Li, X., Sharma, M., Sasaki, C. Y., Longo, D. L., Lim, B. und Sun, Z. (2002). Linking  $\beta$ -catenin to androgen signaling pathway. *J Biol Chem* 277, 11336-11344.
- Yost, C., Farr, G. H., Pierce, S. B., Ferkey, D. M., Chen, M. M. und Kimelman, D. (1998). GBP, an inhibitor of GSK-3, is implicated in Xenopus development and oncogenesis. *Cell* 93, 1031-1041.
- Yost, C., Torres, M., Miller, J. R., Huang, E., Kimelman, D. und Moon, R. T. (1996). The axis-inducing activity, stability, and subcellular distribution of  $\beta$ -catenin is regulated in *Xenopus* embryos by glycogen synthase kinase 3. *Genes Dev* 10, 1443-1454.
- Zhang, Y., Qiu, W. J., Chan, S. C., Han, J., He, X. und Lin, S. C. (2002). Casein kinase I and casein kinase II differentially regulate axin function in Wnt and JNK pathways. *J Biol Chem* 277, 17706-17712.
- Zorn, A. M., Barish, G. D., O., W. B., Lavender, P., Klymkowsky, M. W. und Varmus, H. E. (1999). Regulation of Wnt signaling by Sox proteins: XSox17  $\alpha/\beta$  and XSox3 physically interact with  $\beta$ -catenin. *Mol Cell* 4, 487-498.