

8. Zitierte Literatur

- Abraham WC, Bear MF (1996) Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci* 19:126-130.
- Adler CM, Goldberg TE, Malhotra AK, Pickar D, Breier A (1998) Effects of ketamine on thought disorder, working memory, and semantic memory in healthy volunteers. *Biol Psychiatry* 43:811-816.
- Aghajanian GK, Marek GJ (2000) Serotonin model of schizophrenia: emerging role of glutamate mechanisms. *Brain Res Brain Res Rev* 31:302-312.
- Akbarian S, Huntsman MM, Kim JJ, Tafazzoli A, Potkin SG, Bunney WE, Jr., Jones EG (1995) GABAA receptor subunit gene expression in human prefrontal cortex: comparison of schizophrenics and controls. *Cereb Cortex* 5:550-560.
- Akers RF, Lovinger DV, Colley PA, Linden DJ, Routtenberg A (1986) Translocation of protein kinase C activity may mediate hippocampal long-term potentiation. *Science* 231:587-589.
- Alger BE, Megela AL, Teyler TJ (1978) Transient heterosynaptic depression in the hippocampal slice. *Brain Res Bull* 3:181-184.
- Alger BE, Teyler TJ (1976) Long-term and short-term plasticity in CA1, CA3 and dentate region of the rat hippocampal slice. *Brain Res* 110:463-480.
- Amaral DG, Witter MP (1989) The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 31:571-591.
- Amsterdam JD, Winokur A, Dyson W, Herzog S, Gonzalez F, Rott R, Koprowski H (1985) Borna disease virus. A possible etiologic factor in human affective disorders? *Arch Gen Psychiatry* 42:1093-1096.
- Andersen P, Holmqvist B, Voorhoeve PE (1966) Excitatory synapses on hippocampal apical dendrites activated by entorhinal stimulation. *Acta Physiol Scand* 66:461-472.
- Andersen P, Lomo T (1967) Control of hippocampal output by afferent volley frequency. *Prog Brain Res* 27:400-412.
- Andersen P, Sundberg SH, Sveen O, Wigstrom H (1977) Specific long-lasting potentiation of synaptic transmission in hippocampal slices. *Nature* 266:736-737.
- Andine P, Widermark N, Axelsson R, Nyberg G, Olofsson U, Martensson E, Sandberg M (1999) Characterization of MK-801-induced behavior as a putative rat model of psychosis. *J Pharmacol Exp Ther* 290:1393-1408.
- Anis NA, Berry SC, Burton NR, Lodge D (1983) The dissociative anaesthetics, ketamine and phencyclidine, selectively reduce excitation of central mammalian neurones by N-methyl-aspartate. *Br J Pharmacol* 79:565-575.
- Aradi I, Santhakumar V, Chen K, Soltesz I (2002) Postsynaptic effects of GABAergic synaptic diversity: regulation of neuronal excitability by changes in IPSC variance. *Neuropharmacology* 43:511-522.
- Baimbridge KG, Miller JJ (1981) Calcium uptake and retention during long-term potentiation of neuronal activity in the rat hippocampal slice preparation. *Brain Res* 221:299-305.
- Bear MF, Abraham WC (1996) Long-term depression in hippocampus. *Annu Rev Neurosci* 19:437-462.
- Beck H, Goussakov IV, Lie A, Helmstaedter C, Elger CE (2000) Synaptic plasticity in the human dentate gyrus. *J Neurosci* 20:7080-7086.
- Behrens CJ, van den Boom LP, de HL, Friedman A, Heinemann U (2005) Induction of sharp wave-ripple complexes in vitro and reorganization of hippocampal networks. *Nat Neurosci* 8:1560-1567.
- Benes FM, Vincent SL, Alsterberg G, Bird ED, SanGiovanni JP (1992) Increased GABAA receptor binding in superficial layers of cingulate cortex in schizophrenics. *J Neurosci* 12:924-929.

- Bienenstock EL, Cooper LN, Munro PW (1982) Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J Neurosci* 2:32-48.
- Birse EF, Eaton SA, Jane DE, Jones PL, Porter RH, Pook PC, Sunter DC, Udvarhelyi PM, Wharton B, Roberts PJ, . (1993) Phenylglycine derivatives as new pharmacological tools for investigating the role of metabotropic glutamate receptors in the central nervous system. *Neuroscience* 52:481-488.
- Bliss TV, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361:31-39.
- Bliss TVP, Gardner-Medwin AR (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the unanaesthetized rabbit following stimulation of the perforant path. *J Physiol (Lond)* 232:357-374.
- Bliss TVP, Lømo T (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol (Lond)* 232:331-356.
- Bortolotto ZA, Clarke VR, Delany CM, Parry MC, Smolders I, Vignes M, Ho KH, Miu P, Brinton BT, Fantaske R, Ogden A, Gates M, Ornstein PL, Lodge D, Bleakman D, Collingridge GL (1999) Kainate receptors are involved in synaptic plasticity. *Nature* 402:297-301.
- Boulton CL, von Haebler D, Heinemann U (1992) Tracing of axonal connections by rhodamine-dextran-amine in the rat hippocampal-entorhinal cortex slice preparation. *Hippocampus* 2:99-106.
- Braunewell KH, Manahan-Vaughan D (2001) Long-term depression: a cellular basis for learning? *Rev Neurosci* 12:121-140.
- Breier A, Su TP, Saunders RC, Carson RE, Kolachana BS, De Bartolomeis A, Weinberger DR, Weisenfeld N, Malhotra AK, Eckelman WC, Pickar D (1997) Schizophrenia is associated with elevated amphetamine-induced synaptic dopamine concentrations: Evidence from a novel positron emission tomography method. *Proc Natl Acad Sci U S A* 94:2569-2574.
- Brun VH, Otnass MK, Molden S, Steffenach HA, Witter MP, Moser MB, Moser EI (2002) Place cells and place recognition maintained by direct entorhinal-hippocampal circuitry. *Science* 296:2243-2246.
- Buck N, Cali S, & Behr J (2006). Enhancement of long-term potentiation at CA1-subiculum synapses in MK-801-treated rats. *Neurosci Lett* 392: 5-9
- Buzsaki G (1980) Long-term potentiation of the commissural path-CA1 pyramidal cell synapse in the hippocampus of the freely moving rat. *Neurosci Lett* 19:293-296.
- Buzsaki G (1986) Hippocampal sharp waves: their origin and significance. *Brain Res* 398:242-252.
- Buzsaki G, Horvath Z, Urioste R, Hetke J, Wise K (1992) High-frequency network oscillation in the hippocampus. *Science* 256:1025-1027.
- Buzsaki G, Leung LW, Vanderwolf CH (1983) Cellular bases of hippocampal EEG in the behaving rat. *Brain Res* 287:139-171.
- Carlsson A (1988) The current status of the dopamine hypothesis of schizophrenia. *Neuropsychopharmacology* 1:179-186.
- Castillo PE, Malenka RC, Nicoll RA (1997) Kainate receptors mediate a slow postsynaptic current in hippocampal CA3 neurons. *Nature* 388:182-186.
- Chapman CA, Perez Y, Lacaille JC (1998) Effects of GABA(A) inhibition on the expression of long-term potentiation in CA1 pyramidal cells are dependent on tetanization parameters. *Hippocampus* 8:289-298.
- Church SM, Murray RM, Bramon E, Cotter D (2002) Does schizophrenia result from developmental or degenerative processes? *J Neural Transm Suppl*.
- Cohen AS, Abraham WC (1996) Facilitation of long-term potentiation by prior activation of metabotropic glutamate receptors. *J Neurophysiol* 76:953-962.
- Colbert CM, Levy WB (1992) Electrophysiological and pharmacological characterization of perforant path synapses in CA1: mediation by glutamate receptors. *J Neurophysiol* 68:1-8.

- Colbert CM, Levy WB (1993) Long-term potentiation of perforant path synapses in hippocampal CA1 *in vitro*. *Brain Res* 606:87-91.
- Corso TD, Sesma MA, Tenkova TI, Der TC, Wozniak DF, Farber NB, Olney JW (1997) Multifocal brain damage induced by phencyclidine is augmented by pilocarpine. *Brain Res* 752:1-14.
- Coussens CM, Kerr DS, Abraham WC (1997) Glucocorticoid receptor activation lowers the threshold for NMDA-receptor-dependent homosynaptic long-term depression in the hippocampus through activation of voltage-dependent calcium channels. *J Neurophysiol* 78:1-9.
- Creese I, Burt DR, Snyder SH (1976) Dopamine receptor binding predicts clinical and pharmacological potencies of antischizophrenic drugs. *Science* 192:481-483.
- Davies CH, Clarke VRJ, Jane DE, Collingridge GL (1995) Pharmacology of postsynaptic metabotropic glutamate receptors in rat hippocampal CA1 pyramidal neurones. *Br J Pharmacol* 116:1859-1869.
- Davies CH, Pozza MF, Collingridge GL (1993) CGP 55845A: A potent antagonist of GABA_B receptors in the CA1 region of rat hippocampus. *Neuropharmacology* 32:1071-1073.
- Davies SN, Martin D, Millar JD, Aram JA, Church J, Lodge D (1988) Differences in results from *in vivo* and *in vitro* studies on the use-dependency of N-methylaspartate antagonism by MK-801 and other phencyclidine receptor ligands MK-801 neurotoxicity in male mice: histologic effects and chronic impairment in spatial learning. *Eur J Pharmacol* 145:141-151.
- De No (1934) Studies on the Structure of the cerebral Cortex. *Journal f Psychologie und Neurologie*.
- Deisz RA, Prince DA (1989) Frequency-dependent depression of inhibition in guinea pig neocortex *in vitro* by GABA_B receptor feed-back on GABA release. *J Physiol (Lond)* 412:513-542.
- Doller HJ, Weight FF (1982) Perforant pathway activation of hippocampal CA1 stratum pyramidale neurons: electrophysiological evidence for a direct pathway. *Brain Res* 237:1-13.
- Doller HJ, Weight FF (1985) Perforant pathway-evoked long-term potentiation of CA1 neurons in the hippocampal slice preparation. *Brain Res* 333:305-310.
- Douglas RM (1977) Long lasting synaptic potentiation in the rat dentate gyrus following brief high frequency stimulation. *Brain Res* 126:361-365.
- Douglas RM, Goddard GV (1975) Long-term potentiation of the perforant path-granule cell synapse in the rat hippocampus. *Brain Res* 86:205-215.
- Dudek SM, Bear MF (1992) Homosynaptic long-term depression in area CA1 of hippocampus and effects of N-methyl-D-aspartate receptor blockade. *Proc Natl Acad Sci U S A* 89:4363-4367.
- Dugladze T, Lepsveridze E, Breustedt J, Kehrer C, Heinemann U, Gloveli T (2004) Effects of phencyclidines on signal transfer from the entorhinal cortex to the hippocampus in rats. *Neurosci Lett* 354:185-188.
- Dvorak-Carbone H, Schuman EM (1999a) Long-term depression of temporoammonic-CA1 hippocampal synaptic transmission. *J Neurophysiol* 81:1036-1044.
- Dvorak-Carbone H, Schuman EM (1999b) Patterned activity in stratum lacunosum moleculare inhibits CA1 pyramidal neuron firing. *J Neurophysiol* 82:3213-3222.
- Dworkin RH, Opler LA (1992) Simple schizophrenia, negative symptoms, and prefrontal hypodopaminergia. *Am J Psychiatry* 149:1284-1285.
- Eichenbaum H (2000) A cortical-hippocampal system for declarative memory. *Nat Rev Neurosci* 1:41-50.
- Empson RM, Heinemann U (1995) Perforant path connections to area CA1 are predominantly inhibitory in the rat hippocampal-entorhinal cortex combined slice preparation. *Hippocampus* 5:104-107.
- Erchova I, Kreck G, Heinemann U, Herz AV (2004) Dynamics of rat entorhinal cortex layer II and III cells: characteristics of membrane potential resonance at rest predict oscillation properties near threshold. *J Physiol* 560:89-110.

- Farber NB, Kim SH, Dikranian K, Jiang XP, Heinkel C (2002) Receptor mechanisms and circuitry underlying NMDA antagonist neurotoxicity. *Mol Psychiatry* 7:32-43.
- Farber NB, Wozniak DF, Price MT, Labruyere J, Huss J, St Peter H, Olney JW (1995) Age-specific neurotoxicity in the rat associated with NMDA receptor blockade: potential relevance to schizophrenia? *Biol Psychiatry* 38:788-796.
- Ferraguti F, Cobden P, Pollard M, Cope D, Shigemoto R, Watanabe M, Somogyi P (2004) Immunolocalization of metabotropic glutamate receptor 1alpha (mGluR1alpha) in distinct classes of interneuron in the CA1 region of the rat hippocampus. *Hippocampus* 14:193-215.
- Fix AS, Horn JW, Wightman KA, Johnson CA, Long GG, Storts RW, Farber N, Wozniak DF, Olney JW (1993) Neuronal vacuolization and necrosis induced by the noncompetitive N-methyl-D-aspartate (NMDA) antagonist MK(+)-801 (dizocilpine maleate): a light and electron microscopic evaluation of the rat retrosplenial cortex. *Exp Neurol* 123:204-215.
- Frey U, Krug M, Reymann KG, Matthies H (1988) Anisomycin, an inhibitor of protein synthesis, blocks late phases of LTP phenomena in the hippocampal CA1 region in vitro. *Brain Res* 452:57-65.
- Fuster JM (1997) *The Prefrontal Cortex: Anatomy, Physiology, and Neuropsychology of the Frontal Lobe*. Lippincott-Raven.
- Gao XM, Sakai K, Roberts RC, Conley RR, Dean B, Tamminga CA (2000) Ionotropic glutamate receptors and expression of N-methyl-D-aspartate receptor subunits in subregions of human hippocampus: effects of schizophrenia. *Am J Psychiatry* 157:1141-1149.
- Gloveli T, Schmitz D, Empson RM, Dugladze T, Heinemann U (1997a) Morphological and electrophysiological characterization of layer III cells of the medial entorhinal cortex of the rat. *Neuroscience* 77:629-648.
- Gloveli T, Schmitz D, Empson RM, Heinemann U (1997b) Frequency-dependent information flow from the entorhinal cortex to the hippocampus. *J Neurophysiol* 78:3444-3449.
- Goldman-Rakic PS (1996) Regional and cellular fractionation of working memory. *Proc Natl Acad Sci U S A* 93:13473-13480.
- Goldman-Rakic PS, Selemon LD (1997) Functional and anatomical aspects of prefrontal pathology in schizophrenia. *Schizophr Bull* 23:437-458.
- Grover LM, Teyler TJ (1990) Two components of long-term potentiation induced by different patterns of afferent activation. *Nature* 347:477-479.
- Grunze H, Gallinat J, Haag C, Hegerl U (2000) Psychopathological and auditory evoked potential correlates of ketamine psychosis--a single case report. *Schizophr Res* 41:386-388.
- Grunze HC, Rainnie DG, Hasselmo ME, Barkai E, Hearn EF, McCarley RW, Greene RW (1996) NMDA-dependent modulation of CA1 local circuit inhibition. *J Neurosci* 16:2034-2043.
- Gunther-Genta F, Bovet P, Hohlfeld P (1994) Obstetric complications and schizophrenia. A case-control study. *Br J Psychiatry* 164:165-170.
- Harris EW, Ganong AH, Cotman CW (1984) Long-term potentiation in the hippocampus involves activation of N-methyl-D-aspartate receptors. *Brain Res* 323:132-137.
- Harrison PJ, Weinberger DR (2005) Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Mol Psychiatry* 10:40-68.
- Hebb DO (1949) *The Organization of Behavior*. New York: Wiley.
- Heckers S (2001) Neuroimaging studies of the hippocampus in schizophrenia. *Hippocampus* 11:520-528.
- Heynen AJ, Abraham WC, Bear MF (1996) Bidirectional modification of CA1 synapses in the adult hippocampus *in vivo*. *Nature* 381:163-166.
- Horvath ZC, Czopf J, Buzsaki G (1997) MK-801-induced neuronal damage in rats. *Brain Res* 753:181-195.

- Hucker HB, Hutt JE, White SD, Arison BH, Zacchei AG (1983) Disposition and metabolism of (+)-5-methyl-10,11-dihydro-5H-dibenzo[a,d] cyclohepten-5,10-imine in rats, dogs, and monkeys. *Drug Metab Dispos* 11:54-58.
- Huettner JE, Bean BP (1988) Block of N-methyl-D-aspartate-activated current by the anticonvulsant MK-801: selective binding to open channels. *Proc Natl Acad Sci U S A* 85:1307-1311.
- Huh KH, Wenthold RJ (1999) Turnover analysis of glutamate receptors identifies a rapidly degraded pool of the N-methyl-D-aspartate receptor subunit, NR1, in cultured cerebellar granule cells. *J Biol Chem* 274:151-157.
- Impey S, Obrietan K, Wong ST, Poser S, Yano S, Wayman G, Deloulme JC, Chan G, Storm DR (1998) Cross talk between ERK and PKA is required for Ca²⁺ stimulation of CREB-dependent transcription and ERK nuclear translocation. *Neuron* 21:869-883.
- Izquierdo I, Medina JH (1993) Role of the amygdala, hippocampus and entorhinal cortex in memory consolidation and expression. *Braz J Med Biol Res* 26:573-589.
- Japha K, Koch M (1999) Picrotoxin in the medial prefrontal cortex impairs sensorimotor gating in rats: reversal by haloperidol. *Psychopharmacology (Berl)* 144:347-354.
- Jarrard LE (1993) On the role of the hippocampus in learning and memory in the rat. *Behav Neural Biol* 60:9-26.
- Javitt DC, Zukin SR (1991) Recent advances in the phencyclidine model of schizophrenia. *Am J Psychiatry* 148:1301-1308.
- Jay TM, Glowinski J, Thierry A-M (1989) Selectivity of the hippocampal projection to the prelimbic area of the prefrontal cortex in the rat. *Brain Res* 505:337-340.
- Jentsch JD, Redmond DE, Jr., Elsworth JD, Taylor JR, Youngren KD, Roth RH (1997) Enduring cognitive deficits and cortical dopamine dysfunction in monkeys after long-term administration of phencyclidine. *Science* 277:953-955.
- Joels M (2001) Corticosteroid actions in the hippocampus. *J Neuroendocrinol* 13:657-669.
- Kegeles LS, Abi-Dargham A, Zea-Ponce Y, Rodenhiser-Hill J, Mann JJ, Van Heertum RL, Cooper TB, Carlsson A, Laruelle M (2000) Modulation of amphetamine-induced striatal dopamine release by ketamine in humans: implications for schizophrenia. *Biol Psychiatry* 48:627-640.
- Kemp A, Manahan-Vaughan D (2004) Hippocampal long-term depression and long-term potentiation encode different aspects of novelty acquisition. *Proc Natl Acad Sci U S A* 101:8192-8197.
- Kemp N, Bashir ZI (1999) Induction of LTD in the adult hippocampus by the synaptic activation of AMPA/kainate and metabotropic glutamate receptors. *Neuropharmacology* 38:495-504.
- Khawaled R, Bruening-Wright A, Adelman JP, Maylie J (1999) Bicuculline block of small-conductance calcium-activated potassium channels. *Pflügers Archiv European Journal of Physiology*. 438, 3 / 1999: 314-321
- Kim JJ, Diamond DM (2002) The stressed hippocampus, synaptic plasticity and lost memories. *Nat Rev Neurosci* 3:453-462.
- Kim JJ, Foy MR, Thompson RF (1996) Behavioral stress modifies hippocampal plasticity through N-methyl-D-aspartate receptor activation. *Proc Natl Acad Sci U S A* 93:4750-4753.
- Kim JS, Kornhuber HH, Schmid-Burgk W, Holzmüller B (1980) Low cerebrospinal fluid glutamate in schizophrenic patients and a new hypothesis on schizophrenia. *Neurosci Lett* 20:379-382.
- Kiss J, Buzsáki G, Morrow JS, Glantz SB, Leranth C (1996) Entorhinal cortical innervation of parvalbumin-containing neurons (Basket and Chandelier cells) in the rat Ammon's horn. *Hippocampus* 6:239-246.
- Kokkinidis L, Anisman H (1980) Amphetamine models of paranoid schizophrenia: an overview and elaboration of animal experimentation. *Psychol Bull* 88:551-579.
- Krystal JH, Karper LP, Seibyl JP, Freeman GK, Delaney R, Bremner JD, Heninger GR, Bowers MB, Jr., Charney DS (1994) Subanesthetic effects of the noncompetitive NMDA antagonist, ketamine, in humans. Psychotomimetic, perceptual, cognitive, and neuroendocrine responses. *Arch Gen Psychiatry* 51:199-214.

- Lacaille J-C, Schwartzkroin PA (1988) Stratum lacunosum-moleculare interneurons of hippocampal CA1 region. I. intracellular response characteristics, synaptic responses, and morphology. *J Neurosci* 8:1400-1410.
- Lapointe V, Morin F, Ratte S, Croce A, Conquet F, Lacaille JC (2004) Synapse-specific mGluR1-dependent long-term potentiation in interneurons regulates mouse hippocampal inhibition. *J Physiol* 555:125-135.
- Laruelle M (1998) Imaging dopamine transmission in schizophrenia. A review and meta-analysis. *Q J Nucl Med* 42:211-221.
- Laruelle M, bi-Dargham A, Gil R, Kegeles L, Innis R (1999) Increased dopamine transmission in schizophrenia: relationship to illness phases. *Biol Psychiatry* 46:56-72.
- Levy WB, Steward O (1979) Synapses as associative memory elements in the hippocampal formation. *Brain Res* 175:233-245.
- Lewis DA (2000) GABAergic local circuit neurons and prefrontal cortical dysfunction in schizophrenia. *Brain Res Brain Res Rev* 31:270-276.
- Lewis DA, Pierri JN, Volk DW, Melchitzky DS, Woo TU (1999) Altered GABA neurotransmission and prefrontal cortical dysfunction in schizophrenia. *Biol Psychiatry* 46:616-626.
- Li H, Rogawski MA (1998) GluR5 kainate receptor mediated synaptic transmission in rat basolateral amygdala in vitro. *Neuropharmacology* 37:1279-1286.
- Li P, Wilding TJ, Kim SJ, Calejesan AA, Huettner JE, Zhuo M (1999) Kainate-receptor-mediated sensory synaptic transmission in mammalian spinal cord. *Nature* 397:161-164.
- Li X-G, Somogyi P, Tepper JM, Buzsáki G (1992) Axonal and dendritic arborization of an intracellularly labeled chandelier cell in the CA1 region of rat hippocampus. *Exp Brain Res* 90:519-525.
- Lipska BK, Weinberger DR (2000) To model a psychiatric disorder in animals: schizophrenia as a reality test. Implications of normal brain development for the pathogenesis of schizophrenia. *Neuropsychopharmacology* 23:223-239.
- Little Z, Grover LM, Teyler TJ (1995) Metabotropic glutamate receptor antagonist, (R,S)-alpha-methyl-4-carboxyphenylglycine, blocks two distinct forms of long-term potentiation in area CA1 of rat hippocampus. *Neurosci Lett* 201:73-76.
- Lodge D, Anis NA (1982) Effects of phencyclidine on excitatory amino acid activation of spinal interneurons in the cat. *Eur J Pharmacol* 77:203-204.
- Lopes da Silva FH, Witter MP (1990) Anatomic Organization and Physiology of the Limbic Cortex. *Physiological Review* 70:453-511.
- Lu YF, Kandel ER, Hawkins RD (1999) Nitric oxide signaling contributes to late-phase LTP and CREB phosphorylation in the hippocampus. *J Neurosci* 19:10250-10261.
- Luby ED, Cohen BD, Rosenbaum G, Gottlieb JS, Kelly R (1959) Study of a new schizophrenic-like drug: Sernyl. *Arch Neurol Psychiat* 363-369.
- Lynch G, Larson J, Kelso S, Barrionuevo G, Schottler F (1983) Intracellular injections of EGTA block induction of hippocampal long term potentiation. *Nature* 305:719-723.
- Lynch GS, Dunwiddie T, Gribkoff V (1977) Heterosynaptic depression: a postsynaptic correlate of long-term potentiation. *Nature* 266:737-739.
- Lynch MA (2004) Long-term potentiation and memory. *Physiol Rev* 84:87-136.
- Maguire EA, Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RS, Frith CD (2000) Navigation-related structural change in the hippocampi of taxi drivers. *Proc Natl Acad Sci U S A* 97:4398-4403.
- Malenka RC, Kauer JA, Perkel DJ, Mauk MD, Kelly PT, Nicoll RA, Waxham MN (1989) An essential role for postsynaptic calmodulin and protein kinase activity in long-term potentiation. *Nature* 340:554-557.
- Malinow R, Malenka RC (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* 25:103-126.

- Malinow R, Schulman H, Tsien RW (1989) Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science* 245:862-866.
- Manahan-Vaughan D, Kulla A, Frey JU (2000) Requirement of translation but not transcription for the maintenance of long-term depression in the CA1 region of freely moving rats. *J Neurosci* 20:8572-8576.
- Martin SJ, Grimwood PD, Morris RG (2000) Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* 23:649-711.
- McGrath J, Castle D (1995) Does influenza cause schizophrenia? A five year review. *Aust N Z J Psychiatry* 29:23-31.
- McNaughton BL, Barnes CA, Meltzer J, Sutherland RJ (1989) Hippocampal granule cells are necessary for normal spatial learning but not for spatially-selective pyramidal cell discharge. *Exp Brain Res* 76:485-496.
- McNaughton BL, Douglas RM, Goddard GV (1978) Synaptic enhancement in fascia dentata: cooperativity among coactive afferents. *Brain Res* 157:277-293.
- McNaughton N, Miller JJ (1984) Medial septal projections to the dentate gyrus of the rat: electrophysiological analysis of distribution and plasticity. *Exp Brain Res* 56:243-256.
- Miyamoto E (2006) Molecular mechanism of neuronal plasticity: induction and maintenance of long-term potentiation in the hippocampus. *J Pharmacol Sci* 100:433-442.
- Mody I, Baimbridge KG, Miller JJ (1984) Blockade of tetanic- and calcium-induced long-term potentiation in the hippocampal slice preparation by neuroleptics. *Neuropharmacology* 23:625-631.
- More JC, Nistico R, Dolman NP, Clarke VR, Alt AJ, Ogden AM, Buelens FP, Troop HM, Kelland EE, Pilato F, Bleakman D, Bortolotto ZA, Collingridge GL, Jane DE (2004) Characterisation of UBP296: a novel, potent and selective kainate receptor antagonist. *Neuropharmacology* 47:46-64.
- Moriyoshi K, Masu M, Ishii T, Shigemoto R, Mizuno N, Nakanishi S (1991) Molecular cloning and characterization of the rat NMDA receptor. *Nature* 354:31-37.
- Morris RG, Anderson E, Lynch GS, Baudry M (1986) Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5 MK-801 neurotoxicity in male mice: histologic effects and chronic impairment in spatial learning. *Nature* 319:774-776.
- Morris RG, Garrud P, Rawlins JN, O'Keefe J (1982) Place navigation impaired in rats with hippocampal lesions. *Nature* 297:681-683.
- Morris RGM (2003) Long-term potentiation and memory. *Philos Trans R Soc Lond [Biol]* 358:643-647.
- Morris RGM, Moser EI, Riedel G, Martin SJ, Sandin J, Day M, O'Carroll C (2003) Elements of a neurobiological theory of the hippocampus: the role of activity-dependent synaptic plasticity in memory. *Philos Trans R Soc Lond [Biol]* 358:773-786.
- Moser EI, Krobot KA, Moser MB, Morris RG (1998) Impaired spatial learning after saturation of long-term potentiation. *Science* 281:2038-2042.
- Mott DD, Lewis DV (1991) Facilitation of the induction of long-term potentiation by GABA_B receptors. *Science* 252:1718-1720.
- Mulkey RM, Malenka RC (1992) Mechanisms underlying induction of homosynaptic long-term depression in area CA1 of the hippocampus. *Neuron* 9:967-975.
- Naber PA, Witter MP, Lopez da Silva FH (1999) Perirhinal cortex input to the hippocampus in the rat: evidence for parallel pathways, both direct and indirect. A combined physiological and anatomical study. *Eur J Neurosci* 11:4119-4133.
- Nakao K, Ikegaya Y, Yamada MK, Nishiyama N, Matsuki N (2002) Hippocampal long-term depression as an index of spatial working memory. *Eur J Neurosci* 16:970-974.
- Newberry NR, Nicoll RA (1984) Direct hyperpolarizing action of baclofen on hippocampal pyramidal cells. *Nature* 308:450-452.

- Newcomer JW, Krystal JH (2001) NMDA receptor regulation of memory and behavior in humans. *Hippocampus* 11:529-542.
- Nicoll RA, Oliet SH, Malenka RC (1998) NMDA receptor-dependent and metabotropic glutamate receptor-dependent forms of long-term depression coexist in CA1 hippocampal pyramidal cells. *Neurobiol Learn Mem* 70:62-72.
- Nowak L, Bregestovski P, Ascher P, Herbet A, Prochiantz A (1984) Magnesium gates glutamate-activated channels in mouse central neurons. *Nature* 307:462-465.
- O'Keefe J, Dostrovsky J (1971) The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Research* 34:171-175.
- O'Keefe J, Nadel L (1978) *The hippocampus as a cognitive map*. Oxford: Oxford University Press.
- Olney JW, Farber NB (1995) Glutamate receptor dysfunction and schizophrenia. *Arch Gen Psychiatry* 52:998-1007.
- Olney JW, Labruyere J, Price MT (1989) Pathological changes induced in cerebrocortical neurons by phencyclidine and related drugs. *Science* 244:1360-1362.
- Olney JW, Labruyere J, Wang G, Wozniak DF, Price MT, Sesma MA (1991) NMDA antagonist neurotoxicity: Mechanism and prevention. *Science* 254:1515-1518.
- Olney JW, Newcomer JW, Farber NB (1999) NMDA receptor hypofunction model of schizophrenia. *J Psychiatr Res* 33:523-533.
- Olpe H-R, Karlsson G (1990) The effects of baclofen and two GABA_B-receptor antagonists on long-term potentiation. *Naunyn Schmiedebergs Arch Pharmacol* 342:194-197.
- Otani S, Connor JA (1998) Requirement of rapid Ca²⁺ entry and synaptic activation of metabotropic glutamate receptors for the induction of long-term depression in adult rat hippocampus. *J Physiol* 511 (Pt 3):761-770.
- Ouardouz M, Lacaille JC (1995) Mechanisms of selective long-term potentiation of excitatory synapses in stratum oriens/alveus interneurons of rat hippocampal slices. *J Neurophysiol* 73:810-819.
- Papez JW (1937) A proposed mechanism of emotion. *Arch Neurol Psychiatry* 38:725-743.
- Park S, Holzman PS (1992) Schizophrenics show spatial working memory deficits. *Arch Gen Psychiatry* 49:975-982.
- Patenaude C, Chapman CA, Bertrand S, Congar P, Lacaille JC (2003) GABAB receptor- and metabotropic glutamate receptor-dependent cooperative long-term potentiation of rat hippocampal GABA_A synaptic transmission. *J Physiol (Lond)* 553:155-167.
- Pearce BD, Valadi NM, Po CL, Miller AH (2000) Viral infection of developing GABAergic neurons in a model of hippocampal disinhibition. *Neuroreport* 11:2433-2438.
- Pearson G, Robinson F, Beers GT, Xu BE, Karandikar M, Berman K, Cobb MH (2001) Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev* 22:153-183.
- Penington NJ, Fox AP (1994) Effects of LSD on Ca⁺⁺ currents in central 5-HT-containing neurons: 5-HT_{1A} receptors may play a role in hallucinogenesis. *J Pharmacol Exp Ther* 269:1160-1165.
- Perez Y, Morin F, Lacaille JC (2001) A hebbian form of long-term potentiation dependent on mGluR1a in hippocampal inhibitory interneurons. *Proc Natl Acad Sci U S A* 98:9401-9406.
- Pozza MF, Manuel NA, Steinmann M, Froestl W, Davies CH (1999) Comparison of antagonist potencies at pre- and post-synaptic GABA(B) receptors at inhibitory synapses in the CA1 region of the rat hippocampus. *Br J Pharmacol* 127:211-219.
- Pulver AE (2000) Search for schizophrenia susceptibility genes. *Biol Psychiatry* 47:221-230.
- Racine RJ, Milgram NW, Hafner S (1983) Long-term potentiation phenomena in the rat limbic forebrain. *Brain Res* 260:217-231.

- Ramón y Cajal S (1911) Histologie du Systeme Nerveux de l'Homme et des Vertebres. Vol.II. Maloine, Paris. - 1960 Studies on Vertebrate Neurogenesis. Translated by L. Guth. Charles C Thomas, Springfield, Illinois. Paris: Maloine.
- Ramon y Cajal SR (1893) Estructura del Asta de Ammon y Fascial Dentata. Anal soc esp hist nat 22.
- Remondes M, Schuman EM (2002) Direct cortical input modulates plasticity and spiking in CA1 pyramidal neurons. Nature 416:736-740.
- Remondes M, Schuman EM (2004) Role for a cortical input to hippocampal area CA1 in the consolidation of a long-term memory. Nature 431:699-703.
- Remondes M, Schuman EM (2003) Molecular Mechanisms Contributing to Long-Lasting Synaptic Plasticity at the Temporoammonic-CA1 Synapse. Learning Memory 10:247-252.
- Rempel-Clower NL, Zola SM, Squire LR, Amaral DG (1996) Three cases of enduring memory impairment after bilateral damage limited to the hippocampal formation. J Neurosci 16:5233-5255.
- Robinson TE, Becker JB (1986) Enduring changes in brain and behavior produced by chronic amphetamine administration: a review and evaluation of animal models of amphetamine psychosis. Brain Res 396:157-198.
- Rujescu D, Bender A, Keck M, Hartmann AM, Ohl F, Raeder H, Giegling I, Genius J, McCarley RW, Moller HJ, Grunze H (2006) A pharmacological model for psychosis based on N-methyl-D-aspartate receptor hypofunction: molecular, cellular, functional and behavioral abnormalities. Biol Psychiatry 59:721-729.
- Sakurai SY, Cha JH, Penney JB, Young AB (1991) Regional distribution and properties of [3H]MK-801 binding sites determined by quantitative autoradiography in rat brain. Neuroscience 40:533-543.
- Sams-Dodd F (1996) Phencyclidine-induced stereotyped behaviour and social isolation in rats: A possible animal model of schizophrenia. Behav Pharmacol 7:3-23.
- Schulz PE (1997) Long-term potentiation involves increases in the probability of neurotransmitter release. Proc Natl Acad Sci U S A 94:5888-5893.
- Schulz PE, Cook EP, Johnston D (1994) Changes in paired-pulse facilitation suggest presynaptic involvement in long-term potentiation. J Neurosci 14:5325-5337.
- Schwartzkroin PA, Wester K (1975) Long-lasting facilitation of a synaptic potential following tetanization in the *in vitro* hippocampal slice. Brain Res 89:107-119.
- Scoville WB, Milner B (2000) Loss of recent memory after bilateral hippocampal lesions. 1957. J Neuropsychiatry Clin Neurosci 12:103-113.
- Seeman P (1987) Dopamine receptors and the dopamine hypothesis of schizophrenia. Synapse 1:133-152.
- Selten JP, van VK, Pleyte W, Herzog S, Hoek HW, van Loon AM (2000) Borna disease virus and schizophrenia in Surinamese immigrants to the Netherlands. Med Microbiol Immunol (Berl) 189:55-57.
- Siekmeier PJ, Hasselmo ME, Howard MW, Coyle J (2007) Modeling of context-dependent retrieval in hippocampal region CA1: implications for cognitive function in schizophrenia. Schizophr Res 89:177-190.
- Simpson MD, Slater P, Deakin JF, Royston MC, Skan WJ (1989) Reduced GABA uptake sites in the temporal lobe in schizophrenia. Neurosci Lett 107:211-215.
- Spencer WA, Kandel ER (1969) Synaptic inhibition in seizures. In: Basic mechanisms of epilepsies (Jasper HH, Ward AA, Pope A, eds), pp 575-603. Boston: Little Brown and Company.
- Squire LR, Zola-Morgan SM (1991) The medial temporal lobe memory system. Science 253:1380-1386.
- Stell BM, Mody I (2002) Receptors with different affinities mediate phasic and tonic GABA(A) conductances in hippocampal neurons. J Neurosci 22:RC223.
- Steullet P, Neijt HC, Cuenod M, Do KQ (2006) Synaptic plasticity impairment and hypofunction of NMDA receptors induced by glutathione deficit: relevance to schizophrenia. Neuroscience 137:807-819.

- Steward O, Scoville SA (1976) Cells of origin of entorhinal cortical afferents to the hippocampus and fascia dentata of the rat. *J Comp Neurol* 169:347-370.
- Subramaniam S, McGonigle P (1991) Quantitative autoradiographic characterization of the binding of (+)-5-methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5, 10-imine ($[^3\text{H}]$ MK-801) in rat brain: regional effects of polyamines. *J Pharmacol Exp Ther* 256:811-819.
- Suzuki SS, Smith GK (1987) Spontaneous EEG spikes in the normal hippocampus. I. Behavioral correlates, laminar profiles and bilateral synchrony. *Electroencephalogr Clin Neurophysiol* 67:348-359.
- Swartzwelder HS, Ferrari CM, Anderson WW, Wilson WA (1989) The drug MK-801 attenuates the development, but not the expression, of long-term potentiation and stimulus train-induced bursting in hippocampal slices. *Neuropharmacology* 28:441-445.
- Swerdlow NR, Shoemaker JM, Bongiovanni MJ, Neary AC, Tochen LS, Saint Marie RL (2005) Reduced startle gating after D1 blockade: effects of concurrent D2 blockade. *Pharmacol Biochem Behav* 82:293-299.
- Tordjman S, Drapier D, Bonnot O, Graignic R, Fortes S, Cohen D, Millet B, Laurent C, Roubertoux PL (2007) Animal models relevant to schizophrenia and autism: validity and limitations. *Behav Genet* 37:61-78.
- Torrey EF, Yolken RH (2003) *Toxoplasma gondii* and schizophrenia. *Emerg Infect Dis* 9:1375-1380.
- Tremblay R, Chakravarthy B, Hewitt K, Tauskela J, Morley P, Atkinson T, Durkin JP (2000) Transient NMDA receptor inactivation provides long-term protection to cultured cortical neurons from a variety of death signals. *J Neurosci* 20:7183-7192.
- Tsai G, Coyle JT (2002) Glutamatergic mechanisms in schizophrenia. *Annu Rev Pharmacol Toxicol* 42:165-179.
- Turrigiano GG, Nelson SB (2004) Homeostatic plasticity in the developing nervous system. *Nat Rev Neurosci* 5:97-107.
- van Welie I, van Hooff JA, Wadman WJ (2004) Homeostatic scaling of neuronal excitability by synaptic modulation of somatic hyperpolarization-activated Ih channels. *Proc Natl Acad Sci U S A* 101:5123-5128.
- Volk DW, Austin MC, Pierri JN, Sampson AR, Lewis DA (2000) Decreased glutamic acid decarboxylase67 messenger RNA expression in a subset of prefrontal cortical gamma-aminobutyric acid neurons in subjects with schizophrenia. *Arch Gen Psychiatry* 57:237-245.
- Wagner JJ, Alger BE (1995) GABAergic and developmental influences on homosynaptic LTD and depotentiation in rat hippocampus. *J Neurosci* 15:1577-1586.
- Wang HL, Wang GH, Li QY, Shu C, Jiang MS, Guo Y (2006) Prevalence of *Toxoplasma* infection in first-episode schizophrenia and comparison between *Toxoplasma*-seropositive and *Toxoplasma*-seronegative schizophrenia. *Acta Psychiatr Scand* 114:40-48.
- Weinberger DR (1987) Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry* 44:660-669.
- Weinberger DR (1999) Cell biology of the hippocampal formation in schizophrenia: Implications of normal brain development for the pathogenesis of schizophrenia. *Biol Psychiatry* 45:395-402.
- Willems J, Balster RL, Leander JD (1990) The behavioral pharmacology of NMDA receptor antagonists. *Trends Pharmacol Sci* 11:423-428.
- Wilsch VW, Behnisch T, Jager T, Reymann KG, Balschun D (1998) When are class I metabotropic glutamate receptors necessary for long-term potentiation? *J Neurosci* 18:6071-6080.
- Witter MP (1993) Organization of the entorhinal-hippocampal system: A review of current anatomical data. *Hippocampus* 3 Suppl.:33-44.
- Witter MP, Groenewegen HJ, Lopes da Silva FH, Lohman AH (1989) Functional organization of the extrinsic and intrinsic circuitry of the parahippocampal region. *Prog Neurobiol* 33:161-253.

- Witter MP, Wouterlood FG, Naber PA, Van Haefen T (2000) Anatomical organization of the parahippocampal-hippocampal network. *Ann N Y Acad Sci* 911:1-24.
- Wöhrl R., Eisenach S., Manahan-Vaughan D.; Heinemann U., von Haebler D. (2007), Acute and long-term effects of MK-801 on direct cortical input evoked homo- and heterosynaptic plasticity in the CA1 region of the female rat. *Eur J Neurosci.* (akzeptiert 9/2007)
- Wong EH, Kemp JA, Priestley T, Knight AR, Woodruff GN, Iversen LL (1986) The anticonvulsant MK-801 is a potent N-methyl-D-aspartate antagonist. *Proc Natl Acad Sci U S A* 83:7104-7108.
- Xu L, Anwyl R, Rowan MJ (1997) Behavioural stress facilitates the induction of long-term depression in the hippocampus. *Nature* 387:497-500.
- Yamada S, Harano M, Annoh N, Nakamura K, Tanaka M (1999) Involvement of serotonin 2A receptors in phencyclidine-induced disruption of prepulse inhibition of the acoustic startle in rats. *Biol Psychiatry* 46:832-838.
- Yamamoto C, Chujo T (1978) Long-term potentiation in thin hippocampal sections studied by intracellular and extracellular recordings. *Exp Neurol* 58:242-250.
- Ylinen A, Bragin A, Nadasdy Z, Jando G, Szabo I, Sik A, Buzsaki G (1995) Sharp wave-associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular mechanisms. *J Neurosci* 15:30-46.
- Zhang XL, Zhou ZY, Winterer J, Muller W, Stanton PK (2006) NMDA-dependent, but not group I metabotropic glutamate receptor-dependent, long-term depression at Schaffer collateral-CA1 synapses is associated with long-term reduction of release from the rapidly recycling presynaptic vesicle pool. *J Neurosci* 26:10270-10280.
- Zucker RS (1989) Short-term synaptic plasticity. *Annu Rev Neurosci* 12:13-31.