

5 Bibliographie

- (1) Jeffrey FMH, Mallow CR, Radda GK, 1987. Influence of intracellular acidosis on contractile function in the working rat heart. *Am J Physiol* 253: H1499-H1505
- (2) Mason MJ, Smith JD, Garcia-Soto JJ, Grinstein S. Internal pH-sensitive site couples $\text{Cl}^-/\text{HCO}_3^-$ exchange to the Na^+/H^+ antiport in lymphocytes. *Am J Physiol* 1989; 256: C428-33
- (3) Lagadic-Gossmann D, Buckler KJ, Vaughan-Jones RD. Role of bicarbonate in pH recovery from intracellular acidosis in the guinea-pig ventricular myocyte. *J Physiol (Lond)* 1992; 458: 361-84
- (4) Wallert MA, Fröhlich O. Na^+-H^+ exchange in isolated myocytes from rat heart. *Am J Physiol* 1989; 257: C207-13
- (5) Lucchesi PA, Berk BC. Regulation of sodium-hydrogen exchange in vascular smooth muscle. *Cardiovascular Research* 1995; 29: 172-177
- (6) Burckhardt B-C, Burckhardt G. Cellular mechanisms of proximal tubular acidification. In: Häussinger D, ed. *pH Homeostasis*. New York: Academic Press, 1988: 233-57
- (7) Sardet C, Franchi A, Pouyssegur J. Molecular cloning, primary structure, and expression of the human growth factor-activatable Na^+/H^+ antiporter. *Cell* 1989; 56: 271-80
- (8) Orlowski J, Kandasamy RA, Shull GE. Molecular cloning of putative members of the Na^+/H^+ exchanger gene family. *J Biol Chem* 1992; 267: 9331-9
- (9) Tse C-M, Brant SR, Walker MS, Pouyssegur J, Donowitz M. Cloning and sequencing of a rabbit cDNA encoding an intestinal and kidney-specific Na^+/H^+ exchanger isoform (NHE-3). *J Biol Chem* 1992; 267: 9340-6

- (10) Wang Z, Orlowski J, Schull GE. Primary structure and functional expression of a novel gastrointestinal isoform of the rat Na^+/H^+ exchanger. *J Biol Chem* 1993; 11925-8
- (11) Counillon L, Pouyssegur J, Reithmeier R. The Na^+/H^+ exchanger NHE-1 possesses N- and O-linked glycosylation restricted to the first N-terminal extracellular domain. *Biochemistry* 1994; 33:10463-9
- (12) Tani M, Neely JR. Role of intracellular Na^+ in Ca^{2+} overload and depressed recovery of ventricular function of reperfused ischemic rat hearts. Possible involvement of Na^+/H^+ and $\text{Na}^+/\text{Ca}^{2+}$ exchange. *Circ Res* 1989; 65: 1045-56
- (13) Pike MM, Lou CS, Clark MD, et al. NMR measurements of Na^+ and cellular energy in ischemic rat heart: Role of Na^+/H^+ exchange. *Am J Physiol* 1993; 265: H2017-26
- (14) Dennis SC, Gevers W, Opie LH. Protons in ischemia: Where do they come from; Where do they go to ? *J Mol Cell Cardiol* 1991; 23:1077-86
- (15) Karmazyn M. 1996. Role of sodium-hydrogen exchange in mediating myocardial ischemic and reperfusion injury. Mechanisms and therapeutic implications. In: Fliegel L (ed.). *The Na^+/H^+ Exchanger*. Austin. Texas: Springer/R.G. Landes Company. 189-215
- (16) Avkiran M, Yasutake M. 1996. Role of the sarcolemmal Na^+/H^+ Exchanger in arrhythmogenesis during reperfusion of ischemic myocardium. In: Fliegel L (ed.) *The Na^+/H^+ Exchanger*. Austin. Texas: Springer/R.G. Landes Company. 173-187
- (17) Murphy E, Perlman M, London RE, Steenbergen. Amiloride delays the ischemia-induced rise in cytosolic free calcium. *Circ Res* 1991; 68: 1250-8
- (18) Frelin C, Vigne P, Lazdunski M. The role of Na^+/H^+ exchange system in cardiac cells in relation to the control of the internal Na^+ concentration. *J Biol Chem* 1984; 259: 8880-5
- (19) Lazdunski M, Frelin C, Vigne P. The sodium/hydrogen exchange system in cardiac cells: its biochemical and pharmacological properties and its role in regulation internal concentrations of sodium and internal pH. *J Mol Cell Cardiol* 1985; 17:1029-42

- (20) Vaughn-Jones RD, Wu ML. Extracellular H⁺ inactivation of Na⁺-H⁺ exchange in the sheep cardiac Purkinje fibre. *J Physiol (Lond)* 1990; 428: 441-66
- (21) Ibuki C, Hearse DJ, Avkiran M. Mechanisms of antifibrillatory effect of acidic reperfusion: role of perfusate bicarbonate concentration. *Am J Physiol* 1993; 264: H783-90
- (22) Becker LC, Ambrosio G. Myocardial consequences of reperfusion. *Prog Cardiovasc Dis* 1987; 30: 23-44
- (23) Pierce GN, Meng H. The role of sodium-proton exchange in ischemic-reperfusion injury in the heart. *Am J Cardiovasc Pathol* 1992; 4: 91-102
- (24) Opie LH. Reperfusion injury and its pharmacologic modification. *Circulation* 1989; 80: 1049-62
- (25) Shen AC, Jennings RB. Kinetics of calcium accumulation in acute myocardial ischemic injury. *Am J Pathol* 1972; 67: 441-52
- (26) Grinwald PM, Nayler WG. Calcium entry in the calcium paradox. *J Mol Cell Cardiol* 1981; 13: 867-80
- (27) Ward C, Moffat M. Modulation of sodium-hydrogen exchange activity in cardiac myocytes during acidosis and realkalisation: effects on calcium, pH_i, and cell shortening. *Cardiovascular Research*; 1995 29: 247-253
- (28) Lee JA, Allen DG. Changes in intracellular free calcium concentration during long exposures to stimulated ischemia in isolated mammalian ventricular muscle. *Circ Res* 1992; 71: 58-69
- (29) Shine KI, Douglas AM. Low calcium reperfusion of ischemic myocardium. *J Mol Cell Cardiol* 1983; 15: 251-60
- (30) Miura Y, Kimura J. Sodium-Calcium exchange current. Dependence on internal Ca²⁺ and Na⁺ and competitive binding site of external Na⁺ and Ca²⁺. *J Gen Physiol* 1989; 93: 1129-45

- (31) Gores GJ, Nieminen A-L, Fleishman KE, Dawson TL, Herman B, Lemasters JJ. Extracellular acidosis delays onset of cell death in ATP-deleted hepatocytes. *Am J Physiol* 1988; 255: C315-22
- (32) Bond JM, Herman B, Lemasters JJ. Protection by acidotic pH against anoxia/reoxygenation injury to rat neonatal cardiac myocytes. *Biochem Biophys Res Commun* 1991; 179:798-803
- (33) Bond JM, Chacon E, Herman B, Lemasters JJ. Intracellular pH and Ca^{2+} homeostasis in the pH paradox of reperfusion injury to neonatal rat cardiac myocytes. *Am J Physiol* 1993; 265: C129-37
- (34) Meng H-P, Lonsberry BB, Pierce GN. Influence of perfusate pH on the postischemic recovery of cardiac contractile function: involvement of sodium-hydrogen exchange. *J Pharmacol Exp Ther* 1991; 258: 772-7
- (35) Hearse DJ, Bolli R. Reperfusion induced injury: manifestations, mechanisms, and clinical relevance. *Cardiovasc Res* 1992; 26: 101-8
- (36) Karmazyn M. Ischemic and reperfusion injury in the heart. Cellular mechanisms pharmacological interventions. *Can J Physiol Pharmacol* 1991; 69: 719-30
- (37) Godt RE. A simple electrostatic model can explain the effect of pH upon the force-pCa relation of skinned frog skeletal muscle fibres. *Biophys J* 1981; 35: 385-92
- (38) Jeffray FMH, Mallow CR, Radda GK. Influence of intracellular acidosis on contractile function in the working rat heart. *Am J Physiol* 1987; 253: H1499-505
- (39) Malloy C, Matthews P, Smith M, Radda GK. In vivo ^{31}P NMR study of the regional metabolic responses to cardiac ischemia. *Adv Myocardiol* 1985; 6: 461-4
- (40) van den Bosch H. Intracellular phospholipase A. *Biochim Biophys Acta* 1980; 604: 191-246
- (41) Kinsella JL, Aronson PS. Amiloride inhibition of the Na-H exchanger in renal microvillus membrane vesicles. *Am J Physiol* 1981; 241: F374-9
- (42) Duarte CG, Chomety F, Giebisch G. Effect of amiloride, ouabain and furosemide on distal tubular function in the rat. *Am J Physiol* 1971; 221: 632-9

- (43) Baer, JE, Jones CB, Spitzer SA, Russo HF. The potassium sparing and natriuretic activity of N-aminidino-3,5-diamino-6-chlorpyrazinecarboxamine hydrochloride dihydrate (amiloride hydrochloride). *J Pharmacol Exp Ther* 1967; 157: 472-85
- (44) Benos DJ. Amiloride; a molecular probe of sodium transport in tissues and cells. *Am J Physiol* 1982; 242: C131-45
- (45) Scholz W, Albus U, Lang HJ, et al. Hoe 694, a new Na^+/H^+ exchange inhibitor and its effects in cardiac ischemia. *Br J Pharmacol* 1993; 109: 562-8
- (46) Counillon LT, Scholz W, Lang HJ, Pouyssegur J. Pharmacological characterization of stably transfected Na^+/H^+ antiporter isoforms using amiloride analogs and a new inhibitor exhibiting antiischemic properties. *Mol Pharmacol* 1993; 44: 1041-5
- (47) Hendrikx M, Mubagawa K, Veronck F, et al. New Na^+/H^+ exchange inhibitor HOE 694 improves postischemic function and high energy phosphate resynthesis and reduces Ca^{2+} overload in isolated perfused rabbit heart. *Circulation* 1994; 89: 2787-98
- (48) Scholz W, Albus U, Counillon L, Gögelein H, Lang H-J, Linz W, Weichert A, Schölkens B. Protective effects of HOE 642, a selective sodium-hydrogen exchange subtype 1 inhibitor, on cardiac ischemia and reperfusion. *Cardiovascular Research* 1995; 29: 260-268
- (49) Orlowski J, Kandasamy RA, Shull GE. Molecular cloning of putative members of the Na/H exchanger gene family. *J Biol Chem* 1992; 267: 9331-9
- (50) Scholz W, Albus U. Potential of selective sodium-hydrogen exchange inhibitors in cardiovascular therapy. *Cardiovascular Research* 1995; 29: 184-188
- (51) Khandoudi N, Bernard, Cozzone P, Feuvray D. Intracellular pH and role of Na/H exchange during ischemia and reperfusion of normal and diabetic rat hearts. *Cardiovasc Res* 1990; 24: 873-8
- (52) Vandenberg JI, Metcalfe JC, Grace AA. Mechanisms of pH_i recovery after global ischemia in the perfused heart. *Circ Res* 1993; 72: 993-1003
- (53) E. Bugge, K. Ytrehus. Inhibition of sodium-hydrogen exchange reduces infarct size in the isolated rat heart – a protective additive to ischemic preconditioning. *Cardiovascular Research* 1995; 29: 269-274

- (54) Koike A. 1996. Protective effects of dimethyl amiloride against postischemic myocardial dysfunction in rabbit hearts. *The Journal of Thoracic and Cardiovascular Surgery*, Volume 112, Number 3, Page 765-775
- (55) ML Myers, S Mathur, GH Li, M Karmazyn. Sodium-hydrogen exchange inhibitors improve postischaemic recovery of function in the perfused rabbit heart. *Cardiovascular Research* 1995; 29: 209-214
- (56) Meng HP, Pierce GN. Protective effects of 5-(N₁ N-dimethyl) amiloride on ischemia-reperfusion injury in hearts. *Am J Physiol* 1990; 258: H1615-9
- (57) Duff HJ. Clinical and in vivo antiarrhythmic potential of sodium-hydrogen exchange inhibitors. *Cardiovascular Research* 1995; 29: 189-193
- (58) Scholz W, Albus U, Hropot M, Klaus E, Linz W, Scholkens BA. Zunahme des Na⁺/H⁺-Austausches an Kaninchenerthrozyten unter atherogener Diät. In: Assmann G, Betz E, Heinle H, Schulte H, eds. *Neue Aspekte aus Zellbiologie und Molekulargenetik, Epidemiologie und Klinik*. Wiesbaden: Viehweg, Braunschweig, 1990; 296-302
- (59) Sack S, Mohri M, Schwarz ER, et al. Effects of a new Na⁺/H⁺ antiporter inhibitor on postischemic reperfusion in pig heart. *J Cardiovasc Pharmacol* 1994; 23: 72-8
- (60) Harada K, Franklin A, Johnson RG, Grossman W, Morgan JP. Acidemia and hypernatremia enhance postischemic recovery of excitation-contraction coupling. *Circ Res* 1994; 74: 1197-209
- (61) Moffat MP, Karmazyn M. Protective effects of the potent Na/H exchange inhibitor methylisobutyl amiloride against post-ischemic contractile dysfunction in rat and guinea pig hearts. *J Mol Cell Cardiol* 1993; 25: 959-71
- (62) Khandoudi N, Bernard M, Cozzone P, Feuvray D. Intracellular pH and role of Na⁺/H⁺ exchange during ischaemia and reperfusion of normal and diabetic rat hearts. *Cardiovasc Res* 1990; 24: 873-8
- (63) Pierce GN, Ramjiawann B, Dhalla NS, Ferrari R. Na⁺-H⁺ exchange in cardiac sarcolemmal vesicles isolated from diabetic rats. *Am J Physiol* 1990; 258: H255-61

- (64) Kusama Y, Hearse DJ, Avkiran M. Diabetes and susceptibility to reperfusion-induced ventricular arrhythmias. *J Mol Cell Cardiol* 1992; 24: 411-21
- (65) Meno H, Jarmakani JJ, Philipson KD. Developmental changes of sarcolemmal Na^+/H^+ exchange. *J Mol Cell Cardiol* 1989; 21: 1179-85
- (66) Avkiran M, Goss MW, McGill C, Brooks G. Developmental changes in cardiac Na^+/H^+ exchanger (NHE-1) mRNA expression in the rat. *Circulation* 1993; 88: 2349
- (67) Fliegel L, Haworth RS, Dyck JRB. Characterization of the placental brush border membrane Na^+/H^+ exchanger: identification of a thiol dependent transition in apparent molecular size. *Biochem J* 1993; 289: 101-7
- (68) Krapf R, Pearce D, Lynch C, et al. Expression of rat renal Na^+/H^+ antiporter mRNA levels in response to respiratory and metabolic acidosis. *J Clin Invest* 1991; 87: 747-51
- (69) Horie S, Moe O, Tejedor A, Alpern RJ. Preincubation in acid medium increases Na/H antiporter activity in cultured renal proximal tubule cells. *Proc Natl Acad Sci USA* 1990; 87: 4742-5
- (70) Livne A, Grinstein S, Rothstein A. Characterization of Na^+/H^+ exchange in platelets. *Thromb Haemost* 1987; 58: 971-7
- (71) Karmazyn M, Moffat MP. Role of Na^+/H^+ exchange in cardiac physiology and pathophysiology: mediation of myocardial reperfusion injury by the pH paradox. *Cardiovasc Res* 1993; 27: 915-24
- (72) Fafournoux P, Noel J, Pouyssegur J. Evidence that Na^+/H^+ exchanger isoforms NHE-1 and NHE-3 exists as stable dimers in membranes with a high degree of specificity for homodimers. *J Biol Chem* 1994; 269:2589-96
- (73) Fliegel L, Fröhlich O. The Na^+/H^+ exchanger: an update on structure, regulation and cardiac physiology. *Biochem J* 1993; 296: 273-85
- (74) Wakabayashi S, Sardet C, Fafournoux P, Pouyssegur J. The Na^+/H^+ antiporter cytoplasmic domain mediates growth factors signals and controls „ H^+ sensing“. *Proc Natl Acad Sci USA* 1992; 89: 2424-8

- (75) Tse M, Levine S, Yun C, et al. Structure/function studies of the epithelial isoforms of the mammalian Na^+/H^+ exchanger gene family. *J Membr Biol* 1993; 135: 93-108
- (76) Counillon L, Pouyssegur J. Structure-function studies and molecular regulation of the growth factor activatable sodium-hydrogen exchanger (NHE-1). *Cardiovascular Research* 1995; 29: 147-154
- (77) Kaiser S, Curthoys NP. Effect of pH and bicarbonate on phosphoenolpyruvate carboxykinase and glutaminase mRNA levels in cultured renal epithelial cells. *J Biol Chem* 1991; 266: 9397-402
- (78) Jason R B Dyck, T G Maddaford, G N Pierce, L Fliegel. Induction of expression of the sodium-hydrogen exchanger in rat myocardium. *Cardiovascular Research* 1995; 29: 203-208
- (79) Igarashi P, Freed MI, Ganz MB, Reilly RF. Effects of chronic metabolic acidosis on Na^+/H^+ exchangers in LLC-PK₁ renal epithelial cells. *Am J Physiol* 1992; 263: F83-8
- (80) Fliegel L, Sardet C, Pouyssegur J, Barr A. Identification of the protein and cDNA of the cardiac Na^+/H^+ exchanger. *FEBS Lett* 1991; 279: 25-9
- (81) Moe OW, Tejedor A, Levi M, Seldin DW, Preisig PA, Alpern RJ. Dietary NaCl modulates Na^+/H^+ antiporter activity in renal cortical apical membrane vesicles. *Am J Physiol* 1991; 260: F130-7
- (82) Wallert MA, Fröhlich O. Alpha 1-adrenergic stimulation of Na^+/H^+ exchange in cardiac myocytes. *Am J Physiol* 1992; 263: C1096-102
- (83) Breen TE, Pressler ML. α_1 -Adrenergic stimulation and phorbol esters alter intracellular pH in cardiac Purkinje fibres. (Abstract) *Clin Res* 1988; 36: 226A
- (84) Puceat M, Clement-Chomienne O, Terzic A, Vassort G. α_1 -Adrenoceptor and purinoceptor agonists modulate Na-H antiport in single cardiac cells. *Am J Physiol* 1993; 264: H310-9
- (85) Gambassi G, Lakatta EG, Capogrossi M. α_{1b} -Adrenoceptors decrease cytosolic pH via protein-kinase C mediated inhibition of Na^+/H^+ exchange. (Abstract) *Circulation* 1993; 88: I-382

- (86) Kramer BK, Smith TW, Kelly RA. Endothelin and increased contractility in adult rat ventricular myocytes. Role of intracellular alkalosis induced by activation of the protein-kinase C-dependent Na^+/H^+ exchanger. *Circ Res* 1991; 68: 269-79
- (87) Lagadic-Gossmann D, Vaughan-Jones RD, Buckler KJ. Adrenaline and extracellular ATP switch between two modes of acid extrusion in the guinea-pig ventricular myocyte. *J Physiol (Lond)* 1992; 458: 385-407
- (88) Lagadic-Gossmann D, Vaughan-Jones RD. Coupling of dual acid extrusion in the guinea-pig isolated ventricular myocyte to α_1 -and β -adrenoceptor. *J Physiol (Lond)* 1993; 464: 49-73
- (89) Grace AA, Vandenberg JI. Angiotensin II enhances cardiac pH regulation in the presence of HCO_3^- via three acid-base transport mechanisms. (Abstract) *Circulation* 1993; 88: I-331
- (90) Wehling M. Aldosterone specific membrane receptors, rapid activation of the sodium-hydrogen exchanger, and cardiovascular implications. *Cardiovascular Research* 1995; 29: 167-171
- (91) Yamaji Y, Hayashi M, Iyori M, Katajima W, Saruta T. Chronic DOC treatment enhances Na^+/H^+ exchanger activity of betaintercalated cells in rabbit CCD. *Am J Physiol* 1992; 265: F712-7
- (92) Schömig A, Richardt G. The role of catecholamines in ischemia. *J Cardiovasc Pharmacol* 1990; 16: S105-12
- (93) Karmazyn M. Na^+/H^+ exchange inhibitors reverse lactate-induced depression in postischaemic ventricular recovery. *Br J Pharmacol* 1993; 108: 50-6
- (94) Ferrier GR, Moffat MP, Lukas A. Possible mechanisms of ventricular arrhythmias elicited by ischemia followed by reperfusion. *Circ Res* 1985; 56:184-94
- (95) Mitsuka M, Berk BC. Long-term regulation of Na^+-H^+ exchange in vascular smooth muscle cells – of protein kinase -C. *Am J Physiol* 1991; 260: C562-9

- (96) Berk BC, Elder E, Mitsuka M. Hypertrophy and hyperplasia cause differing effects on vascular smooth muscle cell Na^+/H^+ exchange and intracellular pH. *J Biol. Chem* 1990; 265: 19632-7
- (97) Grinstein S, Rotin D, Mason J. Na^+/H^+ exchange and growth factor-induced cytosolic pH changes. Role in cellular proliferation. *Biochim Biophys Acta* 1989; 988: 73-97
- (98) Mitsuka M, Nagae M, Berk BC. Na^+/H^+ exchange inhibitors decrease neointimal formation after rat carotid injury. Effects on smooth muscle cell migration and proliferation. *Circ Res* 1993; 73: 269-75
- (99) Kranzhofer R, Schirmer J, Schomig A, et al. Suppression of neointimal thickening and smooth muscle cell proliferation after arterial injury in the rat by inhibitors of Na^+/H^+ exchange. *Circ Res* 1993; 73: 264-8
- (100) Grinstein S, ed. *Na^+/H^+ exchange*. Boca Raton: CRC Press, 1988
- (101) Kinsella JL, Aronson P. Interaction of NHE_4^+ and Li^+ with the renal microvillus membrane Na^+/H^+ exchanger. *Am J Physiol* 1981; 241: C220-6
- (102) Dyck JRB, Lopaschuk GD, Fliegel L. Identification of a small Na^+/H^+ exchanger-like message in the rabbit myocardium. *FEBS Lett* 1992; 310: 255-9
- (103) Takaichi K, Wang D, Balkovetz DF, Warnock DG. Cloning sequencing, and expression of Na-H antiporter cDNAs from human tissues. *Am J Physiol* 1992; 267: C1069-76
- (104) Stringham JC, Southard JH, Hegge J, Triemstra L, Fields BL, Belzer FO. Limitation of heart preservation by cold storage. *Transplantation* 1992; 53: 287-94
- (105) Schofer J, Lampe F, Müller-Hansen S, Voigt KD. Anstieg der Kreatinkinase und ihrer mm-Isoformen nach erfolgreicher und unkomplizierter Koronarangioplastie. *Z Kardiol* 1992; 81: 326-30
- (106) Eisenberg JD, Sobel BE, Geltman EM. Differentiation of ischemic from non-ischemic cardiomyopathy with positron emission tomography. *Am J Cardiol* 1987; 59: 1410-14
- (107) Fliegel L, Wang H. Regulation of the Na^+ / H^+ Exchanger in the Mammalian Myocardium. *J Mol Cell Cardiol* 1997; 29: 1991-1999

-
- (108) Pitt Bertram, M.D., Faiez Zannad and Witten Janet, Ph.D. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *The New England Journal of Medicine*, 1999. Volume 341, Number 10, Page 709-717

Das Versuchsvorhaben wurde im Rahmen einer Arzneimittelprüfung beim Ministerium für Natur und Umwelt des Landes Schleswig Holstein in Kiel am 17.05.1995 bekannt gegeben.