

- Allsopp, T.E., S. Wyatt, H.F. Paterson, and A.M. Davies. **1993**. The proto-oncogene bcl-2 can selectively rescue neurotrophic factor- dependent neurons from apoptosis. *Cell* 73, 295-307.
- Alnemri, E.S., D.J. Livingston, D.W. Nicholson, G. Salvesen, N.A. Thornberry, W.W. Wong, and J. Yuan. **1996**. Human ICE/CED-3 protease nomenclature. *Cell* 87, 171.
- Alvarado, C., M.E. Alarcon-Riquelme, J. Alcocer-Varela, and C. Fernandez. **1997**. Participation of the Bcl-2 and Fas molecules in experimental apoptosis of spleen B lymphocytes. *Rev Invest Clin* 49, 171-8.
- Angelidis, C.E., I. Lazaridis, and G.N. Pagoulatos. **1991**. Constitutive expression of heat-shock protein 70 in mammalian cells confers thermoresistance. *Eur J Biochem* 199, 35-9.
- Antonsson, B., F. Conti, A. Ciavatta, S. Montessuit, S. Lewis, I. Martinou, L. Bernasconi, A. Bernard, J.J. Mermod, G. Mazzei, K. Maundrell, F. Gambale, R. Sadoul, and J.C. Martinou. **1997**. Inhibition of Bax channel-forming activity by Bcl-2. *Science* 277, 370-2.
- Arends, M.J., and A.H. Wyllie. **1991**. Apoptosis: mechanisms and roles in pathology. *Int Rev Exp Pathol* 32:223-54.
- Arrigo, A.P. **1998**. Small stress proteins: chaperones that act as regulators of intracellular redox state and programmed cell death. *Biol Chem* 379, 19-26.
- Baffy, G., T. Miyashita, J.R. Williamson, and J.C. Reed. **1993**. Apoptosis induced by withdrawal of interleukin-3 (IL-3) from an IL-3- dependent hematopoietic cell line is associated with repartitioning of intracellular calcium and is blocked by enforced Bcl-2 oncogene production. *J Biol Chem* 268, 6511-9.
- Baldini, N. **1997**. Multidrug resistance--a multiplex phenomenon. *Nat Med* 3, 378-80.
- Bargou, R.C., K. Bommert, P. Weinmann, P.T. Daniel, C. Wagener, M.Y. Mapara, and B. Dorken. **1995**. Induction of Bax-alpha precedes apoptosis in a human B lymphoma cell line: potential role of the bcl-2 gene family in surface IgM-mediated apoptosis. *Eur J Immunol* 25, 770-5.
- Bargou, R.C., C. Wagener, K. Bommert, M.Y. Mapara, P.T. Daniel, W. Arnold, M. Dietel, H. Guski, A. Feller, H.D. Royer, and B. Dorken. **1996**. Overexpression of the death-promoting gene bax-alpha which is downregulated in breast cancer restores sensitivity to different apoptotic stimuli and reduces tumor growth in SCID mice. *J Clin Invest* 97, 2651-9.
- Barry, M.A., C.A. Behnke, and A. Eastman. **1990**. Activation of programmed cell death (apoptosis) by cisplatin, other anticancer drugs, toxins and hyperthermia. *Biochem Pharmacol* 40, 2353-62.
- Bates, S., and K.H. Vousden. **1999**. Mechanisms of p53-mediated apoptosis. *Cell Mol Life Sci* 55, 28-37.
- Benjamin, C.W., R.R. Hiebsch, and D.A. Jones. **1998**. Caspase activation in MCF7 cells responding to etoposide treatment. *Mol Pharmacol* 53, 446-50.
- Bissonnette, R.P., F. Echeverri, A. Mahboubi, and D.R. Green. **1992**. Apoptotic cell death induced by c-myc is inhibited by bcl-2. *Nature* 359, 552-4.

- Bittl, A., M. Nap, W. Jager, F. Cornillie, and N. Lang. **1995**. Immunohistochemical detection of P-glycoprotein on frozen and paraffin- embedded tissue sections of normal and malignant tissues. *Anticancer Res* 15, 1007-14.
- Blagosklonny, M.V. **2000**. Cell death beyond apoptosis. *Leukemia* 14, 1502-8.
- Boise, L.H., M. Gonzalez-Garcia, C.E. Postema, L. Ding, T. Lindsten, L.A. Turka, X. Mao, G. Nunez, and C.B. Thompson. **1993**. bcl-x, a bcl-2-related gene that functions as a dominant regulator of apoptotic cell death. *Cell* 74, 597-608.
- Boldin, M.P., T.M. Goncharov, Y.V. Goltsev, and D. Wallach. **1996**. Involvement of MACH, a novel MORT1/FADD-interacting protease, in Fas/APO-1- and TNF receptor-induced cell death. *Cell* 85, 803-15.
- Boyd, J.M., G.J. Gallo, B. Elangovan, A.B. Houghton, S. Malstrom, B.J. Avery, R.G. Ebb, T. Subramanian, T. Chittenden, R.J. Lutz, and et al. **1995**. Bik, a novel death-inducing protein shares a distinct sequence motif with Bcl-2 family proteins and interacts with viral and cellular survival-promoting proteins. *Oncogene* 11, 1921-8.
- Bradford, M.M. **1976**. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-54.
- Buchholz, T.A., M.M. Weil, M.D. Story, E.A. Strom, W.A. Brock, and M.D. McNeese. **1999**. Tumor suppressor genes and breast cancer. *Radiat Oncol Investig* 7, 55-65.
- Bukau, B., and A.L. Horwich. **1998**. The Hsp70 and Hsp60 chaperone machines. *Cell* 92, 351-66.
- Bump, N.J., M. Hackett, M. Hugunin, S. Seshagiri, K. Brady, P. Chen, C. Ferenz, S. Franklin, T. Ghayur, P. Li, and et al. **1995**. Inhibition of ICE family proteases by baculovirus antiapoptotic protein p35. *Science* 269, 1885-8.
- Cai, J., and D.P. Jones. **1998**. Superoxide in apoptosis. Mitochondrial generation triggered by cytochrome c loss. *J Biol Chem* 273, 11401-4.
- Casciola-Rosen, L., D.W. Nicholson, T. Chong, K.R. Rowan, N.A. Thornberry, D.K. Miller, and A. Rosen. **1996**. Apopain/CPP32 cleaves proteins that are essential for cellular repair: a fundamental principle of apoptotic death. *J Exp Med* 183, 1957-64.
- Cash, C.D. **1997**. Are the reactive oxygen-derived species (ROS) interactive properties of the many therapeutic drugs from various categories pertinent to their beneficial effects? *Gen Pharmacol* 28, 169-75.
- Chinnaiyan, A.M., and V.M. Dixit. **1996**. The cell-death machine. *Curr Biol* 6, 555-62.
- Chinnaiyan, A.M., K. Orth, K. O'Rourke, H. Duan, G.G. Poirier, and V.M. Dixit. **1996**. Molecular ordering of the cell death pathway. Bcl-2 and Bcl-xL function upstream of the CED-3-like apoptotic proteases. *J Biol Chem* 271, 4573-6.
- Chittenden, T., C. Flemington, A.B. Houghton, R.G. Ebb, G.J. Gallo, B. Elangovan, G. Chinnadurai, and R.J. Lutz. **1995**. A conserved domain in Bak, distinct from BH1 and BH2, mediates cell death and protein binding functions. *Embo J* 14, 5589-96.

- Ciocca, D.R., G.M. Clark, A.K. Tandon, S.A. Fuqua, W.J. Welch, and W.L. McGuire. **1993.** Heat shock protein hsp70 in patients with axillary lymph node-negative breast cancer: prognostic implications. *J Natl Cancer Inst* 85, 570-4.
- Cohen, G.M. **1997.** Caspases: the executioners of apoptosis. *Biochem J* 326 (Pt 1), 1-16.
- Cossarizza, A., M. Baccarani-Contri, G. Kalashnikova, and C. Franceschi. **1993.** A new method for the cytofluorimetric analysis of mitochondrial membrane potential using the J-aggregate forming lipophilic cation 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolcarbocyanine iodide (JC-1). *Biochem Biophys Res Commun* 197, 40-5.
- Crompton, M. **1999.** The mitochondrial permeability transition pore and its role in cell death. *Biochem J* 341, 233-49.
- Daniel, P.T., K.T. Pun, S. Ritschel, I. Sturm, J. Holler, B. Dorken, and R. Brown. **1999.** Expression of the death gene Bik/Nbk promotes sensitivity to drug- induced apoptosis in corticosteroid-resistant T-cell lymphoma and prevents tumor growth in severe combined immunodeficient mice. *Blood* 94, 1100-7.
- Daniel, P.T., I. Sturm, S. Ritschel, K. Friedrich, B. Dorken, P. Bendzko, and T. Hillebrand. **1999.** Detection of genomic DNA fragmentation during apoptosis (DNA ladder) and the simultaneous isolation of RNA from low cell numbers. *Anal Biochem* 266, 110-5.
- Daniel, P. T. **2000.** Dissecting the pathways to death. *Leukemia* 14(12), 2035-44.
- Daniel, P. T., T. Wieder, et al. **2001.** The kiss of death: promises and failures of death receptors and ligands in cancer therapy. *Leukemia* 15(7), 1022-32.
- Darzynkiewicz, Z., S. Bruno, G. Del Bino, W. Gorczyca, MA. Hotz, P. Lassota, F. Traganos. **1992.** Features of apoptotic cells measured by flow cytometry. *Cytometry* 13, 795-808.
- Datta, R., Y. Manome, N. Taneja, L.H. Boise, R. Weichselbaum, C.B. Thompson, C.A. Slapak, and D. Kufe. **1995.** Overexpression of Bcl-XL by cytotoxic drug exposure confers resistance to ionizing radiation-induced internucleosomal DNA fragmentation. *Cell Growth Differ* 6, 363-70.
- Datta, S. R., A. Brunet, et al. **1999.** Cellular survival: a play in three Akts. *Genes Dev* 13(22), 2905-27.
- De Jong, D., F.A. Prins, D.Y. Mason, J.C. Reed, G.B. van Ommen, and P.M. Kluin. **1994.** Subcellular localization of the bcl-2 protein in malignant and normal lymphoid cells. *Cancer Res* 54, 256-60.
- Decaudin, D., S. Geley, T. Hirsch, M. Castedo, P. Marchetti, A. Macho, R. Kofler, and G. Kroemer. **1997.** Bcl-2 and Bcl-XL antagonize the mitochondrial dysfunction preceding nuclear apoptosis induced by chemotherapeutic agents. *Cancer Res* 57, 62-7.
- Desagher, S., A. Osen-Sand, et al. **1999.** Bid-induced conformational change of Bax is responsible for mitochondrial cytochrome c release during apoptosis. *J Cell Biol* 144(5), 891-901.

- Deveraux, Q.L., R. Takahashi, G.S. Salvesen, and J.C. Reed. **1997**. X-linked IAP is a direct inhibitor of cell-death proteases. *Nature* 388, 300-4.
- Di Marco, A., T. Dasdia, and W. Pastori. **1984**. Interaction of calcium ions and cAMP on the cytotoxic effect of doxorubicin. *Tumori* 70, 217-21.
- Droin, N., L. Dubrez, et al. **1998**. Upregulation of CASP genes in human tumor cells undergoing etoposide-induced apoptosis. *Oncogene* 16(22), 2885-94.
- Du, C., M. Fang, et al. **2000**. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell* 102(1), 33-42.
- Dubrez, L., I. Savoy, et al. **1996**. Pivotal role of a DEVD-sensitive step in etoposide-induced and Fas-mediated apoptotic pathways. *Embo J* 15(20), 5504-12.
- Dyson, J.E., D.M. Simmons, J. Daniel, J.M. McLaughlin, P. Quirke, and C.C. Bird. **1986**. Kinetic and physical studies of cell death induced by chemotherapeutic agents or hyperthermia. *Cell Tissue Kinet* 19, 311-24.
- Elangovan, B., and G. Chinnadurai. **1997**. Functional dissection of the pro-apoptotic protein Bik. Heterodimerization with anti-apoptosis proteins is insufficient for induction of cell death. *J Biol Chem* 272, 24494-8.
- Ellerby, L.M., H.M. Ellerby, S.M. Park, A.L. Holleran, A.N. Murphy, G. Fiskum, D.J. Kane, M.P. Testa, C. Kayalar, and D.E. Bredesen. **1996**. Shift of the cellular oxidation-reduction potential in neural cells expressing Bcl-2. *J Neurochem* 67, 1259-67.
- Ellis, R.J., and F.U. Hartl. **1996**. Protein folding in the cell: competing models of chaperonin function. *Faseb J* 10, 20-6.
- Ellis, R.J. **1996**. Discovery of molecular chaperones. *Cell Stress Chaperones* 1, 155-60.
- Eskes, R., S. Desagher, B. Antonsson, and J.C. Martinou. **2000**. Bid induces the oligomerization and insertion of Bax into the outer mitochondrial membrane. *Mol Cell Biol* 20, 929-35.
- Essmann, F., T. Wieder, A. Otto, EC. Müller, B. Dörken, PT: Daniel. **2000**. GDP dissociation inhibitor D4-GDI (Rho-GDI 2), but not the homologous rho-GDI 1, is cleaved by caspase-3 during drug induced apoptosis. *Biochem J* 346, 777-83.
- Fairbairn, J.J., M.W. Khan, K.J. Ward, B.W. Loveridge, D.W. Fairbairn, and K.L. O'Neill. **1995**. Induction of apoptotic cell DNA fragmentation in human cells after treatment with hyperthermia. *Cancer Lett* 89, 183-8.
- Farrow, S.N., J.H. White, I. Martinou, T. Raven, K.T. Pun, C.J. Grinham, J.C. Martinou, and R. Brown. **1995**. Cloning of a bcl-2 homologue by interaction with adenovirus E1B 19K. *Nature* 374, 731-3.
- Franke, T. F. and L. C. Cantley. **1997**. Apoptosis. A Bad kinase makes good. *Nature* 390(6656), 116-7.
- Freeman, B.C., M.P. Myers, R. Schumacher, and R.I. Morimoto. **1995**. Identification of a regulatory motif in Hsp70 that affects ATPase activity, substrate binding and interaction with HDJ-1. *Embo J* 14, 2281-92.

- Friedrich, K., T. Wieder, C. Von Haefen, S. Radetzki, R. Janicke, K. Schulze-Osthoff, B. Dorken, and P.T. Daniel. **2001**. Overexpression of caspase-3 restores sensitivity for drug-induced apoptosis in breast cancer cell lines with acquired drug resistance. *Oncogene* 20, 2749-60.
- Friess, H., Z. Lu, H.U. Graber, A. Zimmermann, G. Adler, M. Korc, R.M. Schmid, and M.W. Buchler. **1998**. bax, but not bcl-2, influences the prognosis of human pancreatic cancer. *Gut* 43, 414-21.
- Fuller, K.J., R.D. Issels, D.O. Slosman, J.G. Guillet, T. Soussi, and B.S. Polla. **1994**. Cancer and the heat shock response. *Eur J Cancer* 12, 1884-91.
- Gabai, V.L., I.V. Zamulaeva, A.F. Mosin, Y.M. Makarova, V.A. Mosina, K.R. Budagova, Y.V. Malutina, and A.E. Kabakov. **1995**. Resistance of Ehrlich tumor cells to apoptosis can be due to accumulation of heat shock proteins. *FEBS Lett* 375, 21-6.
- Galen, W.P. **1990**. Hyperthermia and chemotherapy. *Adv Exp Med Biol* 267:209-16.
- Gantchev, T.G., and D.J. Hunting. **1997**. Enhancement of etoposide (VP-16) cytotoxicity by enzymatic and photodynamically induced oxidative stress. *Anticancer Drugs* 8, 164-73.
- Golstein, P. **1997**. Controlling cell death. *Science* 275, 1081-2.
- Gonzalez, D., J.D. van Dijk, and L.E. Blank. **1995**. Radiotherapy and hyperthermia. *Eur J Cancer* 31A, 1351-5.
- Gorman, A.M., A. Samali, A.J. McGowan, and T.G. Cotter. **1997**. Use of flow cytometry techniques in studying mechanisms of apoptosis in leukemic cells. *Cytometry* 29, 97-105.
- Gottesman, M.M., and I. Pastan. **1993**. Biochemistry of multidrug resistance mediated by the multidrug transporter. *Annu Rev Biochem* 62, 385-427.
- Griffon-Etienne, G., J.L. Merlin, and C. Marchal. **1997**. Evaluation of Taxol in head and neck squamous carcinoma multicellular tumor spheroids. *Anticancer Drugs* 8, 48-55.
- Gross, A., X.M. Yin, K. Wang, M.C. Wei, J. Jockel, C. Milliman, H. Erdjument-Bromage, P. Tempst, and S.J. Korsmeyer. **1999**. Caspase cleaved BID targets mitochondria and is required for cytochrome c release, while BCL-XL prevents this release but not tumor necrosis factor-R1/Fas death. *J Biol Chem* 274, 1156-63.
- Guan, R.J., S.F. Moss, N. Arber, S. Krajewski, J.C. Reed, and P.R. Holt. **1996**. 30 KDa phosphorylated form of Bcl-2 protein in human colon. *Oncogene* 12, 2605-9.
- Guo, B., A. Godzik, and J.C. Reed. **2001**. Bcl-G, a novel pro-apoptotic member of the Bcl-2 family. *J Biol Chem* 276, 2780-5.
- Haim, N., J. Roman, J. Nemec, and B.K. Sinha. **1986**. Peroxidative free radical formation and O-demethylation of etoposide(VP- 16) and teniposide(VM-26). *Biochem Biophys Res Commun* 135, 215-20.
- Haldar, S., N. Jena, and C.M. Croce. **1995**. Inactivation of Bcl-2 by phosphorylation. *Proc Natl Acad Sci U S A* 92, 4507-11.
- Hall, P.A., D. Meek, and D.P. Lane. **1996**. p53--integrating the complexity. *J Pathol* 180, 1-5.

- Han, J., P. Sabbatini, and E. White. **1996**. Induction of apoptosis by human Nbk/Bik, a BH3-containing protein that interacts with E1B 19K. *Mol Cell Biol* 16, 5857-64.
- Harmon, B.V., A.M. Corder, R.J. Collins, G.C. Gobe, J. Allen, D.J. Allan, and J.F. Kerr. **1990**. Cell death induced in a murine mastocytoma by 42-47 degrees C heating in vitro: evidence that the form of death changes from apoptosis to necrosis above a critical heat load. *Int J Radiat Biol* 58, 845-58.
- Hecker, D., G. Page, M. Lohrum, S. Weiland, and K.H. Scheidtmann. **1996**. Complex regulation of the DNA-binding activity of p53 by phosphorylation: differential effects of individual phosphorylation sites on the interaction with different binding motifs. *Oncogene* 12, 953-61.
- Hertzberg, R.P., and P.B. Dervan. **1984**. Cleavage of DNA with methidiumpropyl-EDTA-iron(II): reaction conditions and product analyses. *Biochemistry* 23, 3934-45.
- Hettinga, J.V., A.W. Konings, and H.H. Kampinga. **1997**. Reduction of cellular cisplatin resistance by hyperthermia--a review. *Int J Hyperthermia* 13, 439-57.
- Hockenberry, D.M., Z.N. Oltvai, X.M. Yin, C.L. Milliman, and S.J. Korsmeyer. **1993**. Bcl-2 functions in an antioxidant pathway to prevent apoptosis. *Cell* 75, 241-51.
- Hofmann, K., P. Bucher, and J. Tschopp. **1997**. The CARD domain: a new apoptotic signalling motif. *Trends Biochem Sci* 22, 155-6.
- Hollstein, M., B. Shomer, M. Greenblatt, T. Soussi, E. Hovig, R. Montesano, and C.C. Harris. **1996**. Somatic point mutations in the p53 gene of human tumors and cell lines: updated compilation. *Nucleic Acids Res* 24, 141-6.
- Hsu, S.Y., A. Kaipia, E. McGee, M. Lomeli, and A.J. Hsueh. **1997**. Bok is a pro-apoptotic Bcl-2 protein with restricted expression in reproductive tissues and heterodimerizes with selective anti-apoptotic Bcl-2 family members. *Proc Natl Acad Sci U S A* 94, 12401-6.
- Hsu, Y.T., K.G. Wolter, and R.J. Youle. **1997**. Cytosol-to-membrane redistribution of Bax and Bcl-X(L) during apoptosis. *Proc Natl Acad Sci U S A* 94, 3668-72.
- Hu, Y., M.A. Benedict, D. Wu, N. Inohara, and G. Nunez. **1998**. Bcl-XL interacts with Apaf-1 and inhibits Apaf-1-dependent caspase-9 activation. *Proc Natl Acad Sci U S A* 95, 4386-91.
- Hunter, J.J., and T.G. Parslow. **1996**. A peptide sequence from Bax that converts Bcl-2 into an activator of apoptosis. *J Biol Chem* 271, 8521-4.
- Ikeda, K., K. Kajiwara, E. Tanabe, S. Tokumaru, E. Kishida, Y. Masuzawa, and S. Kojo. **1999**. Involvement of hydrogen peroxide and hydroxyl radical in chemically induced apoptosis of HL-60 cells. *Biochem Pharmacol* 57, 1361-5.
- Inohara, N., L. Ding, S. Chen, and G. Nunez. **1997**. harakiri, a novel regulator of cell death, encodes a protein that activates apoptosis and interacts selectively with survival-promoting proteins Bcl-2 and Bcl-X(L). *Embo J* 16, 1686-94.
- Irmler, M., M. Thome, M. Hahne, P. Schneider, K. Hofmann, V. Steiner, J.L. Bodmer, M. Schroter, K. Burns, C. Mattmann, D. Rimoldi, L.E. French, and J. Tschopp. **1997**. Inhibition of death receptor signals by cellular FLIP. *Nature* 388, 190-5.

- Jäättela, M., D. Wissing, K. Kokholm, T. Kallunki, and M. Egeblad. **1998.** Hsp70 exerts its anti-apoptotic function downstream of caspase-3-like proteases. *Embo J* 17, 6124-34.
- Jäättela, M. **1999.** Escaping cell death: survival proteins in cancer. *Exp Cell Res* 248, 30-43.
- Jackson, S., C. Harwood, M. Thomas, L. Banks, and A. Storey. **2000.** Role of Bak in UV-induced apoptosis in skin cancer and abrogation by HPV E6 proteins. *Genes Dev* 14, 3065-73.
- Jacobson, M.D., and G.I. Evan. **1994.** Apoptosis. Breaking the ICE. *Curr Biol* 4, 337-40.
- Johnston, R.N., and B.L. Kucey. **1988.** Competitive inhibition of hsp70 gene expression causes thermosensitivity. *Science* 242, 1551-4.
- Jones, N.A., J. Turner, A.J. McIlwrath, R. Brown, and C. Dive. **1998.** Cisplatin- and paclitaxel-induced apoptosis of ovarian carcinoma cells and the relationship between bax and bak up-regulation and the functional status of p53. *Mol Pharmacol* 53, 819-26.
- Jürgensmeier, J.M., Z. Xie, Q. Devereaux, L. Ellerby, D. Bredesen, and J.C. Reed. **1998.** Bax directly induces release of cytochrome c from isolated mitochondria. *Proc Natl Acad Sci U S A* 95, 4997-5002.
- Kane, D.J., T.A. Sarafian, R. Anton, H. Hahn, E.B. Gralla, J.S. Valentine, T. Ord, and D.E. Bredesen. **1993.** Bcl-2 inhibition of neural death: decreased generation of reactive oxygen species. *Science* 262, 1274-7.
- Kasahara, Y., K. Iwai, A. Yachie, K. Ohta, A. Konno, H. Seki, T. Miyawaki, and N. Taniguchi. **1997.** Involvement of reactive oxygen intermediates in spontaneous and CD95 (Fas/APO-1)-mediated apoptosis of neutrophils. *Blood* 89, 1748-53.
- Kastan, M.B., O. Onyekwere, D. Sidransky, B. Vogelstein, and R.W. Craig. **1991.** Participation of p53 protein in the cellular response to DNA damage. *Cancer Res* 51, 6304-11.
- Kastan, M.B., Q. Zhan, W.S. el-Deiry, F. Carrier, T. Jacks, W.V. Walsh, B.S. Plunkett, B. Vogelstein, and A.J. Fornace, Jr. **1992.** A mammalian cell cycle checkpoint pathway utilizing p53 and GADD45 is defective in ataxia-telangiectasia. *Cell* 71, 587-97.
- Kataoka, T., N. Holler, O. Micheau, F. Martinon, A. Tinel, K. Hofmann, and J. Tschopp. **2001.** Bcl-rambo, a novel Bcl-2 homologue that induces apoptosis via its unique C-terminal extension. *J Biol Chem* 276, 19548-54.
- Kerr, J.F., A.H. Wyllie, and A.R. Currie. **1972.** Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer* 26, 239-57.
- Kerr, J.F., C.M. Winterford, and B.V. Harmon. **1994.** Apoptosis. Its significance in cancer and cancer therapy. *Cancer* 73, 2013-26.
- Kiefer, M.C., M.J. Brauer, V.C. Powers, J.J. Wu, S.R. Umansky, L.D. Tomei, and P.J. Barr. **1995.** Modulation of apoptosis by the widely distributed Bcl-2 homologue Bak. *Nature* 374, 736-9.

- Kischkel, F.C., S. Hellbardt, I. Behrmann, M. Germer, M. Pawlita, P.H. Krammer, and M.E. Peter. **1995**. Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *Embo J* 14, 5579-88.
- Kitanaka, C., T. Namiki, K. Noguchi, T. Mochizuki, S. Kagaya, S. Chi, A. Hayashi, A. Asai, Y. Tsujimoto, and Y. Kuchino. **1997**. Caspase-dependent apoptosis of COS-7 cells induced by Bax overexpression: differential effects of Bcl-2 and Bcl-xL on Bax-induced caspase activation and apoptosis. *Oncogene* 15, 1763-72.
- Kluck, R.M., E. Bossy-Wetzel, D.R. Green, and D.D. Newmeyer. **1997**. The release of cytochrome c from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science* 275, 1132-6.
- Knudson, C.M., K.S. Tung, W.G. Tourtellotte, G.A. Brown, and S.J. Korsmeyer. **1995**. Bax-deficient mice with lymphoid hyperplasia and male germ cell death. *Science* 270, 96-9.
- Knudson, C.M., and S.J. Korsmeyer. **1997**. Bcl-2 and Bax function independently to regulate cell death. *Nat Genet* 16, 358-63.
- Knutsen, T., V.K. Rao, T. Ried, L. Mickley, E. Schneider, K. Miyake, B.M. Ghadimi, H. Padilla-Nash, S. Pack, L. Greenberger, K. Cowan, M. Dean, T. Fojo, and S. Bates. **2000**. Amplification of 4q21-q22 and the MXR gene in independently derived mitoxantrone-resistant cell lines. *Genes Chromosomes Cancer* 27, 110-6.
- Kondo, S., Y. Shinomura, Y. Miyazaki, T. Kiyohara, S. Tsutsui, S. Kitamura, Y. Nagasawa, M. Nakahara, S. Kanayama, and Y. Matsuzawa. **2000**. Mutations of the bak gene in human gastric and colorectal cancers. *Cancer Res* 60, 4328-30.
- Korsmeyer, S.J. **1995**. Regulators of cell death. *Trends Genet* 11, 101-5.
- Korsmeyer, S.J., J.R. Shutter, D.J. Veis, D.E. Merry, and Z.N. Oltvai. **1993**. Bcl-2/Bax: a rheostat that regulates an anti-oxidant pathway and cell death. *Semin Cancer Biol* 4, 327-32.
- Krajewska, M., S.F. Moss, S. Krajewski, K. Song, P.R. Holt, and J.C. Reed. **1996**. Elevated expression of Bcl-X and reduced Bak in primary colorectal adenocarcinomas. *Cancer Res* 56, 2422-7.
- Krajewski, S., C. Blomqvist, K. Franssila, M. Krajewska, V.M. Wasenius, E. Niskanen, S. Nordling, and J.C. Reed. **1995**. Reduced expression of proapoptotic gene BAX is associated with poor response rates to combination chemotherapy and shorter survival in women with metastatic breast adenocarcinoma. *Cancer Res* 55, 4471-8.
- Krajewski, S., A. Hugger, M. Krajewska, J.C. Reed, and J.K. Mai. **1998**. Developmental expression patterns of Bcl-2, Bcl-x, Bax, and Bak in teeth. *Cell Death Differ* 5, 408-15.
- Krajewski, S., M. Krajewska, L.M. Ellerby, K. Welsh, Z. Xie, Q.L. Devereaux, G.S. Salvesen, D.E. Bredesen, R.E. Rosenthal, G. Fiskum, and J.C. Reed. **1999a**. Release of caspase-9 from mitochondria during neuronal apoptosis and cerebral ischemia. *Proc Natl Acad Sci U S A* 96, 5752-7.

- Krajewski, S., M. Krajewska, B.C. Turner, C. Pratt, B. Howard, J.M. Zapata, V. Frenkel, S. Robertson, Y. Ionov, H. Yamamoto, M. Peruchó, S. Takayama, and J.C. Reed. **1999b**. Prognostic significance of apoptosis regulators in breast cancer. *Endocr Relat Cancer* 6, 29-40.
- Krishan, A., K. Paika, and E. Frei, III. **1975**. Cytofluorometric studies on the action of podophyllotoxin and epipodophyllotoxins (VM-26, VP-16-213) on the cell cycle traverse of human lymphoblasts. *J Cell Biol* 66, 521-30.
- Kroemer, G., N. Zamzami, and S.A. Susin. **1997**. Mitochondrial control of apoptosis. *Immunol Today* 18, 44-51.
- Kroemer, G., and J.C. Reed. **2000**. Mitochondrial control of cell death. *Nat Med* 6, 513-9.
- Lämmli, UK. **1970**. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-5.
- Lane, D.P., and P.A. Hall. **1997**. MDM2--arbiter of p53's destruction. *Trends Biochem Sci* 22, 372-4.
- Lazebnik, Y.A., A. Takahashi, R.D. Moir, R.D. Goldman, G.G. Poirier, S.H. Kaufmann, and W.C. Earnshaw. **1995**. Studies of the lamin proteinase reveal multiple parallel biochemical pathways during apoptotic execution. *Proc Natl Acad Sci U S A* 92, 9042-6.
- Lehnert, M. **1998**. Chemotherapy resistance in breast cancer. *Anticancer Res* 18, 2225-6.
- Li, H., H. Zhu, C.J. Xu, and J. Yuan. **1998**. Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell* 94, 491-501.
- Li, P., D. Nijhawan, I. Budihardjo, S.M. Srinivasula, M. Ahmad, E.S. Alnemri, and X. Wang. **1997**. Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell* 91, 479-89.
- Lin, E.Y., A. Orlofsky, H.G. Wang, J.C. Reed, and M.B. Prystowsky. **1996**. A1, a Bcl-2 family member, prolongs cell survival and permits myeloid differentiation. *Blood* 87, 983-92.
- Lindquist, S., and E.A. Craig. **1988**. The heat-shock proteins. *Annu Rev Genet* 22, 631-77.
- Linn, S.C., A.H. Honkoop, K. Hoekman, P. van der Valk, H.M. Pinedo, and G. Giaccone. **1996**. p53 and P-glycoprotein are often co-expressed and are associated with poor prognosis in breast cancer. *Br J Cancer* 74, 63-8.
- Liu, X., H. Zou, C. Slaughter, and X. Wang. **1997**. DFF, a heterodimeric protein that functions downstream of caspase-3 to trigger DNA fragmentation during apoptosis. *Cell* 89, 175-84.
- Lown, J.W., S.K. Sim, K.C. Majumdar, and R.Y. Chang. **1977**. Strand scission of DNA by bound adriamycin and daunorubicin in the presence of reducing agents. *Biochem Biophys Res Commun* 76, 705-10.
- Luo, X., I. Budihardjo, H. Zou, C. Slaughter, and X. Wang. **1998**. Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* 94:481-90.

- Mancini, M., D.W. Nicholson, S. Roy, N.A. Thornberry, E.P. Peterson, L.A. Casciola-Rosen, and A. Rosen. **1998**. The caspase-3 precursor has a cytosolic and mitochondrial distribution: implications for apoptotic signaling. *J Cell Biol* 140, 1485-95.
- Mandal, M., L. Adam, J. Mendelsohn, and R. Kumar. **1998**. Nuclear targeting of Bax during apoptosis in human colorectal cancer cells. *Oncogene* 17, 999-1007.
- Mandic, A., K. Viktorsson, M. Molin, G. Akusjarvi, H. Eguchi, S.I. Hayashi, M. Toi, J. Hansson, S. Linder, and M.C. Shoshan. **2001**. Cisplatin induces the proapoptotic conformation of Bak in a deltaMEKK1-dependent manner. *Mol Cell Biol* 21, 3684-91.
- Martinez, J.D., M.T. Craven, E. Joseloff, G. Milczarek, and G.T. Bowden. **1997**. Regulation of DNA binding and transactivation in p53 by nuclear localization and phosphorylation. *Oncogene* 14, 2511-20.
- Matsushima, M., T. Fujiwara, E. Takahashi, T. Minaguchi, Y. Eguchi, Y. Tsujimoto, K. Sukumari and Y. Nakamura. **1998**. Isolation, mapping, and functional analysis of a novel human cDNA (BNIP3) encoding a protein homologous to human Nip3. *Genes Chrom Cancer* 21, 230-235.
- Matsumoto, H., A. Takahashi, X. Wang, K. Ohnishi, and T. Ohnishi. **1997**. Transfection of p53-knockout mouse fibroblasts with wild-type p53 increases the thermosensitivity and stimulates apoptosis induced by heat stress. *Int J Radiat Oncol Biol Phys* 38, 1089-95.
- McCarthy, N.J., M.K. Whyte, C.S. Gilbert, and G.I. Evan. **1997**. Inhibition of Ced-3/ICE-related proteases does not prevent cell death induced by oncogenes, DNA damage, or the Bcl-2 homologue Bak. *J Cell Biol* 136, 215-27.
- McDonnell, T.J., and S.J. Korsmeyer. **1991**. Progression from lymphoid hyperplasia to high-grade malignant lymphoma in mice transgenic for the t(14; 18). *Nature* 349, 254-6.
- McKay, D.B. **1993**. Structure and mechanism of 70-kDa heat-shock-related proteins. *Adv Protein Chem* 44, 67-98.
- Miller, A.B. **1991**. Epidemiological approaches to primary and secondary prevention of cancer. *J Cancer Res Clin Oncol* 117, 177-85.
- Miller, T.M., K.L. Moulder, C.M. Knudson, D.J. Creedon, M. Deshmukh, S.J. Korsmeyer, and E.M. Johnson, Jr. **1997**. Bax deletion further orders the cell death pathway in cerebellar granule cells and suggests a caspase-independent pathway to cell death. *J Cell Biol* 139, 205-17.
- Minn, A.J., P. Velez, S.L. Schendel, H. Liang, S.W. Muchmore, S.W. Fesik, M. Fill, and C.B. Thompson. **1997**. Bcl-x(L) forms an ion channel in synthetic lipid membranes. *Nature* 385, 353-7.
- Miyashita, T., and J.C. Reed. **1993**. Bcl-2 oncoprotein blocks chemotherapy-induced apoptosis in a human leukemia cell line. *Blood* 81, 151-7.
- Miyashita, T., M. Harigai, M. Hanada, and J.C. Reed. **1994**. Identification of a p53-dependent negative response element in the bcl-2 gene. *Cancer Res* 54, 3131-5.

- Morimoto, R.I., K.D. Sarge, and K. Abravaya. **1992**. Transcriptional regulation of heat shock genes. A paradigm for inducible genomic responses. *J Biol Chem* 267, 21987-90.
- Mosmann, T. **1983**. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65, 55-63.
- Mosser, D.D., A.W. Caron, L. Bourget, C. Denis-Larose, and B. Massie. **1997**. Role of the human heat shock protein hsp70 in protection against stress- induced apoptosis. *Mol Cell Biol* 17, 5317-27.
- Mouridsen, H.T., C. Alftahan, L. Bastholt, J. Bergh, M. Dalmark, S. Eksborg, S. Hellsten, M. Kjaer, C. Peterson, T. Skovsgard, and et al. **1990**. Current status of epirubicin (Farmorubicin) in the treatment of solid tumours. *Acta Oncol* 29, 257-85.
- Muchmore, S.W., M. Sattler, H. Liang, R.P. Meadows, J.E. Harlan, H.S. Yoon, D. Nakagawa, T., H. Zhu, N. Morishima, E. Li, J. Xu, B.A. Yankner, and J. Yuan. **2000**. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 403, 98-103.
- Nagata, S. **1997**. Apoptosis by death factor. *Cell* 88:355-65.
- Nakagawa, T., H. Zhu, N. Morishima, E. Li, J. Xu, B.A. Yankner, and J. Yuan. **2000**. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 403, 98-103.
- Narita, M., S. Shimizu, T. Ito, T. Chittenden, R.J. Lutz, H. Matsuda, and Y. Tsujimoto. **1998**. Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria. *Proc Natl Acad Sci U S A* 95, 14681-6.
- Naundorf, H., E.C. Rewasowa, I. Fichtner, B. Buttner, M. Becker, and M. Gorlich. **1992**. Characterization of two human mammary carcinomas, MT-1 and MT-3, suitable for in vivo testing of ether lipids and their derivatives. *Breast Cancer Res Treat* 23, 87-95.
- Nelson, W.G., and M.B. Kastan. **1994**. DNA strand breaks: the DNA template alterations that trigger p53- dependent DNA damage response pathways. *Mol Cell Biol* 14, 1815-23.
- Ng, F.W., M. Nguyen, T. Kwan, P.E. Branton, D.W. Nicholson, J.A. Cromlish, and G.C. Shore. **1997**. p28 Bap31, a Bcl-2/Bcl-XL- and procaspase-8-associated protein in the endoplasmic reticulum. *J Cell Biol* 139, 327-38.
- Nicholson, D.W., A. Ali, N.A. Thornberry, J.P. Vaillancourt, C.K. Ding, M. Gallant, Y. Gareau, P.R. Griffin, M. Labelle, Y.A. Lazebnik, and et al. **1995**. Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature* 376, 37-43
- Nicholson, D.W., and N.A. Thornberry. **1997**. Caspases: killer proteases. *Trends Biochem Sci* 22, 299-306.
- Nicolaou, K.C., Z. Yang, J.J. Liu, H. Ueno, P.G. Nantermet, R.K. Guy, C.F. Claiborne, J. Renaud, E.A. Couladouros, K. Paulvannan, and et al. **1994**. Total synthesis of taxol. *Nature* 367, 630-4.

- Nicoletti, I., G. Migliorati, MC. Pagliacci, F. Grignani, C. Riccardi. **1991**. A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J Immunol Methods* 139, 271-9.
- Nollen, E.A., J.F. Brunsting, H. Roelofsen, L.A. Weber, and H.H. Kampinga. **1999**. In vivo chaperone activity of heat shock protein 70 and thermotolerance. *Mol Cell Biol* 19, 2069-79.
- Nunez, G., R. Merino, D. Grillot, and M. Gonzalez-Garcia. **1994**. Bcl-2 and Bcl-x: regulatory switches for lymphoid death and survival. *Immunol Today* 15, 582-8.
- O'Connor, L., A. Strasser, L.A. O'Reilly, G. Hausmann, J.M. Adams, S. Cory, and D.C. Huang. **1998**. Bim: a novel member of the Bcl-2 family that promotes apoptosis. *Embo J* 17, 384-95.
- Oda, E., R. Ohki, H. Murasawa, J. Nemoto, T. Shibue, T. Yamashita, T. Tokino, T. Taniguchi, and N. Tanaka. **2000**. Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science* 288, 1053-8.
- Oltvai, Z.N., C.L. Milliman, and S.J. Korsmeyer. **1993**. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 74, 609-19.
- Orth, K., and V.M. Dixit. **1997**. Bik and Bak induce apoptosis downstream of CrmA but upstream of inhibitor of apoptosis. *J Biol Chem* 272, 8841-4.
- Osman, Y., Y. Moriyama, and A. Shibata. **1995**. Enhanced elimination of Ph+ chromosome cells in vitro by combined hyperthermia and other drugs (AZT, IFN-alpha, TNF, and quercetin): its application to autologous bone marrow transplantation for CML. *Exp Hematol* 23, 444-52.
- Packer, M.A., and M.P. Murphy. **1995**. Peroxynitrite formed by simultaneous nitric oxide and superoxide generation causes cyclosporin-A-sensitive mitochondrial calcium efflux and depolarisation. *Eur J Biochem* 234, 31-9.
- Pan, S.S., L. Pedersen, and N.R. Bachur. **1981**. Comparative flavoprotein catalysis of anthracycline antibiotic. Reductive cleavage and oxygen consumption. *Mol Pharmacol* 19, 184-6.
- Pastan, I.H., and M.M. Gottesman. **1988**. Molecular biology of multidrug resistance in human cells. *Important Adv Oncol*, 3-16.
- Pierce, J., and C.H. Suelter. **1977**. An evaluation of the Coomassie brilliant blue G-250 dye-binding method for quantitative protein determination. *Anal Biochem* 81, 478-80.
- Pulleyblank, D.E. **1997**. Of topo and Maxwell's dream. *Science* 277, 648-9.
- Puthalakath, H., D.C. Huang, L.A. O'Reilly, S.M. King, and A. Strasser. **1999**. The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol Cell* 3, 287-96.
- Rampino, N., H. Yamamoto, Y. Ionov, Y. Li, H. Sawai, J.C. Reed, and M. Perucho. **1997**. Somatic frameshift mutations in the BAX gene in colon cancers of the microsatellite mutator phenotype. *Science* 275, 967-9.

- Ray, C.A., R.A. Black, S.R. Kronheim, T.A. Greenstreet, P.R. Sleath, G.S. Salvesen, and D.J. Pickup. **1992**. Viral inhibition of inflammation: cowpox virus encodes an inhibitor of the interleukin-1 beta converting enzyme. *Cell* 69, 597-604.
- Reed, J.C. **1994**. Bcl-2 and the regulation of programmed cell death. *J Cell Biol* 124, :1-6.
- Reed, J.C. **1995**. Bcl-2 family proteins: regulators of chemoresistance in cancer. *Toxicol Lett* 82-83, 155-8.
- Reed, J.C. **1999**. Dysregulation of apoptosis in cancer. *J Clin Oncol* 17, 2941-53.
- Reed, J.C. **2000**. Mechanisms of apoptosis. *Am J Pathol* 157, 1415-30.
- Reers, M., S.T. Smiley, C. Mottola-Hartshorn, A. Chen, M. Lin, and L.B. Chen. **1995**. Mitochondrial membrane potential monitored by JC-1 dye. *Methods Enzymol* 260, 406-17.
- Robertson, J.D., K. Datta, and J.P. Kehrer. **1997**. Bcl-xL overexpression restricts heat-induced apoptosis and influences hsp70, bcl-2, and Bax protein levels in FL5.12 cells. *Biochem Biophys Res Commun* 241, 164-8.
- Rotonda, J., D.W. Nicholson, K.M. Fazil, M. Gallant, Y. Gareau, M. Labelle, E.P. Peterson, D.M. Rasper, R. Ruel, J.P. Vaillancourt, N.A. Thornberry, and J.W. Becker. **1996**. The three-dimensional structure of apopain/CPP32, a key mediator of apoptosis. *Nat Struct Biol* 3, 619-25.
- Rotter, V., R. Aloni-Grinstein, D. Schwartz, N.B. Elkind, A. Simons, R. Wolkowicz, M. Lavigne, P. Beserman, A. Kapon, and N. Goldfinger. **1994**. Does wild-type p53 play a role in normal cell differentiation? *Semin Cancer Biol* 5, 229-36.
- Rowe, T.C., G.L. Chen, Y.H. Hsiang, and L.F. Liu. **1986**. DNA damage by antitumor acridines mediated by mammalian DNA topoisomerase II. *Cancer Res* 46, 2021-6.
- Sakaguchi, Y., L.C. Stephens, M. Makino, T. Kaneko, F.R. Strelbel, L.L. Danhauser, G.N. Jenkins, and J.M. Bull. **1995**. Apoptosis in tumors and normal tissues induced by whole body hyperthermia in rats. *Cancer Res* 55, 5459-64.
- Samali, A., B. Zhivotovsky, D.P. Jones, and S. Orrenius. **1998**. Detection of pro-caspase-3 in cytosol and mitochondria of various tissues. *FEBS Lett* 431, 167-9.
- Sanfilippo, O., E. Ronchi, C. De Marco, G. Di Fronzo, and R. Silvestrini. **1991**. Expression of P-glycoprotein in breast cancer tissue and in vitro resistance to doxorubicin and vincristine. *Eur J Cancer* 27, 155-8.
- Sarafian, T.A., and D.E. Bredesen. **1994**. Is apoptosis mediated by reactive oxygen species? *Free Radic Res* 21, 1-8.
- Schendel, S.L., R. Azimov, K. Pawlowski, A. Godzik, B.L. Kagan, and J.C. Reed. **1999**. Ion channel activity of the BH3 only Bcl-2 family member, BID. *J Biol Chem* 274, 21932-6.
- Schendel, S.L., Z. Xie, M.O. Montal, S. Matsuyama, M. Montal, and J.C. Reed. **1997**. Channel formation by antiapoptotic protein Bcl-2. *Proc Natl Acad Sci U S A* 94:5113-8.
- Schinkel, A.H., S. Kemp, M. Dolle, G. Rudenko, and E. Wagenaar. **1993**. N-glycosylation and deletion mutants of the human MDR1 P-glycoprotein. *J Biol Chem* 268, 7474-81.

- Schlesinger, P.H., A. Gross, X.M. Yin, K. Yamamoto, M. Saito, G. Waksman, and S.J. Korsmeyer. **1997.** Comparison of the ion channel characteristics of proapoptotic BAX and antiapoptotic BCL-2. *Proc Natl Acad Sci U S A* 94, 11357-62.
- Schmitt, E., A. Steyaert, G. Cimoli, and R. Bertrand. **1998.** Bax-alpha promotes apoptosis induced by cancer chemotherapy and accelerates the activation of caspase 3-like cysteine proteases in p53 double mutant B lymphoma Namalwa cells. *Cell Death Differ* 5, 506-16.
- Schmitz, I., H. Walczak, P.H. Krammer, and M.E. Peter. **1999.** Differences between CD95 type I and II cells detected with the CD95 ligand. *Cell Death Differ* 6, 821-2.
- Schulze-Osthoff, K., D. Ferrari, M. Los, S. Wesselborg, and M.E. Peter. **1998.** Apoptosis signaling by death receptors. *Eur J Biochem* 254, 439-59.
- Schulze-Osthoff, K., P.H. Krammer, and W. Droge. **1994.** Divergent signalling via APO-1/Fas and the TNF receptor, two homologous molecules involved in physiological cell death. *Embo J* 13, 4587-96.
- Shchepotin, I.B., V. Soldatenkov, R.R. Buras, R.J. Nauta, M. Shabahang, and S.R. Evans. **1994.** Apoptosis of human primary and metastatic colon adenocarcinoma cell lines in vitro induced by 5-fluorouracil, verapamil, and hyperthermia. *Anticancer Res* 14, 1027-31.
- Shimizu, S., M. Narita, and Y. Tsujimoto. **1999.** Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. *Nature* 399, 483-7.
- Shimizu, S., and Y. Tsujimoto. **2000.** Proapoptotic BH3-only Bcl-2 family members induce cytochrome c release, but not mitochondrial membrane potential loss, and do not directly modulate voltage-dependent anion channel activity. *Proc Natl Acad Sci U S A* 97, 577-82.
- Simonian, P.L., D.A.M. Grillot, R. Merino, and G. Nunez. **1996.** Bax can antagonize Bcl-XL during etoposide and cisplatin-induced cell death independently of its heterodimerization with Bcl-XL. *J Biol Chem* 271, 2764-72.
- Simonian, P.L., D.A. Grillot, and G. Nunez. **1997.** Bak can accelerate chemotherapy-induced cell death independently of its heterodimerization with Bcl-XL and Bcl-2. *Oncogene* 15, 1871-5.
- Skulachev, V.P. **1996.** Why are mitochondria involved in apoptosis? Permeability transition pores and apoptosis as selective mechanisms to eliminate superoxide-producing mitochondria and cell. *FEBS Lett* 397, 7-10.
- Skulachev, V.P. **1998.** Cytochrome c in the apoptotic and antioxidant cascades. *FEBS Lett* 423, 275-80.
- Smith, P.K., R.I. Krohn, G.T. Hermanson, A.K. Mallia, F.H. Gartner, M.D. Provenzano, E.K. Fujimoto, N.M. Goedeke, B.J. Olson, and D.C. Klenk. **1985.** Measurement of protein using bicinchoninic acid. *Anal Biochem* 150, 76-85.
- Solary, E., N. Droin, A. Bettaieb, L. Corcos, M.T. Dimanche-Boitrel, and C. Garrido. **2000.** Positive and negative regulation of apoptotic pathways by cytotoxic agents in hematological malignancies. *Leukemia* 14, 1833-49.

- Somasundaram, K., T.K. MacLachlan, T.F. Burns, M. Sgagias, K.H. Cowan, B.L. Weber, and W.S. el-Deiry. **1999**. BRCA1 signals ARF-dependent stabilization and coactivation of p53. *Oncogene* 18, 6605-14.
- Song, Q., Y. Kuang, V.M. Dixit, and C. Vincenz. **1999**. Boo, a novel negative regulator of cell death, interacts with Apaf-1. *Embo J* 18, 167-78.
- Srinivasula, S.M., M. Ahmad, T. Fernandes-Alnemri, and E.S. Alnemri. **1998**. Autoactivation of procaspase-9 by Apaf-1-mediated oligomerization. *Mol Cell* 1, 949-57.
- Streffer, C. **1985**. Metabolic changes during and after hyperthermia. *Int J Hyperthermia* 1, 305-19.
- Stein, U., W. Walther, M. Lemm, H. Naundorf, and I. Fichtner. **1997**. Development and characterisation of novel human multidrug resistant mammary carcinoma lines in vitro and in vivo. *Int J Cancer* 72, 885-91.
- Strobel, T., S.K. Kraeft, L.B. Chen, and S.A. Cannistra. **1998**. BAX expression is associated with enhanced intracellular accumulation of paclitaxel: a novel role for BAX during chemotherapy-induced cell death. *Cancer Res* 58, 4776-81.
- Strobel, T., Y.T. Tai, S. Korsmeyer, and S.A. Cannistra. **1998b**. BAD partly reverses paclitaxel resistance in human ovarian cancer cells. *Oncogene* 17, 2419-27.
- Strumberg, D., J.L. Nitiss, J. Dong, K.W. Kohn, and Y. Pommier. **1999**. Molecular analysis of yeast and human type II topoisomerases. Enzyme- DNA and drug interactions. *J Biol Chem* 274, 28246-55.
- Sturm, I., C.H. Kohne, G. Wolff, H. Petrowsky, T. Hillebrand, S. Hauptmann, M. Lorenz, B. Dorken, and P.T. Daniel. **1999**. Analysis of the p53/BAX pathway in colorectal cancer: low BAX is a negative prognostic factor in patients with resected liver metastases. *J Clin Oncol* 17, 1364-74.
- Sturm, I., H. Petrowsky, R. Volz, M. Lorenz, S. Radetzki, T. Hillebrand, G. Wolff, S. Hauptmann, B. Dorken, and P.T. Daniel. **2001**. Analysis of p53/BAX/p16(ink4a/CDKN2) in esophageal squamous cell carcinoma: high BAX and p16(ink4a/CDKN2) identifies patients with good prognosis. *J Clin Oncol* 19, 2272-81.
- Stürzbecher, H.W., C. Addison, and J.R. Jenkins. **1988**. Characterization of mutant p53-hsp72/73 protein-protein complexes by transient expression in monkey COS cells. *Mol Cell Biol* 8, 3740-7.
- Sullivan, D.M., K.C. Chow, B.S. Glisson, and W.E. Ross. **1987a**. Role of proliferation in determining sensitivity to topoisomerase II-active chemotherapy agents. *NCI Monogr*, 73-8.
- Sullivan, D.M., M.D. Latham, and W.E. Ross. **1987b**. Proliferation-dependent topoisomerase II content as a determinant of antineoplastic drug action in human, mouse, and Chinese hamster ovary cells. *Cancer Res* 47, 3973-9.
- Susin, S.A., N. Zamzami, M. Castedo, T. Hirsch, P. Marchetti, A. Macho, E. Daugas, M. Geuskens, and G. Kroemer. **1996**. Bcl-2 inhibits the mitochondrial release of an apoptogenic protease. *J Exp Med* 184, 1331-41.

- Susin, S.A., N. Zamzami, M. Castedo, E. Daugas, H.G. Wang, S. Geley, F. Fassy, J.C. Reed, and G. Kroemer. **1997**. The central executioner of apoptosis: multiple connections between protease activation and mitochondria in Fas/APO-1/CD95- and ceramide- induced apoptosis. *J Exp Med* 186, 25-37.
- Susin, S.A., H.K. Lorenzo, N. Zamzami, I. Marzo, C. Brenner, N. Larochette, M.C. Prevost, P.M. Alzari, and G. Kroemer. **1999a**. Mitochondrial release of caspase-2 and -9 during the apoptotic process. *J Exp Med* 189, 381-94.
- Susin, S.A., H.K. Lorenzo, N. Zamzami, I. Marzo, B.E. Snow, G.M. Brothers, J. Mangion, E. Jacotot, P. Costantini, M. Loeffler, N. Larochette, D.R. Goodlett, R. Aebersold, D.P. Siderovski, J.M. Penninger, and G. Kroemer. **1999b**. Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* 397, 441-6.
- Swat, W., L. Ignatowicz, and P. Kisielow. **1991**. Detection of apoptosis of immature CD4+8+ thymocytes by flow cytometry. *J Immunol Methods* 137, 79-87.
- Szabo, I., G. Bathori, F. Tombola, A. Coppola, I. Schmehl, M. Brini, A. Ghazi, V. De Pinto, and M. Zoratti. **1998**. Double-stranded DNA can be translocated across a planar membrane containing purified mitochondrial porin. *Faseb J* 12, 495-502.
- Tai, Y.T., S. Lee, E. Niloff, C. Weisman, T. Strobel, and S.A. Cannistra. **1998**. BAX protein expression and clinical outcome in epithelial ovarian cancer. *J Clin Oncol* 16, 2583-90.
- Takayama, S., Z. Xie, and J.C. Reed. **1999**. An evolutionarily conserved family of Hsp70/Hsc70 molecular chaperone regulators. *J Biol Chem* 274, 781-6.
- Tewari, M., L.T. Quan, K. O'Rourke, S. Desnoyers, Z. Zeng, D.R. Beidler, G.G. Poirier, G.S. Salvesen, and V.M. Dixit. **1995**. Yama/CPP32 beta, a mammalian homolog of CED-3, is a CrmA-inhibitable protease that cleaves the death substrate poly(ADP-ribose) polymerase. *Cell* 81, 801-9.
- Thornberry, N.A., H.G. Bull, J.R. Calaycay, K.T. Chapman, A.D. Howard, M.J. Kostura, D.K. Miller, S.M. Molineaux, J.R. Weidner, J. Aunins, and et al. **1992**. A novel heterodimeric cysteine protease is required for interleukin-1 beta processing in monocytes. *Nature* 356, 768-74.
- Thornberry, N.A., T.A. Rano, E.P. Peterson, D.M. Rasper, T. Timkey, M. Garcia-Calvo, V.M. Houtzager, P.A. Nordstrom, S. Roy, J.P. Vaillancourt, K.T. Chapman, and D.W. Nicholson. **1997**. A combinatorial approach defines specificities of members of the caspase family and granzyme B. Functional relationships established for key mediators of apoptosis. *J Biol Chem* 272, 17907-11.
- Tsujimoto, Y., J. Cossman, E. Jaffe, and C.M. Croce. **1985**. Involvement of the bcl-2 gene in human follicular lymphoma. *Science* 228, 1440-3.
- Tsujimoto, Y. **1997**. Regulation of cell death by Bcl-2 family. *Tanpakushitsu Kakusan Koso* 42, 1644-8.
- Twentyman, P.R. and C.H. Versantvoort. **1996**. Experimental modulation of MRP-mediated resistance. *Eur J Cancer* 32, 1002-9.

- Tyurina, Y.Y., V.A. Tyurin, J.C. Yalowich, P.J. Quinn, H.G. Claycamp, N.F. Schor, B.R. Pitt, and V.E. Kagan. **1995**. Phenoxy radicals of etoposide (VP-16) can directly oxidize intracellular thiols: protective versus damaging effects of phenolic antioxidants. *Toxicol Appl Pharmacol* 131, 277-88.
- Van Bree, C., B. van der Maat, H.M. Ceha, N.A. Franken, J. Haveman, and P.J. Bakker. **1999**. Inactivation of p53 and of pRb protects human colorectal carcinoma cells against hyperthermia-induced cytotoxicity and apoptosis. *J Cancer Res Clin Oncol* 125, 549-55.
- Van der Zee, A.G., S. de Jong, W.N. Keith, H. Hollema, H. Boonstra, and E.G. de Vries. **1994**. Quantitative and qualitative aspects of topoisomerase I and II alpha and beta in untreated and platinum/cyclophosphamide treated malignant ovarian tumors. *Cancer Res* 54, 749-55.
- Vander Heiden, M.G., N.S. Chandel, E.K. Williamson, P.T. Schumacker, and C.B. Thompson. **1997**. Bcl-xL regulates the membrane potential and volume homeostasis of mitochondria. *Cell* 91, 627-37.
- Vander Heiden, M.G., and C.B. Thompson. **1999**. Bcl-2 proteins: regulators of apoptosis or of mitochondrial homeostasis? *Nat Cell Biol* 1, E209-16.
- Vargas-Roig, L.M., F.E. Gago, O. Tello, J.C. Aznar, and D.R. Ciocca. **1998**. Heat shock protein expression and drug resistance in breast cancer patients treated with induction chemotherapy. *Int J Cancer* 79, 468-75.
- Vaux, D.L., S. Cory, and J.M. Adams. **1988**. Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* 335, 440-2.
- Verhagen, A.M., P.G. Ekert, M. Pakusch, J. Silke, L.M. Connolly, G.E. Reid, R.L. Moritz, R.J. Simpson, and D.L. Vaux. **2000**. Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins. *Cell* 102, 43-53.
- Vogelstein, B. **1990**. Cancer. A deadly inheritance. *Nature* 348, 681-2.
- Volm, M. **1998**. Multidrug resistance and its reversal. *Anticancer Res* 18, 2905-17.
- Volm, M., and T. Efferth. **1990**. Relationship of DNA ploidy to chemoresistance of tumors as measured by in vitro tests. *Cytometry* 11, 406-10.
- Wagener, C., R.C. Bargou, P.T. Daniel, K. Bommert, M.Y. Mapara, H.D. Royer, and B. Dorken. **1996**. Induction of the death-promoting gene bax-alpha sensitizes cultured breast-cancer cells to drug-induced apoptosis. *Int J Cancer* 67, 138-41.
- Walker, N.I., R.E. Bennett, and R.A. Axelsen. **1988**. Melanosis coli. A consequence of anthraquinone-induced apoptosis of colonic epithelial cells. *Am J Pathol* 131, 465-76.
- Walker, N.P., R.V. Talanian, K.D. Brady, L.C. Dang, N.J. Bump, C.R. Ferenz, S. Franklin, T. Ghayur, M.C. Hackett, L.D. Hammill, and et al. **1994**. Crystal structure of the cysteine protease interleukin-1 beta-converting enzyme: a (p20/p10)2 homodimer. *Cell* 78, 343-52.
- Walton, M.I., D. Whysong, P.M. O'Connor, D. Hockenberry, S.J. Korsmeyer, and K.W. Kohn. **1993**. Constitutive expression of human Bcl-2 modulates nitrogen mustard and camptothecin induced apoptosis. *Cancer Res* 53, 1853-61.

- Wang, G.Q., B.R. Gastman, E. Wieckowski, L.A. Goldstein, A. Gambotto, T.H. Kim, B. Fang, A. Rabinovitz, X.M. Yin, and H. Rabinowich. **2001**. A role for mitochondrial Bak in apoptotic response to anticancer drugs. *J Biol Chem* 276, 34307-17.
- Wang, Y., and C. Prives. **1995**. Increased and altered DNA binding of human p53 by S and G2/M but not G1 cyclin-dependent kinases. *Nature* 376, 88-91.
- Wang, J.C. **1996**. DNA topoisomerases. *Annu Rev Biochem*, 635-92.
- Wang, K., X.M. Yin, D.T. Chao, C.L. Milliman, and S.J. Korsmeyer. **1996**. BID: a novel BH3 domain-only death agonist. *Genes Dev* 10, 2859-69.
- Wang, Q., G.P. Zambetti, and D.P. Suttle. **1997**. Inhibition of DNA topoisomerase II alpha gene expression by the p53 tumor suppressor. *Mol Cell Biol* 17, 389-97.
- Watt, P.M., and I.D. Hickson. **1994**. Structure and function of type II DNA topoisomerases. *Biochem J* 303, 681-95.
- Wieder, T., F. Essmann, A. Prokop, K. Schmelz, K. Schulze-Osthoff, R. Beyaert, B. Dörken, P.T. Daniel. **2001**. Activation of caspase-8 in drug-induced apoptosis of B-lymphoid cells is independent of CD95/Fas receptor-ligand interaction and occurs downstream of caspase-3. *Blood* 97, 1378-87.
- Wilson, K.P., J.A. Black, J.A. Thomson, E.E. Kim, J.P. Griffith, M.A. Navia, M.A. Murcko, S.P. Chambers, R.A. Aldape, S.A. Raybuck, and et al. **1994**. Structure and mechanism of interleukin-1 beta converting enzyme. *Nature* 370, 270-5.
- Wyllie, A.H., J.F. Kerr, and A.R. Currie. **1980**. Cell death: the significance of apoptosis. *Int Rev Cytol*, 251-306.
- Xanthoudakis, S., S. Roy, D. Rasper, T. Hennessey, Y. Aubin, R. Cassady, P. Tawa, R. Ruel, A. Rosen, and D.W. Nicholson. **1999**. Hsp60 accelerates the maturation of pro-caspase-3 by upstream activator proteases during apoptosis. *Embo J* 18, 2049-56.
- Xiang, J., D.T. Chao, and S.J. Korsmeyer. **1996**. BAX-induced cell death may not require interleukin 1 beta-converting enzyme-like proteases. *Proc Natl Acad Sci U S A* 93, 14559-63.
- Yamamoto, H., H. Sawai, T.K. Weber, M.A. Rodriguez-Bigas, and M. Perucho. **1998**. Somatic frameshift mutations in DNA mismatch repair and proapoptosis genes in hereditary nonpolyposis colorectal cancer. *Cancer Res* 58, 997-1003.
- Yang, E., J. Zha, J. Jockel, L.H. Boise, C.B. Thompson, and S.J. Korsmeyer. **1995**. Bad, a heterodimeric partner for Bcl-XL and Bcl-2, displaces Bax and promotes cell death. *Cell* 80, 285-91.
- Yang, E., and S.J. Korsmeyer. **1996**. Molecular thanatopsis: a discourse on the BCL2 family and cell death. *Blood* 88, 386-401.
- Yang, J., X. Liu, K. Bhalla, C.N. Kim, A.M. Ibrado, J. Cai, T.I. Peng, D.P. Jones, and X. Wang. **1997**. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked [see comments]. *Science* 275, 1129-32.

- Yen, H.C., T.D. Oberley, C.G. Gairola, L.I. Szweda, and D.K. St Clair. **1999**. Manganese superoxide dismutase protects mitochondrial complex I against adriamycin-induced cardiomyopathy in transgenic mice. *Arch Biochem Biophys* 362, 59-66.
- Yin, C., C.M. Knudson, S.J. Korsmeyer, and T. Van Dyke. **1997**. Bax suppresses tumorigenesis and stimulates apoptosis in vivo. *Nature* 385, 637-40.
- Yin, X.M., Z.N. Oltvai, D.J. Veis-Novack, G.P. Linette, and S.J. Korsmeyer. **1994**. Bcl-2 gene family and the regulation of programmed cell death. *Cold Spring Harb Symp Quant Biol* 59, 387-93.
- Yonezawa, M., T. Otsuka, N. Matsui, H. Tsuji, K.H. Kato, A. Moriyama, and T. Kato. **1996**. Hyperthermia induces apoptosis in malignant fibrous histiocytoma cells in vitro. *Int J Cancer* 66, 347-51.
- Yuan, J., and H.R. Horvitz. **1992**. The *Caenorhabditis elegans* cell death gene ced-4 encodes a novel protein and is expressed during the period of extensive programmed cell death. *Development* 116, 309-20.
- Zamzami, N., P. Marchetti, M. Castedo, C. Zanin, J.L. Vayssiere, P.X. Petit, and G. Kroemer. **1995**. Reduction in mitochondrial potential constitutes an early irreversible step of programmed lymphocyte death in vivo. *J Exp Med* 181, 1661-72.
- Zamzami, N., S.A. Susin, P. Marchetti, T. Hirsch, I. Gomez-Monterrey, M. Castedo, and G. Kroemer. **1996**. Mitochondrial control of nuclear apoptosis. *J Exp Med* 183, 1533-44.
- Zamzami, N., P. Marchetti, M. Castedo, T. Hirsch, S.A. Susin, B. Masse, and G. Kroemer. **1996**. Inhibitors of permeability transition interfere with the disruption of the mitochondrial transmembrane potential during apoptosis. *FEBS Lett* 384, 53-7.
- Zamzami, N., C. Brenner, I. Marzo, S.A. Susin, and G. Kroemer. **1998**. Subcellular and submitochondrial mode of action of Bcl-2-like oncoproteins. *Oncogene* 16, 2265-82.
- Zapata, J.M., M. Krajewska, S. Krajewski, R.P. Huang, S. Takayama, H.G. Wang, E. Adamson, and J.C. Reed. **1998**. Expression of multiple apoptosis-regulatory genes in human breast cancer cell lines and primary tumors. *Breast Cancer Res Treat* 47, 129-40.
- Zha, H., C. Aime-Sempe, T. Sato, and J.C. Reed. **1996**. Proapoptotic protein Bax heterodimerizes with Bcl-2 and homodimerizes with Bax via a novel domain (BH3) distinct from BH1 and BH2. *J Biol Chem* 271, 7440-4.
- Zhou, Q., S. Snipas, K. Orth, M. Muzio, V.M. Dixit, G.S. Salvesen. **1997**. Target protease specificity of the viral serpin CRM A; analysis of five caspases. *J Biol Chem* 272, 7797-7800.
- Zou, H., W.J. Henzel, X. Liu, A. Lutschg, and X. Wang. **1997**. Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* 90, 405-13.