

7 Literatur

- Abbiw, D. K. (1990). *Useful plants of Ghana*, Intermediate Technology Publications and the Royal Botanic Gardens Kew, London, S.154–157.
- Aboagye–Nyame, F. (1995). *Systematic steps and procedures for accelerated development of traditional medicine*. Ghana Federation of Traditional Healers. Unpublished seminar report, Accra (Ghana), S. 60–64.
- Adesanya, S. A. (1988). Antisickling activity of *Adansonia digitata*. *Planta Medica* **54**, 374.
- Adesina, S. K., Reisch, J. (1985). A triterpenoid glykoside from *Tetrapleura tetraptera* fruit. *Phytochemistry* **24**, 3003–3006.
- Adewunmi, C. O., Furu, P. (1989). Evaluation of aridanin, a glycoside, and aridan, an aqueous extract of *Tetrapleura tetraptera* fruit, on *Schistosoma mansoni* and *S. bovis*. *Journal of Ethnopharmacology* **27**, 277–283.
- Almeida–Alves, T. M., Nagem, T. J., de Carvalho, L. Z., Krettli, A. U., Zani, C. L. (1997). Antiplasmodial triterpenes from *Vernonia brasiliensis*. *Planta Medica* **63**, 554–555.
- Ang, K. K. H., Holmes, M. J., Higa, T., Hamann, M. T., Kara, U. A. K. (2000). *In vivo* antimalarial activity of the beta–carboline alkaloid manzamine A. *Antimicrobial Agents and Chemotherapy* **44**, 1645–1649.
- Angerhofer, C. K., König, G. M., Wright, A. D., Sticher, O., Milhous, W. K., Cordell, G. A., Farnsworth, N. R., Pezzuto, J. M. (1992). Novel antimalarial compounds. In: *Advances in natural product chemistry* (Herausgeber: C. Atta–ur–Rhaman), Harwood Academic, Chur, S. 311–329.
- Angus, B. J. (2001). Malaria on the world wide web. *Clinical Infectious Diseases* **33**, 651–661.
- Anthonsen, T., Bergland, G. (1971). The diterpenoids of some *Solidago* species. *Acta Chemica Scandinavica* **25**, 1924–1925.
- Aquino, R., D’Agostino, M., De Simone, F., Pizza, C. (1988). 4–Arylcoumarin glycosides from *Coutarea hexandra*. *Phytochemistry* **27**, 1827–1830.
- Azuine, M. A. (1998). *Cancer and Ethnobotany of Nigeria*. Shaker Verlag Aachen.
- Babady–Bila, T., Gedris, E., Herz, W. (1996). Niruroidine, a norsecurinine–type alkaloid from *Phyllanthus niruroides*. *Phytochemistry* **41**, 1441–1443.
- Banerji, A., Chintalwar, G. J., Joshi, N. K., Chadha, M. S. (1971). Isolation of ecdysterone from Indian plants. *Phytochemistry* **10**, 2225–2226.

- Baslas, R. K., Agarwal, R. (1980). Chemical investigation of some anticancer plants of *Euphorbia* genus. *Indian Journal of Chemistry* **19**, 717–718.
- Baytelman, B. (1979). *Etnobotanica en el estado de Morelos*. Instituto Nacional de Antropología e Historia, Centro Regional de Morelos, Cuernavaca.
- Berendsohn, W. G. (2001). Botanischer Garten und Botanisches Museum Berlin–Dahlem, Berlin, persönliche Mitteilung.
- Belmain, S. R., Neal, G. E., Ray, D. E., Golob, P. (2001). Insecticidal and vertebrate toxicity associated with ethnobotanicals used as post-harvest protectants in Ghana. *Food and Chemical Toxicology* **39**, 287–291.
- Benoit, F., Valentin, A., Péliissier, Y., Marion, C., Dakuyo, Z., Mallié, M., Bastide, J. M. (1995). Antimalarial activity *in vitro* of *Cochlospermum tinctorium* tubercle extracts. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **89**, 217–218.
- Beuze, R. (1973). La santé par les plantes des antilles francaises. Emile Desormeaux, Pointe-a-pitre, Guadeloupe, S. 80.
- Bhacca, N. S. (1963). *High resolution NMR–spectra catalog*. Varian Associates, Palo Alto, S. 267.
- Bharel, S., Gulati, A., Abdin, M. Z., Srivastava P. S., Jain, S. K. (1996). Structure, biosynthesis and functions of artemisinin. *Fitoterapia* **67**, 387–402.
- Boeker, R., Bohlmann, F., King, R. M. (1986). Umbelliferone derivatives from *Heterothalamus spartioides*. *Revista Latinoamericana de Química* **17**, 47–48.
- Bohlmann, F., Niedballa, U., Rode, K. M. (1966). Über neue Polyine mit C₁₇–Kette. *Chemische Berichte* **99**, 3552–3553.
- Bohlmann, F., Zdero, C. (1969). Über ein neues Coniferylalkohol–Derivat aus *Blumea lacera*. *Tetrahedron Letters* **2**, 69–70.
- Bohlmann, F., Zdero, C. (1977). Neue Germacranolide aus *Calea zacatechichi*. *Phytochemistry* **16**, 1065–1068.
- Bohlmann, F., Fritz, U. (1978). Neue Diterpene und Acetylenverbindungen aus *Nidorella*–Arten. *Phytochemistry* **17**, 1769–1772.
- Bohlmann, F., Fritz, U., King, R. M., Robinson, H. (1981a). Fourteen heliangolides from *Calea* species. *Phytochemistry* **20**, 743–749.
- Bohlmann, F., Zhong–Liang, C., Schuster, A. (1981b). Aromatic esters from *Solidago decurrens*. *Phytochemistry* **20**, 2601–2602.

- Bokkenheuser, V. D., Shackleton, C. H. L., Winter, J. (1987). Hydrolysis of dietary flavonoid glycosides by strains of intestinal *Bacteroides* from humans. *Biochemical Journal* **248**, 953–956.
- Bork, P. M., Schmitz, M. L., Kuhnt, M., Escher, C., Heinrich, M. (1997). Sesquiterpene lactones containing Mexican Indian medicinal plants and pure sesquiterpene lactones as potent inhibitors of transcription factor NF- κ B. *FEBS Letter* **402**, 85–90.
- Bray, D. H., Warhurst, D. C., Connolly, J. D., O'Neill, M. J., Phillipson, J. D. (1990). Plants as sources of antimalarial drugs. Part 7. Activity of some species of Meliaceae plants and their constituent limonoids. *Phytotherapy Research* **4**, 29–34.
- Bray, P. G., Boulter, M. K., Ritchie, G. Y., Howells, R. E., Ward, S. A. (1994). Relationship of global chloroquine transport and reversal of resistance in *Plasmodium falciparum*. *Molecular and Biochemical Parasitology* **63**, 87–94.
- Brieskorn, C. H., Michel, H. (1968). Flavone aus dem Blatt von *Rosmarinus officinalis* L. *Tetrahedron Letters* **30**, 3447–3448.
- Bringmann, G., Saeb, W., Aké-Assi, L., François, G., Narayanan, A. S. S., Peters, K., Peters, E. M. (1997). Betulinic acid: isolation from *Triphyophyllum peltatum* and *Ancistrocladus heyneanus*, antimalarial activity, and crystal structure of the benzyl ester. *Planta Medica* **64**, 54–57.
- Brown, R. E., Stancato, F. A., Wolfe, A. D. (1979). The effects of mefloquine on *Escherichia coli*. *Life Sciences* **25**, 1857–1864.
- Bruce, T. B. F. (1998, 2000, 2001). Phytotherapeut in Accra–North Kaneshie, Accra (Ghana), persönliche Mitteilung.
- Bruce-Chwatt, L. J. (1963). A longitudinal survey of malaria infection in a group of West African adults. *West African Medicinal Journal* **12**, 141–173.
- Bunnag, D., Karbwang, J., Na-Bangchang, K., Thanavibul, A., Chittamas, S., Harinasuta, T. (1996). Quinine-tetracycline for multidrug resistant falciparum malaria. *Southeast Asian Journal of Tropical Medicine and Public Health* **27**, 15–18.
- Cabantchik, Z. I., Glickstein, H., Golenser, J., Loyevsky, M., Tsafack, A. (1996). Iron chelators: mode of action as antimalarials. *Acta Haematologica* **95**, 70–77.
- Calera, M. R., Mata, R., Lotina-Hennsen, B., Anaya, A. L. (1996). Uncoupling behavior of the 4-phenylcoumarins in spinach chloroplasts: structure–activity relationships. *Journal of Agricultural and Food Chemistry* **44**, 2966–2969.
- Campos-Neves, M. T., Campos-Neves, A. (1966). Composition of the bark of the trunk and branches of *Hymenodictyon floribundum* of Angola. *Garcia Orta* **14**, 401–410.

- Cheng, D., Röder, E. (1986). Pyrrolizidin–Alkaloide aus *Emilia sonchifolia*. *Planta Medica* **6**, 484–486.
- Cocker, W., Moore, L., Pratt, A. C. (1965). Dextrorotatory hardwickiic acid. An extractive of *Copaifera officinalis*. *Tetrahedron Letters* **24**, 1983–1985.
- Crabbe, P., Azpeitia, E. A., Djerassi, C. (1961). Steroids and terpenoids of different species of *Iresine*. *Bulletin des Sociétés Chimiques Belges* **70**, 168–188.
- Crew, M. C., Melgar, M. D., George, S., Greenough, R. C., Szpiech, J. M., DiCarlo, F. J. (1976). Metabolism of 3–(hydroxymethyl)–8–methoxychromone in the rat. II. Classification and identification of urinary drug metabolites. *Xenobiotica* **6**, 89–97.
- Cubukcu, B., Bray, D. H., Warhurst, D. H., Mericli, A. H., Ozhatay, N., Sariyar, G. (1990). In vitro antimalarial activity of crude extracts and compounds from *Artemisia abrotanum*. *Phytotherapy Research* **4**, 203–204.
- D'Agostino, M., De Feo, V., De Simone, F., Pizza, C. (1989). 4–Arylcoumarins from *Coutarea hexandra*. *Phytochemistry* **28**, 1773–1774.
- Delle Monache, G., Botta, B., Neto, A. S., De Lima, R. A. (1983). 4–Arylcoumarins from *Coutarea hexandra*. *Phytochemistry* **22**, 1657–1658.
- Delle Monache, G., Botta, B., De Lima, R. A. (1984). A 4–Arylcoumarin from *Coutarea hexandra*. *Phytochemistry* **23**, 1813.
- Delle Monache, G., Botta, B., Vinciguerra, V., Pinheiro, R. M. (1990). 4–Arylcoumarins from *Coutarea latiflora*. *Phytochemistry* **29**, 3984–3986.
- Desjardins, R. E., Canfield, C. J., Chulay, J. D. (1979). Quantitative assessment of anti-malarial activity *in vitro* by a semiautomated microdilution technique. *Antimicrobial Agents and Chemotherapy* **16**, 710–718.
- Desjardins, R. E., Pamplin, C. L. 3rd, von Bredow, J., Barry, K. G., Canfield, C. J. (1979). Kinetics of a new antimalarial, mefloquine. *Clinical Pharmacology and Therapeutics* **26**, 372–379.
- De Vries, P. J., Dien, T. K. (1996). Clinical pharmacology and therapeutic potential of artemisinin and its derivatives in the treatment of malaria. *Drugs* **52**, 818–836.
- Divo, A. A., Geary, T. G., Jensen, J. B., Ginsburg, H. (1985). The mitochondrion of *Plasmodium falciparum* visualized by rhodamine 123 fluorescence. *Journal of Protozoology* **32**, 442–446.
- Dokosi, O. B. (1998). *Herbs of Ghana*. Ghana Universities Press Accra, Accra (Ghana), S. 313–314.

- Eich, E., Eichberg, D. (1982). Zur antibakteriellen Wirkung von Clavinalkaloiden und deren partialsynthetischen Derivaten. *Planta Medica* **45**, 146–147.
- Eich, E., Becker, C., Müller, W. E. G. (1984). Zur zytostatischen Wirkung von Festuclavin und seinen Derivaten. *Farmaceutisch Tijdschrift voor België* **61**, 267.
- Eich, E., Becker, C., Eichberg, D., Maidhof, A., Müller, W. E. G. (1986). Clavines as antitumor agents, 5: *In vitro* activity of semisynthetic festuclavines. *Pharmazie* **41**, 156–157.
- Eich, E. (1992). Ergolin-Derivate—»schmutzige«, spezifische und selektive Arzneistoffe. *Pharmazeutische Zeitung* **22**, 9–22.
- Eiden, F. (1998). Ausflug in die Vergangenheit: Chinin und andere Chinaalkaloide. 1. Teil: Von der Isolierung der Chinolinalkaloide bis zur Konstitutionsaufklärung. *Pharmazie in unserer Zeit* **6**, 257–271.
- Eiden, F. (1999). Ausflug in die Vergangenheit: Chinin und andere Chinaalkaloide. 3. Teil: Vom Weg zur Totalsynthese der Chinolin-Chinaalkaloide über die Herstellung besser wirksamer Malariamittel bis zur Erforschung der Indol-Chinaalkaloide. *Pharmazie in unserer Zeit* **2**, 74–86.
- Elford, B. C., Roberts, M. F., Phillipson, J. D., Wilson, R. J. M. (1987). Potentiation of the antimalarial activity of qinghaosu by methoxylated flavones. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **81**, 434–436.
- El Izzi, A., Benie, T., Thieulant, M. L., Duval, J. (1990). Inhibitory effects of saponins from *Tetrapleura tetraptera* on the LH release by cultured rat pituitary cells. *Planta Medica* **56**, 357–359.
- El-Seedi, H., Ghia, F., Torssell, K. B. G. (1994). Cadinane sesquiterpenes from *Siparuna macrotrepala*. *Phytochemistry* **35**, 1495–1497.
- Faizi, S., Siddiqui, B. S., Saleem, R., Siddiqui, S., Aftab, K., Gilani A. (1995). Fully acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa oleifera*. *Phytochemistry* **38**, 957–963.
- Fletcher, K. A., Evans, D. A., Gilles, H. M., Greaves, J., Bunnag, D., Harinasuta, T. (1981). Studies on the pharmacokinetics of primaquine. *Bulletin of the World Health Organization* **59**, 407–12.
- François, G., Aké Assi, L., Holenz, J., Bringmann, G. (1996). Constituents of *Picralima nitida* display pronounced inhibitory activities against asexual erythrocytic forms of *Plasmodium falciparum* *in vitro*. *Journal of Ethnopharmacology* **54**, 113–117.
- François, G., Timperman, G., Eling, W., Aké-Assi L., Holenz, J., Bringmann, G. (1997).

- Naphthylisoquinoline alkaloids against malaria: evaluation of the curative potential of dioncophylline C and dioncopeltine A against *Plasmodium berghei* *in vivo*. *Antimicrobial Agents and Chemotherapy* **41**, 2533–2539.
- Gilles, H. M. (Hrsg.) (1991). *Management of severe and complicated malaria*. WHO, Genf.
- Ginsburg, H., Ward, S.A., Bray, P.G. (1999). An integrated model of chloroquine action. *Parasitology Today* **15**, 357–360.
- Graefe, E. U., Derendorf, H., Veit, M. (1999). Pharmacokinetics and bioavailability of the flavonol quercetin in humans. *International Journal of Clinical Pharmacology and Therapeutics* **37**, 219–233.
- Greenman, J. M. (1907). *New and noteworthy spermatophytes from Mexico, Central America, and the West Indies*. Field Columbian Museum–Botany, Band 2. S. 247–287.
- Griffiths, L. A., Smith, G. E. (1972). Metabolism of apigenin and related compounds in the rat. *Biochemical Journal* **128**, 901–911.
- Gunstone, F. D., Harwood, J. L., Padley, F. P. (1995). *The Lipid Handbook*. Chapman & Hall, London.
- Halazy, S., Perez, M., Fourrier, C., Pallard, I., Pauwels, P. J., Palmier, C., Joun, G. W., Valentin, J., Bonnafous, R., Martinez, J. (1996). Serotonin dimers: application of the bivalent ligand approach to the design of new potent and selective 5-HT_{1B/1D} agonists. *Journal of Medicinal Chemistry* **39**, 4920–4927.
- Harborne, J. B. (1967). *Comparative Biochemistry of the flavonoids*. Academic Press, London, New York.
- Harborne, J. B., Williams, C. A. (1975). Flavone and flavonol glycosides. In: *The flavonoids* (Herausgeber: J. B. Harborne, T. J. Mabry, H. Mabry), Chapman & Hall, London, S. 376–441.
- Heilmann, J., Merfort, I. (1998). Aktueller Kenntnisstand zum Metabolismus von Flavonoiden I. Resorption und Metabolismus von Flavonolen. *Pharmazie in unserer Zeit* **2**, 58–64.
- Heinrich, M. (1989). Ethnobotanik der Tieflandmixe (Oaxaca, Mexico) und phytochemische Untersuchung von *Capraria biflora* L. (Scrophulariaceae). Dissertationes Botanicae No. 144. J. Cramer in Gebr. BorntraegerVerlagsbuchhandlung, Berlin, Stuttgart.
- Hellwig, B. (2001). Artemether / Lumefantrin. *Neue Arzneimittel (Deutsche Apotheker*

- Zeitung–Beilage*) **48**, 94–97.
- Heymann, H., Tezuka, Y., Kikuchi, T., Supriyatna, S. (1994). Constituents of *Sindora sumatrana* MIQ. III.: New *trans*-clerodane diterpenoids from the dried pods. *Chemical & Pharmaceutical Bulletin* **42**, 1202–1207.
- Herz, W., Narendra, K. (1980) Sesquiterpene lactones of *Calea zacatechichi* and *C. urticiflora*. *Phytochemistry* **19**, 593–597.
- Herz, W., Kulanthaivel, P. (1985). Diterpenes and sesquiterpene lactones from *Mikania congesta*. *Phytochemistry* **24**, 1761–1768.
- Hider, R. C., Liu, Z. (1997). The treatment of malaria with iron chelators. *Journal of Pharmacy and Pharmacology* **49**, 59–64.
- Hildebrand, M. R., Willuhn, G., Schmidt, T. J. (2000). New Dihydrobenzofurans and further constituents from roots of *Microglossa pyrifolia*. Poster, International Congress and 48th Annual Meeting of the Society for Medicinal Plant Research (GA), Zürich, Abstract Book, P1A/14.
- Hill, A. V., Allsopp, C. E., Kwiatkowski, D., Anstey, N. M., Twumasi, P., Rowe, P. A., Bennett, S., Brewster, D., McMichael, A. J., Greenwood, B. M. (1991). Common west African HLA antigens are associated with protection from severe malaria. *Nature* **352**, 595–600.
- Houghton, P. J., Lala, P. K., Shellard, E. J., Spong, K. (1976). The alkaloids of *Mitragyna stipulosa* (D.C.) O. Kuntze. *Journal of Pharmacy and Pharmacology* **28**, 664.
- Hudson, A. T. (1993). Atovaquone—a novel broad-spectrum anti-infective drug. *Parasitology Today* **9**, 66–68.
- Ito, A., Chai, H., Shin, Y. G., García, R., Mejía, M., Gao, Q., Fairchild, C. R., Lane, K. E., Menendez, A. T., Farnsworth, N. R., Cordell, A. G., Pezzuto, J. M., Kinghorn, A. D. (2000). Cytotoxic constituents of the roots of *Exostema acuminatum*. *Tetrahedron* **56**, 6401–6405.
- Irvine, F. R. (1961). *Woody plants of Ghana*. Oxford University Press, London, S. 678.
- Iwu, M. M., Obidoa, O., Anazodo, M. (1986). Biochemical mechanism of the antimalarial activity of *Azadirachta indica*. *Pharmacological Research Communications* **18**, 81–90.
- Jain, S. C., Jain, R., Mascolo, N., Capasso, F., Vijayvergia, R., Sharma, R. A., Mittal, C. (1998). Ethnopharmacological evaluation of *Pergularia daemia* (Forsk.) Chiov. *Phytotherapy Research* **12**, S. 378–380.
- Jenett–Siems, K. (2002). Untersuchung ethnomedizinisch genutzter afrikanischer und

- mittelamerikanischer Arzneipflanzen im Hinblick auf Phytochemie und *in-vitro*-Aktivität gegen *Plasmodium falciparum*. Habilitation, Fachbereich Pharmazie, Freie Universität Berlin.
- Johns, T., Faubert, G. M., Kokwaru, J. O., Mahunnah, R. L. A., Kimanami, E. K. (1995). Anti-giardical activity of gastrointestinal remedies of the Luo of East-Africa. *Journal of Ethnopharmacology* **46**, 17–23.
- Kamchonwongpaisan, S., Nilanonta, C., Tarnchompoo, B., Thebtaranonth, C., Thebtaranonth, Y., Yuthavong, Y., Kongsaree, P., Clardy, J. (1995). An antimalarial peroxide from *Amomum krervanh* Pierre. *Tetrahedron Letters* **36**, 1821–1824.
- Kapadia, G. J., Angerhofer, C. K., Ansa-Asamoah, R. (1993). Akuammine: an antimalarial indolemonoterpene alkaloid of *Picralima nitida* seeds. *Planta Medica* **59**, 565–566.
- Kasper, R. (1991). Zur Entwicklung onkostatisch wirksamer Festuclavin-Derivate. Dissertation, Institut für Pharmazie, Freie Universität Berlin.
- Keawpradub, N., Kirby, G. C., Steele, J. C. P., Houghton, P. J. (1999). Antiplasmodial activity of extracts and alkaloids of three *Alstonia* species from Thailand. *Planta Medica* **65**, 690–694.
- Khalid, S. A., Farouk, A., Geary, T. G., Jensen, J. B. (1986). Potential antimalarial candidates from African plants: an *in vitro* approach using *Plasmodium falciparum*. *Journal of Ethnopharmacology* **15**, 201–209.
- Khoo, S. F. (1973). Structure and stereochemistry of the diterpenes of *Hymenaea courbaril* (Caesalpinioideae) seed pod resin. *Tetrahedron* **29**, 3379–3388.
- Kirby, G. C., Paine, A., Warhurst, D. C. (1995). *In vitro* and *in vivo* antimalarial activity of cryptolepine, a plant-derived indoloquinoline. *Phytotherapy Research* **9**, 359–363.
- Köhler, I. (1999). Traditionelle Medizin in Ghana. *Zeitschrift für Phytotherapie* **20**, 22–24.
- Köhler, I., Jenett-Siems, K., Bienzle, U., Eich, E. (2000). *In vitro* antiplasmodial activity of natural and semi-synthetic ergoline derivatives. Poster, International Congress and 48th Annual Meeting of the Society for Medicinal Plant Research (GA), Zürich, Abstract Book, P2A/51.
- Kokwaro, J. O. (1976). *Medicinal plants of East Africa*. East African Literature Bureau, Nairobi, Kenia, S. 134.
- Koranteng, G. D. (1998). Phytotherapeut in Accra North, Accra (Ghana), persönliche Mitteilung.

- Kraft, C., Jenett-Siems, K., Siems, K., Jakupovic, J., Mavi, S., Bienzle, U., Eich, E. (2002). *In vitro* antiplasmodial evaluation of medicinal plants from Zimbabwe. *Phytotherapy Research*, in press.
- Krämer, S. (1999). Absorption prediction from physicochemical parameters. *Pharmaceutical Sciences & Technology Today* **2**, 373–380.
- Křen, V., Weignerová, L., Kuzma, M., Jegorov, A., Sedmera, P. (2001). Semisynthetic dimers of antiparkinsonic ergot alkaloids. *Heterocycles* **55**, 1045–1056.
- Křen, V., Fišerová, A., Weignerová, L., Stibor, I., Halada, P., Sedmera, P., Pospíšil, M. (2002). Clustered ergot alkaloids modulate cell-mediated cytotoxicity. *Bioorganic Medicinal Chemistry* **10**, 415–424.
- Kretschmer, H., Bienzle, U., Klauß, V., Kreamsner, P. G., Leichsenring, M. (1996). Malaria. In: *Tropen- und Reisemedizin* (Herausgeber: J. Knobloch), Gustav Fischer Verlag, Jena, 134–163.
- Krugliak, M., Deharo, E., Shalmiev, G., Sauvain, M., Moretti, C., Ginsburg, H. (1995). Antimalarial effects of C18 fatty acids on *Plasmodium falciparum* in culture and on *Plasmodium vinckei petteri* and *Plasmodium yoelii nigeriensis* in vivo. *Experimental Parasitology* **81**, 97–105.
- Kuiate, J. R., Zollo, P. H. A., Nguefa, E. H., Bessiere, J. M., Lamaty, G., Menut, C. (1999). Composition of the essential oils from leaves of *Microglossa pyrifolia* (Lam.) O. Kuntze and *Helichrysum odoratissimum* (L.) Less. growing in Cameroon. *Flavour & Fragrance Journal* **14**, 82–84.
- Kumaratilake, L. M., Robinson, B. S., Ferrante, A., Poulos, A. (1992). Antimalarial properties of n-3 and n-6 polyunsaturated fatty acids: *in vitro* effects on *Plasmodium falciparum* and *in vivo* effects on *P. berghei*. *Journal of Clinical Investigation* **89**, 961–967.
- Kusano, A., Nikaido, T., Kuge, T., Ohmoto, T., Monache, G. D., Botta, B., Botta, M., Saitoh, T. (1991). Inhibition of adenosine 3',5'-cyclic monophosphate phosphodiesterase by flavonoids from licorice roots and 4-arylcoumarins. *Chemical & Pharmaceutical Bulletin* **39**, 930–933.
- Lipinsky, C. A., Lombardo, F., Dominy, B. W., Feeney, P. J. (1997). Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Advanced Drug Delivery Reviews* **23**, 3–25.
- Liu, K. C. S., Yang, S. L., Roberts, M. F., Elford, B. C., Phillipson, J. D. (1989). The contribution of flavonoids to the antimalarial activity of *Artemisia annua*. *Planta*

- Medica* **55**, 654–655.
- Liu, S. Y., Sporer, F., Wink, M., Jourdane, J., Hennig, R., Li, Y. L., Ruppel, A. (1997). Anthraquinones in *Rheum palmatum* and *Rumex dentatus* (Polygonaceae), and phorbol esters in *Jatropha curcas* (Euphorbiaceae) with molluscicidal activity against the schistosome vector snails *Oncomelania*, *Biomphalaria* and *Bulinus*. *Tropical Medicine & International Health* **2**, 179–188.
- Lucius, R. (1997). *Parasitologie: Grundlagen für Biologen, Mediziner, Veterinärmediziner*. Spektrum Akademischer Verlag GmbH Heidelberg, Berlin, S. 97–112.
- Luzzi, G. A., Peto, T. E. A. (1993). Adverse effects of antimalarials. An update. *Drug Safety* **8**, 295–311.
- Mambu, L., Martin, M., Razafimahefa, D., Ramanitrahasimbola, D., Rasoanaivo, P., Frappier, F. (2000). Spectral characterisation and antiplasmodial activity of bisbenzylisoquinolines from *Isolona ghesquiereina*. *Planta Medica* **66**, 537–540.
- Martinez, M. (1959). *Las Plantas Medicinales de Mexico*. 4th ed. Ediciones Botas, Mexico City, S. 347–349.
- Mata, R., Calzada, F., Garcia, M. R., Reguero, M. T. (1987). Chemical studies on Mexican plants used in traditional medicine, III: New 4-phenylcoumarins from *Exostema caribaeum*. *Journal of Natural Products* **50**, 866–871.
- Mata, R., Calzada, F., Garcia, M. R. (1988). Chemical studies on Mexican plants used in traditional medicine, VI. Additional new 4-phenylcoumarins from *Exostema caribaeum*. *Journal of Natural Products* **51**, 851–856.
- Mata, R., Albor, C., Pereda-Miranda, R., McLaughlin, J. (1990). Cytotoxic constituents of *Exostema mexicanum*. *Planta Medica*, **56**, 241.
- Matsuda, H., Li, Y., Murakami, T., Matsumura, N., Yamahara, J., Yoshikawa, M. (1998). Antidiabetic principles of natural medicines. III. Structure-related inhibitory activity and action mode of oleanolic acid glycosides on hypoglycemic activity. *Chemical & Pharmaceutical Bulletin* **46**, 1399–1403.
- Mavi, S. (1997). Department of Pharmacy, University of Zimbabwe, Harare (Zimbabwe), persönliche Mitteilung.
- Mayagoitia, L., Díaz, J. L., Contreras, C. M. (1986). Psychopharmacologic analysis of an alleged oneirogenic plant: *Calea zacatechichi*. *Journal of Ethnopharmacology* **18**, 229–243.
- Mc Dowell, T., Bremer B., (1998). Implications of morphological and molecular analyses.

- Plant Systematics & Evolution* **212**, 215–246.
- Melzig, M. F. (2002). Institut für Pharmazie, Humboldt Universität zu Berlin, Berlin, persönliche Mitteilung.
- Menzies, J. R., Paterson, S. J., Duwiejua, M., Corbett, A. D. (1998). Opioid activity of alkaloids extracted from *Picralima nitida* (fam. Apocynaceae). *European Journal of Pharmacology* **29**, 101–108.
- Meshnick, S. R., Taylor, T. E., Kamchonwongpaisan, S. (1996). Artemisinin and the antimalarial endoperoxides: from herbal remedy to targeted chemotherapy. *Microbiological Reviews* **60**, 301–315.
- Metz, G. (2000). Strukturvielfalt aus Obst und Gemüse. *Pharmazeutische Zeitung* **26**, 24–26.
- Milliken, W. (1997). Malaria and antimalarial plants in Roraima, Brazil. *Tropical Doctor* **27** (Supplement I), 20–25.
- Mizokawa, T., Akai, T., Nakata, Y., Yamaguchi, M., Nakagawa, H., Hasan, S., Rattig, K. J., Wachtel, H. (1993). Terguride as a new anti-hyperprolactinemic agent: characterization in rats and dogs in comparison with bromocriptine. *Japanese Journal of Pharmacology* **63**, 269–278.
- Mockenhaupt, F.P. (1995). Reviews: mefloquine resistance in *Plasmodium falciparum*. *Parasitology Today* **11**, 248–253.
- Mockenhaupt, F.P., Thies, F.L., Meyer, C.G. (1998). Malaria im Kindesalter. *Pädiatrische Praxis* **55**, 305–318.
- Mockenhaupt, F. P. (2000). Institut für Tropenmedizin, Medizinische Fakultät Charité, Humboldt Universität zu Berlin, Berlin, persönliche Mitteilung.
- Morton, J. F., (1981). *Atlas of Medicinal Plants of Middle America. Bahamas to Yucatan* (Herausgeber: C. Thomas), Springfield (USA).
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* **65**, 55–63.
- Müller, M., Heide, L. (2001). Malaria in Afrika: Wie hilfreich sind Arzneipflanzen? *Zeitschrift für Phytotherapie* **22**, 77–81.
- Muñoz, V., Sauvain, M., Bourdy, G., Callapa, J., Bergeron, S., Rojas, I., Bravo, J. A., Balderrama, L., Ortiz, B., Gimez, A. Deharo E. (2000a). The search for natural bioactive compounds through a multidisciplinary approach in Bolivia. Part I. Evaluation of the antimalarial activity of plants used by the Chacobo Indians. *Journal*

- of Ethnopharmacology* **69**, 127–137.
- Muñoz, V., Sauvain, M., Bourdy, G., Callapa, J., Rojas, I., Vargas, L., Tae, A. Deharo E. (2000b). The search for natural bioactive compounds through a multidisciplinary approach in Bolivia. Part II. Antimalarial activity of some plants used by Mosekene indians. *Journal of Ethnopharmacology* **69**, 139–155.
- Murakami, H., Kitazono, Y., Jiwajinda, S., Koshimizu, K., Ohigashzi, H. (1998). Niaziminin, a thiocarbamate from the leaves of *Moringa oleifera*, holds a strict structural requirement for inhibition of tumor–promoter–induced Epstein–Barr virus activation. *Planta Medica* **64**, 319–323.
- Mustofa, Valentin, A., Benoit–Vical, F., Péliissier, Y., Koné–Bamba, D., Mallié, M. (2000). Antiplasmodial activity of plant extracts used in west African traditional medicine. *Journal of Ethnopharmacology* **73**, 145–151.
- Mutschler, E. (1996). *Arzneimittelwirkungen*. 6. Auflage, Wissenschaftliche Verlagsgesellschaft mbH Stuttgart.
- Nakanishi, T., Ogaki, I. A., Murata, H., Nishi, M. (1985). Flavonoids of *Striga asiatica*. *Journal of Natural Products* **48**, 491–493.
- Neuwinger, H. D. (1998). *Afrikanische Arzneipflanzen und Jagdgifte*. 2. Auflage, Wissenschaftliche Verlagsgesellschaft mbH Stuttgart.
- Neuwinger, H. D. (2000). *African Traditional Medicine*. Medpharm GmbH, Stuttgart.
- Nkunya, M. H. H., Weenen, H., Bray, D. H., Mgani, Q. A., Mwasumbi, L. B. (1991). Antimalarial activity of Tanzanian plants and their active constituents: The genus *Uvaria*. *Planta Medica* **57**, 341–343.
- Noster, S., Kraus, L. (1990). *In vitro* antimalarial activity of *Coutarea latiflora* and *Exostema caribaeum* extracts on *Plasmodium falciparum*. *Planta Medica* **56**, 63–65.
- Nurse, G.T. (1979). Iron, the thalassaemias, and malaria. *The Lancet* **3**, 938–940.
- Olliaro, P. L., Yuthavong, Y. (1999). An overview of chemotherapeutic targets of antimalarial drug discovery. *Pharmacology & Therapeutics* **1**, 91–110.
- Olliaro, P. L., Haynes, R. K., Meunier, B., Yuthavong, Y. (2001). Possible modes of action of the artemisinin–type compounds. *Trends in Parasitology* **17**, 122–126.
- Oliver–Bever, B. (1982). Medicinal plants in tropical West Africa I. Plants acting on the cardiovascular system. *Journal of Ethnopharmacology* **5**, 1–71.
- Onegi, B. (1999). Department of Pharmacy (Pharmacognosy Unit), Makerere University, Kampala (Uganda), persönliche Mitteilung.

- O'Neill, M. J., Bray, D. H., Boardmann, P., Phillipson, J. D., Warhurst, D. C. (1985). Plants as sources of antimalarial drugs part 1. *In vitro* method for the evaluation of crude extracts from plants. *Planta Medica* **51**, 394–398.
- O'Neill, M. J., Phillipson, J. D. (1989). Plants as sources of antimalarial compounds. *Revista Latinoamericana de Química* **20**, 111–118.
- Ortega, A., Vivar, R., Díaz, E., Romo, J. (1970). Determinación de las estructuras de la calaxina y de la ciliarina, nuevos germacranólidos furanónicos. *Revista Latinoamericana de Química* **1**, 81–85.
- Osifo, N. G. (1984). Chloroquine-induced pruritus among patients with malaria. *Archives of Dermatology* **120**, 80–82.
- Panchagnula, R., Thomas, N. S. (2000). Biopharmaceutics and pharmacokinetics in drug research. *International Journal of Pharmaceutics* **201**, 131–150.
- Paulo, A., Gomes, E. T., Steele, J., Warhurst, D. C., Houghton, P. J. (2000). Antiplasmodial activity of *Cryptolepis sanguinolenta* alkaloids from leaves and roots. *Planta Medica* **66**, 30–34.
- Peters, W. (1998). Drug resistance in malaria parasites of animals and man. *Advances in Parasitology* **41**, 1–62.
- Petersen, E., Hogh, B., Marbiah, N. T., Dolopaie, E., Gottschau, A., Hanson, A. P., Bjorkman, A. (1991). Clinical and parasitological studies on malaria in Liberian adults living under intense malaria transmission. *Annals of Tropical Medicine and Parasitology* **85**, 577–84.
- Pongprayoon, U., Baeckström, P., Jacobsson, U., Lindström, M., Bohlin, L. (1992). Antispasmodic activity of β -damascenone and *E*-phytol isolated from *Ipomoea pes-caprae*. *Planta Medica* **58**, 19–21.
- Portoghese, P.S. (1992). The role of concepts in structure–activity relationship studies of opioid ligands. *Journal of Medicinal Chemistry* **35**, 1927–1937.
- Pradines, B., Ramiandrasoa, F., Basco, L. K., Bricard, L., Kunesch, G., Le Bras, J. (1996). *In vitro* activities of novel catecholate siderophores against *Plasmodium falciparum*. *Antimicrobial Agents and Chemotherapy* **40**, 2094–2098.
- Quadri-Spinelli, T., Heilmann, J., Rali, T., Sticher, O. (2000). Bioactive coumarin derivatives from the fern *Cyclosorus interruptus*. *Planta Medica* **66**, 728–733.
- Quijano, L., Calderón, J. S., Ríos, T. (1977). Los componentes de la *Calea zacatechichi* estructura de los caleocromenos A y B. *Revista Latinoamericana de Química* **8**, 90–93.

- Quijano, L., Vivar, R., Ríos, T. (1978). Los componentes de la *Calea zacatechichi* estructura de los caleinas A y B. *Revista Latinoamericana de Química* **9**, 86–89.
- Quijano, L., Vivar, R., Ríos, T. (1979). Revision of the structures of caleine A and B, germacranolide sesquiterpenes from *Calea zacatechichi*. *Phytochemistry* **18**, 1745–1747.
- Radloff, P. D., Phillips, J., Nkeyi, M., Hutchinson, D., Kremsner, P. G. (1996). Atovaquone and proguanil for *Plasmodium falciparum* malaria. *The Lancet* **347**, 1511–1513.
- Ramesh, D., Dennis, T. J., Shingare, M. S. (1992). Constituents of *Adansonia digitata* root bark. *Fitoterapia* **63**, 278–279.
- Ramos, R. R., Alarcon–Aguilar, F., Lara–Lemus, A., Flores–Saenz, J. L. (1992). Hypoglycemic effect of plants used in Mexico as antidiabetics. *Archives of Medical Research* **23**, 59–64.
- Rao, C. V. K., Ganapaty, S. (1983). Investigation on *Euphorbia pilulifera* L. *Fitoterapia* **6**, 265–269.
- Rašková, H., Čelada, L., Lavický, J., Vaneček, J., Urbanová, Z., Křeček, J., Priborska, Z., Elis, J., Krejčí, I. (1987). Pharmacological interventions to antagonize stress–induced immune consequences. *Annals of the New York Academy of Sciences* **496**, 436–446.
- Rasoanaivo, P., Ratsimamanga–Urverg, S., Milijaona, R., Rafatro, H., Rakoto–Ratsimamanga, A., Galeffi, C., Nicoletti, M. (1994). *In vitro* and *in vivo* chloroquine–potentiating action of *Strychnos myrtoides* alkaloids against chloroquine–resistant strains of *Plasmodium* Malaria. *Planta Medica* **60**, 13–16.
- Rätsch, C. (1998). *Enzyklopädie der psychoaktiven Pflanzen*. 3. Auflage, AT Verlag Aarau, S. 116–118.
- Ratsimamanga–Urverg, S., Rasoanaivo, P., Ramiaramanana, L., Milijaona, R., Rafatro, H., Verdier, F., Rakoto–Ratsimamanga, A., Le Bras, J. (1992). *In vitro* antimalarial activity and chloroquine–potentiating action of two bisbenzylisoquinoline enantiomer alkaloids isolated from *Strychnos thouarsii* and *Spirospermum penduliflorum*. *Planta Medica* **58**, 540–543.
- Ravelonjato, B., Libot, F., Ramiandrasoa, F., Kunesch, N., Gayral, P., Poisson, J. (1992). Molluscicidal constituents of *Calophyllum* from Madagascar: activity of some natural and synthetic neoflavonoids and khellactones. *Planta Medica* **58**, 51–55.
- Reddy, K. L. N., Reddy, S. P., Sharma, G. V. M. (1994). Synthesis of 13–Hydroxy–9Z,11E,15Z–octadecatrienoic acid. *Synthetic Communications* **24**, 1395–1401.

- Reher, G., Kraus, L., (1982). A neoflavonoid from *Coutarea latiflora* (Rubiaceae). *Phytochemistry* **22**, 1524–1525.
- Reher, G., Kraus, L., (1984). New neoflavonoids from *Coutarea latiflora*. *Journal of Natural Products* **47**, 172–174.
- Reinhard, E. (2001). *Pharmazeutische Biologie*. 6. Auflage, Wissenschaftliche Verlagsgesellschaft mbH Stuttgart. S. 549.
- Rinne, U.K. (1986). Dopamine agonists as primary treatment in Parkinson's disease. *Advances in Neurology* **45**, 519–523.
- Ross R.(1910) *The prevention of malaria*. John Murray London.
- Rücker, G., Kehrenbaum, S., Sakulas, H., Lawong, B., Goeltenboth, F. (1992). Acetylenic glycosides from *Microglossa pyrifolia*. *Planta Medica* **58**, 266–269.
- Rücker, G., Kehrenbaum, S., Sakulas, H., Lawong, B., Goeltenboth, F. (1994). Acetylated aurone glycosides from *Microglossa pyrifolia*. *Planta Medica* **60**, 288–289.
- Ruwende, C., Khoo, S. C., Snow, R. W., Yates, S. N., Kwiatkowski, D., Gupta, S., Warn, P., Allsopp, C. E., Gilbert, S. C., Peschu, N. (1995). Natural selection of hemi- and heterozygotes for G6PD deficiency in Africa by resistance to severe malaria. *Nature* **376**, 246–249.
- Sam, G. H. (1998). Pharmakologie am Centre for Scientific Research into Plant Medicine (CSRPM) Mampong–Akwapuem, Mampong–Akwapuem (Ghana), persönliche Mitteilung.
- Sánchez–Viesca, F. (1969). The structure of exostemin, a new 4–phenylcoumarin isolated from *Exostema caribaeum*. *Phytochemistry* **8**, 1821–1823.
- Sanz, J. F., Marco, J. A. (1990). Synthesis of sinapyl alcohol diisovalerate, a new phenylpropanoid from *Artemisia assoana*. *Journal of Natural Products* **53**, 1034–1035.
- Scheline, R. R. (1991). *CRC Handbook of mammalian metabolism of plant compounds*. CRC Press, Boca Raton, Ann Arbor, Boston, S. 255–287.
- Seshadri, T. R., Videeswaran, S. (1971). Chemical constituents of *Daemia extensa* roots. *Current Science* **40**, 594–595.
- Silva, M., Mundaca, J. M., Sammes, P. G. (1971). Flavonoid and triterpene constituents of *Baccharis rhomboidalis*. *Phytochemistry* **10**, 1942–1943.
- Singh, P., Jain, S., Jakupovic, J. (1988). Clerodane derivatives from *Grangea maderaspatana*. *Phytochemistry* **27**, 1537–1539.
- Sittie, A. A., Lemmich, E., Olsen, C. E., Hviid, L., Kharazmi, A., Nkrumah, F. K.,

- Christensen, S. B. (1999). Structure–activity studies: *In vitro* antileishmanial and antimalarial activities of anthraquinones from *Morinda lucida*. *Planta Medica* **65**, 259–261.
- Slater, T. F., Sawyer, B., Strauli, U. D. (1963). Studies on succinate–tetrazolium reductase systems III. Points of coupling of four different tetrazolium salts. *Biochimica et Biophysica Acta* **77**, 383–393.
- Srinivasan, K. K., Sankara Subbramanian, S. (1980). Chemical investigation of *Emilia sonchifolia*. *Fitoterapia* **51**, 241–243.
- Stahl, E. (1967). *Dünnschichtchromatographie*. Springer Verlag, Basel, Heidelberg, New York, S. 87.
- Standley, P. C. (1926). *Trees and shrubs of Mexico*. United States National Herbarium Part 5, Number 23. Smithsonian Institute Washington D.C., 1313–1721
- Standley, W., Williams, C. O. (1975). Flora of Guatemala–Part XI, Numbers 1 to 3. *Fieldiana: Botany* **24**, 1–274.
- Stauffacher, D., Tschertter, H., Hofmann, A. (1965). Isolierung von Ergosin und Ergosinin neben Agroclavin aus den Samen von *Ipomoea argyrophylla* VATKE (Convolvulaceae). *Helvetica Chimica Acta* **48**, 1379–1380.
- Steele, J. C. P., Warhurst, D. C., Kirby, G. C., Simmonds, M. S. J. (1999). *In Vitro* and *in vivo* evaluation of betulinic acid as an antimalarial. *Phytotherapy Research* **13**, 115–119.
- Subramanian S. S., Nagarajan, S. (1968). Phytochemical studies on *Crotalaria laburnifolia*. *Planta Medica* **16**, 432–435.
- Talapatra, B., Basak, A., Goswami, S., Talapatra, S. K. (1981). Terpenoids and related compounds. XIX. 3- β -Hydroxyfriedelan-27-on, a new triterpene from *Pergularia extensa* N. E. Brown. *Indian Journal of Chemistry* **20B**, 249–250.
- Tan, R. X., Zheng, W. F., Tang, H. Q. (1998). Biologically active substances from the genus *Artemisia*. *Planta Medica* **64**, 295–302.
- Tandon, S., Rastogi, R. P. (1979). Strictic acid, a novel diterpene from *Conyza stricta*. *Phytochemistry* **18**, 494–495.
- Teuschner, E. (1997). *Biogene Arzneimittel*. 5. Auflage, Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart.
- Thebtaranonth, C., Thebtaranonth, Y., Wanauppathamkul, S., Yuthavong, Y. (1995). Antimalarial sesquiterpenes from tubers of *Cyperus rotundus*: structure of 10,12-peroxycalamenene, a sesquiterpene endoperoxide. *Phytochemistry* **40**, 125–128.

- Tchuendem, M. H. K., Mbah, J. A., Tsopmo, A., Ayafor, J. F., Sterner, O., Okunjic, C. C., Iwu, M. M., Schuster, B. M. (1999). Antiplasmodial sesquiterpenoids from the African *Reneilmia cincinnata*. *Phytochemistry* **52**, 1095–1099.
- The International Plant Names Index, Collaboration of The Royal Botanic Gardens Kew, The Harvard University Herbaria, and the Australian National Herbarium (2001). <http://www.ipni.org/>, Zugriff am 15. 03. 2002.
- Tona, L., Ngimbi, N. P., Tsakala, M., Messia, K., Cimaga, K., Apers, S., De Bruyne, T., Pieters, L., Totte, J., Vlietnick, A. J. (1999). Antimalarial activity of 20 crude extracts from nine African medicinal plants used in Kinshasa, Congo. *Journal of Ethnopharmacology* **68**, 193–203.
- Toon, S., Rowland, M. (1983). Structure–pharmacokinetic relationships among the barbiturates in the rat. *Journal of Pharmacology and Experimental Therapeutics* **225**, 752–763.
- Trager, W., Jensen, J. B. (1976). Human malaria parasites in continuous culture. *Science* **193**, 673–675.
- Traore, F., Gasquet, M., Laget, M., Guiraud, H., Di Giorgio, C., Azas, N., Doumbo, O., Timon-David, P. (2000). Toxicity and genotoxicity of antimalarial alkaloid rich extracts derived from *Mitragyna inermis* O. Kuntze and *Nauclea latifolia*. *Phytotherapy Research* **14**, 608–611.
- Umlawski, J. (1998). Malariatherapie durch vorzeitige Eliminierung Falciparum–infizierter normaler und genetisch veränderter Erythrozyten aus dem Blutkreislauf, ausgelöst durch gezielte oxidative Schädigung. Dissertation, Medizinische Fakultät, Humboldt–Universität zu Berlin, Berlin.
- Valentin, A., Benoit–Vical, F., Moulis, C., Stanislas, E., Mallié, M., Fouraste, I., Bastide, J. (1997). In vitro antimalarial activity of penduline, a bisbenzylisoquinoline from *Isopyrum thalictroides*. *Antimicrobial Agents and Chemotherapy* **41**, 2305–2307.
- Varshney, I. P., Jain, D. C., Beg, M. F. A., Srivastava, H. C. (1985). Samanin–C and Samanin–E new saponins from *Pithecolobium saman*. *Fitoterapia* **56**, 281–283.
- Vongchareonsathit, A., De–Eknankul, W. (1998). Rapid TLC–densitometric analysis of plaunotol from *Croton sublyratus* leaves. *Planta Medica* **64**, 279–280.
- Watt, J. M., Breyer–Brandwijk, M. G. (1962). *The Medicinal and Poisonous Plants of Southern and Eastern Africa*. E.& S. Livingstone, Edinburgh, London.
- White, N. J. (1996). Malaria. In: *Manson's Tropical Diseases* (Herausgeber: G. C. Cook), W. B. Saunders Company London, London, S. 1087–1165.

- WHO (1997). World malaria situation in 1994. *Weekly Epidemiological Record* **72**, 269–292.
- WHO (2000). *World Health Report 2000*. WHO Press, Geneva.
- Wolters, B. (1996). *Agave bis Zaubernuß. Heilpflanzen der Indianer Nord- und Mittelamerikas*. Urs Freund Verlag, Greifenberg, S. 284–285.
- Wussow, J. R., Urbatsch, L. E., Sullivan, G. A. (1985). *Calea* (Asteraceae) in Mexico, Central America, and Jamaica. *Systematic Botany* **10**, 241–267.
- Yadav, J. S., Deshpande, P. K., Sharma, G. V. M. (1990). Synthesis of biologically active compounds of agricultural interest. *Pure & Applied Chemistry* **62**, 1333–1338.
- Yoshida, T., Namba, O., Chen, L., Okuda, T. (1990). Tannins and related polyphenols of euphorbiaceous plants V. Euphorbin C and equilibrated dimeric dehydroellagttannin having a new tetrameric galloyl group. *Chemical & Pharmaceutical Bulletin* **38**, 86–93.
- Zdero, C, Ahmed, A. A., Bohlmann, F., Mungai, G. M. (1990). Diterpenes and sesquiterpene xylosides from East African *Conyza* species. *Phytochemistry* **29**, 3167–3172.