

---

## 7 Literaturverzeichnis

- Aberle,H., Bauer,A., Stappert,J., Kispert,A., and Kemler,R. (1997). Beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J.* 16, 3797.
- Ahumada,A., Slusarski,D.C., Liu,X., Moon,R.T., Malbon,C.C., and Wang,H.Y. (2002). Signaling of rat Frizzled-2 through phosphodiesterase and cyclic GMP. *Science* 298, 2006.
- Amit,S., Hatzubai,A., Birman,Y., Andersen,J.S., and Ben Shushan,E. (2002). Axin-mediated CKI phosphorylation of beta-catenin at Ser 45: a molecular switch for the Wnt pathway. *Genes Dev.* 16, 1066.
- Angers,S., Thorpe,C.J., Biechele,T.L., Goldenberg,S.J., Zheng,N., MacCoss,M.J., and Moon,R.T. (2006). The KLHL12-Cullin-3 ubiquitin ligase negatively regulates the Wnt-beta-catenin pathway by targeting Dishevelled for degradation. *Nat. Cell Biol.* 8, 348-357.
- Ausubel,F.M., Brent,R., Kingston,R.E., Moore,D.D., Seidman,I.G., Smith,J.A., and Struhl,K. (1987). Current protocols in molecular biology. (Greene Publishing Associates and Wiley-Interscience, New York).
- Axelrod,J.D., Miller,J.R., Shulman,J.M., Moon,R.T., and Perrimon,N. (1998). Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes Dev.* 12, 2610-2622.
- Baeg,G.H., Lin,X., Khare,N., Baumgartner,S., and Perrimon,N. (2001). Heparan sulfate proteoglycans are critical for the organization of the extracellular distribution of Wingless. *Development* 128, 87.
- Bakkers,J., Kramer,C., Pothof,J., Quaedvlieg,N.E., Spaink,H.P., and Hammerschmidt,M. (2004). Has2 is required upstream of Rac1 to govern dorsal migration of lateral cells during zebrafish gastrulation. *Development* 131, 525.
- Basler,K., Christen,B., and Hafen,E. (1991). Ligand-independent activation of the sevenless receptor tyrosine kinase changes the fate of cells in the developing Drosophila eye. *Cell* 64, 1069-1081.
- Behrens,J., Jerchow,B.A., Wurtele,M., Grimm,J., Asbrand,C., Wirtz,R., Kuhl,M., Wedlich,D., and Birchmeier,W. (1998). Functional interaction of an axin homolog, conductin, with beta-catenin, APC, and GSK3beta. *Science* 280, 596-599.

- Behrens,J., von Kries,J.P., Kuhl,M., Bruhn,L., and Wedlich,D. (1996). Functional interaction of beta-catenin with the transcription factor LEF-1. *Nature* 382, 638.
- Bilder,D. and Perrimon,N. (2000). Localization of apical epithelial determinants by the basolateral PDZ protein Scribble. *Nature* 403, 676.
- Blader,P., Strahle,U., and Ingham,P.W. (1996). Three Wnt genes expressed in a wide variety of tissues during development of the zebrafish, *Danio rerio*: developmental and evolutionary perspectives. *Dev. Genes Evol.* 206, 3-13.
- Bogenrieder,T. and Herlyn,M. (2003). Axis of evil: molecular mechanisms of cancer metastasis. *Oncogene* 22, 6524-6536.
- Boutros,M. and Mlodzik,M. (1999). Dishevelled: at the crossroads of divergent intracellular signaling pathways. *Mech. Dev.* 83, 27-37.
- Boutros,M., Paricio,N., Strutt,D.I., and Mlodzik,M. (1998). Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and wingless signaling. *Cell* 94, 109-118.
- Brand,A.H. and Perrimon,N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118, 401-415.
- Brivanlou,A.H. and Darnell,J.E., Jr. (2002). Signal transduction and the control of gene expression. *Science* 295, 813-818.
- Bryja,V., Schulte,G., Rawal,N., Grahn,A., and Arenas,E. (2007). Wnt-5a induces Dishevelled phosphorylation and dopaminergic differentiation via a CK1-dependent mechanism. *J. Cell Sci.* 120, 586-595.
- Cabrera,C.V., Alonso,M.C., Johnston,P., Phillips,R.G., and Lawrence,P.A. (1987). Phenocopies Induced with Antisense Rna Identify the Wingless Gene. *Cell* 50, 659-663.
- Cadigan,K.M. and Liu,Y.I. (2006). Wnt signaling: complexity at the surface. *J. Cell Sci.* 119, 395-402.
- Cafferty,P., Yu,L., and Rao,Y. (2004). The receptor tyrosine kinase Off-track is required for layer-specific neuronal connectivity in *Drosophila*. *Development* 131, 5287-5295.
- Cantley,L.C. (2002). The phosphoinositide 3-kinase pathway. *Science* 296, 1655-1657.

- Carreira-Barbosa,F., Concha,M.L., Takeuchi,M., Ueno,N., Wilson,S.W., and Tada,M. (2003). Prickle 1 regulates cell movements during gastrulation and neuronal migration in zebrafish. *Development* 130, 4037.
- Chae,J., Kim,M.J., Goo,J.H., Collier,S., Gubb,D., Charlton,J., Adler,P.N., and Park,W.J. (1999). The Drosophila tissue polarity gene starry night encodes a member of the protocadherin family. *Development* 126, 5421-5429.
- Chan,D.W., Chan,C.Y., Yam,J.W., Ching,Y.P., and Ng,I.O. (2006). Prickle-1 negatively regulates Wnt/beta-catenin pathway by promoting Dishevelled ubiquitination/degradation in liver cancer. *Gastroenterology* 131, 1218-1227.
- Cheng,J.C., Miller,A.L., and Webb,S.E. (2004). Organization and function of microfilaments during late epiboly in zebrafish embryos. *Dev. Dyn.* 231, 313.
- Christodoulides,C., Scarda,A., Granzotto,M., Milan,G., Nora,E.D., Keogh,J., Pergola,G.D., Stirling,H., Pannacciulli,N., Sethi,J.K., Federspil,G., Vidal-Puig,A., Farooqi,I.S., O'Rahilly,S., and Vettor,R. (2006). WNT10B mutations in human obesity. *Diabetologia* V49, 678-684.
- Clarke,A.R. (2006). Wnt signalling in the mouse intestine. *Oncogene* 25, 7512-7521.
- Cooper,M.T. and Bray,S.J. (1999). Frizzled regulation of Notch signalling polarizes cell fate in the Drosophila eye. *Nature* 397, 526-530.
- Copp,A.J., Greene,N.D., and Murdoch,J.N. (2003). The genetic basis of mammalian neurulation. *Nat. Rev. Genet.* 4, 784-793.
- Corey,D.R. and Abrams,J.M. (2001). Morpholino antisense oligonucleotides: tools for investigating vertebrate development. *Genome Biol.* 2, REVIEWS1015.
- Curtin,J.A., Quint,E., Tsipouri,V., Arkell,R.M., and Cattanach,B. (2003). Mutation of Celsr1 disrupts planar polarity of inner ear hair cells and causes severe neural tube defects in the mouse. *Curr. Biol.* 13, 1129.
- Dabdoub,A., Donohue,M.J., Brennan,A., Wolf,V., and Montcouquiol,M. (2003). Wnt signaling mediates reorientation of outer hair cell stereociliary bundles in the mammalian cochlea. *Development* 130, 2375.
- Daggett,D.F., Boyd,C.A., Gautier,P., Bryson-Richardson,R.J., and Thisse,C. (2004). Developmentally restricted actin-regulatory molecules control morphogenetic cell movements in the zebrafish gastrula. *Curr. Biol.* 14, 1632.

- Dallas,S.L., Chen,Q., and Sivakumar,P. (2006). Dynamics of assembly and reorganization of extracellular matrix proteins. *Curr. Top. Dev. Biol.* 75, 1-24.
- Das,G., Jenny,A., Klein,T.J., Eaton,S., and Mlodzik,M. (2004). Diego interacts with Prickle and Strabismus/Van Gogh to localize planar cell polarity complexes. *Development* 131, 4467-4476.
- De Ferrari,G.V. and Moon,R.T. (2006). The ups and downs of Wnt signaling in prevalent neurological disorders. *Oncogene* 25, 7545-7553.
- De Robertis,E.M., Larrain,J., Oelgeschlager,M., and Wessely,O. (2000). The establishment of Spemann's organizer and patterning of the vertebrate embryo. *Nat. Rev. Genet.* 1, 171-181.
- Dierick,H. and Bejsovec,A. (1999). Cellular mechanisms of wingless/Wnt signal transduction. *Curr. Top. Dev. Biol.* 43, 153-190.
- Divecha,N. and Irvine,R.F. (1995). Phospholipid Signaling. *Cell* 80, 269-278.
- Fanto,M. and McNeill,H. (2004). Planar polarity from flies to vertebrates. *J. Cell Sci.* 117, 527-533.
- Fanto,M. and Mlodzik,M. (1999). Asymmetric Notch activation specifies photoreceptors R3 and R4 and planar polarity in the Drosophila eye. *Nature* 397, 523-526.
- Fanto,M., Weber,U., Strutt,D.I., and Mlodzik,M. (2000). Nuclear signaling by Rac and Rho GTPases is required in the establishment of epithelial planar polarity in the Drosophila eye. *Curr. Biol.* 10, 979.
- Feigui,F., Hannus,M., Mlodzik,M., and Eaton,S. (2001). The ankyrin repeat protein Diego mediates Frizzled-dependent planar polarization. *Dev. Cell* 1, 93-101.
- Fields,S. and Song,O.K. (1989). A Novel Genetic System to Detect Protein Protein Interactions. *Nature* 340, 245-246.
- Formstone,C.J. and Mason,I. (2005). Combinatorial activity of Flamingo proteins directs convergence and extension within the early zebrafish embryo via the planar cell polarity pathway. *Dev. Biol.* 282, 320.
- Giles,R.H., van Es,J.H., and Clevers,H. (2003). Caught up in a Wnt storm: Wnt signaling in cancer. *Biochim. Biophys. Acta* 1653, 1.

- Gong,Y., Mo,C., and Fraser,S.E. (2004). Planar cell polarity signalling controls cell division orientation during zebrafish gastrulation. *Nature* 430, 689-693.
- Gonzalez-Sancho,J.M., Brennan,K.R., Castelo-Soccio,L.A., and Brown,A.M. (2004). Wnt proteins induce dishevelled phosphorylation via an LRP5/6- independent mechanism, irrespective of their ability to stabilize beta-catenin. *Mol. Cell Biol.* 24, 4757-4768.
- Gordon,M.D. and Nusse,R. (2006). Wnt signaling: multiple pathways, multiple receptors, and multiple transcription factors. *J. Biol. Chem.* 281, 22429-22433.
- Grant,S.F.A., Thorleifsson,G., Reynisdottir,I., Benediktsson,R., Manolescu,A., Sainz,J., Helgason,A., Stefansson,H., Emilsson,V., Helgadottir,A., Styrkarsdottir,U., Magnusson,K.P., Walters,G.B., Palsdottir,E., Jonsdottir,T., Gudmundsdottir,T., Gylfason,A., Saemundsdottir,J., Wilensky,R.L., Reilly,M.P., Rader,D.J., Bagger,Y., Christiansen,C., Gudnason,V., Sigurdsson,G., Thorsteinsdottir,U., Gulcher,J.R., Kong,A., and Stefansson,K. (2006). Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet* 38, 320-323.
- Gubb,D., Green,C., Huen,D., Coulson,D., Johnson,G., Tree,D., Collier,S., and Roote,J. (1999). The balance between isoforms of the prickle LIM domain protein is critical for planar polarity in Drosophila imaginal discs. *Genes Dev.* 13, 2315-2327.
- Guder,C., Philipp,I., Lengfeld,T., Watanabe,H., Hobmayer,B., and Holstein,T.W. (2006). The Wnt code: cnidarians signal the way. *Oncogene* 25, 7450-7460.
- Guo,N., Hawkins,C., and Nathans,J. (2004). Frizzled6 controls hair patterning in mice. *Proc. Natl. Acad. Sci. USA* 101, 9277.
- Habas,R., Dawid,I.B., and He,X. (2003). Coactivation of Rac and Rho by Wnt/Frizzled signaling is required for vertebrate gastrulation. *Genes Dev.* 17, 295-309.
- Habas,R., Kato,Y., and He,X. (2001). Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell* 107, 843-854.
- Hakeda-Suzuki,S., Ng,J., Tzu,J., Dietzl,G., Sun,Y., Harms,M., Nardine,T., Luo,L., and Dickson,B.J. (2002). Rac function and regulation during Drosophila development. *Nature* 416, 438-442.
- Hamblet,N.S., Lijam,N., Ruiz-Lozano,P., Wang,J., Yang,Y., Luo,Z., Mei,L., Chien,K.R., Sussman,D.J., and Wynshaw-Boris,A. (2002). Dishevelled 2 is essential for cardiac outflow tract development, somite segmentation and neural tube closure. *Development* 129, 5827-5838.

- Hammerschmidt,M., Pelegri,F., Mullins,M.C., Kane,D.A., Brand,M., van Eeden,F.J., Furutani-Seiki,M., Granato,M., Haffter,P., Heisenberg,C.P., Jiang,Y.J., Kelsh,R.N., Odenthal,J., Warga,R.M., and Nusslein-Volhard,C. (1996). Mutations affecting morphogenesis during gastrulation and tail formation in the zebrafish, *Danio rerio*. *Development* 123, 143-151.
- Heisenberg,C.P. and Nusslein-Volhard,C. (1997). The function of silberblick in the positioning of the eye anlage in the zebrafish embryo. *Dev. Biol.* 184, 85.
- Heisenberg,C.P., Tada,M., Rauch,G.J., Saude,L., Concha,M.L., Geisler,R., Stemple,D.L., Smith,J.C., and Wilson,S.W. (2000). Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* 405, 76-81.
- Henderson,D.J., Phillips,H.M., and Chaudhry,B. (2006). Vang-like 2 and noncanonical Wnt signaling in outflow tract development. *Trends in Cardiovascular Medicine* 16, 38-45.
- Huang,H.C. and Klein,P.S. (2004). The Frizzled family: receptors for multiple signal transduction pathways. *Genome Biol.* 5, 234.
- Huber,M.A., Kraut,N., and Beug,H. (2005). Molecular requirements for epithelial-mesenchymal transition during tumor progression. *Current Opinion in Cell Biology* 17, 548-558.
- Huelsken,J. and Birchmeier,W. (2001). New aspects of Wnt signaling pathways in higher vertebrates. *Curr. Opin. Genet. Dev.* 11, 547-553.
- Ikeda,S., Kishida,S., Yamamoto,H., Murai,H., Koyama,S., and Kikuchi,A. (1998). Axin, a negative regulator of the Wnt signaling pathway, forms a complex with GSK-3beta and beta-catenin and promotes GSK-3beta-dependent phosphorylation of beta-catenin. *EMBO J.* 17, 1371-1384.
- Ito,H., Fukuda,Y., Murata,K., and Kimura,A. (1983). Transformation of Intact Yeast-Cells Treated with Alkali Cations. *Journal of Bacteriology* 153, 163-168.
- James,P., Halladay,J., and Craig,E.A. (1996). Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast. *Genetics* 144, 1425-1436.
- Jenny,A., Darken,R.S., Wilson,P.A., and Mlodzik,M. (2003). Prickle and Strabismus form a functional complex to generate a correct axis during planar cell polarity signaling. *EMBO J.* 22, 4409-4420.
- Jenny,A. and Mlodzik,M. (2006). Planar cell polarity signaling: a common mechanism for cellular polarization. *Mt. Sinai J. Med.* 73, 738-750.

- Jenny,A., Reynolds-Kenneally,J., Das,G., Burnett,M., and Mlodzik,M. (2005). Diego and Prickle regulate Frizzled planar cell polarity signalling by competing for Dishevelled binding. *Nat. Cell Biol.* 7, 691-697.
- Jessen,J.R., Topczewski,J., Bingham,S., Sepich,D.S., Marlow,F., Chandrasekhar,A., and Solnica-Krezel,L. (2002). Zebrafish trilobite identifies new roles for Strabismus in gastrulation and neuronal movements. *Nat. Cell Biol.* 4, 610-615.
- Karner,C., Wharton,K.A., and Carroll,T.J. (2006a). Apical-basal polarity, Wnt signaling and vertebrate organogenesis. *Semin. Cell Dev. Biol.* 17, 214-222.
- Karner,C., Wharton,K.A., Jr., and Carroll,T.J. (2006b). Planar cell polarity and vertebrate organogenesis. *Semin. Cell Dev. Biol.* 17, 194-203.
- Keegan,B.R., Meyer,D., and Yelon,D. (2004). Organization of cardiac chamber progenitors in the zebrafish blastula. *Development* 131, 3081.
- Keller,R. (2002). Shaping the vertebrate body plan by polarized embryonic cell movements. *Science* 298, 1950-1954.
- Kibar,Z., Vogan,K.J., Groulx,N., Justice,M.J., Underhill,D.A., and Gros,P. (2001). Ltap/Lpp1, a mammalian homolog of Drosophila Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat. Genet.* 28, 251.
- Kiefer,J.C. (2005). Planar cell polarity: heading in the right direction. *Dev. Dyn.* 233, 695-700.
- Kilian,B., Mansukoski,H., Barbosa,F.C., Ulrich,F., Tada,M., and Heisenberg,C.P. (2003). The role of Ppt/Wnt5 in regulating cell shape and movement during zebrafish gastrulation. *Mech. Dev.* 120, 467-476.
- Kimelman,D. and Xu,W. (2006). beta-Catenin destruction complex: insights and questions from a structural perspective. *Oncogene* 25, 7482-7491.
- Kimmel,C.B., Ballard,W.W., Kimmel,S.R., Ullmann,B., and Schilling,T.F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* 203, 253.
- Kitagawa,M., Hatakeyama,S., Shirane,M., Matsumoto,M., Ishida,N., Hattori,K., Nakamichi,I., Kikuchi,A., Nakayama,K., and Nakayama,K. (1999). An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of beta-catenin. *EMBO J.* 18, 2401-2410.
- Klein,T.J., Jenny,A., Djiane,A., and Mlodzik,M. (2006). CKlepsilon/discs overgrown promotes both Wnt-Fz/beta-catenin and Fz/PCP signaling in Drosophila. *Curr. Biol.* 16, 1337-1343.

- Klein,T.J. and Mlodzik,M. (2005). Planar cell polarization: An emerging model points in the right direction. *Annu. Rev. Cell. Dev. Biol.* 21, 155.
- Klingensmith,J., Nusse,R., and Perrimon,N. (1994). The *Drosophila* segment polarity gene *dishevelled* encodes a novel protein required for response to the wingless signal. *Genes Dev.* 8, 118.
- Klingensmith,J., Yang,Y., Axelrod,J.D., Beier,D.R., Perrimon,N., and Sussman,D.J. (1996). Conservation of *dishevelled* structure and function between flies and mice: isolation and characterization of Dvl2. *Mech. Dev.* 58, 15-26.
- Kockel,L., Zeitlinger,J., Staszewski,L.M., Mlodzik,M., and Bohmann,D. (1997). Jun in *Drosophila* development: redundant and nonredundant functions and regulation by two MAPK signal transduction pathways. *Genes Dev.* 11, 1748-1758.
- Krieg,P.A. and Melton,D.A. (1984). Functional messenger RNAs are produced by SP6 in vitro transcription of cloned cDNAs. *Nucleic Acids Res.* 12, 7057-7070.
- Kuan,C.Y., Yang,D.D., Samanta Roy,D.R., Davis,R.J., Rakic,P., and Flavell,R.A. (1999). The Jnk1 and Jnk2 protein kinases are required for regional specific apoptosis during early brain development. *Neuron* 22, 667-676.
- Kuhl,M., Sheldahl,L.C., Malbon,C.C., and Moon,R.T. (2000a). Ca(2+)/calmodulin-dependent protein kinase II is stimulated by Wnt and Frizzled homologs and promotes ventral cell fates in *Xenopus*. *J. Biol. Chem.* 275, 12701-12711.
- Kuhl,M., Sheldahl,L.C., Park,M., Miller,J.R., and Moon,R.T. (2000b). The Wnt/Ca<sup>2+</sup> pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet.* 16, 279.
- Kupperman,E., An,S., Osborne,N., Waldron,S., and Stainier,D.Y. (2000). A sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature* 406, 192-195.
- Laemmli,U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.
- Lawrence,P.A., Casal,J., and Struhl,G. (2002). Towards a model of the organisation of planar polarity and pattern in the *Drosophila* abdomen. *Development* 129, 2749-2760.
- Lee,P.N., Pang,K., Matus,D.Q., and Martindale,M.Q. (2006). A WNT of things to come: Evolution of Wnt signaling and polarity in cnidarians. *Seminars in Cell & Developmental Biology* 17, 157-167.

- Lele,Z., Bakkers,J., and Hammerschmidt,M. (2001). Morpholino phenocopies of the swirl, snailhouse, somitabun, minifin, silberblick, and pipetail mutations. *Genesis*. **30**, 190-194.
- Lepley,D., Paik,J.H., Hla,T., and Ferrer,F. (2005). The G protein-coupled receptor S1P2 regulates Rho/Rho kinase pathway to inhibit tumor cell migration. *Cancer Res.* **65**, 3788-3795.
- Li,L., Yuan,H., Xie,W., Mao,J., Caruso,A.M., McMahon,A., Sussman,D.J., and Wu,D. (1999). Dishevelled proteins lead to two signaling pathways. Regulation of LEF-1 and c-Jun N-terminal kinase in mammalian cells. *J. Biol. Chem.* **274**, 129-134.
- Lijam,N., Paylor,R., McDonald,M.P., Crawley,J.N., Deng,C.X., Herrup,K., Stevens,K.E., Maccaferri,G., McBain,C.J., Sussman,D.J., and Wynshaw-Boris,A. (1997). Social interaction and sensorimotor gating abnormalities in mice lacking Dvl1. *Cell* **90**, 895-905.
- Lin,X. and Perrimon,N. (1999). Dally cooperates with Drosophila Frizzled 2 to transduce Wingless signalling. *Nature* **400**, 281.
- Logan,C.Y. and Nusse,R. (2004). THE WNT SIGNALING PATHWAY IN DEVELOPMENT AND DISEASE. *Annual Review of Cell and Developmental Biology* **20**, 781-810.
- Lu,X., Borchers,A.G., Jolicoeur,C., Rayburn,H., Baker,J.C., and Tessier-Lavigne,M. (2004). PTK7/CCK-4 is a novel regulator of planar cell polarity in vertebrates. *Nature* **430**, 93-98.
- Machingo,Q.J., Fritz,A., and Shur,B.D. (2006). A beta1,4-galactosyltransferase is required for convergent extension movements in zebrafish. *Dev. Biol.* **297**, 471-482.
- Makita,R., Mizuno,T., Koshida,S., Kuroiwa,A., and Takeda,H. (1998). Zebrafish wnt11: pattern and regulation of the expression by the yolk cell and No tail activity. *Mech. Dev.* **71**, 165-176.
- Marlow,F., Topczewski,J., Sepich,D., and Solnica-Krezel,L. (2002). Zebrafish Rho kinase 2 acts downstream of Wnt11 to mediate cell polarity and effective convergence and extension movements. *Curr. Biol.* **12**, 876.
- Matsui,T., Raya,A., Kawakami,Y., Callol-Massot,C., Capdevila,J., Rodriguez-Esteban,C., and Izpisua Belmonte,J.C. (2005). Noncanonical Wnt signaling regulates midline convergence of organ primordia during zebrafish development. *Genes Dev.* **19**, 164-175.
- Miller,J.R. (2002). The Wnts. *Genome Biol.* **3**, REVIEWS3001.

- Mlodzik,M. (2002). Planar cell polarization: do the same mechanisms regulate Drosophila tissue polarity and vertebrate gastrulation? *Trends Genet.* **18**, 564-571.
- Moeller,H., Jenny,A., Schaeffer,H.J., Schwarz-Romond,T., Mlodzik,M., Hammerschmidt,M., and Birchmeier,W. (2006). Diversin regulates heart formation and gastrulation movements in development. *Proc. Natl. Acad. Sci. U. S. A* **103**, 15900-15905.
- Molenaar,M., van de,W.M., Oosterwegel,M., Peterson-Maduro,J., and Godsake,S. (1996). XTcf-3 transcription factor mediates beta-catenin-induced axis formation in Xenopus embryos. *Cell* **86**, 391.
- Montcouquiol,M., Rachel,R.A., Lanford,P.J., Copeland,N.G., Jenkins,N.A., and Kelley,M.W. (2003). Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. *Nature* **423**, 173.
- Moon,R.T., Brown,J.D., and Torres,M. (1997). WNTs modulate cell fate and behavior during vertebrate development. *Trends Genet.* **13**, 157-162.
- Morgan,D., Turnpenny,L., Goodship,J., Dai,W., Majumder,K., Matthews,L., Gardner,A., Schuster,G., Vien,L., Harrison,W., Elder,F.F., Penman-Splitt,M., Overbeek,P., and Strachan,T. (1998). Inversin, a novel gene in the vertebrate left-right axis pathway, is partially deleted in the inv mouse. *Nat. Genet.* **20**, 149-156.
- Moriguchi,T., Kawachi,K., Kamakura,S., Masuyama,N., Yamanaka,H., Matsumoto,K., Kikuchi,A., and Nishida,E. (1999). Distinct domains of mouse dishevelled are responsible for the c-Jun N-terminal kinase/stress-activated protein kinase activation and the axis formation in vertebrates. *J. Biol. Chem.* **274**, 30957-30962.
- Mosavi,L.K., Cammett,T.J., Desrosiers,D.C., and Peng,Z.Y. (2004). The ankyrin repeat as molecular architecture for protein recognition. *Protein Sci.* **13**, 1435-1448.
- Moser,A.R., Luongo,C., Gould,K.A., McNeley,M.K., Shoemaker,A.R., and Dove,W.F. (1995). ApcMin: a mouse model for intestinal and mammary tumorigenesis. *Eur. J. Cancer* **31A**, 1061-1064.
- Myers,D.C., Sepich,D.S., and Solnica-Krezel,L. (2002). Convergence and extension in vertebrate gastrulae: cell movements according to or in search of identity? *Trends Genet.* **18**, 447.
- Nusse,R., Brown,A., Papkoff,J., Scambler,P., Shackleford,G., McMahon,A., Moon,R., and Varmus,H. (1991). A new nomenclature for int-1 and related genes: the Wnt gene family. *Cell* **64**, 231.

- Nusse,R., van Ooyen,A., Rijsewijk,F., van Lohuizen,M., Schuuring,E., and van't Veer,L. (1985). Retroviral insertional mutagenesis in murine mammary cancer. Proc. R. Soc. Lond B Biol. Sci. 226, 3-13.
- Nusse,R. and Varmus,H.E. (1982). Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. Cell 31, 99-109.
- Nüsslein-Volhard,C. and Wieschaus,E. (1980). Mutations Affecting Segment Number and Polarity in Drosophila. Nature 287, 795-801.
- Ohkawara,B., Yamamoto,T.S., Tada,M., and Ueno,N. (2003). Role of glypican 4 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. Development 130, 2129-2138.
- Pan,W.J., Pang,S.Z., Huang,T., Guo,H.Y., Wu,D., and Li,L. (2004). Characterization of function of three domains in dishevelled-1: DEP domain is responsible for membrane translocation of dishevelled-1. Cell Res. 14, 324-330.
- Pandur,P., Lasche,M., Eisenberg,L.M., and Kuhl,M. (2002). Wnt-11 activation of a non-canonical Wnt signalling pathway is required for cardiogenesis. Nature 418, 636-641.
- Park,M. and Moon,R.T. (2002). The planar cell-polarity gene stbm regulates cell behaviour and cell fate in vertebrate embryos. Nat. Cell Biol. 4, 20.
- Park,T.J., Gray,R.S., Sato,A., Habas,R., and Wallingford,J.B. (2005). Subcellular localization and signaling properties of dishevelled in developing vertebrate embryos. Curr. Biol. 15, 1039-1044.
- Peters,J.M., Mckay,R.M., Mckay,J.P., and Graff,J.M. (1999). Casein kinase I transduces Wnt signals. Nature 401, 345-350.
- Puech,P.H., Taubenberger,A., Ulrich,F., Krieg,M., Muller,D.J., and Heisenberg,C.P. (2005). Measuring cell adhesion forces of primary gastrulating cells from zebrafish using atomic force microscopy. J. Cell Sci. 118, 4199-4206.
- Radtke,F., Clevers,H., and Riccio,O. (2006). From Gut Homeostasis to Cancer. Current Molecular Medicine 6, 275-289.
- Rauch,G.J., Hammerschmidt,M., Blader,P., Schauerte,H.E., Strahle,U., Ingham,P.W., McMahon,A.P., and Haffter,P. (1997). Wnt5 is required for tail formation in the zebrafish embryo. Cold Spring Harb. Symp. Quant. Biol. 62, 227-234.

- Riesgo-Escovar,J.R., Jenni,M., Fritz,A., and Hafen,E. (1996). The Drosophila Jun-N-terminal kinase is required for cell morphogenesis but not for DJun-dependent cell fate specification in the eye. *Genes Dev.* 10, 2759-2768.
- Rijsewijk,F., Schuermann,M., Wagenaar,E., Parren,P., Weigel,D., and Nusse,R. (1987). The Drosophila Homolog of the Mouse Mammary Oncogene Int-1 Is Identical to the Segment Polarity Gene Wingless. *Cell* 50, 649-657.
- Rossmann,K.L., Der,C.J., and Sondek,J. (2005). GEF means go: turning on RHO GTPases with guanine nucleotide-exchange factors. *Nat. Rev. Mol. Cell Biol.* 6, 167-180.
- Rosso,S.B., Sussman,D., Wynshaw-Boris,A., and Salinas,P.C. (2005). Wnt signaling through Dishevelled, Rac and JNK regulates dendritic development. *Nat. Neurosci.* 8, 34-42.
- Saburi,S. and McNeill,H. (2005). Organising cells into tissues: new roles for cell adhesion molecules in planar cell polarity. *Curr. Opin. Cell Biol.* 17, 482-488.
- Sambrook,E.F., Fritsch,E.F., and Maniatis,T. (1989). Molecular cloning: a laboratory handbook. (Cold Spring Harbor Laboratory Press, New York).
- Saneyoshi,T., Kume,S., Amasaki,Y., and Mikoshiba,K. (2002). The Wnt/calcium pathway activates NF-AT and promotes ventral cell fate in Xenopus embryos. *Nature* 417, 295-299.
- Schier,A.F. (2001). Axis formation and patterning in zebrafish. *Curr. Opin. Genet. Dev.* 11, 393-404.
- Schier,A.F. and Talbot,W.S. (2005). Molecular genetics of axis formation in zebrafish. *Annu. Rev. Genet.* 39, 561-613.
- Schwarz-Romond,T., Asbrand,C., Bakkers,J., Kuhl,M., Schaeffer,H.J., Huelsken,J., Behrens,J., Hammerschmidt,M., and Birchmeier,W. (2002). The ankyrin repeat protein Diversin recruits Casein kinase Iepsilon to the beta-catenin degradation complex and acts in both canonical Wnt and Wnt/JNK signaling. *Genes Dev.* 16, 2073-2084.
- Seifert,J.R. and Mlodzik,M. (2007). Frizzled/PCP signalling: a conserved mechanism regulating cell polarity and directed motility. *Nat. Rev. Genet.* 8, 126-138.
- Sheldahl,L.C., Park,M., Malbon,C.C., and Moon,R.T. (1999). Protein kinase C is differentially stimulated by Wnt and Frizzled homologs in a G-protein-dependent manner. *Curr. Biol.* 9, 695.

- Sheldahl,L.C., Slusarski,D.C., Pandur,P., Miller,J.R., Kuhl,M., and Moon,R.T. (2003). Dishevelled activates Ca<sup>2+</sup> flux, PKC, and CamKII in vertebrate embryos. *J. Cell Biol.* **161**, 769-777.
- Slusarski,D.C., Corces,V.G., and Moon,R.T. (1997a). Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* **390**, 410.
- Slusarski,D.C., Yang-Snyder,J., Busa,W.B., and Moon,R.T. (1997b). Modulation of embryonic intracellular Ca<sup>2+</sup> signaling by Wnt-5A. *Dev. Biol.* **182**, 114.
- Solnica-Krezel,L. and Driever,W. (1994). Microtubule arrays of the zebrafish yolk cell: organization and function during epiboly. *Development* **120**, 2443.
- Spemann,H. and Mangold,H. (2001). Induction of embryonic primordia by implantation of organizers from a different species. 1923. *Int. J. Dev. Biol.* **45**, 13-38.
- Stainier,D.Y. (2001). Zebrafish genetics and vertebrate heart formation. *Nat. Rev. Genet.* **2**, 39-48.
- Strutt,D. (2003). Frizzled signalling and cell polarisation in *Drosophila* and vertebrates. *Development* **130**, 4501-4513.
- Strutt,D.I., Weber,U., and Mlodzik,M. (1997). The role of RhoA in tissue polarity and Frizzled signalling. *Nature* **387**, 292-295.
- Strutt,H., Price,M.A., and Strutt,D. (2006). Planar polarity is positively regulated by casein kinase epsilon in *Drosophila*. *Curr. Biol.* **16**, 1329-1336.
- Sumanas,S., Kim,H.J., Hermanson,S., and Ekker,S.C. (2001). Zebrafish frizzled 2 morphant displays defects in body axis elongation. *Genesis* **30**, 114.
- Sussman,D.J., Klingensmith,J., Salinas,P., Adams,P.S., Nusse,R., and Perrimon,N. (1994). Isolation and characterization of a mouse homolog of the *Drosophila* segment polarity gene dishevelled. *Dev. Biol.* **166**, 73.
- Tada,M., Concha,M.L., and Heisenberg,C.P. (2002). Non-canonical Wnt signaling and regulation of gastrulation movements. *Semin. Cell Dev. Biol.* **13**, 251.
- Tam,P.P., Loebel,D.A., and Tanaka,S.S. (2006). Building the mouse gastrula: signals, asymmetry and lineages. *Curr. Opin. Genet. Dev.* **16**, 419-425.
- Taylor,J., Abramova,N., Charlton,J., and Adler,P.N. (1998). Van Gogh: a new *Drosophila* tissue polarity gene. *Genetics* **150**, 199.

- Tetsu,O. and McCormick,F. (1999). Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* 398, 422.
- Theisen,H., Purcell,J., Bennett,M., Kansagara,D., Syed,A., and Marsh,J.L. (1994). dishevelled is required during wingless signaling to establish both cell polarity and cell identity. *Development* 120, 347-360.
- Thisse,C. and Zon,L.I. (2002). Organogenesis--heart and blood formation from the zebrafish point of view. *Science* 295, 457-462.
- Tomlinson,A. (1988). Cellular interactions in the developing Drosophila eye. *Development* 104, 183-193.
- Topczewski,J., Sepich,D.S., Myers,D.C., Walker,C., Amores,A., Lele,Z., Hammerschmidt,M., Postlethwait,J., and Solnica-Krezel,L. (2001). The zebrafish glypcan knypek controls cell polarity during gastrulation movements of convergent extension. *Dev. Cell* 1, 251-264.
- Torban,E., Wang,H.J., Groulx,N., and Gros,P. (2004). Independent mutations in mouse Vangl2 that cause neural tube defects in looptail mice impair interaction with members of the Dishevelled family. *J. Biol. Chem.* 279, 52703-52713.
- Tree,D.R., Ma,D., and Axelrod,J.D. (2002). A three-tiered mechanism for regulation of planar cell polarity. *Semin. Cell Dev. Biol.* 13, 217-224.
- Trinh,L.A. and Stainier,D.Y. (2004). Fibronectin regulates epithelial organization during myocardial migration in zebrafish. *Dev. Cell* 6, 371-382.
- Tsang,M., Lijam,N., Yang,Y., Beier,D.R., Wynshaw-Boris,A., and Sussman,D.J. (1996). Isolation and characterization of mouse dishevelled-3. *Dev. Dyn.* 207, 253.
- Turner,D.L. and Weintraub,H. (1994). Expression of Achaete-Scute Homolog-3 in Xenopus-Embryos Converts Ectodermal Cells to A Neural Fate. *Genes & Development* 8, 1434-1447.
- Usui,T., Shima,Y., Shimada,Y., Hirano,S., and Burgess,R.W. (1999a). Flamingo, a seven-pass transmembrane cadherin, regulates planar cell polarity under the control of Frizzled. *Cell* 98, 585.
- Usui,T., Shima,Y., Shimada,Y., Hirano,S., and Burgess,R.W. (1999b). Flamingo, a seven-pass transmembrane cadherin, regulates planar cell polarity under the control of Frizzled. *Cell* 98, 585.

- van Amerongen,R. and Berns,A. (2006). Knockout mouse models to study Wnt signal transduction. *Trends Genet.* 22, 678-689.
- Veeman,M.T., Axelrod,J.D., and Moon,R.T. (2003a). A second canon. Functions and mechanisms of beta-catenin-independent Wnt signaling. *Dev. Cell* 5, 367-377.
- Veeman,M.T., Slusarski,D.C., Kaykas,A., Louie,S.H., and Moon,R.T. (2003b). Zebrafish Prickle, a modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements. *Curr. Biol.* 13, 680.
- Vinson,C.R. and Adler,P.N. (1987). Directional non-cell autonomy and the transmission of polarity information by the frizzled gene of Drosophila. *Nature* 329, 549.
- Vinson,C.R., Conover,S., and Adler,P.N. (1989). A Drosophila tissue polarity locus encodes a protein containing seven potential transmembrane domains. *Nature* 338, 263.
- Wada,H., Iwasaki,M., Sato,T., Masai,I., and Nishiwaki,Y. (2005). Dual roles of zygotic and maternal Scribble1 in neural migration and convergent extension movements in zebrafish embryos. *Development* 132, 2273.
- Wallingford,J.B. (2005). Neural tube closure and neural tube defects: studies in animal models reveal known knowns and known unknowns. *Am. J. Med. Genet. C Semin. Med. Genet.* 135, 59.
- Wallingford,J.B., Fraser,S.E., and Harland,R.M. (2002). Convergent extension: the molecular control of polarized cell movement during embryonic development. *Dev. Cell* 2, 695-706.
- Wallingford,J.B. and Habas,R. (2005). The developmental biology of Dishevelled: an enigmatic protein governing cell fate and cell polarity. *Development* 132, 4421-4436.
- Wang,J., Hamblet,N.S., Mark,S., Dickinson,M.E., Brinkman,B.C., Segil,N., Fraser,S.E., Chen,P., Wallingford,J.B., and Wynshaw-Boris,A. (2006a). Dishevelled genes mediate a conserved mammalian PCP pathway to regulate convergent extension during neurulation. *Development* 133, 1767-1778.
- Wang,J., Mark,S., Zhang,X., Qian,D., Yoo,S.J., Radde-Gallwitz,K., Zhang,Y., Lin,X., Collazo,A., Wynshaw-Boris,A., and Chen,P. (2005). Regulation of polarized extension and planar cell polarity in the cochlea by the vertebrate PCP pathway. *Nat. Genet.* 37, 980-985.
- Wang,Y., Guo,N., and Nathans,J. (2006b). The role of Frizzled3 and Frizzled6 in neural tube closure and in the planar polarity of inner-ear sensory hair cells. *J. Neurosci.* 26, 2147-2156.

- Wang,Y. and Nathans,J. (2007). Tissue/planar cell polarity in vertebrates: new insights and new questions. *Development* 134, 647-658.
- Warga,R.M. and Nusslein-Volhard,C. (1999). Origin and development of the zebrafish endoderm. *Development* 126, 827.
- Watanabe,D., Saijoh,Y., Nonaka,S., Sasaki,G., Ikawa,Y., Yokoyama,T., and Hamada,H. (2003). The left-right determinant Inversin is a component of node monocilia and other 9+0 cilia. *Development* 130, 1725-1734.
- Weber,U., Paricio,N., and Mlodzik,M. (2000). Jun mediates Frizzled induced R3/R4 cell fate distinction and planar polarity determination in the Drosophila eye. *Development* 127, 3619.
- Westfall,T.A., Brimeyer,R., Twedt,J., Gladon,J., Olberding,A., Furutani-Seiki,M., and Slusarski,D.C. (2003a). Wnt-5/pipetuin functions in vertebrate axis formation as a negative regulator of Wnt/beta-catenin activity. *J. Cell Biol.* 162, 889-898.
- Westfall,T.A., Hjertos,B., and Slusarski,D.C. (2003b). Requirement for intracellular calcium modulation in zebrafish dorsal-ventral patterning. *Dev. Biol.* 259, 380-391.
- Wharton,K.A., Jr. (2003). Runnin' with the Dvl: proteins that associate with Dsh/Dvl and their significance to Wnt signal transduction. *Dev. Biol.* 253, 1-17.
- Winter,C.G., Wang,B., Ballew,A., Royou,A., and Karess,R. (2001). Drosophila Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton. *Cell* 105, 81.
- Witzel,S., Zimyanin,V., Carreira-Barbosa,F., Tada,M., and Heisenberg,C.P. (2006). Wnt11 controls cell contact persistence by local accumulation of Frizzled 7 at the plasma membrane. *J. Cell Biol.* 175, 791-802.
- Wodarz,A. and Nusse,R. (1998). Mechanisms of Wnt signaling in development. *Annual Review of Cell and Developmental Biology* 14, 59-88.
- Wolff,T. and Rubin,G.M. (1998). strabismus, a novel gene that regulates tissue polarity and cell fate decisions in Drosophila. *Development* 125, 1149.
- Wong,H.C., Bourdelas,A., Krauss,A., Lee,H.J., and Shao,Y. (2003). Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in the C-terminal region of Frizzled. *Mol. Cell.* 12, 1251.

- Wong,H.C., Mao,J., Nguyen,J.T., Srinivas,S., Zhang,W., Liu,B., Li,L., Wu,D., and Zheng,J. (2000). Structural basis of the recognition of the dishevelled DEP domain in the Wnt signaling pathway. *Nat. Struct. Biol.* 7, 1178-1184.
- Yamanaka,H., Moriguchi,T., Masuyama,N., Kusakabe,M., Hanafusa,H., Takada,R., Takada,S., and Nishida,E. (2002). JNK functions in the non-canonical Wnt pathway to regulate convergent extension movements in vertebrates. *EMBO Rep.* 3, 69-75.
- Yoneda,A., Ushakov,D., Multhaupt,H.A., and Couchman,J.R. (2007). Fibronectin matrix assembly requires distinct contributions from Rho kinases I and -II. *Mol. Biol. Cell* 18, 66-75.
- Zhu,S., Liu,L., Korzh,V., Gong,Z., and Low,B.C. (2006). RhoA acts downstream of Wnt5 and Wnt11 to regulate convergence and extension movements by involving effectors Rho kinase and Diaphanous: use of zebrafish as an in vivo model for GTPase signaling. *Cell Signal.* 18, 359-372.