

8. Literaturverzeichnis

- Alcaini M, Giard MH, Thevenet M, Pernier J (1994): Two separate frontal components in the N1 wave of the human auditory evoked response. *Psychophysiology* 31: 611-615
- Almasy L, Porjesz B, Blangero J, Chorlian DB, O'Conner SJ, Kuperman S, Rohrbaugh J, Bauer LO, Reich T, Polich J, Begleiter H (1988): Heritability of event-related brain potentials in families with a history of alcoholism. *Am.J.Med.Genet.*: 383-390
- Amati F, Conti E, Novelli A, Bengala M, Diglio MC, Marino B, Gianotti A, Gabrielli O, Novelli G, Dallapiccola B (1999): Atypical deletions suggest five 22q11.2 critical regions related to the DiGeorge/velo-cardio-facial syndrome. *Eur J Hum Genet* 7(8): 903-9
- Arinami T, Gao M, Hamaguchi H, Toru M (1997): A functional polymorphism in the promoter region of the dopamine D2 receptor gene is associated with schizophrenia. *Hum Mol Genet*. Apr.6(4): 577-82.
- Badner JA, Gershon ES (2002): Meta-analysis of whole-genome linkage scans of bipolar disorder and schizophrenia. *Mol Psychiatry*. 7(4): 405-11.
- Barden N, Morissette J (1999): Chromosome 13 workshop report. *Am J Med Genet*; 88 (3): 260-2
- Bassett AS (1992) Chromosomal aberrations and schizophrenia Autosomes. *Br J Psychiatry* 161: 323-34
- Begleiter H, Porjesz B (1995): Neurophysiological phenotypic factors in the development of alcoholism. In Begleiter H, Kissin B (Eds.), *Alcohol and Alcoholism*, Vol. 1, The Genetics of Alcoholism. Oxford University Press, New York: 269-293
- Birbaumer N, Schmidt RF (1996): Biologische Psychologie Springer Verlag Berlin Heidelberg: 534
- Blackwood DHR, St. Clair DM, Muir WJ, Duffy JC (1991): Auditory P300 and eye tracking dysfunktion in schizophrenic pedigrees. *Arch Gen Psychiatry* 48: 899-909
- Bleuler E. (1911): Dementia praecox oder Gruppe der Schizophrenien. Deutike Wien

- Breier A, Su TP, Saunders R, Carson RE, Kolachana BS, de Bartolomeis A, Weinberger DR, Weisenfeld N, Malhotra AK, Eckelman WC, Pickar D (1997): Schizophrenia is associated with elevated amphetamine induced synaptic dopamine concentrations: evidence from a novel positron emission tomography method. *Proc Natl Acad Sci USA* 94(6): 2569-74
- Bruder GE, Kayser J, Tenke CE, Friedman M, Malaspina D, Gorman JM (2001): Event-related potentials in schizophrenia during tonal and phonetic oddball tasks: relations to diagnostic subtype, symptom features and verbal memory. *Biol Psychiatry* 50: 447-452
- Burnet PW, Eastwood SL, Harrison PJ (1997): (3H)WAY-100635 for 5-HT1A receptor autoradiography in human brain: a comparison with (3H)8-OH-DPAT and demonstration of increased binding in the frontal cortex in schizophrenia. *Neurochem Int* 30(6): 565-74
- Cannon TD, Gasperoni TL, Van Erp TGM, Rosso IM (2001) Quantitative neural indicators of liability to schizophrenia: implications for molecular genetic studies. *Am J Med Genet* 105: 16-19
- Carlsson A (1988): The current status of the dopamine hypothesis of schizophrenia. *Neuropharmacology* 1(3): 179-86 Review
- Carlsson A, Hansson LO, Waters N, Carlsson ML (1999): A glutamatergic deficiency model of schizophrenia. *Br J Psychiatry Suppl*: 2-6.
- Carter CS, Mintun M, Nichols T, Cohen JD (1997): Anterior cingulate gyrus dysfunction and selective attention deficits in schizophrenia: [15O]H₂O PET study during single-trial Stroop task performance. *Am J Psychiatry* 154: 1670-1675.
- Chen WJ, Faraone SV (2000): Sustained attention deficits as markers of genetic susceptibility to schizophrenia. *Am J Med Genet* 97: 52-57.
- Cleghorn JM, Garnett ES, Nahmias C, Brown GM, Kaplan RD, Szechtman H, Szechtman B, Franco S, Dermer SW, Cook P (1990): Regional brain metabolism during auditory hallucinations in chronic schizophrenia. *Br J Psychiatry Oct*; 157: 562-70
- Cornblatt BA, Malhotra AK (2001) Impaired attention as an endophenotype for molecular genetic studies of schizophrenia. *Am J Med Genet* 105: 11-15

- Craddock N, Lendon C (1999): Chromosome workshop: chromosomes 11, 14 and 15. *Am J Med Genet*; 88 (3): 244-54
- Crowe RR, Vieland V (1999): Report of the Chromosome 5 Workshop of the Sixth World Congress on Psychiatric Genetics. *Am J Med Genet*; 88 (3): 229-32
- Curtis D (1999): Chromosome 21 workshop. *Am J Med Genet*; 88 (3): 272-5
- Davis KL, Kahn RS, Ko G, Davidson M (1991): Dopamine in schizophrenia: a review and reconceptualization. *Am J Psychiatry*. Nov; 148(11): 1474-86. Review.
- Detera-Wadleigh SD (1999): Chromosomes 12 and 16 workshop. *Am J Med Genet*; 88 (3): 255-9
- Dolan RJ, Fletcher P, Frith CD, Friston KJ, Frackowiak RS, Grasby PM (1995): Dopaminergic modulation of impaired cognitive activation in the anterior cingulate cortex in schizophrenia. *Nature* 378: 180-182
- Dubertret C, Gorwood P, Gouya L, Deybach JC, Ades J (2001): Association and excess of transmission of a DRD2 haplotype in a sample of French schizophrenic patients. *Schizophr Res.* 49(1-2): 203-12.
- East SZ, Burnet PW, Leslie RA, Roberts JC, Harrison PJ (2002): 5-HT₆ receptor binding sites in schizophrenia and following antipsychotic drug administration: autoradiographic studies with (125I)SB-258585. *Synapse* 45(3): 191-9
- Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE, Goldman D, Weinberger DR (2001): Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proc Natl Acad Sci USA* 98(12): 6917-22
- Eischen SE, Polich J (1994): P300 from families. *Electroencephalogr Clin Neurophysiol*. Jul.92(4): 369-372
- Emri M, Glaub T, Berecz R, Lengyel Z, Mikecz P, Repa I, Bartok E, Degrell I, Tron L (2006): Brain blood flow changes measured by positron emission tomography during an auditory cognitive task in healthy volunteers and in schizophrenic patients. *Prog Neuropsychopharmacol Biol Psychiatry*. Jan 9
- Faraone SV, Seidman LJ, Kremen WS, Toomey R, Pepple JR, Tsuang MT (2000): Neuropsychologic functioning among the nonpsychotic relatives of schizophrenic patients: the effect of genetic loading. *Biol Psychiatry* 48: 120-126

- Ford JM, Mathalon DH, Kalba S, Marsh L, Pfefferbaum A (2001): N100 and P300 abnormalities in patients with schizophrenia, epilepsy, and epilepsy with schizophrenialike features. *Biol Psychiatry* 49: 848-860
- Ford JM, White PM, Csernansky JG, Faustman WO, Roth WT, Pfefferbaum A (1994): ERPs in schizophrenia: Effects of antipsychotic medication. *Biol Psychiatry* 36: 153-170
- Franke P, Maier W, Hain C, Klingler T (1992) Wisconsin Card Sorting Test: an indicator of vulnerability to schizophrenia? *Schizophr Res* 6: 243-249
- Friedman D, Brown C, Vaughan HG, Cornblatt B, Kimling LE (1984): Cognitive brain potential components in adolescents. *Psychophysiology* 21: 83-96
- Gallinat J, Hegerl U (1994): Dipole source analysis. Linking scalp potentials to their generating neuronal structures. *Pharmacopsychiatry* 27: 52-53.
- Gallinat J, Mulert C, Bajbouj M, Herrmann WM, Schunter J, Senkowski D, Moukhtieva R, Kronfeldt D, Winterer G (2002): frontal and temporal dysfunction of auditory stimulus processing in schizophrenia. *NeuroImage* 17: 110-127.
- Gejman PV (1999): Chromosomes 19 and 20 report. *Am J Med Genet* 88(3):271
- Giard MH, Perrin F, Echallier JF, Thevenet M, Froment JC, Pernier J (1994): Dissociation of temporal and frontal components in the human auditory N1 wave: a scalp current density and dipole model analysis. *Electroencephalogr Clin Neurophysiol* 92: 238-252.
- Gill M, Hawi Z, O'Neill FA, Walsh D, Straub RE, Kendler KS (1996) Neurotrophin-3 gene polymorphisms and schizophrenia: no evidence for linkage or association. *Psychiatr Genet* 6(4): 183-6
- Goodin DS, Squires KC, Henderson BH, Starr A (1978): Age-related variations in evoked potentials to auditory stimuli in normal human subjects. *Electroencephal Clin Neurophysiol* 44: 447-458
- Gottesman II. (1991) Schizophrenia genesis: the origin of madness. New York: W.H. Freeman
- Gottesman L, Shields J (1972): Schizophrenia and Genetics: A Twin Study Vantage Point. Academic Press, New York.

- Grandy DK, Zhang Y, Civelli O (1993): PCR detection of the TaqA RFLP at DRD2 locus. *Hum Mol Genet* 2: 2197
- Hallmayer J (1999) Chromosomes 1, 2 and 7 workshop. *Am J Med Genet*; 88 (3): 219-23
- Harrison PJ (1999): Neurochemical alterations in schizophrenia affecting the putative receptor targets of atypical antipsychotics. Focus on dopamin (D1, D3, D4) and 5-HT2a receptors. *Br J Psychiatry Suppl* (38): 12-22
- Hauge XY, Crandy DK, Eubank JH, Evans GA, Civelli O, Litt M (1991): Detection and characterization of additional DNA polymorphisms in the D2 receptor gene. *Genomics* 10: 527-530
- Hawi Z, Straub RE, O'Neill A, Kendler KS, Walsh D, Gill M (1998): No linkage or linkage disequilibrium between brain-derived neurotrophic factor (BDNF) dinucleotide repeat polymorphism and schizophrenia in irish families. *Psychiatry Res* 81(2): 111-6
- Hegerl U (1998): Ereigniskorrelierte Potentiale. In U. Hegerl (Ed.), *Neurophysiologische Untersuchungen in der Psychiatrie: EEG, EKP, Schlafpolygraphie, Motorik, autonome Funktionen*. Wien, New York: Springer: 95-140
- Hegerl U, Klotz S, Ulrich G (1985): Späte akustisch evozierte Potentiale- Einfluss von Alter, Geschlecht und unterschiedlichen Untersuchungsbedingungen. *Z EEG-EMG* 16: 129-139
- Hetrick WP, Sandman CA, Bunney WE jr, Jin Y, Potkin SG, White MH (1996): Gender differences in gating of the auditory evoked potential in normal subjects. *Biol Psychiatry* 39 (1): 51-8
- Hillyard SA, Hink RF, Schwent VL, Picton TW (1973): Electrical signs of selective attention in the human brain. *Science* 182: 177-180
- Holcomb PJ, Ackerman PT, Dykman RA (1986): Auditory event-related potentials in attention and reading disabled boys. *Int. J. Psychophysiol.* 3: 263-273
- Holzman PS, Kringlen E, Levy DL, Proctor LR, Haberman SJ, Yasillo NJ (1977): Abnormal-pursuit eye movements in schizophrenia. Evidence for a genetic indicator. *Arch Gen Psychiatry* 34: 802-805
- Hovatta I, Lichtermann D, Juvonen H, Suvisaari J, Terwilliger JD, Arajarvi R, Kokko-Sahin ML, Ekelund J, Lonnqvist J, Peltonen L (1998): Linkage analysis of putative schizophrenia gene candidate regions on chromosomes 3p, 5q,

6p, 8p, 20p and 22q in a population-based sampled finnish family set. Mol Psychiatry. Sep; 3(5): 452-7.

Itokawa M, Arinami T, Futamura N, Hamaguchi H, Toru M (1993): A structural polymorphism of human dopamin D2 receptor, D2(Ser311-Cys). Biochem Biophys Res Commun 196 (Suppl 3): 1369-1375

Jasper HH (1958): The ten- twenty electrode system of the International Federation. J. Electroenceph. Clin. Neurophysiol. 20: 371-375

Jerger J, Weikers NJ, Sharbrough FW, Jerger S (1969): Bilateral lesions of the temporal lobe: A case study. Acta Otolaryngologica 258: 1-51

Johnstone SJ, Barry RJ, Anderson JW, Stephen FC (1996): Age-related changes in child and adolescent event related potential component morphology, amplitude and latency to standard and target stimuli in an auditory oddball task. International Journal of Psychophysiology 24: 223-238

Katsanis J, Iacono WG, McGue MK, Carlson SR (1998): P300 event-related potentials heritability in monozygotic and dizygotic twins. Psychophysiology Jan.34(1): 47-58

Kayser J, Bruder GE, Tenke CE, Stuart BK, Amador XF, Gormann JM (2001): Event-related brain potentials (ERPs) in schizophrenia for tonal and phonetic oddball tasks. Biol Psychiatry 49: 832-847

Kendler KS, Gruenberg AM, Kinney DK (1994): Independent diagnoses of adoptees and relatives as defined by DSM-3 in the provincial and national samples of the Danish Adoption Study of Schizophrenia. Arch Gen Psychiatry 51(6): 456-68

Kendler KS, Myers JM, O'Neill FA, Martin R, Murphy B, MacLean CJ, Walsh D, Straub RE (2000): Clinical features of schizophrenia and linkage to chromosomes 5q, 6p, 8p, and 10p in the Irish Study of High-Density Schizophrenia Families. Am J Psychiatry. Mar; 157(3): 402-8.

Kendler KS, Robinette CD (1983): Schizophrenia in the National Academy of Sciences-National Research Council Twin Registry: a 16-year update. Am J Psychiatry 140(12): 1551-63

Kennedy JL, Basile VS, Macciardi FM (1999): Chromosome 4 Workshop Summary: Sixth World Congress on Psychiatric Genetics, Bonn, Germany, October 6-10, 1998. Am J Med Genet; 88 (3): 224-8

Kohn Y, Lerer B (2002): Genetics of schizophrenia: a review of linkage findings. *Isr J Psychiatry Relat Sci.*; 39(4): 340-51

Krabbendam L, Marcelis M, Delespaul P, Jolles J, van Os J (2001): Single or multiple familial cognitive risk factors in schizophrenia? *Am J Med Genet* 105: 183-188

Krabbendam L, van Os J (2005): Schizophrenia and urbanicity: a major environmental influence-conditional on genetic risk. *Schizophr Bull.* Oct;31(4):795-9.

Kraepelin, E (1906): Lehrbuch der Psychiatrie. Barth Leipzig

Laruelle M (1998): Imaging dopamine transmission in schizophrenia. A review and meta analysis. *Q J Nucl med* 42(3): 211-21

Lerer B, Segman RH, Hamdan A, Kanyas K, Karni O, Kohn Y, Korner M, Lanktree M, Kaadan M, Turetsky N, Yakir A, Kerem B, Macciardi F (2003): Genome scan of Arab Israeli families maps a schizophrenia susceptibility gene to chromosome 6q23 and supports a locus at chromosome 10q24. *Mol Psychiatry*. May; 8(5): 488-98.

Lowitzsch K, Maurer K, Hopf HC (1983): Evozierte Potentiale in der klinischen Diagnostik. Thieme Stuttgart 180-195, 270-279

Massat I, Souery D, Del-Favero J, Van Gestell S, Serretti A, Macciardi F, Smeraldi E, Kaneva R, Adolfsson R, et al. (2002): Positive association of dopamin D2 receptor polymorphism with bipolar affective disorder in a european multicenter association study of affective disorders. *American Journal of Medical Genetics (Neuropsychiatric Genetics)* 114: 177-185

Maurer K, Leitner H, Schäfer E (1982): Akustisch Evozierte Potentiale. Enke Stuttgart 1-56, 211-242

Maziade M, Roy MA, Rouillard E, Bissonnette L, Fournier JP, Roy A, Garneau Y, Montgrain N, Potvin A, Cliche D, Dion C, Wallot H, Fournier A, Nicole L, Lavallee JC, Merette C (2001): A search for specific and common susceptibility loci for schizophrenia and bipolar disorder: a linkage study in 13 target chromosomes. *Mol Psychiatry*. Nov; 6(6): 684-93.

Mc Guffin P, Farmer AE, Gottesman II, Murray RM, Reveley AM (1984): Twin concordance for operationally defined schizophrenia. Confirmation of familiality and heritability. *Arch Gen Psychiatry* 41(6): 541-5

Mega MS, Cummings JL (1997): The Cingulate and Cingulate Syndromes. In M.R. Trimble & J.L. Cummings (Eds.) Contemporary Behavioral Neurology: 189-214

Michel F, Peronnet F (1982): L'hemianacusie, un deficit auditif dans un hemisphere.

Michie PT, Bearpark HM, Crawford JM, Glue LC (1990): The nature of selective attention effects on auditory event-related potentials. Biol Psychol 30: 219-250

Miller SA, Dykes DD, Polesky HF (1988): A simple salting out procedure for extracting DNA from human nucleated cells. Nucleic Acids. Res. 16: 1215

Mulert C, Gallinat J, Pascual-Marqui R, Dorn H, Frick K, Schlattmann P, Mientus S, Herrmann WM, Winterer G (2001): Reduced event-related current density in the anterior cingulate cortex in schizophrenia. Neuroimage 13: 589-600

Näätänen R (1987): Event related brain potentials in research of cognitive process - a classification of components. In: Van der Meer E, Hofmann J: Knowledge aided information processing Elsevier Amsterdam

Näätänen R, Picton T (1987): The N1 wave of the human electric and magnetic response to sound: a review and an analysis of the component structure. Psychophysiology 24: 375-425.

Nordahl TE, Carter CS, Salo RE, Kraft L, Baldo J, Salamat S, Robertson L, Kusubov N (2001): Anterior cingulate metabolism correlates with stroop errors in paranoid schizophrenia patients. Neuropsychopharmacology Jul; 25(1): 139-48

Nordstrom AL, Farde L, Eriksson L, Halldin C (1995): No elevated D2 dopamine receptors in neuroleptic-naive schizophrenic patients revealed by positron emission tomography and (11C)N-methylspiperone. Psychiatry Res 61 (2): 67-83

Nüchterlein KH, Dawson ME (1984): Information processing and attentional functioning in the developmental course of schizophrenic disorders. Schizophr.Bull.10: 160-203

Nurnberger Jr JI, Foroud T (1999): Chromosome 6 workshop report. Am J Med Genet; 88 (3): 233-8

O'Connor S, Morzorati S, Christian JC, Li TK (1994): Heritable features of the auditory oddball event-related potential: peaks latencies, morphology and topography. Electroencephalography Clin. Neurophysiol. 92: 115-125

O'Donnell BF, Shenton ME, Mc Carley RW, Faux SF, Smith RS, Salisbury DF, et al. (1993): The auditory N2 component in schizophrenia: Relationship to MRI temporal lobe gray matter and to other ERP abnormalities. *Biol Psychiatry* 34: 26-40

Okubo Y, Olsson H, Ito H, Lofti M, Suhara T, Halladin C, Farde L (1999) : PET mapping of extrastriatal D2-like dopamine receptors in the human brain using an anatomic standardization technique and [11C]FLB 457. *Neuroimage* Dec; 10(6): 666-74

Olney JW, Farber NB (1995): Glutamate receptor dysfunction and schizophrenia [see comments]. *Arch Gen Psychiatry* 52: 998-1007

Onstad S, Skre I, Torgersen S, Kringlen E (1991): Twin concordance for DSM-III-R schizophrenia. *Acta Psychiatr Scand.* 83(5): 395-401

Oruc L, Furac I, Croux C, Jakovljevic M, Kracun I, Folnegovic and Van Broeckhoven C (1996): Association study of bipolar disorder with candidate genes involved in dopamine-serotonin metabolism and GABAergic neurotransmission: a preliminary report. *Psychiatr Genet* 6: 213-217

Papanicolaou AC, Rogers RL, Baumann S, Saydjadi C, Eisenberg, HM (1990): Source localization of two evoked magnetic field components using two alternative procedures. *Exp Brain Res* 80: 44-48.

Parsian A, Chakraverty S, Todd RD (1995): Possible association between the dopamin D3 receptor gene and bipolar affective disorder. *Am J Med Genet* 60: 234-237

Paterson AD (1999): Sixth World Congress of Psychiatric Genetics X Chromosome Workshop. *Am J Med Genet*; 88 (3): 279-86

Pfefferbaum A, Ford JM, Roth WT, Kopell BS (1980): Age related Changes in Auditory Event- Related Potentials. *Electroencephal Clin Neurophysiol* 49: 266-267

Pfefferbaum A, Ford JM, White PM, Roth WT (1989): P3 in schizophrenia is affected by stimulus modality, response requirements, medication status and negative symptoms. *Arch Gen Psychiatry* 46: 1035-1046

Picton TW, Hillyard SA, Krausz HI, Galambos R (1974): Human auditory evoked potentials. I. Evaluation of components. *Electroencephalogr Clin Neurophysiol* 36: 179-190.

- Picton TW, Stuss DT, Champagne SC, Nelson RF (1984): The effects of age on human event-related potentials. *Psychophysiology* 21: 312-325
- Polich J, Burns T (1984): P300 from identical twins. *Neuropsychologia* 25: 299-304
- Polich J, Ladish C, Burns T (1990): Normal variation of P300 in children: age, memory span, and head size. *Int. J. Psychophysiol.* 9: 237-248
- Posner MI, Desimone R (1998): Beyond images [editorial]. *Curr Opin Neurobiol* 8: 175-177
- Pulver AE, Mulle J, Nestadt G, Schwartz KL, Blouin JL, Dombroski B, Liang KY, Housman DE, Kazazian HH, Antonarakis SE, Lasseter VK, Wolyniec PS, Thornquist MH, McGrath JA (2000): Genetic heterogeneity in schizophrenia: stratification of genome scan data using co-segregating related phenotypes. *Mol Psychiatry*. Nov; 5(6): 650-3.
- Raymond JR, Mukhin YV, Gelasco A, Turner J, Collinsworth G, Gettys TW, Grewal JS, Garnovskaya MN (2001): Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacol Ther* 92(2/3): 179-212
- Risch N (1990): Linkage strategies for genetically complex traits. I. Multilocus models. *Am J Hum Genet* 46: 222-8
- Risch N, Merikangas K (1996): The future of genetic studies of complex human diseases. *Science* 273: 1516-7
- Rockstroh B (1989) Slow cortical potentials and behaviour. Urban and Schwarzenberg Baltimore
- Roth WT, Horvath TB, Pfefferbaum A, Kopell BS (1980) Event related potentials in schizophrenics. *Electroencephalog Clin Neurophysiol* 48: 127-139
- Sachs L (1978): *Angewandte Statistik* Springer Verlag Berlin Heidelberg New York S.219
- Sandman CA, Patterson JV (2000): The auditory event-related potential is a stable and reliable measure in elderly subjects over a 3 year period. *Clinical Neurophysiology* 111 : 1427-1437
- Sarkar G, Kapelner S, Grandy DK, Marchionni M, Civelli O Sobell J, Heston L, Sommer S (1991): Direct sequencing of the dopamin D2 receptor (DRD2) in schizophrenics reveals three polymorphisms but no structural change in the receptor. *Genomics* 11: 8-14

- Scherg M, Picton TW (1991): Separation and identification of event-related potential components by brain electric source analysis. *Electroencephalogr Clin Neurophysiol Suppl* 42: 24-37
- Scherg M, Berg P (1991): Use of prior knowledge in brain electromagnetic source analysis. *Brain Topogr* 4: 143-150
- Schneider K (1939) *Psychischer Befund und psychiatrische Diagnose*. Thieme Leipzig
- Schreiber H, Born GS, Kornhuber HH, Born J (1992): Event-related potential correlates of impaired selective attention in children at high risk for schizophrenia. *Biol. Psychiatry* 32: 634-651
- Schultz SK, O'Leary DS, Boles Ponto LL, Arndt S, Magnotta V, Watkins GL, Hichwa RD, Andreasen NC (2002): Age and regional cerebral blood flow in schizophrenia: age effects in anterior cingulate, frontal and parietal cortex. *J Neuropsychiatry Clin Neurosci.* Winter; 14: 19-24
- Schwab SG, Wildenauer DB (1999): Chromosome 22 workshop report. *Am J Med Genet*; 88 (3): 276-8
- Seidman LJ, Buka SL, Goldstein JM, Horton NJ, Rieder RO, Tsuang MT (2000): The relationship of prenatal and perinatal complications to cognitive functioning at age 7 in the New England Cohorts of the National Collaborative Perinatal Project. *Schizophr Bull.* ;26(2):309-21.
- Seidman LJ, Faraone SV, Goldstein JM et al. (2002): Left hippocampal volume as a vulnerability indicator for schizophrenia: a magnetic resonance imaging morphometric study of nonpsychotic first-degree relatives. *Arch Gen Psychiatry* 59: 839-849
- Seretti A, Macchiardi F, Smeraldi E (1998): Dopamine receptor D2 Ser/Cys311 variant associated with disorganized symptomatology of schizophrenia. *Schizophr Res.* Nov 30; 34(3): 207-10
- Shagass C, Straumanis JJ Jr, Roemer RA, Amadeo M (1977): Evoked potentials of schizophrenics in several sensory modalities. *Biol Psychiatry* 12: 221-235
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavas J, Weiller E, Hergueta T, Baker R, Dunbar GC (1998): The mini-international neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 59 Suppl 20: 22-33; quiz 34-57

- Shelley AM, Silipo G, Javitt DC (1999): Diminished responsiveness of ERPs in schizophrenic subjects to changes in auditory stimulation parameters: Implications for theories of cortical dysfunction. *Schizophr Res* 37(1): 65-79
- Shelley AM, Ward PB, Cats SV, Michie PT, Andrews S, McConaghy N (1991): Mismatch negativity: An index of a preattentive processing deficit in schizophrenia. *Biol Psychiatry* 30: 1059-1062
- Siedenberg R, Goodin DS, Aminoff MJ, Rowley HA, Roberts TP (1996): Comparison of late components in simultaneously recorded event-related electrical potentials and event-related magnetic fields. *Electroencephalogr Clin Neurophysiol* 99: 191-197
- Sklar P (2002): Linkage analysis in psychiatric disorders: the emerging picture. *Annu Rev Genomics Hum Genet.* 3: 371-413.
- Souery D, Lipp O, Mahieu B, Mendelbaum K, De Martelaer V, Van Broeckhoven C, Mendlewicz J (1996): Association study of bipolar disorder with candidate genes involved in catecholamine neurotransmission: DRD2, DRD3, DAT1, and TH genes. *Am J Med Genet* 67: 551-555
- Squires KC, Goodin DS, Starr A (1978): Auditory Evoked Potentials in Ageing and Dementia. In: Barber C(Hrsg): *Evoked Potentials. International Evoked Potentials Symp.*, Nottingham, England. Lancaster MIP Press (1980): 353-355
- Stefansson H, Sigurdsson E, Steinthorsdottir V, Bjornsdottir S, Sigmundsson T, Ghosh S, Brynjolfsson J, Gunnarsdottir S et al. (2002): Neuregulin 1 and susceptibility to schizophrenia. *Am J Hum Genet* 71(4): 877-92
- Suhara T, Okubo Y, Yasuno F, Sudo Y, Inoue M, Ichimiya T, Nakashima Y, Nakayama K, Tanada S, Suzuki K, Halldin C, Farde L (2002): Decreased dopamine D2 receptor binding in anterior cingulated cortex in schizophrenia. *Arch Gen Psychiatry* Jan; 59(1): 25-30
- Tauscher J, Kapur S, Verhoeff NP, Hussey DF, Daskalakis ZJ, Wisniewski ST, Wilson AA, Houle S, Kasper S, Zipursky RB (2002): Brain serotonin 5-HT(1A) receptor binding in schizophrenia measured by positron emission tomography and (11C)WAY-100635. *Arch Gen Psychiatry* 59(6): 514-20
- Tienari P (1991): Interaction between genetic Vulnerability and family environment: the Finnish adoptive family study of schizophrenia. *Acta Psychiatr Scand* 84(5): 460-5

- Van Beijsterveldt CEM, Molenaar PCM, de Geus EJC, Boomsma DI (1998): Individual differences in P300 amplitude: A genetic study in adolescent twins. *Biol. Psychol.* 47: 97-120
- Van Broeckhoven C, Verheyen G (1999): Report of the chromosome 18 workshop. *Am J Med Genet*; 88 (3): 263-70
- Vawter MP, Barrett T, Cheadle C, Sokolov BP, Wood WH, Donovan DM, Webster M, Freed WJ, Becker KG (2001): Application of cDNA microarrays to examine gene expression differences in schizophrenia. *Brain Res Bull* 55(5): 641-50
- Virgos C, Martorell L, Valero J, Figuera L, Civeira F, Joven J, Labad A, Vitella E (2001): Association study of schizophrenia with polymorphisms at six candidate genes. *Schizophr Res.* 49(1-2): 65-71
- Vogeley K, Axmann TS, Pfeiffer U et al. (2000): Disturbed gyration of the prefrontal region in male schizophrenic patients: A morphometric postmortem study. *Am J Psychiatry* 157: 34-39
- Vollenweider FX, Leenders KL, Oye I, Hell D, Angst J (1997): Differential psychopathology and patterns of cerebral glucose utilisation produced by (S)- and (R)-ketamine in healthy volunteers using positron emission tomography (PET). *Eur Neuropsychopharmacol.* Jan; 7(1): 25-38
- Wagner M, Rendtorff N, Kathmann N, Engel RR (1996): CNV, Pinv and probe-evoked potentials in schizophrenics. *Electroencephalogr Clin Neurophysiol* 98: 130-143
- Walhovd KB , Fjell AM (2002): One year test-retest reliability of auditory ERPs in young and old adults. *International Journal of Psychophysiology* 46: 29-40
- Wildenauer DB, Schwab SG (1999): Chromosomes 8 and 10 workshop. *Am Jmed Genet*; 88 (3): 239-43
- Williams J, Spurlock G, Mc Guffin P, Mallet J, Nothen MM, Gill M, Aschauer H, Nylander PO, Macciardi F, Owen MJ (1996): Association between schizophrenia and T102C polymorphism of the 5-hydroxytryptamine type 2a-receptor gene. European multicentre Association Study of schizophrenia (EMASS) Group. *Lancet* 347(9011): 1294-6
- Wong AHC, Hubert HM Van Tol. (2003): Schizophrenia: from phenomenology to neurobiology. *Neuroscience and Biobehavioral Reviews* 27: 269-306
- Wright P, Murray RM (1993): Schizophrenia: prenatal influenza and autoimmunity. *Ann Med. Oct*;25(5):497-502. Review.

Yasuno F, Suhara T, Okubo Y, Ichimiya T, Takano A, Sudo Y, Inoue M (2005): Abnormal effective connectivity of dopamine D2 receptor binding in schizophrenia. Psychiatry Res. Apr 30; 138(3): 197-207

Yucel M, Pantelis C, Stuart GW, Wood SJ, Maruff P, Velakoulis D, Pipingas A, Crowe SF, Tochon-Danguy HJ, Egan GF (2002): Anterior cingulated activation during stroop task performance: a PET to MRI coregistration study of individual patients with schizophrenia. Am J Psychiatry Feb; 159(2): 251-4

Zobel A, Maier W (2003): Endophänotypen ein neues Konzept zur biologischen Charakterisierung psychischer Störungen. Der Nervenarzt Dez; Springer Verlag