

9 Literaturverzeichnis

A

Adachi, K.; Kohara, T.; Nakao, N.: Design, synthesis and structure activity relationships of 2-substituted 2-amino-1,3-propanediols: discovery of a novel immunosuppressant, FTY 720. *Bioorg. Med. Chem. Lett.* 5 (1995) 853-856.

Adrian, M.; Dubochet, J.; Lepault, J.; McDowell, A.W.: Cryo-electron microscopy of viruses, *Nature* 308 (1984) 32 – 36.

Ahmed, I., Gokhale, R.D., Shah, M.V., Patton, T.F., Physicochemical determinants of drug diffusion across the conjunctiva, sclera, and cornea, *J. Pharm. Sci.* 76 (1987) 583-586.

Allen, T. M.; Redemann, C.; Martin, F.; Hansen, C.: Liposomes containing synthetic lipid derivatives of polyethylene glycol show prolonged circulation half-lives in vivo. *Biochim. Biophys. Acta, Biomembranes* 1066 (1991) 29-36.

AMK Mitteilungen 377/47/05: Arzneimittelkommission der Deutschen Apotheker, in: *Pharmazeutische Zeitung* 150 (2005) 7 103.

AMK Mitteilungen 71/05: Arzneimittelkommission der Deutschen Apotheker, in: *Deutsche Apothekerzeitung* 145 (2005) 3245.

AMK Mitteilungen 117/14/06, in: *Pharmazeutische Zeitung* 151 (2006) 1382.

Arndt, D; Fichtner, I: *Liposomen: Darstellung, Eigenschaften, Anwendung*, Akad. Verlag Berlin, 1986.

Arnoux, P.; Morrison, R.: Drug analysis of biological samples. A survey of validation approaches in chromatography in the UK pharmaceutical industry, *Xenobiotica* 22 (1992) 757.

Arzneimitteltelegramm 16.2.05, Tusann Press, Rue de Provence 4, F-67116 Reichstett.

Arzneistoffprofile – Basisinformationen über Arzneiliche Wirkstoffe, GOVI-Verlag GmbH, Frankfurt/Main, 18. Ergänzungslieferung 2003.

Ashcroft, D. M.; Dimmock, P.; Garside, R.; Stein, K.; Williams, H. C.: Efficacy and tolerability of topical pimecrolimus and tacrolimus in the treatment of atopic dermatitis: meta-analysis of randomised controlled trials, *BMJ [British Medical Journal]* (2005) 516-522.

Ashton, P., Podder, S.K., Lee, V.H.L.: Formulation influence on conjunctival penetration of four beta-blockers in the pigmented rabbit: a comparison with corneal penetration, *Pharm. Res.* 8 (1991) 1166-1174.

B

Baban, D. F.; Seymour L. W.: Control of tumour vascular permeability, *Adv. Drug Del. Rev.* 34 (1998) 109-119.

Badwan, A. A.; El-Khordagui, L. K.; Saleh, A. M.; Khalil, S. A.: The solubility of benzodiazepines in sodium salicylate solution and a proposed mechanism for hydrotropic solubilization, *Int. J. Pharm.* 13 (1982) 67-74.

Bahri A.M., Heyne B.J., Hans P., Secret A.E., Mouithys-Mickalad A.A., Hoebeke M.D.: Quantification of lipid bilayer effective microviscosity and fluidity effect induced by propofol, *Biophys. Chem.* 114 (2005) 53-61.

Balasubramanian, D.; Srinivas, V.; Gaikar, V. G.; Sharma, M. M: Aggregation behavior of hydrotropic compounds in aqueous solution, *J. Phys. Chem.* 93 (1989) 3865-3870.

Baldelli, S., Murgia, S., Merlini, S.; Zenoni, S.; Perico, N.; Remuzzi, G.; Cattaneo, D.: High-performance liquid chromatography with ultraviolet detection for therapeutic drug monitoring of everolimus, *J. Chromat. B* 816 (2005) 99-105.

Bangham, A. D.; Horne, R. W.: Negative staining of phospholipids and their structural modification by surface-active agents as observed in the electron microscope, *J. Mol. Biol.* 8 (1964) 660-668.

Bangham, A. D.; Standish, M. M.; Watkins, J. C.: Diffusion of univalent ions across the lamellae of swollen phospholipids, *J. Mol. Biol.* 13 (1965) 238-252.

Barry, B.W.: Mode of action of penetration enhancers in human skin, *J. Control. Rel.* 6 (1987a) 85-97.

Barry, B.W.: Penetration enhancers. In: Shroot, B.; Schaefer, H. (Hg.): *Skin Pharmacokinetics*, Karger, Basel, 1987b.

Bartlett, J.D.; Jaanus, S.D.: *Clinical Ocular Pharmacology*, Butterworth Publishers Stoneham, USA, 1989.

Bartolone J.; Santhanam U.; Penksa, C.; Lang B.: Hydroxy acids modulate skin cell biology, *J. Invest. Dermatol.* 104 (1995) 609.

Bauduin, P.: Characterization of short polypropylene glycol monoalkylethers and design of enzymatic reaction media, *Dissertation, Universität Regensburg*, 2005.

Bauer, S.: Interaktionen im oxidativen Stoffwechsel von Cyclosporin A, *Dissertation, Humboldt Universität zu Berlin*, 1993.

Bentley, M.V.L.B.; Vianna, R. F.; Wilson, S.; Collet, J.H.: Characterisation of the influence of some cyclodextrins on the stratum corneum from hairless mouse, *J. Pharm. Pharmacol.* 49 (1997) 397-402.

Berdel, W.; Panse, J.: Stellenwert der Angiogenese-Inhibition bei Tumorerkrankungen – präklinische Daten. *Onkol.* 28 Suppl. 4 (2005) 25-28.

- van den Bergh, B. A. I.; Salomons-de Vries, I.; Bouwstra, J. A.: Interactions between liposomes and human stratum corneum studied by freeze-substitution electron microscopy, *Int. J. Pharm.* 167 (1998) 57-67.
- van den Bergh, B.A.I.; Vroom, J.; Gerritsen, H.; Junginger, H. E.; Bouwstra, J. A.: Interactions of elastic and rigid vesicles with human skin in vitro: electron microscopy and two-photon excitation microscopy, *Biochim. Biophys. Acta* 1461 (1999) 155-173.
- van den Berg, B. A. I., Wertz, P. W., Junginger, H. E., Bouwstra, J. A.: Elasticity of vesicles assessed by electron spin resonance, electron microscopy and extrusion measurements, *Int. J. Pharm.* 217 (2001) 13-24.
- Bergethon, P. R.: *The physical basis of biochemistry: the foundations of molecular biophysics*, Springer, New York, 1998, pp. 342.
- Betz, G.; Imboden, R.; Imanidis, G.: Interactions of liposome formulation with human skin in vitro, *Int. J. Pharm.* 229 (2001) 117-129.
- Bietti G. B., Virno, M., und Pecori-Giraldi, J. Propylenglycol. A new osmotic agent for ophthalmic uses. *Documenta Ophthalmol.* 34 (1973) 77-92.
- Billich, A.; Aschauer, H.; Aszodi, A.; Stuetz, A.: Percutaneous absorption of drugs used in atopic eczema: pimecrolimus permeates less trough skin than corticosteroids and tacrolimus, *Int. J. Pharm.* 269 (2004) 29-34.
- Blume, A.; Jansen, M.; Ghyczy, M.; Gareiss, J.: Interaction of phospholipid liposomes with lipid modell mixtures for stratum corneum lipids, *Int. J. Pharm.* 99 (1993) 219-228.
- Blume, G.; Cevc, G: Drug-carrier and stability properties of the long-lived lipid vesicles, cryptosomes, in vitro and in vivo. *J. Liposome Res.* 2, 355-368 (1992).
- Bodde, H.E.; van den Brink, I.; Koerten, H.K.; de Haan, F.H.N.: Visualization of in vitro percutaneous penetration of mercuric chloride; transport trough intercellular space versus cellular uptake trough desmosomes, *J. Control. Rel.* 15 (1991) 1155-1159.
- Böhler, T.; Waiser, J.; Budde, K.; Neumayer, H.H.: SDZ RAD und FTY 720 Zukunftsperspektiven in der Transplantationsimmunologie, *innovartis* 4 (1998) 20-24.
- Böhler T; Waiser, J.; Schütz, M.; Budde, K.: FTY 720: Development of a novel and unique, immunomodulating drug that has the potential to revolutionize transplantation medicine, *Transplant* 4 (2002) 116-123.
- Bornhövd, E. C.; Schuller, E.; Bieber, T.; Wollenberg, A.: Immunsuppressive Makrolide und ihr Einsatz in der Dermatologie, *Hautarzt* 51 (2000) 646-654.
- Borel, J. F.; Feurer, Camille; Gubler, H. U.; Staehelin, H. Biological effects of cyclosporin A: a new antilymphocytic agent, *Agents Actions* 6 (1976) 468-475.

Bousema, M.T.; Tank, B.; Heule, F.; Naafs, B.; Stolz, E.; van Joost, T.: Placebo-controlled study of psoriasis patients treated topically with a 10 % cyclosporine gel, *J. Am. Acad. Dermatol.* 22 (1990) 126-127.

Bouwstra, J.A.; Gooris, G.S. ; Dubbelaar, F.E.R. ; Ponc, M.: Phase behaviour of lipid mixtures based on human ceramides: coexistence of crystalline and liquid phases, *J. Lipid Res.* 42 (2001) 1759–1770.

Bouwstra, J.A., Ponc, M.: The skin barrier in healthy and diseased state, *Biochim. Biophys. Acta* 2006, Artikel ist noch im Druck.

Brechue, W.F., Maren, T.H.: pH and drug ionization affects ocular pressure lowering of topical carbonic anhydrase inhibitors, *Invest. Ophthalmol. Vis. Sci.* 34 (1993) 2581-2587.

Brinkmann, V.: FTY 720: Mechanism of action and potential benefit in organ transplantation, *Yonsei Med. J.* 45 (2004) 991-994.

Brinkmann, V.; Cyster, J. G.; Hla, T.: FTY 720: Sphingosin 1-phosphate receptor in the control of lymphocyte egress and endothelial barrier function, *Am. J. Transpl.* 4 (2004) 1019-1025.

Bucke, W. E.: Liposomen zur Behandlung von Mykobakterieninfektionen, Dissertation, Freie Universität Berlin, Berlin, 1996.

Buckton, G.; Armstrong, J. K.; Chowdry, B. Z.; Leharne, S. Beezer, E.: The use of high sensitivity differential scanning calorimetry to characterise dilute aqueous dispersions of surfactants: 2. Further studies on polyoxyethylene alkyl ethers, *Int. J. Pharm.* 110 (1994) 179–187.

Bullingham R.E.; Nicholls A.J.; Kamm B.R.: Clinical pharmacokinetics of mycophenolate mofetil, *Clin. Pharmacokinet.* 34 (1998) 429-455.

Burns, M.K.; Ellis, C.N.; Eisen, D.; Duell, E.; Griffiths, C.E.M.; Annesley, T.M.; Hammilton, T.A.; Birnbaum, J.E.; Voorhees, J.J.: Intralesional cyclosporine for psoriasis. Relationship of dose, tissue levels, and efficacy, *Arch. Dermatol.* 128 (1992) 786-90.

C

Canet D.: NMR-Konzepte und Methoden, Springer Verlag Berlin Heidelberg, 1994.

Carney, L.G.; Hill, R.M.: Human tear pH/ Diurnal variations, *Arch. Ophthalmol.* 94 (1976) 821-824.

Cevc, G.; Blume, G.: Lipid vesicles penetrate into skin owing to the transdermal osmotic gradients and hydration force, *Biochim. Biophys. Acta* 1104 (1992) 226-232.

Cevc, G.; Blume, G.; Schätzlein, A.: Transfersomes-mediated transepidermal delivery improves the regiospecificity and biological activity of corticosteroids in vivo, *J. Control. Rel.* 45 (1997) 211-226.

Cevc G., Gebauer D., Stieber J., Schätzlein A., Blume G.: Ultraflexible vesicles, Transfersomes, have an extremely low pore resistance and transport therapeutic amounts of insulin across the intact mammalian skin, *Biochim. Biophys. Acta* 1368 (1998) 201-215.

Chamlin, S. L.; Frieden, I. J.; Fowler, A.; Williams, M.; Kao, J.; Sheu, M.; Elias, P. M.: Ceramide-Dominant, Barrier-repair lipids improve childhood atopic dermatitis, *Arch. Dermatol.* 137 (2001) 1110-1112.

Chapman D.: Lipid dynamics in cell membranes, In: Weissman, G; Claiborne R (eds), *Cell membranes, biochemistry, cell biology and pathology*, HP Publishing, New York (1975).

Chapman T. M; Perry, C. M.: Everolimus, *Drugs* 64 (2004) 861-872.

Cheeks L.; Kaswan R. L.; Green K.: Influence of vehicle and anterior chamber protein concentration on cyclosporine penetration through the isolated rabbit cornea, *Cur. Eye Res.* 11 (1992) 641-649.

Chetoni, P.; Panichi, L.; Burgalassi, S.; Benelli, U.; Saettone, M. F.: Pharmacokinetics and anti-inflammatory activity in rabbits of a novel indomethacin ophthalmic solution, *J. Ocul. Pharmacol. Ther.* 16 (2000) 363-372.

Chetoni, P.; Rossi, S.; Burgalassi, S.; Monti, D.; Mariotti, S.; Saettone, M. F.: Comparison of Liposome -Encapsulated Acyclovir with Acyclovir Ointment: Ocular Pharmacokinetics in Rabbits, *J. Ocul. Pharmacol. Ther.* 20 (2004) 169-177.

Chrai, S. S.; Patton, T. F.; Mehta, A.; Robinson, J. R.: Lacrimal and instilled fluid dynamics in rabbit eyes, *J. Pharm. Sci.* 62 (1973) 1112-1121.

Coderch, L.; Fonollosa, J.; De Pera, M.; Estelrich, J.; De La Maza, A.; Parra, J.L.: Influence of cholesterol on liposome bilayer fluidity by EPR Relationship with percutaneous absorption, *J. Control. Rel.* 68 (2000) 85-95.

Collins, J.F.; Augustin, A.J.: *Augenheilkunde*, Springer Verlag Berlin Heidelberg New York, 1996.

Conquet, P., Durand, G., Laillier, J., et al.: Evaluation of ocular eye irritation in the rabbit. Objective versus subjective assessment. *Tox. App. Pharmacol.* 39 (1977) 29-139.

Coupland, S.E.; Klebe, S.; Karow, A. C.; Krause, L.; Kruse, H.; Bartlett, R.R.; Hoffmann, F.: Leflunomide therapy following penetrating keratoplasty in the rat, *Graefes Arch. Clin. Exp. Ophthalm.* 232 (1994) 622-627.

D

Daniels, R.: Galenische Prinzipien moderner Hautpflegeprodukte, Skin Care Forum online 25 (2001), www.scf-online.com/german/25_d/galenik_25_d.htm.

Danielsson, I.; Stenius, P.: Anion association and micelle formation in solutions of hydrotropic and short-chain carboxylates, *J. Coll. Interf. Sci.* 37 (1971) 264-280.

Danielsson, I.; Lindman, B.: The definition of microemulsion, *Coll. Surf. B* 3 (1981) 391-392.

Danino, D. et al.: Microstructures in the aqueous solutions of a hybrid anionic fluorocarbon/ hydrocarbon surfactant, *J. Coll. Interf. Sci.* 259, 382 – 390 (2003).

Davies, J. E. D.: *Cyclodextrin Technology*, Kluwer Academic Publishers, AA Dordrecht (Niederlande), 1988.

DAZ 12/2002: *Deutsche Apothekerzeitung* 142 (2002) 1485-1486.

DAZ 7/2004: Beilage Neue Arzneimittel, Everolimus, *Deutsche Apothekerzeitung* 51 (2004) 57-63.

DAZ 36/2005: *Deutsche Apothekerzeitung* 145 (2005) 3759-3760.

Dejnika, N. S., Kuroki, A. M., Fosnot, J., Tang, W.; Tolentino, M. J.; Bennett, J.: Systemic rapamycin inhibits retinal and choroidal neovascularization in mice, *Mol. Vis.* 10 (2004) 964-972.

De Paepe, K.; Roseeuw, D.; Rogiers, V.: Repair of acetone- and sodium lauryl sulphate-damaged human skin barrier function using topically applied emulsions containing barrier lipids, *J. Eur. Acad. Dermatol. Venereol.* 16 (2002) 587 – 594.

DIN 32645: Nachweis-, Erfassungs- und Bestimmungsgrenze, Beuth Verlag, Berlin, 1994.

Dintest online: www.rzuser.uni-heidelberg.de/~df6/themenpark/kalibration.htm

Doane, M.G.; Jensen, A.D.; Dohlmann, C.H.: Penetration routes of topically applied eye medication, *Am. J. Ophthalmol.* 85 (1978) 383-386.

Dolder, R.: *Ophthalmika: Pharmakologie, Biopharmazie und Galenik der Augenarzneimittel*, Wiss. Verlagsges. Stuttgart 1990.

Döppenschmitt, S.: Entwicklung und Validierung analytischer Methoden zur Bestimmung verschiedener Glucocortikoide in Humanplasma und Humanserum, Dissertation, Humboldt-Universität zu Berlin, 1996.

van Doorne, H.: Interactions between cyclodextrins and ophthalmic drugs, *Eur. J. Pharm. Biopharm.* 39 (1993) 133-139.

Downing, D. T., Wertz, P. W., Stewart, M. E.: The role of sebum and epidermal lipids in the cosmetic properties of skin, *Int. J. Cosm. Sci.* 8 (1986) 115-123.

Downing, D. T., Stewart, M. E., Wertz, P. W., Colton, S. W., Abraham, W., Strauss, J. S.: Skin Lipids: An Update, *J. Invest. Dermatol.* 88 (1987) 2s-6s.

Dubochet, J. et al.: Cryo-electron microscopy of vitrified specimens, *Quart. Rev. Biophys.* 21, 129-228 (1988).

Durrani, A.M.; Davies, N.M.; Thomas, M.; Kellaway, I.W.: Pilocarpine bioavailability from a mucoadhesive liposomal ophthalmic drug delivery system, *Int. J. Pharm.* 88 (1992) 409 – 415.

E

Ecclestone, G.M.: Microemulsions. In: Swarbrick, J. and Boylan J.C. (eds) *Encyclopedia of pharmaceutical technology*, Vol. 9, Marcel Dekker Inc., New York, Basel und Hong Kong, 1994, 375-421.

Edsman, K.; Carlfors, J.; Petersson, R.: Rheological evaluation of Poloxamer as an in situ gel for ophthalmic use, *Eur. J. Pharm. Sci.* 6 (1998) 105-112.

Egbaria, K.; Ramachandran, C.; Kittayanond, D.; Weiner, N.: Topical delivery of liposomally encapsulated interferon evaluated by in vitro diffusion studies, *Antimicrob. Agents Chemother.* 34 (1990a) 107-110.

Egbaria, K.; Ramachandran, C.; Weiner, N.: Topical delivery of ciclosporin: evaluation of various formulations using in vitro diffusion studies in hairless mouse skin, *Skin Pharmacol.* 3 (1990b) 21-28.

Egbaria, K.; Ramachandran, C.; Weiner, N.: Topical application of liposomally entrapped cyclosporin evaluated by in vitro diffusion studies with human skin, *Skin Pharmacol.* 4 (1991a), 21-28.

Egbaria, K.; Weiner, N. Liposomes as a topical drug delivery system evaluated by in vitro diffusion studies, *Cosmet. Pharm. Appl. Polym.* [Proc. Am. Chem. Soc. Symp. Polym. Cosmet. Pharm. Appl.] (1991b) 215-24.

Ehrenstein, G. W.; Riedel, G.; Trawiel, P.: *Praxis der thermischen Analyse von Kunststoffen*, Carl Hanser Verlag, München, 2003.

Elias, P. M., Friend, D. S.: The permeability barrier, *J. Cell Biol.* 65 (1975) 180-191.

Elias, P.M., Brown, B.E.; Ziboh, V.A.: The permeability barrier in essential fatty acid deficiency: evidence for direct role of linoleic acid in barrier function, *J. Invest. Dermatol.* 74 (1980) 230-233.

Elias, P. M.: Epidermal lipids, membranes, and keratinisation, *Int. J. Dermatol.* 20 (1981a) 1-19.

Elias, P. M.: Lipids and the epidermal permeability barrier, *Arch. Dermatol. Res.* 270 (1981b) 95-117.

Elias, P.M.: Epidermal lipids, barrier function and desquamation, *J. Invest. Dermatol.* 80 (1983) 44-49.

Elias, P.M.; Williams, M.L.; Maloney, M.E.; Bonifas, J.A.; Brown, B.E.; Grayson, S.; Epstein, E.H.: Stratum corneum lipids in disorders of cornification. Steroid sulfatase and cholesterol sulfate in normal desquamation and the pathogenesis of recessive X-linked ichthyosis, *J. Clin. Invest.* 74 (1984) 1414-1421.

Erdogan, M.; Wright, J.R.; McAlister, V.C.: Liposomal tacrolimus lotion as a novel topical agent for treatment of immune-mediated skin disorders: experimental studies in a murine model, *Br. J. Dermatol.* 146 (2002) 964-967.

Evans R.G., Wain A.J., Hardacre C., Compton R.G.: An electrochemical and ESR Spectroscopic study on the molecular dynamics of TEMPO in room temperature ionic liquid solvents, *Chem. Phys. Chem.* 6 (2005) 526-533.

F

Fahr, A.; Holz, M.; Fricker, G.: Liposomal Formulation of Cyclosporin A: Influence of lipid type and dose on Pharmacokinetics, *Pharm. Res.* 12 (1995) 1189-1197.

Firman, P.; Haase, D.; Jen, J.; Kahlweit, M.; Strey, R.: On the effect of electrolytes on the mutual solubility between water and nonionic amphiphiles, *Langmuir* 1 (1985) 718-724.

Foldvari, M.; Gesztes, A.; Mezei, M.: Dermal drug delivery by liposome encapsulation: clinical and electron microscopic studies, *J. Microencapsul.* 7 (1990) 479-89.

Foldvari, M.; Moreland, A.: Clinical observations with topical liposome-encapsulated interferon alpha for the treatment of genital papilloma virus infections, *J. Lipos. Res.* 7 (1997) 115-126.

Forslind, B., Engström, S., Engblom, J., Norlén, L. A novel approach to the understanding of human skin barrier function, *J. Dermatol. Sci.* 14 (1997) 115-125.

Friberg, S. G., Kayali, I., Beckerman, W., Rhein, L. D., Simion, A.: Water permeation of reaggregated stratum corneum with model lipids, *J. Invest. Dermatol.* 94 (1990) 377-380.

Friis, G. J., Bundgaard, H.: Kinetics and degradation of cyclosporin A in acidious aqueous solution and its implication in its oral absorption, *Int. J. Pharm.* 82 (1992) 79-83.

Fröhlich, M.: Elektronenmikroskopische und physikalische Untersuchungen an Hautmodellen aus humanen Stratum corneum Lipiden und topischen phospholipidhaltigen Wirkstoff-Formulierungen, Dissertation, Albert-Ludwigs-Universität zu Freiburg i. Br., Deutschland, 2000.

Fujita, T.; Inoue, K.; Yamamoto, S.; Ikumoto, T.; Sasaki, S.; Toyama, R.: Fungal metabolites. Part 11. A potent immunosuppressive activity found in *Isaria sinclarii* metabolite, *J. Antibiot.* 47 (1994) 208-215.

Fulton, B.; Markham, A.: Mycophenolate mofetil: a review of its pharmacodynamic and pharmacokinetic properties and clinical efficacy in renal transplantation, *Drugs* 51 (1996) 278-298.

G

Gallarate, M.; Gasco, M. R.; Trotta, M.: Influence of octanoic acid on membrane permeability of timolol from solutions and from microemulsions, *Acta Pharm. Technol.* 34 (1988) 102-105.

Gallarate, M.; Gasco, M. R.; Trotta, M.; Chetoni, P.; Saettone, M. F.: Preparation and evaluation in vitro of solutions and o/w microemulsions containing levobunol as ion pair, *Int. J. Pharm.* 100 (1993) 219 – 225.

Galla H.J.: *Spektroskopische Methoden in der Biochemie*, Georg Thieme Verlag Stuttgart New York, 1988.

Garidel, P.: The thermotropic behaviour of phyto-ceramide 1 as investigated by ATR-FTIR and DSC, *Phys. Chem. Chem. Phys.* 4 (2002) 2714-2720.

Gasco, M. R.; Gallarate, M.; Trotta, M.; Bauchiero, L.; Gremmo, E.; Chiappero, O.: Microemulsions as topical delivery vehicles: ocular administration of timolol, *J. Pharm. Biomed. Anal.* 7 (1989) 433-9.

Gebler, H.: *Pharmazie für die Praxis*, Deutscher Apotheker-Verlag Stuttgart, 2000.

Geilen, C. C., Orfanos-Boeckel, H.; Offermann, G.; Orfanos, C. E.: Mycophenolatmofetil: Ein neues Immunsuppressivum in der Dermatologie und seine Einsatzmöglichkeiten, *Hautarzt* 51 (2000) 63-69.

Gesztes, A.; Mezei, M.: Topical anesthesia of the skin by liposome encapsulated tetracaine, *Anesth. Analg.* 67 (1988) 1079-1081.

Ghyczy, H.: Arzneimittel mit Phosphatidylcholin und Liposomen: Entwicklung, Bewertung, Perspektiven, in: Müller, R.H.; Hildebrand, G.E.: *Pharmazeutische Technologie: Moderne Arzneiformen*, Wiss. Verlagsges. Stuttgart 1998.

Gierer, A., Wirtz, K.: Molekulare Theorie der Mikrorreibung, *Z. Naturf.* 8A (1953) 532-538.

Gillette, T.E., Chandler, J.W. et al.: Langerhans cells on the ocular surface, *Ophthalmol.* 89 (1982) 700-711.

Glombitza, B.: Lipidsysteme als stratum corneum Modelle Charakterisierung und Eignung für Permeationsuntersuchungen, Dissertation, Technische Universität Carolo-Wilhelmina zu Braunschweig, Deutschland, 2001.

Gluck, R.; Metcalfe, I. C.: New technology platforms in the development of vaccines for the future, *Vaccine* 20 Suppl. 5 (2002) B10-B16.

Grant M.S.: Toxicology of the eye, Charles C. Thomas, Springfield, Illinois, USA, 1993.

Gregoriadis, G.; Buckland, R. A.: Enzyme-containing liposomes alleviate a model for storage disease, *Nature* 244 (1973) 170-172.

Griffiths, C.E.M; Powles, A.V.; Baker, B.S.; Fry, L. Vladimirsson, H.: Topical cyclosporin and psoriasis, *Lancet* 1 (1987) 806.

Grisanti, S.: Das Immunprivileg des Auges, *Ophthalmologie* 95 (1998) 124-135.

Grubauer, G., Feingold, K. R., Harris, R. M., Elias, P. M.: Lipid content and lipid type as determinants of the epidermal permeability barrier, *J. Lipid. Res.* 30 (1989) 89-96.

Gruber, F.: Untersuchungen zur Encapsulierung von Paclitaxel in kationischen Liposomen, Dissertation, Ludwig Maximilians Universität München, 2004.

Guo, J. X.; Ping, Q. N.; Sun, G. Q.; Jiao, C. H.: Lecithin vesicular carriers for transdermal delivery of cyclosporin A, *Int. J. Pharm.* 194 (2000) 201-207.

H

Hasse, A.; Keipert, S.: Development and characterization of microemulsions for ocular application, *Eur. J. Pharm. Biopharm.* 43 (1997) 179 – 183.

Hauptmann, S.: Einführung in die organische Chemie, Deutscher Verlag für Grundstoffindustrie, Leipzig, 1982.

Henge, U.; Ruzicka, T.; Schubert, R: Topische Calcineurin-Inhibitoren in der Dermatologie, *Pharm. Unserer Zeit* 34 (2005) 296-303.

Hoar, T.P.; Schulman, J.H.: Transparent water in oil dispersions: the oleopathic hydro-micelle, *Nature* 152 (1943) 102-103.

Hochberg, M.; Enk, C. D.: Partial protection against epidermal IL-10 transcription and Langerhans cell depletion by sunscreens after exposure of human skin to UVB, *Photochem. Photobiol.* 70 (1999) 766-772.

- Ho, P. C.; Kraus, K. A.: The effect of hydrotropic salts on phase relationships involving hydrocarbons, water, and alcohols, SPEJ (Society of Petroleum Engineers Journal) 22 (1982) 363-370.
- Holopainen, J. M., Lemmich, J.; Richter, F.; Mouritsen, O. G. ; Rapp, G.; Kinnunen, P. K. J.: Dimyristoylphosphatidylcholine/C16:0-Ceramide binary Liposomes studied by Differential Scanning Calorimetry and Wide- and Small-Angle X-Ray Scattering, Biophys. J. 78 (2000) 2459-2469.
- Hooijmaaijer, E.; Brandl, M. ; Nelson, J.; Lustig, D.: Degradation products of mycophenolate mofetil in aqueous solution, Drug Dev. Ind. Pharm. 25 (1999) 361-365.
- Honeywell-Nguyen, P.L.; Wouter Groenink, H.W.; de Graeff, A.M.; Bouwstra, J.A.: The in vivo transport of elastic vesicles into human skin: effects of occlusion, volume and duration of application, J. Control. Rel. 90 (2003) 243-255.
- Hofner, A. F.: Sirolimus, ein neues Immunsuppressivum, Pharmazeutische Zeitung 148 (2003) 28-34.
- Hoogstrate, A.J.; Verhoef, J.; Brussee, J.; Ijzerman, A.P.; Spies, F.; Bodde, H.E.: Kinetics, ultrastructural aspects and molecular modelling of transepidermal peptide flux enhancement by N-allylazacycloheptanones, Int. J. Pharm. 76 (1991) 37-47.
- Hou, S.Y.; Mitra, A.K.; White, S.H.; Menon, G.K.; Ghadially, R. Elias, P.: Membrane structures in normal and essential fatty acid-deficient stratum corneum: characterization by ruthenium tetroxide staining and X-ray diffraction, J. Invest. Dermatol. 6 (1991) 215-223.
- Huang, C.: Studies on phosphatidylcholine vesicles. Formation and physical characterization, Biochemistry 8 (1969) 344-351.
- Huang, C.: Highly efficient target binding of immunoliposomes, Liposomes in Drug Delivery, 21 years on, London, 12. – 15. 12. 1990.
- Huang, H.-S., Schoenwald, R.D., Lach, J.L.: Corneal penetration behavior of betablocking agents II: assessment of barrier contributions, J. Pharm. Sci. 72 (1983) 1272-1286.
- Hubbell W. L.; McConnell H. M.: Molecular motion in spin-labeled phospholipids and membranes, J. Amer. Chem. Soc. 93 (1971) 314-26.
- Hueber, F.; Schaefer, H.; Wepierre, J.: Role of transepidermal and transfollicular routes in percutaneous absorption of steroids: In vitro studies on human skin, Skin Pharmacol. 7 (1994) 237-244.
- Hultsch, T.; Kapp, A.; Spengel, J.: Immunomodulation and safety of topical calcineurin inhibitors for the treatment of atopic dermatitis, Dermatol. 211 (2005) 174-187.

I

Imokawa, G., Akasaki, S., Hattori, M., Yoshizuka, M.: Selective recovery of deranged water-holding properties by stratum corneum lipids, *J. Invest. Dermatol.* 87 (1986) 758-761.

Imokawa, G., Akasaki, S., Minematsu, Y., Kawai, M.: Importance of intercellular lipids in water-retention properties of the stratum corneum: induction and recovery study of surfactant dry skin, *Arch. Dermatol. Res.* 281 (1989) 45-51.

Imokawa, G., Abe, A., Jin, K., Higaki, Y., Kawashima, M., Hidano, A.: Decreased level of ceramides in human stratum corneum in atopic dermatitis: an etiologic factor in atopic dry skin? *J. Invest. Dermatol.* 96 (1991) 523-26.

Imura, T.; Sakai, H.; Yamauchi, H.; Kaise, C.; Kozowa, K.; Yokoyama, S.; Abe, M.: Preparation of liposomes containing Ceramide 3 and their membrane characteristics, *Coll. Surf. B: Biointerfaces* 20 (2001) 1-8.

Irie, T.; Uekama, K.: Pharmaceutical applications of cyclodextrins. III. Toxicological Issues and safety evaluation, *J. Pharm. Sci.* 86 (1997) 147-162.

Ishida-Yamamoto, A.; Iizuka, H.: Structural organization of cornified cell envelopes and alterations in inherited skin disorders, *Exp. Dermatol.* 7 (1998) 1–10.

Ismailos, G.; Reppas, C.; Dressman, J.B.; Macheras, P.: Unusual solubility behaviour of cyclosporin A in aqueous media, *J. Pharm. Pharmacol.* 43 (1991) 287-289.

J

Jacobs, M.; Martin, G. P.; Mariott, C.: Effects of phosphatidylcholine on the topical bioavailability of corticosteroids assessed by the human skin blanching assay, *J. Pharm. Pharmacol.* 40 (1988) 829-833.

Jahn, K.; Fischer, A.; Neubert, R. H. H.; Wohlrab, J.: Investigation of the penetration behaviour of mycophenolate mofetil from a semisolid formulation into human skin ex vivo, *J. Pharm. Pharmacol.* 53 (2001) 1581-1587.

Jahn, K.: Moderne galenische Zubereitungen zur dermalen Anwendung von Ciclosporin A und Mycophenolatmofetil. Dissertation, Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Deutschland, 2002.

Jansen, T.; Xhonneux, B.; Messens, J.; Borgers, M.: Beta-cyclodextrines as vehicles in eye drop formulations: an evaluation of their effects on rabbit corneal epithelium, *Lens Eye Tox. Res.* 7 (1990) 459-468.

Javier, A. F.; Csorgo, Z. B.; Ellis, C. N.; Kang, S.; Voorhees, J. J.; Cooper, K. D.: Rapamycin (Sirolimus) inhibits proliferating cell nuclear antigen expression and blocks cell cycle in the G1 phase in human keratinocyte stem cells, *J. Clin. Invest.* 99 (1997) 2094-2099.

Jores, K.: Lipid nanodispersions as drug carrier systems - a physicochemical characterization, Dissertation, Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Deutschland, 2004.

Jenning, V.; Mäder, K.; Gohla, S. H.: Solid lipid nanoparticles (SLNTM) based on binary mixtures of liquid and solid lipids: a ¹H-NMR study, *Int. J. Pharm.* 205 (2000) 15-21.

Jeschke, G.: Einführung in die ESR-Spektroskopie, Vorlesungsskript, Johannes-Gutenberg-Universität Mainz, 1998.

Jones, J.; Saadat-Lajevardi, S.; Lee, T.; Horwatt, R.; Hicks, D.; Johnston, A.; Holt, D. W.: An Immunoassay for the measurement of sirolimus, *Clin. Therapeut.* 22 Suppl. B (2000) B49-B61.

K

Kaever, V.; Raesch, K.: Prinzipien der Immunsuppression, *Pharm. Unserer Zeit* 34 (2005) 268-274.

Kahan, B.D.; Stepkowski, S.M.; Napoli, K. L.; Katz S. M.; Knight, R. J.; Van Buren, C.: The development of sirolimus: The university of Texas-Houston experience, *Clin. Transpl.* (2000) 145 -158.

Kahan, B.D.; Karlix, J. L.; Ferguson, R. M.; Leichtman, A. B.; Mulgaonkar, S.; Gonwa, T. A.; Skerjanec, A.; Schmouder, R. L.; Chodoff, L.: Pharmacodynamics, pharmacokinetics, and safety of multiple doses of FTY 720 in stable renal transplant patients: a multicenter, randomized, placebo-controlled, Phase I study, *Transplantation* 76 (2003) 1079 – 1084.

Kaltwasser, J.P.; Nash, P.; Gladman, D.; Rosen, C.F.; Behrens, F.; Jones, P.; Wollenhaupt, J.; Falk, F. G.; Mease, P.: Efficacy and safety of leflunomide in the treatment of psoriatic arthritis and Psoriasis, *Arthritis Rheumatism* 50 (2004) 1939-1950.

Kanai, A.; Alba, R. M.; Takano, T.; Kobayashi, C.; Nakajima, A.; Kurihara, K.; Yokoyama, T.; Fukami, M.: The effect on the cornea of alpha cyclodextrin vehicle for cyclosporin eye drops, *Transpl. Proc.* 21 (1989) 3150-3152.

Kauffman, H. M.; Cherikh, W. S.; Cheng, Y; Hanto, D. W.; Kahan, B. D.: Maintenance immunosuppression with target-of-rapamycin inhibitors is associated with a reduced incidence of de novo malignancies, *Transplant.* 80 (2005) 883-889.

Katami, M.: The mechanisms of corneal allograft failure in the rat, *Eye* 9 (1995) 197-207.

Katre, N. V.; Asherman, J.; Schaefer, H.; Hora, M.: Multivesicular Liposome (DepoFoam) Technology for the Sustained Delivery of Insulin-like Growth Factor-I (IGF-I), *J. Pharm. Sci.* 87 (1998) 1341-1346.

Kawai, M.; Gunawanadrana, I. W. K.; Mollison, K. W.; Hsieh, G.C.; Lane, B. C.; Luly, J. R.: Studies on an immunosuppressive Macrolactam, Ascomycin: Synthesis of a C-33 Hydroxyl Derivative, *Bioorg. Med. Chem. Lett.* 8 (1998) 935-938.

Keipert, S., Siebenbrodt, I.; Lüders, F.; Bornschein, M.: Mikroemulsionen und ihre potenzielle pharmazeutische Nutzung, *Pharmazie* 44 (1989) 433-444.

Keipert, S.; Schulz, G.: Mikroemulsionen auf Sacharoseesterbasis, *Pharmazie* 49 (1994) 195 – 197.

Keipert, S.: Ophthalmika: etablierte Arzneiformen und neue Konzepte, in: Müller, R. H., Hildebrandt, G. E. (Hg.): *Pharmazeutische Technologie: Moderne Arzneiformen*, Wiss. Verlagsges. Stuttgart 1997.

Kim, E.; Gao, Z. G.; Park, J.S.; Li, H.; Han, K.: rhEGF/HP- β -CD complex in poloxamer gel for ophthalmic delivery, *Int. J. Pharm.* 233 (2002) 159-167.

Kirjavainen, M.; Urtti, A.; Valjakka-Koskela, R.; Kiesvaara, J.; Mönkkönen, J.: Liposome-skin interactions and their effects on the skin permeation of drugs, *Eur. J. Pharm. Sci.* 7 (1999) 279-286.

Kleuser, B.: Wirkungen und Signalwege von $1\alpha,25$ -Dihydroxyvitamin D₃ und Spingosin-1-phosphat, Habilitation, Freie Universität Berlin, 2001.

Knapp, S.; Bertelmann, E.; Hartmann, C.; Keipert, S.; Pleyer, U.: Intraocular availability of topically applied mycophenolate mofetil in rabbits, *J. Ocul. Pharmacol. Ther.* 19 (2003) 181-192.

Knapp-Ulrich, S. U.: Einfluss von Cyclodextrinen auf die okuläre Verfügbarkeit von Antiglaukomatosa und Immunsuppressiva in vitro/ in vivo, Thesis, Humboldt Universität, Berlin, Deutschland, (2004).

König, K.; Zijilstra, J.: In-vitro-Untersuchung: Liposomen als Arzneistoffträger in der Lokaltherapie, *Deutsche Apothekerzeitung* 145 (2005) 993-994.

Kohno, T.; Tsui, T.; Hirayama, K.; Watabe, K.; Matsumoto, A.; Kohno, T.; Fujita, T.: A novel immunomodulator, FTY 720, prevents spontaneous dermatitis in NC/Nga mice, *Biol. Pharm. Bull.* 27 (2004) 1392-1396.

Kovarik, J. M.; Schmouder, R.L.; Serra, D.; Wang, Y.; Wiegand, H.; Dilzer, S.c.; Lasseter, K. C.: FTY 720 pharmakokinetics in mild to moderate hepatic impairment, *J. Clin. Pharmacol.* 45 (2005) 446-452.

Kreuter, A.; Somemr, A.; Hyun, J., et al.: 1 % Pimecrolimus, 0,005 % calcipotriol, and 0,1 % betamethasone in the treatment of intertriginous psoriasis: a double blind, randomized controlled study, *Arch. Dermatol.* 142 (2006) 1138-1143.

Kromidas, S.: Validierung in der Analytik, Wiley VCH Weinheim, 1999, S. 179.

Krutmann, J.: New developments in photoprotection of human skin, *Skin Pharmacol. Appl. Skin Physiol.* 14 (2001) 401–407.

Kuhn, B.; Jacobsen, W.; Christians, U.; Benet, L. Z.; Kollman, P. A.: Metabolism of sirolimus and its derivative everolimus by cytochrome P450 3A4: Insights from Docking, molecular dynamics, and quantum chemical calculations, *J. Med. Chem.* 44 (2001) 2027-2034.

Kurose, S.; Ikeda, E.; Tokiwa, M.; Hikita, N.; Mochizuki, M.: Effects of FTY 720, a novel immunosuppressant, on experimental autoimmune uveoretinitis in rats, *Exp. Eye Res.* 70 (2000) 7-10.

Kwon, Y. S.; Kim, J. C.; Inhibition of corneal neovascularization by rapamycin, *Experim. Mol. Med.* 38 (2006) 173-179.

L

Ladbrooke, B. D.; Williams, R. M.; Chapman, D.: Studies on lecithin-cholesterol-water interactions by differential scanning calorimetry and x-ray diffraction, *Biochim. Biophys. Acta, Biomembranes* 150 (1968) 333-40.

Laillier, J., Plazonnet, B., Le Douarec, J.C., et al.: Evaluation of an objective method of studying eye irritation, *Proc. Eur. Soc. Toxicol.* 17 (1976) 336-350 (Proc. Meet. 1975).

Landmann, L.: Epidermal permeability barrier: transformation of lamellar granule-disks into intercellular sheets by a freeze fracture study, *J. Invest. Dermatol.* 87 (1986) 202-209.

Landmann, L.: The epidermal permeability barrier, *Anatomy and Embryology*, 178 (1988) 1-13.

Lasic, D. D.: *Liposomes from physics to applications*, Elsevier Science B.V., Amsterdam, 1993.

Lasic, D. D.: Novel applications of liposomes, *Trends Biotechnol.* 16 (1998) 307-321.

Lasch, J.; Wohlrab, W.: Liposome bound cortisol: a new approach to cutaneous therapy, *Biomed. Biochim. Acta* 45 (1986) 1295-1299.

Lasch, J.; Laub, R.; Wohlrab, W.: How deep do intact liposomes penetrate into human skin?, *J. Control. Rel.* 18 (1991) 55-58.

Lauerma, A.I.; Surber, C.; Maibach, H.I.: Absorption of topical Tacrolimus (FK 506) in vitro through human skin: comparison with cyclosporin A; *Skin Pharmacol.* 10 (1997) 230-234.

Lavker, R.M.; Kaidbey, K.; Leyden, J.J.: Effects of topical ammonium lactate on cutaneous atrophy from a potent topical corticosteroid, *J. Am. Acad. Dermatol.* 26 (1992) 535–544.

Law, S. L.; Huang, K.J.; Chiang, C.H.: Acyclovir-containing liposomes for potential ocular delivery Corneal penetration and absorption, *J. Control. Rel.* 63 (2000) 135 – 140.

Lawrence, M.J.: Microemulsions as drug delivery vehicles, *Curr. Opin. Coll. Interface Sci.* 1 (1996) 826-832.

Leaf, C.W. Toxicology of some nonionic surfactants. In: *Soap Chem. Spec.* 43 S. 48 – 51, 106, 108-110, New York, 1967.

Lebwohl, M.; Freeman, A.K.; Chapman, M.; et al.: Tacrolimus ointment is effective for facial and intertriginous psoriasis, *J. Am. Acad. Dermatol.* 5 (2004) 723-730.

Lee, S.; Dausch, S.; Maierhofer, G.; Dausch, D.: A new therapy concept for the treatment of dry eye - the usefulness of phospholipid liposomes, *Klinische Monatsblätter für Augenheilkunde* 221 (2004) 825-836.

Lehmann, O.: Neue Untersuchungen über flüssige Kristalle. II. Teil, Carl Winter's Universitätsbuchhandlung, Heidelberg 1912.

Lehninger, A. L.; Nelson, D. L.; Cox, M. M.: *Lehninger Biochemie*, Springer-Verlag Berlin Heidelberg New York, 2001.

Levine, A.M.; Tulpule, A.; Espina, B; Sherrod, A; Boswell, W.D.; Lieberman, R.D.; Nathwani, B.N.; Welles, L.: Liposome-encapsulated doxorubicin in combination with standard agents (cyclophosphamide, vincristine, prednisone) in patients with newly diagnosed AIDS-related non-Hodgkin's lymphoma: Results of therapy and correlates of response, *J. Clin. Oncol.* 22 (2004) 2662–2670.

Leydecker, W; Grehn, F.: *Augenheilkunde*, Springer-Verlag Berlin Heidelberg New York, 1993.

Liaw, J.; Robinson, J.R.: The effect of polyethylene glycol molecular weight on corneal transport and the related influence of penetration enhancers, *Int. J. Pharm.* 88 (1992) 125-140.

Liaw, J.; Rojanasakul, Y.; Robinson, J.R.: The effect of drug charge density on corneal transport, *Int. J. Pharm.* 88 (1992) 111-124.

Li, X.; Hirsh, Donald, J.; Cabral-Lilly, D.; Zirkel, A.; Gruner, Sol M.; Janoff, A. S.; Perkins, W. R.: Doxorubicin physical state in solution and inside liposomes loaded via a pH gradient, *Biochim. Biophys. Acta, Biomembranes* 1415, (1998) 23-40.

Lidgate, D.; Brandl, M.; Holper, M.; Abubakari, Xiaoyang, W.: Influence of ferrous sulfate on the solubility, partition coefficient, and stability of mycophenolic acid and the ester mycophenolate mofetil, *Drug Develop. Ind. Pharm.* 28 (2002) 1275-1283.

- Lieb, L. M.; Ramachandran, C.; Egbaria, K.; Weiner, N.: Topical delivery enhancement with multilamellar liposomes into pilosebaceous units: I. In vitro evaluation using fluorescent techniques with the hamster ear model, *J. Inv. Dermatol.* 99 (1992), 108-113.
- Lin, Z. et al. : Comparison of drag reduction, rheology, micelle structure and stress-induced precipitation of dilute cationic surfactant solutions with odd and even alkyl chains, *Rheo. Acta* 41, 483 – 492 (2002).
- Lippert, J.L., Peticolas, W.L.: Laser Raman investigation of the effect of cholesterol on conformational changes in dipalmitoyl lecithin multilayers, *Proc. Natl. Acad. Sci. USA* 68 (1971) 1572-1576.
- Liu, H.; Li, S.; Wang, Y.; Yao, H.; Zhang, Y.: Effect of vehicles and enhancers on the topical delivery of cyclosporin A, *Int. J. Pharm.* 311 (2006) 182-186.
- Löffler, G.; Petrides, P. E.: *Biochemie und Pathobiochemie*, Springer Verlag Berlin Heidelberg New York, 2003.
- Loftsson, T.; Brewster, M. E.: *Pharmaceutical Applications of Cyclodextrins. 1. Drug Solubilization and Stabilization*, *J. Pharm. Sci.* 85 (1996) 1017-1025.
- Loftsson, T.; Stefansson, E.; Kristinsson, J. K.; Fridriksdottir, H.; Sverrisson, T.; Gudmundsdottir, G.; Thorisdottir, S.: Topically effective acetazolamide eye-drop solution in man, *Pharm. Sci.* 2 (1996) 277-279.
- Loftsson, T.: Increasing the cyclodextrin complexation of drugs and drug bioavailability through addition of water-soluble polymers, *Pharmazie* 53 (1998) 733-740.
- Loftsson, T.; Matthiasson, K.; Masson, M.: The effects of organic salts on the cyclodextrin solubilization of drugs, *Int. J. Pharm.* 262 (2003) 101-107.
- Loftsson, T.; Masson, M.: The effects of water-soluble polymers on cyclodextrins and cyclodextrin solubilization of drugs, *J. Drug Deliv. Sci. Technol.* 14 (2004) 35-43.
- Lopes, L.B.; Collet, J. H.; Vitoria, M.; Bentley, L. B.: Topical delivery of cyclosporin A: an in vitro study using monoolein as a penetration enhancer, *Eur. J. Pharm. Biopharm.* 60 (2005) 25-30.
- Lowe, N.J., Stoughton, R. B.: Essential fatty acid deficient hairless mouse: a model of chronic epidermal hyperproliferation, *Br. J. Dermatol.* (1977) 155-162.
- Luepke, N.P.: Hen's egg chorio allantoic membrane test for irritation potential, *Fd. Chem. Toxic.* 23 (1985) 287-291.
- Luepke, N. P.; Kemper, F. H. The HET-CAM test: an alternative to the Draize eye test. *Fd. Chem. Toxic.* 24 (1986) 495-496.
- Luger, T.: *Organ der Gesellschaft für Dermopharmazie*, *DermoTopics* 5 (2005), 23-24.

Lv, F.F.; Li, N.; Zheng, L.Q.; Tung, C. H.: Studies of the chloramphenicol in the microemulsion free of alcohols, *Eur. J. Pharm. Biopharm.* 62 (2006) 288-294.

M

Mäder, K.: Pharmazeutische Anwendungen der Elektronenspinresonanzspektroskopie (ESR), *PZ Prisma* 5, 202 – 212 (1998).

Mäder, K.; Swartz, H.M.; Stößer, R.; Borchert, H.-H.: The application of EPR spectroscopy in the field of pharmacy, *Pharmazie* 49 (1994) 97 – 101.

Madison, K.C.; Swarzendruber, D.C.; Wertz, P.C.; Downing, D.T.: Presence of intact intercellular lipid lamellae in the upper layers of the stratum corneum, *J. Invest. Dermatol.* 88 (1987) 714–718.

Maleki, S.; Graves, S.; Becker, S.; Horwatt, R.; Hicks, D.; Stroshane, R. M.; Kincaid, H.: Therapeutic monitoring of sirolimus in human whole-blood samples by high-performance liquid chromatography, *Clin. Therapeut.* 22 Suppl. B (2000) B25-B37.

Mao-Qiang, M.; Feingold, K. R.; Wang, F.; Thornfeldt, C. R.; Elias, P. M.: Exogenous lipids influence permeability barrier recovery in acetone-treated murine skin, *Archiv. Dermatol.* 129 (1993) 129 728-38.

Mao-Qiang, M.; Feingold, K. R.; Wang, F.; Thornfeldt, C. R.; Elias, P. M.: A natural lipid mixture improves barrier function and hydration in human and murine skin, *J. Soc. Cos. Chem.* 47 (1996) 157-166.

Mastman, G.L.; Blades, E.J.; Henderson, J.W.: The total osmotic pressure of tears in normal and various pathological conditions, *Arch. Ophthalmol.* 65 (1961) 509.

Matero, Anna: Hydrotropes, In: Krister, Holmberg (Hg.): *Handbook of applied surfaces and colloid chemistry*, Part 2, Jon Wiley & Sons, New York, 2001.

Mayer, K.; Birnbaum, F.; Reinhard, T.; Reis, A.; Braunstein, S.; Claas, F.; Sundmacher, R.: FTY 720 prolongs clear allograft survival with a different effect on different lymphocyte populations, *Br. J. Ophthalmol.* 88 (2005) 915-919.

Medawar, P.B.: Immunity to homologous grafted skin III. The fate of skin homografts transplanted to the brain, to subcutaneous tissue, and to the anterior chamber of the eye, *Br. J. Exp. Pathol.* 29 (1948) 58-67.

Meingassner, J. G., Aschauer, H.; Stuetz, A.; Billich, A.: Pimecrolimus permeates less through normal, inflamed, or corticosteroid-pretreated skin, *Experiment. Dermatol.* 14 (2005) 752-757.

Melton, J.L.; Wertz, P.W.; Swarzendruber, D.C.; Downing, D.T.: Effects of essential fatty acid deficiency epidermal O-acylsphingolipids and transepidermal water loss in young pigs, *Biochim. Biophys. Acta* 921 (1987) 191–197.

- Merck Index – an encyclopaedia of chemicals, drugs and biologicals, Merck & Co. Inc., Whitehouse Station, NJ, USA, 12. Aufl. (1996)
- Meisner, D.; Pringle, J.; Mezei, M.: Liposomal ophthalmic drug delivery. III Pharmacodynamic and biodisposition studies of atropine, *Int. J. Pharm.* 55 (1989) 105 – 113.
- Metz, G.: MMF, ein neuer Wirkstoff zur Immunsuppression, *Pharmazeutische Zeitung* 33 (1996) 36 – 42.
- Mezei, M.; Gulasekharam, Y.: Liposomes – a selective drug delivery system for the topical route of administration: gel dosage form, *J. Pharm. Pharmacol.* 34 (1982) 473-474.
- Mezei, M.: Liposomes as a skin drug delivery system, In: *Topics in Pharmaceutical Sciences*, Breimer D. D.; Speiser P. (Hg.), Elsevier Sci. Pub, Amsterdam, 345-358, 1985.
- Mezei, M.: Liposomes and the skin, In: *Gregoriadis, G.; Florence, A. T.; Patel, H. M.: Liposomes in drug delivery*, Harwood academic publishers, Chur, Schweiz, 1993.
- Miller, C. S.; Donovan, M. D.: Effect of poloxamer 407 gel on the miotic activity of pilocarpine nitrate in rabbits, *Int. J. Pharm.* 12 (1982) 147 – 152.
- Milani, J. M.; Pleyer, U.; Dukes, A.; Chou, J.; Lutz, S.; Rueckert, D.; Schmidt, K.-H.; Mondino, B. J.: Prolongation of corneal allograft survival with liposome-encapsulated Cyclosporine in the rat eye, *Ophthalmol.* 100 (1993) 890 – 896.
- Mischler, R.; Metcalfe, I. C.: Inflexal V a trivalent virosome subunit influenza vaccine production, *Vaccine* 20 Suppl. 5 (2002) B17-B23.
- Mistry-Burchardi, N.: Neue Therapieansätze – FTY 720, in: *Rupprecht, H.; Burchardi, Ch.; Mistry-Burchardi, N.; Fischereder, M.; Weidner, S.: Immunsuppression*, Elsevier Urban und Fischer Verlag München Jena, 2005.
- Moll, K. P.: Mobilität von ESR-Spinsonden nach epikutaner Applikation unter besonderer Berücksichtigung der Wechselwirkungen zwischen Liposomen und Haut, *Dissertation, Humboldt-Universität Berlin*, 2004.
- Montalbano, M.; Neff, G.W.; Yamashiki, N.: A retrospective review of liver transplant patients treated with sirolimus from a single center: an analysis of sirolimus related complications, *Transplantation* 78 (2004) 264-268.
- Motta, S., Monti, M., Sesana, S., Caputo, R., Carelli, S., Ghidoni, R.: Ceramide composition of the psoriatic scale, *Biochim. Biophys. Acta.* 1182 (1993) 147-151.
- Mrowietz, U.: *Immunsuppressiva und Immunmodulatoren in der Dermatologie*. Uni-Med Verlag AG, Bremen, Deutschland, 2001.

Müller, B.W.: Topische Mikroemulsionen als neue Wirkstoffträgersysteme, In: Müller, R. H., Hildebrandt, G. E. (Hg.): Pharmazeutische Technologie: Moderne Arzneiformen, Wiss. Verlagsges. Stuttgart 1997.

Müller, H.; Hofer, S.; Kaneider, N.; Neuwirt, H.; Mosheimer, B.; Mayer, G., Konwalinka, G.; Heufler, C.; Tiefenthaler, M.: The immunomodulator FTY 720 interferes with effector functions of human monocyte-derived dendritic cells, Eur. J. Immunol. 35 (2005) 535-545.

Müller, R. H.: Polydispersität und elektroforetische Beweglichkeit hochdispenser Systeme, Dissertation, Christian-Albrechts-Universität zu Kiel, 1983.

Müller, R. H. ; Heinemann, S.: Fat emulsions for parenteral nutrition.I: Evaluation of microscopic and laser light scattering methods for the determination of the physical stability, Clin. Nutr. 11, 223-236 (1992).

Müller, R. H.; Schuhmann, R.: Teilchengrößenmessung in der Laborpraxis, Wiss. Verlagsges. Stuttgart, Deutschland, 1996.

Müller, R. H., Zetapotenziale und Partikelladung in der Laborpraxis, Wiss. Verlagsges. Stuttgart, Deutschland, 1996.

Mutschler, E.: Arzneimittelwirkungen, Wiss. Verlagsges. Stuttgart 2001.

N

Narishetty, S. T. K.; Panchagnula, R.: Effect of L-menthol and 1,8-cineole on phase behavior and molecular organization of SC lipids and skin permeation of zidovudine, J. Control. Rel. 102 (2005) 59-70.

Napoli, K.: A practical guide to the analysis of sirolimus using high performance liquid chromatography with ultraviolet detection, Clin. Therapeut. 22 Suppl. B (2000) B14-B24.

Nash, P.; Thaci, D.; Behrens, F.; Falk, F.; Kaltwasser, P.: Leflunomide improves psoriasis in patients with psoriatic arthritis: an in-depth analysis of data from the TOPAS study, Dermatol. 212 (2006) 238-249.

Navarro, F.; Petit, S.; Stone, G.: Stabilization of polyene macrolides with antioxidants. PCT Int. Appl. WO 2000033878 (2000).

Naveh, N., Muchtar, S., and Benita, S.: Pilocarpine incorporated into a submicron emulsion vehicle causes an unexpectedly prolonged ocular hypotensive effect in rabbits, J. Ocul. Pharmacol. 10 (1994) 509 – 520.

Nell, B.; Walde, I.; Billich, A.; Vit, P.; Meingassner, J. G.: The effect of topical pimecrolimus on keratoconjunctivitis sicca and chronic superficial keratitis in dogs: result from an exploratory study, Vet. Ophthalmol. 8 (2005) 39-46.

Neuberg, C. A.: Hydrotropic phenomena, *Biochemische Zeitschrift*, 76 (1916) 107-176.

Neubert, R.; Wohlrab, W.; Bendas, C.: Modelling of drug penetration into human skin using a multilayer membrane system, *Skin Pharmacol.* 8 (1995) 119 – 129.

Neue, U. D.; Phoebe, C. H.; Tran, K.; Cheng, Y.-F.; Lu, Z.: Dependence of Reversed Phase retention of ionizable analytes on pH, concentration of organic solvents and silanol activity, *J. Chromatogr. A* 925 (2001) 49-67.

New, R. R. C. (Ed.): *Liposomes, a practical approach*, IRC Press, Oxford, 1990.

New York Academy of Science: *The use of liposomes in biology and medicine, Conference proceedings* (1977).

Niemiec, S. M.; Ramachandran, C.; Weiner, N.; Influence of nonionic liposomal composition on topical delivery of peptide drugs into pilosebaceous units: An in vivo study using the hamster ear model, *Pharm. Res.* 12 (1995) 1184-1188.

Nishihata, T.; Kotera, K.; Nakano, Y.; Yamazaki, M.: Rat percutaneous transport of diclofenac and influence of hydrogenated soya phospholipids, *Chem. Pharm. Bull.* 35 (1987) 3807-12.

Nishihata, T.; Kamada, A.; Sakai, K.; Takahashi, K.; Matsumoto, K.; Shinozaki, K.; Tabata, Y.; Keigami, M.; Miyagi, T.; Tatsumi, N.: Percutaneous absorption of diclofenac in rats and humans: aqueous gel formulation, *Int. J. Pharm.* 46 (1988) 1-7.

Nghiem, P.; Pearson, G.; Langley, R. G.: Tacrolimus and pimecrolimus: From clever prokaryotes to inhibiting calcineurin and treating atopic dermatitis, *J. Am. Acad. Dermatol.* 46 (2002) 228-41.

Norn, M.S.: Postmortal staining of the external eye, *Acta Ophthalmol.* 57 (1979) 296-304.

O

Odland, G. F.: A submicroscopic granular component in human epidermis, *J. Invest. Dermatol.* 34 (1960) 11-15.

Ogiso, T.; Hirota, T.; Iwaki, M.; Hino, T.; Tanino, T.: Effect of temperature on percutaneous absorption of terodiline, and relationship between penetration and fluidity of stratum corneum lipids, *Int. J. Pharm.* 176 (1998).

Ohia, E.O., Mancino, M., and Kulkarni, P.S.: Effects of steroids and immunosuppressive drugs on endotoxin-uveitis in rabbits, *J. Ocular. Pharmacol.* 8 (1992) 295-307.

Oldfield, E.; Marsden, J.; Chapman, D.: Proton NMR relaxation study of mobility in lipid water systems, *Chem. Phys. Lipids.* 7 (1971) 1-8.

Oldfield, E.; Chapman, D.: Dynamics of lipids in membranes. Heterogeneity and the role of cholesterol, *FEBS Letters* 23 (1972) 285-97.

Ormerod, A. D.; Shah, S. A. A.; Copeland, P.; Omar, G.; Winfield, A.: Treatment of psoriasis with topical sirolimus: preclinical development and a randomized, double-blind trial, *Br. J. Dermatol.* 152 (2005) 758 – 764.

Olsen, T.W., Benegas, N.M., Joplin, A.C., Evangelista, T. ; Mindrup, E. A.; Holland, E. J.: Rapamycin inhibits corneal allograft rejection and neovascularization, *Arch. Ophthalmol.* 112 (1994) 1471-1475.

Ouyang, C.; Choice, E.; Holland, J.; Meloche, M.; Madden, T. D.: Liposomal Cyclosporine. Characterization of drug incorporation and interbilayer exchange, *Transplantation* 60 (1995) 999-1006.

P

Paller, A. S.; Lebwohl, M.; Fleischer, A. B.; Antaya, R.; Langley, R. G.; Kirsner, R. S.; Blum, R. R.; Rico, M. J.; Jaracz, E.; Croweg, A.; Linowski, G. J.: Tacrolimus ointment is more effective than pimecrolimus cream with a similar safety profile in the treatment of atopic dermatitis: Results from 3 randomized, comparative studies, *J. Amer. Acad. Dermatol.* 52 (2005) 810-822.

Papahadjopoulos D; Miller, N.: Phospholipid model membranes. I. Structural characteristics of hydrated liquid crystals. *Biochim. Biophys. acta* 135, 624-638 (1967).

Papahadjopoulos, D.; Allen, T. M.; Gabizon, A.; Mayhew, E.; Matthay, K.; Huang, S. K.; Lee, K. D.; Woodle, M. C.; Lasic, D. D.: Sterically stabilized liposomes: improvements in pharmacokinetics and antitumor therapeutic efficacy. *Proceedings of the National Academy of Sciences of the United States of America* 88 11460-11464 (1991).

Park, J.; Ha, H.; Ahn, H. J.; Kang, S.-W.; Kim, Y. S.; Seo, J. Y.; Kim, M. S.: Sirolimus inhibits platelet-derived growth factor - induced collagen synthesis in rat vascular smooth muscle cells, *Transplant. Proceed.* 37 (2005), 3459-3462.

Patel H. M.: Liposomes as a controlled release system, *Biochem. Soc. Trans.* 12 (1985) 513-516.

Petrak, K.; Goddard P., Transport of macromolecules across the capillary walls, *Adv. Drug Del. Rev.* 3 (1989) 191-214.

Pfüller, U.: Mizellen – Vesikel – Mikroemulsionen, Tensidassoziate und ihre Anwendung in Analytik und Biochemie, Verlag Volk und Gesundheit, Berlin, 1986.

PharmaForum in Onkologie 27/5/2004: Resistenz umgehen, Prognose verbessern – Myocet[®] beim AIDS-assoziierten Non-Hodgkin-Lymphom.

Ph. Eur. 2005, 5. Ausgabe, 2. Nachtrag: Monographie Mycophenolat Mofetil.

Plätzer, M.: Einsatz moderner Analysemethoden in der Biopharmazie- Kapillarelektrophoretische Untersuchungen und Kopplung zwischen Flüssigchromatographie und Massenspektrometrie, Dissertation, Martin-Luther-Universität Halle-Wittenberg, 2002.

Pilgram, G.S.K.; Vissers, D.C.J.; van der Meulen, H.; Pavel, S.; Lavrijsen, S.P.M.; Bouwstra, J.A.; Koerten, H.K.: Aberrant lipid organization in stratum corneum of patients with atopic dermatitis and lamellar ichthyosis, *J. Invest. Dermatol.* 117 (2001) 710–717.

du Plessis, J.; Egbaria, K.; Weiner, N.; Influence of formulation factors on the deposition of liposomal components into the different strata of the skin, *J. Soc. Cosmet. Chem.* 43 (1992a) 93-100.

du Plessis, J.; Egbaria, K.; Ramachandran, C.; Weiner, N.: Topical delivery of liposomally encapsulated gamma-interferon, *Antiviral Res.* 18 (1992b) 259-65.

Pleyer, U.; Lutz, S.; Jusko, W. J.; Nguyen, K.; Rueckert, D; Mondino, B. J.; Lee, V. H. L. Ocular absorption of topically applied FK506 from liposomal and oil formulations in the rabbit eye, *Inv. Ophthalmol. Vis. Sci.* 34 (1993) 2737-2742.

Poste, G.; Papahadjopoulos, D.: Drug-containing lipid vesicles render drug-resistant cell sensitive to actinomycin D, *Nature* 261 (1976) 699-701.

Price, C.I.; Horton, J.W.; Baxter, C.R.: Topical liposomal delivery of antibiotics in soft tissue infection, *J. Surg. Res.* 49 (1990) 174.

Produktinformation ICI, Wilton, Middlesborough, Cleveland, USA.

Produktinformation Lipoid GmbH, Ludwigshafen, Deutschland, 2002.

Produktinformation Natipide II, Nattermann Phospholipid GmbH, April 1996.

Produktmonographie CellCept®: Roche AG, Grenzach-Wyhlen, 1996.

Produktmonographie Myfortic®: Novartis Pharma GmbH, Nürnberg 2004

Prüfzertifikat Astellas GmbH: Physicochemical properties of FK 506, München, 2006.

Prüfzertifikat Novartis Pharma AG: Use guidelines for RAD, Basel, 2005.

Prüfzertifikat LC Laboratories: Solubility of Rapamycin, Cat. No. R-5000, Woburn, MA, USA, 2004.

PZ 45/2002: Pharmazeutische Zeitung 147 (2002) 4304-4305.

R

Radatz, C.: Untersuchungen zur Lyophilisation von Liposomensuspensionen – Einfluss von Verfahrensparametern, Dissertation, Freie Universität Berlin, Berlin, 1994.

Radomska, A., Dobrucki, R.: The use of some ingredients for microemulsion preparation containing retinol and its esters, *Int. J. Pharm.* 196 (2000) 131 – 134.

Ranby, B; Rabek, J.F.: *ESR spectroscopy in Polymer Research*; Springer-Verlag Berlin, Heidelberg, New York, 1977, pp. 11 – 38.

Rawlings, A.V.; Mayo, A.M.; Rogers, J.; Scott, I.R.: Aging and the seasons influence stratum corneum lipid levels, *J. Invest. Dermatol.* 101 (1993) 483.

Rawlings, A.V.; Watkinson, A.; Rogers, J.; Mayo, A.M.; Hope, J., Scott, I.R.: Abnormalities in stratum corneum structure, lipid composition and desmosome degradation in soap-induced winter xerosis, *J. Soc. Cosmet. Chem.* 45 (1994) 203–220.

Rawlings, A. V.; Davies, A.; Carlomusto, M.; Pillai, S.; Zhang, K.; Kosturko, R.; Verdejo, P.; Feinberg, C.; Nguyen, L.; Chandar, P.: Effect of lactic acid isomers on keratinocyte ceramide synthesis, stratum corneum lipid levels and stratum corneum barrier function, *Arch. Dermatol. Res.* 288 (1996) 383–390.

Raymond, J. P.: *In vitro in vivo modelle zur Absorption von Cyclosporin A*, Dissertation, Universität Basel, 1986.

Rehse, B.: *Tensidhaltige Mehrkomponentensysteme auf der Basis von Polymertensiden*, Diplom, Humboldt-Universität zu Berlin, 1992.

Reim, M.: *Augenheilkunde*, Ferdinand Enke Verlag, Stuttgart, 1990.

Reis, A.; Megahed, M.; Reinhard, T.; Godehardt, E.; Spelsberg, H.; Braunstein, C.; Sundmacher, R.: Coadministration of the new macrolide immunosuppressant RAD and mycophenolate mofetil in experimental corneal transplantation, *Transplant.* 70 (2000) 1397-1401.

Reis, A.; Megahed, M.; Reinhard, T.; Braunstein, C.; Godehardt, E.; Sundmacher, R.: RAD, a new immunosuppressive macrolide in murine corneal transplantation, *Graefe's Arch. Clin. Exp. Ophthalmol.* 239 (2001) 689-692.

Reitamo, S.; Spuls, S.; Sassolas, B.; Lahfa, M.; Claudy, A.; Griffiths, C.E.M.: Efficacy of sirolimus (rapamycin) administered concomitantly with a subtherapeutic dose of cyclosporin in the treatment of severe psoriasis: a randomized controlled trial, *Br. J. Dermatol.* 145 (2001) 438-445.

Richter, T: *In Vitro-Hydratationsdynamik der menschlichen Hautbarriere: Eine Tieftemperatur-REM und TEM Studie von Wassereffekten im Stratum Corneum*, Dissertation, Hamburg, 2004.

Robertson, S.M.; Lang, L.S.: Leflunomide: Inhibition of S-antigen induced autoimmune uveitis in Lewis rats, *Agents Actions* 42 (1994) 3-4.

Robinson, J.C.: *Ocular Anatomy and physiology relevant to ocular drug delivery*, in: Mitra, A. K.: *Ophthalmic drug delivery systems*, Marcelk Decker, New York, 1993.

Robson, K. J.; Stewart, M. E., Michelsen, S.; Lazo, N. D.; Downing, D. T.: 6-OH-4-sphingenine in human epidermal ceramides, *J. Lipid Res.* 35 (1994) 2060-2068.

Röding, J.; Ghyczy, M.: Beeinflussung der Hautfeuchtigkeit der Liposomen. Stabilisierung von pflegenden Ölen und lipophilen Wirkstoffen mit Liposomen, *Seifen-Öle-Fette-Wachse* 10 (1991) 372-378.

Rodrigues, M.M., Rowden, G. et al.: Langerhans cells in the normal conjunctiva and peripheral cornea of selected species, *Invest. Ophthalmol. Vis. Sci.* 21 (1981) 759-65.

Rote Liste 2006, Arzneimittelverzeichnis für Deutschland (einschließlich EU-Zulassungen und bestimmter Medizinprodukte), Verlag Rote Liste Service GmbH, Frankfurt am Main, 2006.

Rupprecht, H.; Burchardi, Ch.; Mistry-Burchardi, N.; Fischereeder, M.; Weidner, S.: Immunsuppression, Elsevier Urban und Fischer Verlag München Jena, 2005.

S

Saettone, M.F.; Giannaccini, B.; Delmonte, G.; Campigli, V.; Tota, G.; La Marca, F. Solubilization of tropicamide by poloxamers: physicochemical data and activity data in rabbits and humans, *Int. J. Pharm.* 43 (1988) 67 – 76.

Saleh, A. M.; El-Khordagui, L. K.: Hydrotropic agents: a new definition, *Int. J. Pharm.* 24 (1985) 231-238.

Sasamoto, Y.; Hirose, S.; Ohno, S.; Onoe, K.; Matsuda, H.: Topical application of ciclosporin ophthalmic solution containing alpha- cyclodextrin in experimental uveitis, *Int. J. Ophthalmol.* 203 (1991) 118-125.

Sato, J.; Denda, M.; Nakanashi, J.; Nomura, J.; Koyama, J.: Cholesterol sulfate inhibits proteases that are involved in desquamation of stratum corneum, *J. Invest. Dermatol.* 111 (1998) 189 – 193.

Schaeffer, H. E.; Krohn, D. L.: Liposomes in topical drug delivery, *Invest. Ophthalmol. Vis. Sci.* 22 (1982) 220 – 227.

Schaefer, H., Redelmeier, T. E. Skin barrier, principles of percutaneous absorption, Verlag Karger, Basel, Freiburg, Paris, 1996.

Scheiffarth, O.F., Gabriel, N. et al.: Immunhistochemische Charakterisierung von Lymphozyten in der Cornea, *Fortschr. Ophthalmol.* 83 (1986) 714-715.

Scheinfeld, N.: The use of topical tacrolimus and pimecrolimus to treat psoriasis: a review, *Dermatol. Online J.* 10 3 (2003)

Scherphof, G., Roerdink, F., Waite, M., and Parks, J., Disintegration of phosphatidylcholine liposomes in plasma as a result of interaction with high-density lipoproteins, *Biochim. Biophys. Acta* 542 (1978) 296-307.

Scheuplein, R.J.; Blank, I.H.: Permeability of the skin, *Physiol. Rev.* 51 (1971) 702-747.

Schleicher, E.: Über den Reaktionsmechanismus der DNA-Photolyasen, Dissertation, Technische Universität München, Deutschland, 2002.

Schmitt, G.: Universität Heidelberg,
<http://www.rzuser.uni-heidelberg.de/~df6/themenpark/kalibration.htm> (29.09.06)

Schmitt-Sody, M. et al, Neovascular targeting therapy: paclitaxel encapsulated in cationic liposomes improves antitumoral efficacy, *Clin. Cancer Res.* 9 (2003) 2335-2341.

Schmölzer, S.: Kinetik der Vesikelbildung in kationischen Tensidsystemen, Dissertation, Universität Bayreuth, 2003.

Schoenwald, R.D, Ward, R.: Relationship between steroid permeability across excised rabbit cornea and octanol-water partition coefficients, *J. Pharm. Sci.* 67 (1978) 786-788.

Schoenwald, R.D., Huang, H.-S.: Corneal penetration behavior of β -blocking agents, I: physicochemical factors, *J. Pharm. Sci.* 72 (1983) 1266-1272.

Scholz, M.: In vitro-Permeationsstudien von hydrophilen und lipophilen Arzneistoffen an okularen Geweben und Zellkulturen, Dissertation, Humboldt Universität Berlin, Deutschland, 2003.

Schote, U.; Ganz, P.; Fahr, A.; Seelig, J.: Interactions of Cyclosporines with lipid membranes as studied by solid-state Nuclear Magnetic Resonance Spectroscopy and High-Sensitivity Titration Calorimetry, *J. Pharm. Sci.* 91 (2002) 856-867.

Schreier, H.; Bouwstra, J.: Liposomes and niosomes as topical drug carriers: dermal and transdermal drug delivery, *J. Control. Rel.* 30 (1994) 1-15.

Schubert, R.; Joos, M.; Deicher, M.; Lasch, J.: Mode of action of topically applied liposomes in dermatology and cosmetics, *Proceedings of MoBBEL (Molekularbiologische und Biotechnologische Entwicklungsliga e.V)* 6 (1992) 20-32.

Schubert, R.; Joos, M.; Deicher, M.; Magerle, R.; Lasch, J.: Destabilization of egg lecithin liposomes on the skin after topical application measured by perturbed angular correlation spectroscopy (PAC) with ^{111}In , *Biochim. Biophys. Acta, Biomembranes* 1150 (1993) 162-164.

Schubert, R.: Liposomen in Arzneimitteln, In: Müller, R. H., Hildebrandt, G. E. (Hg.): *Pharmazeutische Technologie: Moderne Arzneiformen*, Wiss. Verlagsges. Stuttgart 1997.

Schubert, R.: Bioverfügbarkeit und Stabilität von Immunsuppressiva, Entwicklung oraler und parenteraler Arzneiformen, *Pharm. Unserer Zeit* 34 (2005) 296-303.

- Schulman, J.H.; Stoeckenius, W.; Prince, L.M.: Mechanism of formation and structure of micro emulsions by electron microscopy, *J. Phys. Chem.* 63 (1959) 1677-1680.
- Schulze, H.J.; Mahrle, G.; Steigleder, G.K.: Topical cyclosporin A in psoriasis, *Br. J. Dermatol* 122 (1990) 113-114.
- Sedlakova, K.; Muckersie, E.; Robertson, M.; Filipec, M.; Forrester, J. V.: FTY 720 in corneal concordant xenotransplantation, *Transplantation* 79 (2005) 297-303.
- Šentjunc M., Vrhovnik K., Kristl J.: Liposomes as a topical delivery system: the role of size on transport studied by the EPR imaging method, *J. Control. Rel.* 59 (1999) 87-97.
- Sessa, G.; Weissmann, G.: Phospholipid spherules (liposomes) as a model for biological membranes, *J. Lipid Res.* 9 (1968) 310-318.
- Shah, J.; Atienza, J. M.; Duclos Jr., R. I.; Rawlings, A. V.; Dong, Z.; Shipley, G. G.: Structural and thermotropic properties of synthetic C 16:0 (palmitoyl) ceramide: effect of hydration, *J. Lipid Res.* 36 (1995) 1936 – 1944.
- Shi, J.; Kovacs, S.J.; Wang, Y.; Ludden, T.M.; Bhargava, V.O.: Population pharmacokinetics of the active metabolite of leflunomide in pediatric subjects with polyarticular course juvenile rheumatoid arthritis, *J. Pharmacokin. Pharmacodyn.* 32 (2005) 419-439.
- Short, S.M.; Rubas, W.; Paasch, B.D.; Mrsny, R.J.: Transport of biologically active interferon -gamma across human skin in vitro, *Pharm. Res.* 12 (1995) 1140-1145.
- Siebenbrodt, I.: Untersuchungen zur Verfügbarkeitsoptimierung von Ophthalmika, Dissertation, Humboldt-Universität Berlin, 1989.
- Siebenbrodt, I; Keipert, S.: Versuche zur Entwicklung und Charakterisierung ophthalmologisch verwendbarer tensidhaltiger Mehrkomponentensysteme, *Pharmazie* 46 (1991) 435 – 438.
- Siebenbrodt, I.; Keipert, S.: Poloxamer systems as potential ophthalmics. Part 1: Viscous solutions of polymeric surfactants, *PZ Wissenschaft* 5 (1992) 135-141.
- Siebenbrodt, I., Keipert, S. Poloxamer-systems as potential ophthalmics II. microemulsions. *Eur. J. Pharm. Biopharm.* 39 (1993) 25 – 30.
- Sieg, J.W., Robinson, J.R.: Vehicle effects on ocular drug bioavailability, II: evaluation of pilocarpine, *J. Pharm. Sci.* 66 (1977) 1222-1228.
- Siefert, B.: Einfluss von Cyclodextrin auf die Löslichkeit und die okulare Verfügbarkeit von Pilocarpin-HCL bzw. Thalidomid/ Penetrations- und Permeationsstudien an isolierter Cornea, Dissertation, Humboldt Universität zu Berlin, 1998.
- Siefert, B.; Keipert, S.: Influence of alpha-cyclodextrin and hydroxyalkylated beta-cyclodextrin derivatives on the in vitro corneal uptake and permeation of aqueous pilocarpine-HCl solutions, *J. Pharm. Sci.* 86 (1997) 716–720.

Simamora, P.; Alvarez, J.M.; Yalkowsky S. H.: Solubilization of rapamycin, *Int. J. Pharm.* 213 (2001) 25 – 29.

Simoes, S.I.; Marques, C.M.; Cruz, M.E.M.; Cevc, G.; Martins, M.B.F.: The effect of cholate on solubilisation and permeability of simple and protein-loaded phosphatidylcholine/sodium cholate mixed aggregates designed to mediate transdermal delivery of macromolecules, *Eur. J. Pharm. Biopharm.* 58 (2004) 509-519.

K. Singh, M. Mezei, Liposomal ophthalmic drug delivery. I Triamcinolone acetate, *Int. J. Pharm.* 16 (1983) 339 – 344.

Singh, K.; Mezei, M.: Liposomal ophthalmic drug delivery. II Dihydrostreptomycin sulphate, *Int. J. Pharm.* 19 (1984) 263 – 269.

Škalko, N.; Cajkovic, M.; Jelšenjok, I.: Liposomen with clindamycin hydrochloride in the therapy of acne vulgaris, *Int. J. Pharm.* 85 (1992) 97-101.

Song, Y.-K.; Kim, C.-K.: Topical delivery of low-molecular-weight heparin with surface- charged flexible liposomes, *Biomaterials* 27 (2006) 271-280.

Spectral database of organic compounds: National institute of advanced industrial science and technology (AIST). http://www.aist.go.jp/RIODB/SDBS/cgi-bin/cre_index.cgi?lang=eng (28.06.06)

Spernol, A.; Wirtz, K.: Zur Mikrorreibung in Flüssigkeiten, *Z. Naturf.* 8a (1953) 522-532.

Spielmann, H., Liebsch, M., Kalweit, S., et al. Results of a Validation Study in Germany on Two In Vitro Alternatives to the Draize eye irritation test, the HET-CAM Test and the 3T3 NRU Cytotoxicity Test, *ALAT* 24 (1996) 741-858.

Srinivas, V.; Rodley, G. A.; Ravikumar, K.; Robinson, Ward T.; Turnbull, Mark M.; Balasubramanian, D.. Molecular Organization in Hydrotrope Assemblies; *Langmuir* 13 (1997) 3235-3239.

Steele, J.A.; Choi, C.; Kwong, P.C.: Topical tacrolimus in the treatment of inverse psoriasis in children, *J. Am. Acad. Dermatol.* 53 (2005) 713-716.

Steinhilber, D: Pharmakologie der Immunsuppressiva, *Pharm. Unserer Zeit* 34 (2005) 276-283.

Stösser, R.; Mäder, K.; Borchert, H.-H.; Herrmann, W.; Schneider, G.; Liero, A.: Pharmaceutical aspects of ESR investigations on drug delivery systems, tissues and living systems; in: Ohya-Nishiguchi, H.; Packer, L. (Eds.) *Bioradicals detected by ESR spectroscopy*, Birkhäuser, Basel, pp. 301 – 320 (1995)

Stösser, R.; Herrmann, W.; Zehl, A.; Strehmel, V.; Laschewsky, A.: ESR Spin probes in ionic liquids, *Chem. Phys. Chem.* 7 (2006) 1106 – 1111.

Streilein, J.W., Niederkorn, J.Y. and Shadduck, J.A.: Systemic unresponsiveness induced in adult mice by anterior chamber presentation of minor histoincompatibility antigens, *J. Exp. Med.* 152 (1980) 1121-1125.

Strehmel, V.; Laschewsky, A.; Stösser, R.; Zehl, A.; Herrmann, W.: Mobility of spin probes in ionic liquids, *J. Phys. Org. Chem.* 19 (2006) 318-325.

Surber, C.; Itin, P.; Büchner, S.: Clinical controversy on the effect of topical ciclosporin: What is the target side?, *Dermatol.* 185 (1992) 242-245.

Svensson, J.-O.; Brattström, C.; Säwe, J.: Determination of Rapamycin in whole blood by HPLC, *Ther. Drug. Monit.* 19 (1997) 112-116.

T

Tardi, C.; Brandl, M.; Schubert, R.: Erosion and controlled release properties of semisolid vesicular phospholipid dispersions, *J. Control. Rel.* 55 (1998) 261-270.

Taylor, A. L., Watson, C. J. E., Bradley, J. A.: Immunosuppressive agents in solid organ transplantation: Mechanisms of action and therapeutic efficacy, *Crit. Rev. Oncol. Hematol.* 56 (2005) 23-46.

Tanaka, K.; Sekiguchi, M.; Okada, Y.: Restoration of ultraviolet-induced unscheduled DNA synthesis of xeroderma pigmentosum cells by the concomitant treatment with bacteriophage T4 endonuclease V and HVJ (Sendai virus), *Proc. Natl. Acad. Sci.* 72 (1975) 4071-4075.

Tamura, S.; Ohike, A.; Ibuki, R.; Amidon G. L.; Yamashita, S.: Tacrolimus is a Class II low solubility high permeability drug: The effect of p-glycoprotein efflux on regional permeability of tacrolimus in rats, *J. Phar. Sci.* 91 (2002) 719-729.

Theissen U; Luger T A; Schwarz T.: Successful topical administration of ciclosporin A in pyoderma gangraenosum, *Der Hautarzt* 47 (1996) 132-135.

Thiele, B.; Ghyczy, M.; Lunow, C.; Teichert, H.M.; Wolff, H.H.: Influence of phospholipide liposomes (PLL) on UVB – induced erythema formation, *Arch. Dermatol. Res.* 285 (1993) 428-431.

Thompson, P.; Xu, D.; Brunette, I.; Chen, H.: Combined effect of rapamycin and ciclosporine in the prevention of rat corneal allograft rejection, *Transplant. Proceed.* 30 (1998) 1033-1035.

V

Valenta, C.; Steininger, A.; Auner, B. G.: Phloretin and 6-ketocholestanol: membrane interactions studied by a phospholipid/polydiacetylene colorimetric assay and differential scanning calorimetry, *Eur. J. Pharm. Biopharm.* 57 (2004) 329-336.

Vandamme, T. F.: Microemulsions as ocular drug delivery systems: recent developments and future challenges, *Progr. Ret. Eye Res.* 21 (2002) 15 – 34.

Vantrappen, L., Geboes, K. et al.: Lymphocytes and langerhans cells in the normal human cornea, *Invest. Ophthalmol. Vis. Sci.* 26 (1985) 220-225.

Verma D. D.: Invasomes – Novel vesicular carriers for enhanced topical delivery: characterization and skin penetration properties, Dissertation, Philipps-Universität Marburg, 2002.

Verma, D. D.; Verma, S.; Blume, G.; Fahr, A.: Particle size of liposomes influences dermal delivery of substances into skin, *Eur. J. Pharm.* 258 (2003a) 141-151.

Verma, D. D.; Verma, S.; Blume, G.; Fahr, A.: Liposomes increase penetration of entrapped and non-entrapped hydrophilic substances into human skin: a skin penetration and confocal laser scanning microscopy study, *Eur. J. Pharm. Biopharm.* 55 (2003b) 271-277.

Verma, D. D.; Fahr, A.: Synergistic penetration enhancement effect of ethanol and phospholipids on the topical delivery of cyclosporin A, *J. Control. Rel.* 97 (2004) 55-66.

Vermorken, A. J. M.; Hukkelhoven, M. W. A. C.; Vermeesch-Markslag, A. M. G.; Goos, C. M. A. A.; Wirtz, P.; Ziegenmeyer, J. The use of liposomes in the topical application of steroids, *J. Pharm. Pharmacol.* 36 (1984) 334-336.

Vrhovnik, K; Kristl, J; Šentjurc, M; Šmid-Korbar, J.: Influence of liposome bilayer fluidity on the transport of encapsulated substance into the skin as evaluated by EPR, *Pharm. Res.* 15 (1997) 525-530.

W

Wang, D.P.; Lin, C.Y.; Chu, D.L.; Chang, L.C.: Effect of various physical/chemical properties on the transdermal delivery of cyclosporin through topical application, *Drug Dev. Ind. Pharm.* 23 (1997) 99-106.

Wang-Fan W., Küsters, E.; Mak, C.-H., Wang, Y.: Application of centrifugal counter current chromatography to the separation of macrolide antibiotic analogues. II. Determination of partition coefficients in comparison with the shake-flask method, *J. Liq. Chrom. & Rel. Technol.* 23 (2000) 1365-1376.

Wacker Chemie GmbH, Produktinformation (1998)

Wacker Chemie GmbH, EU-Sicherheitsdatenblatt (91/155/EWG) (1999)

Wang, W., Sasaki, H., Chien, D.S., Lee, V.H.L., Lipophilicity influence on conjunctival drug penetration in the pigmented rabbit: a comparison with corneal penetration, *Curr. Eye Res.* 10 (1991) 571-579.

Watsky, M.A., Jablonski, M.M.; Edelhauser, H.F.: Comparison of conjunctival and corneal surface areas in rabbit and human, *Curr. Eye Res.* 7 (1988) 483-486.

Wei, G.; Xu, H.; Ding, P. T.; Li, S. M.; Zheng, J. M.: Thermosetting gels with modulated gelation temperature for ophthalmic use: the rheological and gamma scintigraphic studies, *J. Control. Rel.* 83 (2002) 65-74.

Wertz, P.W.; Downing, D.T.: Acylglucosylceramides of pig epidermis: structure determination, *J. Lipid Res.* 24 (1983) 753-758.

Wertz P.W.; Cho E.I.; Downing D.T.: Effect of essential fatty acid deficiency on the epidermal sphingolipids of the rat; *Biochim. Biophys. Acta* 753 (1983) 350–355.

Wertz, P.W., Miethke, M.C., Long, S.A., Strauss, J.S., Downing, D.T.: The composition of ceramides from human stratum corneum and from comedones, *J. Invest. Dermatol.* 84 (1985) 410–412.

Werfel, T.; Wahn, U.: Black Box Warning für Pimecrolimus und Tacrolimus, *Deutsche Apothekerzeitung* 145 (2005) 1378-1380.

Whitcup, S. M., Pleyer, U., Lai, J.C., et al. Topical liposome-encapsulated FK506 for the treatment of endotoxin induced uveitis, *Ocul. Immunol. Inflam.* 6 (1998) 51-56.

Wohlrab, W.; Lasch, J.: Penetration kinetics of liposomal hydrocortisone in human skin, *Dermatologica* 174 (1987) 18-22.

Wohlrab, W.; Lasch, J.; Taube, K. M.; Wozniak, K. D.: Skin permeation of liposomal incorporated hydrocortisone, *Pharmazie* 44 (1989) 333-335.

Wohlrab, W.; Lasch, J.: The effect of liposomal incorporation of topically applied hydrocortisone on its serum concentration and urinary excretion, *Dermatol. Monatsschr.* 175 (1989) 348-352.

Wohlrab, W.: Aufbau und Funktion der Haut, in: Neubert, R. H. H.; Wohlrab, W.; Marsch, W.: *Dermatopharmazie, Vehikel-Wirkstoffe-Pharmakologie*, Wiss. Verlagsges. Stuttgart, 2001.

Wolka, A. M.; Rytting, J. H.; Reed, B. L.; Finin, B. C.: The interaction of the penetration enhancer DDAIP with a phospholipid model membrane, *Int. J. Pharm.* 271 (2004) 5-10.

Y

Yamabe, K.; Kato, Y.; Onishi, H.; Machida, Y.: In vitro characteristics of liposomes and double liposomes prepared using a novel glass beads method, *J. Control. Rel.* 90 (2003) 71-79.

Yamamoto, A., Serizawa, S., Ito, M., Sato, Y.: Stratum corneum abnormalities in atopic dermatitis, *Arch. Dermatol. Res.* 283 (1991) 219-223.

Yarosh, D. B.; Klein, J.; Kibitel, J.; Alas, L.; O'Connor, A.; Cummings, B.; Grob, D.; Gerstein, D.; Gilchrest, B.A.; Ichihashi, M.; Ogoshi, M.; Ueda, M.; Fernandez, V.; Chadwick, C.; Potten, C.S.; Proby, C.M.; Young, A.R.; Hawk, J.L.: Enzyme therapy of xeroderma pigmentosum: Safety and efficacy testing of T4N5 liposome lotion

containing a prokaryotic DNA repair enzyme. *Photodermatol. Photoimmunol. Photomed.* 12 (1996) 122–130.

Yarosh, D.B.; Klein, J.; O'Connor, A.; Hawk, J.; Rafal, E.; Wolf, P.: Effect of topically applied T4 endonucleaseV in liposomes on skin cancer in xeroderma pigmentosum: A randomised study. *Xeroderma Pigmentosum Study Group, Lancet* 357 (2001) 926–929.

Yarosh, D. B.: Liposomes in investigative dermatology, *Photoderm. Photoimmunol. Photomed.* 17 (2001) 203-212.

Ye, Q.; Asherman, J.; Stevenson, M.; Brownson, E.; Katre, N. V.: DepoFoam technology: a vehicle for controlled delivery of protein and peptide drugs, *J. Control. Rel.* 64 (2000) 155-166.

Z

Zolin, V. V.; Agafonova, O. A.; Kolokoltseva, A. A.; Danilenko, Ye. D.; Fedosova, L. K.; Gamalei, S. G.; Omigov, V. V.; Fadina, V. A.; Masycheva, V. I.: Investigation of the antiviral activity, pharmacokinetics, and toxic properties of human liposomal recombinant α 2b- interferon topically applied. *Vestnik Rossiiskoi Akademii Meditsinskikh Nauk* 5 (2001) 27-31.

Zuberbier, T.: Organ der Gesellschaft für Dermopharmazie, *DermoTopics* 5 (2005), 23-24.