

7 Literatur

- Ackermann, C., Dorresteijn, A. & Fischer, A. (2005): Clonal domains in postlarval *Platynereis dumerilii* (Annelida: Polychaeta). *J. Morphol.*, **266**(3): 258–280.
- Anderson, D. (1959): The embryology of the polychaete *Scoloplos armiger*. *Quart. J. Microscop. Sci.*, **100**(49): 89–166.
- Anderson, D. (1961): The Development of the Polychaete *Haploscoloplos fragilis*. *Quart. J. Microscop. Sci.*, **102**(58): 257–272.
- Anderson, D. (1966a): The comparative early embryology of the Oligochaeta, Hirudinea and Onychophora. *Proc. Linnean Soc. N. S. Wales*, **91**(1): 10–43.
- Anderson, D. (1966b): The comparative embryology of the Polychaeta. *Act. Zool.*, **47**: 1–42.
- Anderson, D. (1973): Embryology and Phylogeny in Annelids and Arthropods, Bd. 50 *International Series of Monographs in pure and applied Biology*. Pergamon Press, Oxford.
- Ax, P. (1999): Das System der Metazoa II. G. Fischer, Stuttgart.
- Bartolomaeus, T. (1989a): Larvale Nierenorgane bei *Lepidochiton cinereus* (Polyplacophora) und *Aeolidia papillosa* (Gastropoda). *Zoomorphology*, **108**(5): 297–307.
- Bartolomaeus, T. (1989b): Ultrastructure and development of the nephridia in *Anaitides mucosa* (Annelida, Polychaeta). *Zoomorphology*, **109**: 15–32.
- Bartolomaeus, T. (1993a): Die Leibeshöhlenverhältnisse und Nephridialorgane der Bilateria - Ultrastruktur, Entwicklung und Evolution. Diss., Göttingen.
- Bartolomaeus, T. (1993b): Ultrastructure of the protonephridia in the larva of *Autolytus prolifer* (Annelida, Syllidae): Implications for annelid phylogeny. *Microfauna mar.*, **8**: 55–64.
- Bartolomaeus, T. (1995): Ultrastructure of the protonephridia in larval *Magelona mirabilis* (Spionida) and *Pectinaria auricoma* (Terebellida): Head kidneys in the ground pattern of the Annelida. *Microfauna mar.*, **10**: 117–141.
- Bartolomaeus, T. (1998): Head Kidneys in Hatchlings of *Scoloplos armiger* (Annelida: Orbiniida): Implications for the Occurrence of Protonephridia in lecithotrophic Larvae. *J. Mar. Biol. Assoc. UK*, **78**: 183–192.
- Bartolomaeus, T. (1999): Structure, function and development of segmental organs in Annelida. *Hydrobiol.*, **402**: 21–37.

- Bartolomaeus, T. & Ax, P. (1992): Protonephridia and Metanephridia - their relation within the Bilateria. *Z. zool. Syst. Evol.-Forsch.*, **30**: 21–45.
- Bartolomaeus, T. & Quast, B. (2005): Structure and development of nephridia in Annelida and related taxa. *Hydrobiol.*, **535/536**: 139–164.
- Bergh, R. (1885a): Über die Metamorphose von *Aulastoma gulo*. *Arb. zool. zootom. Inst. Würzburg*, **7**: 231–291.
- Bergh, R. (1885b): Über die Metamorphose von *Nephelis*. *Z. wiss. Zool.*, **41**: 284–301.
- Bergh, R. (1888): Zur Bildungsgeschichte der Exkretionsorgane bei *Criodrilus*. *Arb. zool. zootom. Inst. Würzburg*, **8**: 223–248.
- Bergrter, A. & Paululat, A. (2007): Pattern of body-wall muscle differentiation during embryonic development of *Enchytraeus coronatus* (Annelida: Oligochaeta; Enchytraeidae). *J. Morphol.*, **268**: 537–549.
- Blake, J. A. (1975): The larval development of Polychaeta from the Northern California Coast. I. *Cirriformia spirabrancha* (Family Cirratulidae). *Trans. Amer. Microscop. Soc.*, **94**(2): 179–188.
- Bleidorn, C., Vogt, L. & Bartolomaeus, T. (2003): New insights into polychaete phylogeny (Annelida) inferred from 18S rDNA sequences. *Mol. Phylogen. Evol.*, **29**(2): 279–288.
- Bonch-Bruevich, E. & Malakhov, V. (1987): Organization of early larve of the polychaete worm *Chaetopterus variopedatus* (Spiromorpha, Chaetopteridae). *Dokl. Akad. Nauk SSSR*, **292**(4): 1013–1015.
- Brandenburg, J. (1966): Die Reusenformen der Cyrtocytten. eine Beschreibung von fünf weiteren Reusengeißelzellen und eine vergleichende Betrachtung. *Zool. Beitr.*, **12**: 345–417.
- Brinkhurst, R. (1971): A Guide for Identification of British Aquatic Oligochaeta. Bd. 22 *Scientific publication / Freshwater Biological Association*, S. 55. Freshw. Biol. Assoc., The Ferry House, Far Sawrey, Ambleside, Cumbria.
- Brinkhurst, R. (1982): Evolution in the Annelida. *Can. J. Zool.*, **60**: 1043–1059.
- Brinkhurst, R. (1994): Evolutionary relationships within the Clitellata: an Update. *Megadrilogica*, **5**(10): 109–112.
- Dohle, W. (1999): The ancestral cleavage pattern of the clitellates and its phylogenetic deviations. *Hydrobiol.*, **402**: 267–283.
- Eisig, H. D. (1898): Zur Entwicklungsgeschichte der Capitelliden. Mittheilungen aus der zoologischen Station zu Neapel, **13**: 1–292.

- Erséus, C. & Källersjö, M. (2004): 18S rDNA phylogeny of Clitellata (Annelida). *Zool. Scri.*, **33**(2): 187–196.
- Fauchald, K. & Rouse, G. (1997): Polychaete systematics: Past and present. *Zool. Scri.*, **26**(2): 71–138.
- Fernández, J. & Stent, G. (1982): Embryonic development of the hirudinid leech *Hirudo medicinalis*: structure, development and segmentation of the germinal plate. *J. Embr. Exp. Morphol.*, **72**: 71–96.
- Gleizer, L. & Stent, G. (1993): Developmental origin of segmental identity in the leech mesoderm. *Development*, **117**(1): 177–189.
- Goodrich, E. (1895): On the coelom, genital ducts, and nephridia. *Quart. J. Microscop. Sci.*, **37**: 477–508.
- Goodrich, E. (1945): The study of Nephridia and genital ducts since 1895. *Quart. J. Microscop. Sci.*, **86**(2): 113–301.
- Gustavsson, L. & Erséus, C. (1999): Morphology and phylogenetic implications of oesophageal modifications in the Limnodriloidinae (Oligochaeta, Tubificidae). *J. Zool.*, **248**(4): 467–482.
- Haszprunar, G. & Ruthensteiner, B. (2000): Microanatomy and ultrastructure of the protonephridial system in the larva of the limpet, *Patella vulgata* L. (Mollusca, Patellogastropoda). *J. Submicroscop. Cyt. Pathol.*, **32**(1): 59–67.
- Hatschek, B. (1886): Zur Entwicklung des Kopfes von *Polygordius*. *Arb. Inst. Univ. Zool. Stat.*, **6**: 109–210.
- Heimler, W. (1981): Untersuchung zur Larvalentwicklung von *Lanice conchilega* (Pallas) 1766 (Polychaeta, Terebellomorpha). Teil II: Bau und Ultrastruktur der Trochophoralarve. *Zool. Jb. Anatom. Ontol.*, **106**(2): 236–277.
- Heimler, W. (1988): Larvae. In: Westheide, W. & Hermans, C. (Hrsg.), *The ultratruucture of the Polychaeta*, Bd. 4 *Microfauna mar.*, S. 353–371. G. Fischer, Stuttgart.
- Hessling, R. (2002): Metameric organisation of the nervous system in developmental stages of *Urechis caupo* (Echiura) and its phylogenetic implications. *Zoomorphology*, **121**(4): 221–234.
- Hessling, R., Müller, C. & Westheide, W. (1999): CLSM analysis of serotonin-immunoreactive neurons in the central nervous system of *Nais variabilis*, *Slavina appendiculata* and *Stylaria lacustris* (Oligochaeta: Naididae). *Hydrobiol.*, **406**: 223–233.

- Hessling, R. & Westheide, W. (1999): CLSM analysis of development and structure of the central nervous system of *Enchytraeus crypticus* ("Oligochaeta", Enchytraeidae). *Zoomorphology*, **119**: 37–47.
- Hill, S. & Boyer, B. (2001): Phalloidin labeling of developing muscle in embryos of the polychaete *Capitella sp. I*. *Biol. Bull.*, **201**(2): 257–258.
- Hoffmann, R. (1899): Beiträge zur Entwicklungsgeschichte der Oligochäten. *Z. wiss. Zool.*, **66**: 335–357.
- Huang, F., Kang, D., Ramirez-Weber, F.-A., Bissen, S. & Weisblat, D. (2002): Micromere lineages in the glossiphoniid leech *Helobdella*. *Development*, **129**: 719–732.
- Iwanoff, P. (1928): Die Entwicklung der Larvalsegmente bei den Anneliden. *Z. Morphol. Ökolog.*, **10**: 62–161.
- Jamieson, B. (1981): The Ultrastructure of the Oligochaeta. Acad. Press, London.
- Kaltenhäuser, U. (1993): Mechanismen der Harnbildung blutsaugender und fleischfressender Egel : *Hirudo medicinalis* (L.) und *Macrobdella decora* (Say), Hirudinea: Gnathobdelliformes. Diss., Konstanz.
- Kleinenberg, N. (1879): The Development of the Earthworm, *Lumbricus trapezoides*, Duges. *Quart. J. Microscop. Sci.*, **19**: 206–244.
- Kuper, M. (2001): Ultrastrukturuntersuchungen der Segmentalorgane, der Spermien und der Brutpflegestrukturen innerhalb der Syllidae (Annelida: Polychaeta). Diss., Osnabrück.
- Lacalli, T. (1984): Structure and Organization of the Nervous System in the Trochophore Larva of *Spirobranchus*. *Phil. Trans. Roy. Soc. London, Ser. B*, **306**(1126): 79–135.
- Lehmann, O. (1887): Beiträge zur Frage von der Homologie der Segmentalorgane und der Ausführgänge der Geschlechtsprodukte bei den Oligochaeten. *Jena. Z. Nat.-Wiss.*, **21**: 322–360.
- Marsden, J. & Anderson, D. (1981): Larval development and metamorphosis of the serpulid polychaete *Galeolaria caespitosa* Lamarck. *Austr. J. Mar. Freshw. Res.*, **32**(4): 667–680.
- Martin, P. (2001): On the origin of the Hirudinea and the demise of the Oligochaeta. *Proc. Roy. Soc. London, Ser. B*, **268**(1471): 1089–1098.
- Martin, P., Kaygorodova, I., Sherbakov, D. & Verheyen, E. (2000): Rapidly Evolving Lineages Impede the Resolution of Phylogenetic Relationships among Clitellata (Annelida). *Mol. Phylogen. Evol.*, **15**(3): 355–368.

- McHugh, D. (1997): Molecular evidence that echiurans and pogonophores are derived annelids. *Proc. Natl. Acad. Sci. U. S. A.*, **94**: 8006–8009.
- Meyer, E. (1901): Studien über den Körperbau der Anneliden. V. Das Mesoderm der Ringelwürmer. *Mittheilungen aus der zoologischen Station zu Neapel*, **14**: 247–585.
- Meyer, F. (1917): Untersuchungen über den Bau und die Entwicklung des Blutgefäßsystems bei *Tubifex tubifex* (MÜLL.). *Jena. Z. Nat.-Wiss., N.F.* **47**: 203–244.
- Neuhaus, B. & Kristensen, R. M. (2007): Ultrastructure of the protonephridia of larval *Rugiloricus* cf. *cauliculus*, male *Armorloricus elegans*, and female *Nanalaricus mysticus* (Loricifera). *J. Morphol.*, **268**(4): 357–370.
- Nielsen, C. (2004): Trochophora larvae: Cell-lineages, ciliary bands, and body regions. 1. Annelida and Mollusca. *J. Exp. Zool.*, **302B**(1): 35–68.
- Nielsen, C. (2005): Trochophora larvae: cell-lineages, ciliary bands and body regions. 2. Other groups and general discussion. *J. Exp. Zool.*, **304B**(5): 401–447.
- Nielsen, C., Haszprunar, G., Ruthensteiner, B. & Wanninger, A. (2007): Early development of the aplacophoran mollusc *Chaetoderma*. *Act. Zool.*, **88**(3): 231–247.
- Okusu, A. (2002): Embryogenesis and development of *Epimenia babai* (Mollusca Neomeniomorpha). *Biol. Bull.*, **203**(1): 87–103.
- Olive, P. (1975): Reproductive biology of *Eulalia viridis* (Müller) (Polychaeta: Phyllodocidae) in the North Eastern U.K. *J. Mar. Biol. Assoc. UK*, **55**: 313–326.
- Pernet, B. (2003): Persistent ancestral feeding structures in nonfeeding annelid larvae. *Biol. Bull.*, **205**: 295–307.
- Purschke, G. (1999): Terrestrial polychaetes - models for the evolution of the Clitellata (Annelida)? *Hydrobiol.*, **406**: 87–99.
- Purschke, G. (2002a): Male genital organs, spermatogenesis and spermatozoa in the enigmatic terrestrial polychaete *Parergodrilus heideri* (Annelida, Parergodrilidae). *Zoology*, **121**(3): 125–138.
- Purschke, G. (2002b): On the ground pattern of Annelida. *Organ. Div. Evol.*, **2**(3): 181–196.
- Purschke, G., Hessling, R. & Westheide, W. (2000): The phylogenetic position of the Clitellata and the Echiura - on the problematic assessment of absent characters. *J. Zool. Syst. Evol. Res.*, **38**: 165–173.

- Purschke, G., Westheide, W., Rohde, D. & Brinkhurst, R. (1993): Morphological reinvestigation and phylogenetic relationship of *Acanthobdella peledina* (Annelida, Clitellata). *Zoomorphology*, **113**: 91–101.
- Quast, B. & Bartolomaeus, T. (2001): Ultrastructure and significance of the transitory nephridia in *Erpobdella octoculata* (Hirudinea, Annelida). *Zoomorphology*, **120**: 205–213.
- Raven, C. P. (1958): Morphogenesis: The Analysis of Molluscan Development, Bd. 2 *International Series of Monographs in pure and applied Biology*. Pergamon Press, Oxford.
- Reynolds, E. S. (1963): The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J. Cell Biol.*, **17**(1): 208–212.
- Reynolds, S. A., French, K. A., Baader, A. & Kristan, W. B. (1998): Staging of middle and late embryonic development in the medicinal leech, *Hirudo medicinalis*. *J. Compar. Neurol.*, **402**: 155–167.
- Rota, E., Martin, P. & Erséus, C. (2001): Soil-dwelling polychaetes: enigmatic as ever? Some hints on their phylogenetic relationships as suggested by a maximum parsimony analysis of 18S rRNA gene sequences. *Contr. Zool.*, **70**(3): 127–138.
- Rouse, G. (2000): The epitome of hand waving? Larval feeding and hypotheses of metazoan phylogeny. *Evol. Develop.*, **2**(4): 222–233.
- Rouse, G. & Fauchald, K. (1995): The Articulation of Annelids. *Zool. Scri.*, **24**(4): 269–301.
- Rouse, G. & Fauchald, K. (1997): Cladistics and polychaetes. *Zool. Scri.*, **26**(2): 139–204.
- Rousset, V., Pleijel, F., Rouse, G., Erséus, C. & Siddall, M. (2007): A molecular phylogeny of annelids. *Cladistics*, **23**(1): 41–63.
- Ruppert, E. & Smith, P. (1988): The functional organisation of filtration nephridia. *Biol. Rev.*, **63**: 231–258.
- Ruthensteiner, B. & Schaefer, K. (1991): On the Protonephridia and 'Larval Kidneys' of *Nassarius (Hinia) reticulatus* (Linaeus) (Caenogastropoda). *J. Mollusc. Stud.*, **57**(3): 323–329.
- Ruthensteiner, B., Wanninger, A. & Haszprunar, G. (2001): The protonephridial system of the tusk shell, *Antalis entalis* (Mollusca, Scaphopoda). *Zoomorphology*, **121**: 19–26.
- Sawyer, R. (1986): Leech Biology and Behaviour. In: Anatomy, Physiology and Behaviour, Bd. 1 *Oxford Sci. Publ.* Clarendon Press, Oxford.

- Schlötzer-Schrehardt, U. (1992): Ultrastrukturelle Untersuchungen zur Reproduktion und Postembryonalentwicklung einschliesslich Adultorganisation von *Pygospio elegans* Claparede, 1863 (Polychaeta, Spionidae). Diss., Konstanz.
- Schmidt, G. A. (1936): Vergleichend embryologische Studien über die Typen der Embryonalanpassungen bei Hirudineen und Nemertinen. *Zoomorphology*, **32**(4): 650–671.
- Seaver, E., Thamm, K. & Hill, S. (2005): Growth patterns during segmentation in the two polychaete annelids, *Capitella sp. I* and *Hydroïdes elegans*: comparisons at distinct life history stages. *Evol. Develop.*, **7**(4): 312–326.
- Segrove, F. (1941): The development of the Serpulid *Pomatoceros triquetus* L. *Quart. J. Microscop. Sci.*, **82**(327): 467–540.
- Shearer, C. (1908): Studies on the Development of Larval Nephridia. Part II. *Polygordius*. *Phil. Trans. Roy. Soc. London, Ser. B*, **199**: 199–230.
- Shearer, C. (1911): On the Development and Structure of the Trochophore of *Hydroïdes uncinatus* (Eupomatus). *Quart. J. Microscop. Sci.*, **56**(223): 543–590.
- Shimizu, T. (1982): Development in the freshwater Oligochaete *Tubifex*. In: Harrison, F. & Cowden, R. (Hrsg.), *Developmental Biology of Freshwater Invertebrates*, S. 283–316. Liss, New York.
- Siddall, M., Apakupakul, K., Burreson, E., Coates, K., Erséus, C., Gelder, S., Källersjö, M. & Trapido-Rosenthal, H. (2001): Validating Livanow: molecular data agree that Leeches, Branchiobdellidans, and *Acanthobdella peledina* form a monophyletic group of Oligochaetes. *Mol. Phylogen. Evol.*, **21**(2): 346–351.
- Sims, R. W. & Gerard, B. M. (1985): Earthworms, Bd. 31 A *Synopsis of the British Fauna*. Brill and Backhuys, London, 171 S.
- Smith, C. & Weisblat, D. (1994): Micromere fate maps in leech embryos: lineage-specific differences in rates of cell proliferation. *Development*, **120**: 3427–3438.
- Smith, P. & Ruppert, E. (1988): Nephridia. In: Westheide, W. & Hermans, C. (Hrsg.), *The ultrastructure of the Polychaeta*, Bd. 4 *Microfauna mar.*, S. 231–262. G. Fischer, Stuttgart.
- Smith, P., Ruppert, E. & Gardiner, S. (1987): A deuterostome-like nephridium in the miraria larva of *Owenia fusiformis* (Polychaeta, Annelida). *Biol. Bull.*, **172**: 315–323.
- Smith, R. (1984): Development and settling of *Spirobranchus giganteus* (Polychaeta; Serpulidae). S. 461–483.

- Stecher, H.-J. (1968): Zur Organisation und Fortpflanzung von *Pisione remota* (Southern) (Polychaeta, Pisionidae). Z. Morphol. Tiere, **61**: 347–410.
- Strathmann, R. R. (1978): The evolution and loss of feeding larval stages of marine invertebrates. Evolution, **32**(4): 894–906.
- Strathmann, R. R., Jahn, T. L. & Fonseca, J. R. C. (1972): Suspension feeding by marine invertebrate larvae: Clearance of particles by ciliated bands of a rotifer, pluteus, and trochophore. Biol. Bull., **142**(3): 505–519.
- Struck, T., Hessling, R. & Purschke, G. (2002): The phylogenetic position of the Aeolosomatidae and Parergodrilidae, two enigmatic oligochaete-like taxa of the 'Polychaeta', based on molecular data from 18S rDNA sequences. J. Zool. Syst. Evol. Res., **40**(3): 155–163.
- Struck, T. & Purschke, G. (2005): The sister group relationship of Aeolosomatidae and Potamodrilidae (Annelida: "Polychaeta") – a molecular phylogenetic approach based on 18S rDNA and cytochrome oxidase I. Zool. Anz., **243**(4): 281–293.
- Struck, T., Schult, N., Kusen, T., Hickman, E., Bleidorn, C., McHugh, D. & Halanych, K. (2007): Annelid phylogeny and the status of Sipuncula and Echiura. Evol. Biol., **7**(1).
- Sukatschoff, B. (1900): Beiträge zur Entwicklungsgeschichte der Hirudineen. I. Zur Kenntnis der Urnieren von *Nephelis vulgaris* Moqu. Tand. und *Aulastomum gulo* Moqu. Tand. Z. wiss. Zool., **67**: 618–639.
- Svetlov, P. (1928): Untersuchungen über die Entwicklungsgeschichte der Regenwürmer. Trav. Lab. zool. Stat. biol. Sébastopol, **13**, Serie 2: 313–329.
- Tannreuther, G. (1915): The Embryology of *Bdellodrilus philadelphicus*. J. Morphol., **26**(2): 143–216.
- Vejdovsky, F. (1887): Das larvale und definitive Excretionssystem. Zool. Anz., **10**: 681–685.
- Vejdovsky, F. (1892): Entwicklungsgeschichtliche Untersuchungen. J. Otto, Prag.
- Wachs, B. (1967a): Die häufigsten hämoglobinführenden Oligochaeten der mitteleuropäischen Binnengewässer. Hydrobiol., **30**(2): 225–247.
- Wachs, B. (1967b): Die Oligochaeten-Fauna der Fließgewässer unter besonderer Berücksichtigung der Beziehungen zwischen der Tubificiden-Besiedlung und dem Substrat. Arch. Hydrobiol., **63**(3): 310–386.
- Weisblat, D., Kim, S. & Stent, G. (1984): Embryonic Origins of Cells in the Leech *Habenularia triserialis*. Develop. Biol., **104**: 65–85.

- Wessing, A. & Polenz, A. (1974): Structure, Development and Function of the Protonephridia in Trochophores of *Pomatoceros triqueter* (Annelida, Polychaeta, Sedentaria). *Cell Tissue Res.*, **156**(1): 21–33.
- Westheide, W. (1986): The nephridia of the interstitial polychaete *Hesionides arenaria* and their phylogenetic significance (Polychaeta, Hesionidae). *Zoomorphology*, **106**: 35–43.
- Westheide, W. (1997): The direction of evolution within the Polychaeta. *J. nat. Hist.*, **31**: 1–15.
- Westheide, W., McHugh, D., Purschke, G. & Rouse, G. (1999): Systematization of the Annelida: different approaches. *Hydrobiol.*, **402**: 291–307.
- Wilcke, D. (1967): Oligochaeta. In: Brohmer, P., Ehrmann, P., Ulmer, G. & Schiemenz, H. (Hrsg.), Urtiere - Hohltiere - Würmer, Bd. 7a *Die Tierwelt Mitteleuropas*, S. 161. Quelle und Meyer, Leipzig.
- Wilson, D. (1932): On the Mitraria Larva of *Owenia fusiformis* Delle Chiaje. *Phil. Trans. Roy. Soc. London, Ser. B*, **221**: 231–334.
- Wilson, D. (1936): The Development of the Sabellid *Branchiomma vesiculosum*. *Quart. J. Microscop. Sci.*, **78**(312): 543–603.
- Wilson, E. (1889): The Embryology of the Earthworm. *J. Morphol.*, **3**: 387–462.
- Woltereck, R. (1905): Zur Kopffrage der Anneliden. *Verh. Dt. Zool. Ges.*, **7**: 154–186.
- Zackson, S. L. (1982): Cell clones and segmentation in leech development. *Cell*, **31**(3): 761–770.