

8. Literaturverzeichnis

Aarestrup, F. M. (1998)

Association between decreased susceptibility to a new antibiotic for treatment of human diseases, everninomycin (SCH 27899), and resistance to an antibiotic used for growth promotion in animals, avilamycin.

Microb. Drug Resist. 4, 137-141

Aarestrup, F. M., Bager, F., Jensen, N.E., Madsen, M., Meyling, A. und Wegener, H.C. (1998)

Surveillance of antimicrobial resistance in bacteria isolated from food animals to antimicrobial growth promoters and related therapeutic agents in Denmark.

APMIS 106, 606-622

Aarestrup, F. M. (2000a)

Characterization of glycopeptide-resistant *Enterococcus faecium* (GRE) from broilers and pigs in Denmark: Genetic evidence that persistence of GRE in pig herds is associated with coselection by resistance to macrolides.

J. Clin. Microbiol. 38, 2774-2777

Aarestrup, F. M. (2000b)

Occurrence, selection and spread of resistance to antimicrobial agents used for growth promotion for food animals in Denmark.

APMIS 108, Suppl. 101, 5-48

Aarestrup, F. M. und Jensen, L. B. (2000)

Presence of variations in ribosomal protein L16 corresponding to susceptibility of enterococci to oligosaccharides (avilamycin and evernimicin).

Antimicrob. Agents Chemother. 44, 3425-3427

Aarestrup, F. M., Kruse, H., Tast, E., Hammerum, A.M. und Jensen, L.B. (2000)

Associations between the use of antimicrobial agents for growth promotion and the occurrence of resistance among *Enterococcus faecium* from broilers and pigs in Denmark, Finland and Norway.

- Microb. Drug Resist. 6, 63-70
- Aarestrup, F. M., Seyfarth, A.M., Emborg, H.-D., Pedersen, K., Hendriksen, R.S. und Bager, F. (2001)
Effect of abolishment of the use of antimicrobial agents for growth promotion on occurrence of antimicrobial resistance in fecal enterococci from food animals in Denmark.
Antimicrob. Agents Chemother. 45, 2054-2059
- Aarestrup, F. M., Hasman, H., Jensen, L. B., Moreno, M., Herrero, I. A., Dominguez, L., Finn, M. und Franklin, A. (2002)
Antimicrobial resistance among enterococci from pigs in three European countries.
Appl. Environ. Microbiol. 68, 4127-4129
- Aarestrup, F. M. und McNicholas, P. M. (2002)
Incidence of high-level evernimicin resistance in *Enterococcus faecium* among food animals and humans.
Antimicrob. Agents Chemother. 46, 3088-3090
- Ambur, O.-H., Reynolds, P.E. und Arias, C. A. (2002)
D-Ala:D-Ala ligase gene flanking the *vanC* cluster: evidence for presence of three ligase genes in vancomycin-resistant *Enterococcus gallinarum* BM4174.
Antimicrob. Agents Chemother. 46, 95-100
- Aminov, R.I., Garrigues-Jeanjean, N. und Mackie, R.I. (2001)
Molecular ecology of tetracycline resistance: development and validation of primers for detection of tetracycline resistance genes encoding ribosomal protection proteins.
Appl. Environ. Microbiol. 67, 22-32
- Anonymus (1997)
Antimicrobial feed additives.
Report from the Commission on Antimicrobial Feed Additives, ministry of agriculture.
Norstedts Tryckeri AB, Stockholm, Schweden
- Arihara, K., Ogihara, S., Sakata, J., Itoh, M. und Kondo, Y. (1991)
Antimicrobial activity of *Enterococcus faecalis* against *Listeria monocytogenes*.

- Lett. Appl. Microbiol. 13, 190-192
- Arihara, K., Cassens, R.G. und Luchansky, J.B. (1993)
 Characterization of bacteriocins from *Enterococcus faecium* with activity against *Listeria monocytogenes*.
 Int. J. Food Microbiology 19, 123-134
- Arizcun, C., Barcina, Y. und Torre, P. (1997)
 Identification and characterization of proteolytic activity of *Enterococcus* ssp. isolated from milk and Roncal and Idiazabal cheese.
 Int. J. Food Microbiology 38, 17-24
- Arthur, M. und Quintiliani, R. Jr. (2001)
 Regulation of van A- and van B- type glycopeptide resistance in enterococci.
 Antimicrob. Agents Chemother. 45, 375-381
- Aymerich, T., Holo, H., Havarstein, L.S., Hugas, M., Garriga, M. und Nes, I.F. (1996)
 Biochemical and genetic characterization of enterocin A from *Enterococcus faecium*, a new antilisterial bacteriocin in the pediocin family of bacteriocins.
 Appl. Environ. Microbiol. 62, 1676-1682
- Aymerich, T., Garriga, M., Ylla, J., Vallier, J., Monfort, J.M. und Hugas, M. (2000)
 Application of enterocins as biopreservatives against *Listeria innocua* in meat products.
 J. Food Protect. 63, 721-726
- Baumgart, J. (1996)
 Enterokokken.
 In: Baumgart, J. (Hrsg.): Mikrobiologische Untersuchung von Lebensmitteln.
 Behr's Verlag, Hamburg, 14-16
- Baumgartner, A., Kueffer, M. und Rohner, P. (2001)
 Occurrence and antibiotic resistance of enterococci in various ready-to-eat foods.
 Arch. Lebensmittelhyg. 52, 16-19
- Bell, R.G. und Gill, C.O. (1982)
 Microbial spoilage of luncheon meat prepared in an impermeable plastic casing.

- J. Appl. Bacteriol. 53, 97-102
- Bell, R.G. und DeLacey, K.M. (1984)
Heat injury and recovery of *Streptococcus faecium* associated with the souring of chub-packed luncheon meat.
J. Appl. Bacteriol. 57, 229-236
- Biavasco, F., Paladini, C., Vignaroli, C., Foglia, G., Manso, E. und Varaldo, P. E. (2001)
Recovery from a single blood culture of two *Enterococcus gallinarum* isolates carrying both *vanC-1* and *vanA* cluster genes and differing in glycopeptide susceptibility.
Eur. J. Clin. Microbiol. Infect. Dis. 20, 309-314
- BMVEL-Bundesministerium für Verbraucherschutz, Ernährung und Landwirtschaft (2002)
Gefährliche Abfälle als Futtermittel in den Handel gebracht- Schnellwarnsystem für Futtermittel dringend erforderlich.
Pressemitteilung Nr.7 vom 14.01.2002
- Boerlin, P., Wissing, A., Aarestrup, F.M., Frey, J. und Nicolet, J. (2001)
Antimicrobial growth promoter ban and resistance to macrolides and vancomycin in enterococci from pigs.
J. Clin. Microbiol. 39, 4193-4195
- Boyd, D. A., Cabral, T., Van Caesele, P., Wylie, J. und Mulvey, M. R. (2002)
Molecular characterization of the *vanE* gene cluster in vancomycin-resistant *Enterococcus faecalis* N00-410 isolated in Canada.
Antimicrob. Agents Chemother. 46, 1977-1979
- Bozdogan, B., Berrezouga, L., Kuo, M.-S., Yurek, D. A., Farley, K. A., Stockman, B. J. und Leclercq, R. (1999)
A new resistance gene, *linB*, conferring resistance to lincosamides by nucleotidylation in *Enterococcus faecium* HM1025.
Antimicrob. Agents Chemother. 43, 925-929
- Bozdogan, B. und Leclercq, R. (1999)
Effects of genes encoding resistance to streptogramins A and B on the activity of quinupristin-dalfopristin against *Enterococcus faecium*.
Antimicrob. Agents Chemother. 43, 2720-2725

- Bridge, P. D. und Sneath, P. H. A. (1982)
Streptococcus gallinarum sp. nov. and *Streptococcus oralis* sp. nov.
Int. J. Syst. Bacteriol. 32, 410-415
- Brisse, S., Fluit, A. C., Wagner, U., Heisig, P., Milatovic, D., Verhoef, J., Scheuring, S., Köhrer, K. und Schmitz, F.-J. (1999)
Association of alterations in ParC and GyrA proteins with resistance of clinical isolates of *Enterococcus faecium* to nine different fluoroquinolones.
Antimicrob. Agents Chemother. 43, 2513-2516
- Bushby, S. R. M. und Hitchings, G. H. (1968)
Trimethoprim, a sulphonamide potentiator.
Br. J. Pharmac. Chemother. 33, 72-90
- Butaye, P., Devriese, L. A. und Haesebrouck, F. (1999)
Phenotypic distinction in *Enterococcus faecium* and *Enterococcus faecalis* strains between susceptibility and resistance to growth-enhancing antibiotics.
Antimicrob. Agents Chemother. 43, 2569-2570
- Butaye, P., Devriese, L. A. und Haesebrouck, F. (2000a)
Influence of different medium components on the *in vitro* activity of the growth-promoting antibiotic flavomycin against enterococci.
J. Antimicrob. Chemother. 46, 713-716
- Butaye, P., Van Damme, K., Devriese, L.A., Van Damme, L., Baele, M., Lauwers, S., Haesebrouck, F. (2000b)
In vitro susceptibility of *Enterococcus faecium* isolated from food to growth-promoting and therapeutic antibiotics.
Int. J. Food Microbiology 54, 181-187
- Butaye, P., Devriese, L. A. und Haesebrouck, F. (2001)
Differences in antibiotic resistance patterns of *Enterococcus faecalis* and *Enterococcus faecium* strains isolated from farm and pet animals.
Antimicrob. Agents Chemother. 45, 1374-1378
- Callewaert, R., Hugas, M. und De Vuyst, L. (2000)
Competitiveness and bacteriocin production of *Enterococci* in the production of spanish-style dry fermented sausages.
Int. J. Food Microbiology 57, 33-42

- Centeno, J.A., Menendez, S. und Rodriguez-Otero, J.L. (1996)
Main microbial flora present as natural starters in Cebreiro raw cow's-milk cheese (Northwest Spain).
Int. J. Food Microbiology 33, 307-313
- Centeno, J.A., Menendez, S., Hermida, M. und Rodriguez-Otero, J.L. (1999)
Effects of the addition of *Enterococcus faecalis* in Cebreiro cheese manufacture.
Int. J. Food Microbiology 48, 97-111
- Chadwick, P.R., Woodford, N., Kaczmarek, E.B., Gray, S., Barrell, R.A. und Oppenheim, B.A. (1996)
Glycopeptide-resistant enterococci isolated from uncooked meat.
J. Antimicrob. Chemother. 38, 908-909
- Chen, D.K., Pearce, L., McGeer, A., Low, D.E. und Willey, B.M. (2000)
Evaluation of D-xylose and 1% methyl- α -D-glucopyranoside fermentation tests for distinguishing *Enterococcus gallinarum* from *Enterococcus faecium*.
J. Clin. Microbiol. 38, 3652-3655
- Chenoweth, C. und Schaberg, D. (1990)
The epidemiology of enterococci.
Eur. J. Clin. Microbiol. Infect. Dis. 9, 80-89
- Chenoweth, C. E., Robinson, K. A. und Schaberg, D. R. (1990)
Efficacy of ampicillin versus trimethoprim-sulfamethoxazole in a mouse model of lethal enterococcal peritonitis.
Antimicrob. Agents Chemother. 34, 1800-1802
- Chow, J. W., Thal, L. A., Perry, M. B., Vazquez, J. A., Donabedian, S. M., Clewell, D. B. und Zervos, M. J. (1993)
Plasmid-associated hemolysin and aggregation substance production contribute to virulence in experimental enterococcal endocarditis.
Antimicrob. Agents Chemother. 37, 2474-2477
- Cintas, L.M., Casaus, P., Havarstein, L.S., Hernandez, P.E. und Nes, I.F. (1997)
Biochemical and genetic characterization of Enterocin P, a novel sec-dependent bacteriocin from *Enterococcus faecium* P13 with a broad antimicrobial spectrum.
Appl. Environ. Microbiol. 63, 4321-4330

- Clark, N. C., Teixeira, L. M., Facklam, R. R. und Tenover, F. C. (1998)
Detection and differentiation of *vanC-1*, *vanC-2*, and *vanC-3* glycopeptide resistance genes in enterococci.
J. Clin. Microbiol. 36, 2294-2297
- Collins, J. E., Bergeland, M. E., Lindeman, C. J. und Duimstra, J. R. (1988)
Enterococcus (Streptococcus) durans adherence in the intestine of a diarrheic pup.
Vet. Path. 25, 396-398
- Collins, M. D., Facklam, R. R., Farrow, J. A. E. und Williamson, R. (1989)
Enterococcus raffinosus sp. nov., *Enterococcus solitarius* sp. nov. and *Enterococcus pseudoavium* sp. nov.
FEMS Microbiol. Lett. 57, 283 - 288
- Collins, M. D., Rodrigues, U. M., Pigott, N. E. und Facklam, R. R. (1991)
Enterococcus dispar sp. nov., a new enterococcus species from human sources.
Lett. Appl. Microbiol. 12, 95-98
- Coque, T. M., Patterson, J. E., Steckelberg, J. M. und Murray, B. E. (1995)
Incidence of hemolysin, gelatinase and aggregation substance among enterococci isolated from patients with endocarditis and other infections and from the feces of hospitalized and community-based persons.
J. Infect. Dis. 171, 1223-1229
- Coque, T. M., Singh, K. V., Weinstock, G. M. und Murray, B. E. (1999)
Characterization of dihydrofolate reductase genes from trimethoprim-susceptible and trimethoprim-resistant strains of *Enterococcus faecalis*.
Antimicrob. Agents Chemother. 43, 141-147
- Coudron, P. E., Markowitz, S. M. und Wong, E. S. (1992)
Isolation of a β -lactamase-producing, aminoglycoside-resistant strain of *Enterococcus faecium*.
Antimicrob. Agents Chemother. 36, 1125-1126
- Crider, S. R. und Colby, S. D. (1985)
Susceptibility of enterococci to trimethoprim and trimethoprim-sulfamethoxazole.
Antimicrob. Agents Chemother. 27, 71-75

- DANMAP-Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (2000)
Consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, foods and humans in Denmark.
Eigenverlag, Kopenhagen, Dänemark
- DANMAP-Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (2001)
Use of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, foods and humans in Denmark.
Eigenverlag, Kopenhagen, Dänemark
- Dargere, S., Vergnaud, M., Verdon, R., Saloux, E., Le Page, O., Leclercq, R. und Bazin, C. (2002)
Enterococcus gallinarum endocarditis occurring on native heart valves.
J. Clin. Microbiol. 40, 2308-2310
- de Baere, T., Claeys, G., Verschraegen, G., Devriese, L. A., Baele, M., van Vlem, B., Vanholder, R., Dequidt, C. und Vaneechoutte, M. (2000)
Continuous ambulatory peritoneal dialysis peritonitis due to *Enterococcus cecorum*.
J. Clin. Microbiol. 38, 3511-3512
- de Vaux, A., Laguerre, G., Divies, C. und Prevost, H. (1998)
Enterococcus asini sp. nov. isolated from the caecum of donkeys (*Equus asinus*).
Int. J. Syst. Bacteriol. 48, 383-387
- Devriese, L. A., Dutta, G. N., Farrow, J. A. E., Van de Kerckhove, A. und Phillips, B. A. (1983)
Streptococcus cecorum, a new species from chickens.
Int. J. Syst. Bacteriol. 73, 421 - 425
- Devriese, L. A., van de Kerckhove, A., Kilpper-Bälz, R., Schleifer, K. H. (1987)
Characterization and identification of *Enterococcus* species isolated from the intestines of animals.
Int. J. Syst. Bacteriol. 37, 257-259
- Devriese, L. A. und Haesebrouck, F. (1991)
Enterococcus hirae in different animal species.
Vet. Record 129, 391-392

- Devriese, L. A., Hommez, J., Wijfels, R. und Haesebrouck, F. (1991a)
Composition of the enterococcal and streptococcal intestinal flora of poultry.
J. Appl. Bacteriol. 71, 46-50
- Devriese, L. A., Ceyskens, K. und Haesebrouck, F. (1991b)
Characteristics of *Enterococcus cecorum* strains from the intestines of different animal species.
Lett. Appl. Microbiol. 12, 137-139
- Devriese, L. A., Collins, M. D. und Wirth, R. (1992a)
The genus *Enterococcus*.
In: Balows, A., H.G. Trüper, M. Dworkin, W. Harder, K. H. Schleifer:
The prokaryotes: a handbook on the biology of bacteria: ecophysiology, isolation, identification, application. Springer, New York, 1465-1481
- Devriese, L. A., Cruz Colque, J. I., De Herdt, P. und Haesebrouck, F. (1992b)
Identification and composition of the tonsillar and anal enterococcal and streptococcal flora of dogs and cats.
J. Appl. Bacteriol. 73, 421-425
- Devriese, L. A., Cruz Colque, J. I., Haesebrouck, F., Desmidt, M., Uyttebroeck, E. und Ducatelle, R. (1992c)
Enterococcus hirae septicaemia of psittacine birds.
Vet. Record 130, 558-559
- Devriese, L. A., Laurier, L., De Herdt, P. und Haesebrouck, F. (1992d)
Enterococcal and streptococcal species isolated from faeces of calves, young cattle and dairy cows.
J. Appl. Bacteriol. 72, 29-31
- Devriese, L. A., Cruz Colque, J. I., De Herdt, P. und Haesebrouck, F. (1992e)
Identification and composition of the tonsillar and anal enterococcal and streptococcal flora of dogs and cats.
J. Appl. Bacteriol. 73, 421-425
- Devriese, L. A., Pot, B. und Collins, M.D. (1993)
Phenotypic identification of the genus *Enterococcus* and differentiation of phylogenetically distinct enterococcal species and species groups.
J. Appl. Bacteriol. 75, 399-408

- Devriese, L. A., Hommez, J., Pot, B. und Haesebrouck, F. (1994)
Identification and composition of the streptococcal and enterococcal flora of tonsils, intestines and faeces of pigs.
J. Appl. Bacteriol. 77, 31-36
- Devriese, L. A. und Pot, B. (1995)
The genus *Enterococcus*.
In: Wood, B. J. B. und Holzapel, W. H. (Hrsg.):
The genera of lactic acid bacteria, Vol. 2.
Chapman & Hall, London, 327 - 367
- Devriese, L. A., Pot, B., Van Damme, L., Kersters, K., Haesebrouck, F. (1995)
Identification of *Enterococcus* species isolated from foods of animal origin.
Int. J. Food Microbiology 26, 187-197
- Devriese, L. A., Pot, B., Kersters, K., Lauwers, S. und Haesebrouck, F. (1996)
Acidification of methyl- α -D-glucopyranoside: a useful test to differentiate *Enterococcus casseliflavus* and *Enterococcus gallinarum* from *Enterococcus faecium* species group and from *Enterococcus faecalis*.
J. Clin. Microbiol. 34, 2607-2608
- Dupont, H., Montravers, P., Mohler, J. und Carbon, C. (1998)
Disparate findings on the role of virulence factors of *Enterococcus faecalis* in mouse and rat models of peritonitis.
Infect. Imm. 66, 2570 - 2575
- Dutka-Malen, S., Blaimont, B., Wauters, G., Courvalin, P. (1994)
Emergence of high-level resistance to glycopeptides in *Enterococcus gallinarum* and *Enterococcus casseliflavus*.
Antimicrob. Agents Chemother. 38, 1675-1677
- Eaton, T. J. und Gasson, M. J. (2001)
Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates.
Appl. Environ. Microbiol. 67, 1628-1635
- El Amin, N., Jalal, S. und Wretling, B. (1999)
Alterations in GyrA and ParC associated with fluoroquinolone resistance in *Enterococcus faecium*.

- Antimicrob. Agents Chemother. 43, 947-949
- El-Samragy, Y.A., Fayed, E.O., Aly, A.A. und Hagrass, A.E.A. (1988)
 Properties of Labneh-like product manufactured using enterococcus starter cultures as novel dairy fermentation bacteria.
 J. Food Protect. 51, 386-390
- Ennahar, S., Asou, Y., Zendo, T., Sonomoto, K. und Ishizaki, A. (2001)
 Biochemical and genetic evidence for production of enterocins A and B by *Enterococcus faecium* WHE 81.
 Int. J. Food Microbiol. 70, 291-301
- Etheridge, M., Yolken, R. H. und Vonderfecht, S. L. (1988)
Enterococcus hirae implicated as a cause of diarrhea in suckling rats.
 J. Clin. Microbiol. 26, 1741-1744
- Facklam, R.R. und Collins, M. D. (1989)
 Identification of *Enterococcus* species isolated from human infections by a conventional test scheme.
 J. Clin. Microbiol. 27, 731-734
- Farrow, J.A.E., Kruze, J., Phillips, B.A., Bramley, A.J. und Collins, M.D. (1984)
 Taxonomic studies on *Streptococcus bovis* and *Streptococcus equinus*:
 Description of *Streptococcus alactolyticus* sp. nov. and *Streptococcus saccharolyticus* sp. nov.
 Syst. Appl. Microbiol. 5, 467-482
- Farrow, J.A.E. und Collins, M.D. (1985)
Enterococcus hirae, a new species that includes amino acid assay strain NCDO 1258 and strains causing growth depression in young chickens.
 Int. J. Syst. Bacteriol. 35, 73-75
- Fines, M., Perichon, B., Reynolds, P., Sahm, D.F. und Courvalin, P. (1999)
 VanE, a new type of acquired glycopeptide resistance in *Enterococcus faecalis* BM4405.
 Antimicrob. Agents Chemother. 43, 2161-2164
- Fontana, R., Canepari, P., Lleo, M. M. und Satta, G. (1990)
 Mechanisms of resistance of enterococci to beta-lactam antibiotics.
 Eur. J. Clin. Microbiol. Infect. Dis. 9, 103-105
- Fontana, R., Aldegheri, M., Ligozzi, M., Lopez, H., Sucari, A. und Satta, G. (1994)
 Overproduction of a low-affinity penicillin-binding protein and high-

- level ampicillin resistance in *Enterococcus faecium*.
Antimicrob. Agents Chemother. 38, 1980-1983
- Franz, C. M. A. P., Holzapfel, W. H. und Stiles, M. E. (1999)
Enterococci at the crossroads of food safety?
Int. J. Food Microbiol. 47, 1-24
- Franz, C. M. A. P., Muscholl-Silberhorn, A. B., Yousif, N. M. K., Vancanneyt, M.,
Swings, J. und Holzapfel, W. H. (2001)
Incidence of virulence factors and antibiotic resistance among
enterococci isolated from food.
Appl. Environ. Microbiol. 67, 4385-4389
- Freitas, A.C., Pais, C., Malcata, X. und Hogg, T.A. (1995)
Microbiological characterization of Picante da Beira Baixa cheese.
J. Food Protect. 59, 155-160
- Fuller, R., Coates, M. E. und Harrison, G. F. (1979)
The influence of specific bacteria and a filterable agent on the growth of
gnotobiotic chicks.
J. Appl. Bacteriol. 46, 335-342
- Gelsomino, R., Vancanneyt, M., Cogan, T.M., Condon, S. und Swings, J. (2002)
Source of enterococci in a farmhouse raw-milk cheese.
Appl. Environ. Microbiol. 68, 3560-3565
- Gevers, D. (2002)
Tetracycline resistance in lactic acid bacteria isolated from fermented dry
sausages.
Belg. Coord. Collect. Microorg. 12, 3-4
- Gilad, J., Borer, A., Riesenber, K., Peled, N., Shnaider, A. und Schlaeffer, F. (1998)
Enterococcus hirae septicemia in a patient with end-stage renal disease
undergoing hemodialysis.
Eur. J. Clin. Microbiol. Infect. Dis. 17, 576-577
- Giraffa, G., Carminati, D. und Tarelli, G.T. (1995)
Inhibition of *Listeria innocua* in milk by bacteriocin-producing *Enterococcus*
faecium 7C5.
J. Food Protect. 58, 621-623
- Giraffa, G. (2002)
Enterococci from foods.

- FEMS Microbiol. Rev. 26, 163-171
- Gray, J. W. und Pedler, S. J. (1992)
Antibiotic-resistant enterococci.
J. Hosp. Infect. 21, 1-14
- Grayson, M. L., Thauvin-Eliopoulos, C., Eliopoulos, G. M., Yao, J. D. C.,
DeAngelis, D. V., Walton, L., Woolley, J. L. und Moellering Jr., R. C.
(1990)
Failure of trimethoprim-sulfamethoxazole therapy in experimental
enterococcal endocarditis.
Antimicrob. Agents Chemother. 34, 1792-1794
- Grayson, M. L., Eliopoulos, G. M., Wennerstern, C., Ruoff, K., de Girolami, P.,
Fenaro, M.-J., Moellering Jr., R. C. (1991)
Increasing resistance to β -lactam antibiotics among clinical isolates of
Enterococcus faecium: a 22-year review at one institution.
Antimicrob. Agents Chemother. 35, 2180-2184
- Greub, G., Devriese, L.A., Pot, B., Dominguez, J. und Bille, J. (1997)
Enterococcus cecorum septicemia in a malnourished adult patient.
Eur. J. Clin. Microbiol. Infect. Dis. 16, 594 - 598
- Grohs, P., Gutmann, L., Legrand, R., Schoot, B. und Mainardi, J. L. (2000)
Vancomycin resistance is associated with serine-containing
peptidoglycan in *Enterococcus gallinarum*.
J. Bacteriol. 182, 6228-6232
- Haroche, J., Allignet, J., Aubert, S., van den Bogaard, A. E. und El Solh, N. (2000)
satG, conferring resistance to streptogramin A, is widely distributed in
Enterococcus faecium strains but not in staphylococci.
Antimicrob. Agents Chemother. 44, 190-191
- Healy, S. P. und Zervos, M. J. (1995)
Mechanisms of resistance of enterococci to antimicrobial agents.
Rev. Med. Microbiol. 6, 70-76
- Henwood, C.J., Livermore, D.M., Johnson, A.P., James, D., Warner, M. und
Gardiner, A. (2000)
Susceptibility of Gram-positive cocci from 25 UK hospitals to antimicrobial
agents including linezolid.
J. Antimicrob. Chemother. 46, 931-940

- Herman, D. J. und Gerding, D. N. (1991)
Antimicrobial resistance among enterococci.
Antimicrob. Agents Chemother. 35, 1-4
- Holzappel, W. und Franz, C. (2002)
Safety and health promotion aspects of enterococci.
EU-FAIR Project CT97-3078, BFE, Karlsruhe
- Houben, J.H. (1982)
Heat resistance of *Streptococcus faecium* in pasteurized ham.
Fleischwirtschaft 62, 490-493
- Hsueh, P.-R., Teng, L.-J., Chen, Y.-C., Yang, P.-C., Ho, S.-W. und Luh, K.-T. (2000)
Recurrent bacteremic peritonitis caused by *Enterococcus cecorum* in a patient with liver cirrhosis.
J. Clin. Microbiol. 38, 2450 - 2452
- Ike, Y., Hashimoto, H., Clewell, D. B. (1987)
High incidence of hemolysin production by *Enterococcus (Streptococcus) faecalis* strains associated with human parenteral infections.
J. Clin. Microbiol. 25, 1524-1528
- Ingham, S.C., Reyes, J. C. N., Schoeller, N.P. und Lang, M.M. (2000)
Potential use of presumptive enterococci and staphylococci as indicators of sanitary condition in plants making hard italian-type cheese.
J. Food Protect. 63, 1697-1701
- Johnson, A.P., Warner, M., Hallas, G. und Livermore, D.M. (2000)
Susceptibility to quinupristin/dalfopristin and other antibiotics of vancomycin-resistant enterococci from the UK, 1997 to mid-1999.
J. Antimicrob. Chemother. 46, 125-128
- Jones, R. N., Marshall, S. A., Pfaller, M. A., Wilke, W. W., Hollis, R. J., Meridith, E.E., Edmond, M. B. und Wenzel, R.P. (1997)
Nosocomial enterococcal blood stream infections in the SCOPE program: antimicrobial resistance, species occurrence, molecular testing results, and laboratory testing accuracy.
Diagn. Microbiol. Infect. Dis. 29, 95-102
- Kamphues, J. (1999)
Leistungsförderer mit antibiotischer Wirkung aus der Sicht der Tierernährung.

- Berl. Münch. Tierärztl. Wschr. 112, 370-379
- Kanematsu, E., Deguchi, T., Yasuda, M., Kawamura, T., Nishino, Y. und Kawada, Y. (1998)
Alterations in the GyrA subunit of DNA gyrase and the ParC subunit of DNA topoisomerase IV associated with quinolone resistance in *Enterococcus faecalis*.
Antimicrob. Agents Chemother. 42, 433-435
- Kaufhold, A. und Ferrieri, P. (1991)
Isolation of *Enterococcus mundtii* from normally sterile body sites in two patients.
J. Clin. Microbiol. 29, 1075-1077
- Kaufhold, A. und Klein, R. (1995)
Species identification and antibiotic susceptibility of enterococci isolated from clinical specimens of hospitalized patients.
Zbl. Bakt. 282, 507-518
- Kawamoto, S., Shima, J., Sato, R., Eguchi, T., Ohmomo, S., Shibato, J., Horikoshi, N., Takeshita, K. und Sameshima, T. (2002)
Biochemical and genetic characterization of Mundticin KS, an antilisterial peptide produced by *Enterococcus mundtii* NFRI 7393.
Appl. Environ. Microbiol. 68, 3830-3840
- Kielwein, G. (1978)
Vorkommen und Bedeutung von Enterokokken in Milch und Milchprodukten.
Arch. Lebensmittelhyg. 29, 127-128
- Klare, I., Heier, H., Claus, H., Böhme, G., Marin, S., Seltmann, G., Hakenbeck, R., Antanassova, V. und Witte, W. (1995)
Enterococcus faecium strains with *vanA*-mediated high-level glycopeptide resistance isolated from animal foodstuffs and fecal samples of humans in the community.
Microb. Drug Resist. 1, 265-272
- Klare, I. und Witte, W. (1997a)
Glykopeptidresistente Enterokokken.
Wien. Klin. Wochenschr. 109, 293-300

- Klare, I. und Witte, W. (1997b)
Glycopeptidresistente Enterokokken: zur Situation in Deutschland.
Hyg. Mikrob. 2, 31-38
- Klare, I. und Reissbrodt, R. (1998)
Glycopeptide-resistant enterococci (GRE).
Biotest Bull. 6, 59-64
- Klare, I. (2001)
unveröffentlichte Mitteilung.
- Klein, G., Pack, A. und Reuter, G. (1998)
Antibiotic resistance patterns of enterococci and occurrence of
vancomycin-resistant enterococci in raw minced beef and pork in
Germany.
Appl. Environ. Microbiol. 64, 1825-1830
- Knudtson, L. M. und Hartman, P. A. (1993)
Antibiotic resistance among enterococcal isolates from environmental
and clinical sources.
J. Food Protect. 56, 489-492
- Kommission der Europäischen Gemeinschaften (2001)
Über eine Strategie der Gemeinschaft zur Bekämpfung der Resistenz
gegen antimikrobielle Mittel.
Mitteilung der Kommission vom 20.06.2001, Brüssel, Belgien
- Kresken, M., Hafner, D., von Rosenstiel, N. (1999)
Zeitliche Entwicklung der Antibiotikaresistenz bei klinisch wichtigen
Bakterienspezies in Mitteleuropa.
Bundesgesundheitsbl.-Gesundheitsforsch.-Gesundheitsschutz 42, 17-25
- Kroker, R. (1991)
Pharmaka zur Behandlung und Verhütung bakterieller Infektionen.
In: Löscher, W., Ungemach, F. R. und Kroker, R.:
Grundlagen der Pharmakotherapie bei Haus- und Nutztieren.
Berlin und Hamburg, 202-237
- Kurup, A., Tee, W. S. N., Loo, L. H. und Lin, R. (2001)
Infection of central nervous system by motile *Enterococcus*: first case
report.
J. Clin. Microbiol. 39, 820-822

- Kühnen, E., Richter, F., Richter, K. und Andries, L. (1988)
Establishment of a typing system for group D streptococci.
Zentralbl. Bakteriol. Hyg. A. 267, 322-330
- Laukova, A. und Marekova, M. (2002)
A bacteriocin-mediated antagonism by *Enterococcus faecium* EK13
against *Listeria innocua* in tofu.
Arch. Lebensmittelhyg. 53, 25-48
- Leclerc, H., Devriese, L.A. und Mossel, D.A.A. (1996)
Taxonomical changes in intestinal (faecal) enterococci and streptococci:
Consequences on their use as indicators of faecal contamination in
drinking water.
J. Appl. Bacteriol. 81, 459-466
- Leclercq, R. und Courvalin, P. (1991)
Bacterial resistance to macrolide, lincosamide, and streptogramin
antibiotics by target modification.
Antimicrob. Agents Chemother. 35, 1267-1272
- Lemcke, R. und Bülte, M. (1999)
Vorkommen von *vanA*-, *vanB*-, *vanC1*-, *vanC2*- und *vanC3*-Genen in
Enterokokken aus Geflügel- und Schweinefleisch.
In: Proceedingband zur 40. Arbeitstagung des Arbeitsgebietes
Lebensmittelhygiene der DVG
Verlag der DVG, Gießen, 389-394
- Lemcke, R. und Bülte, M. (2000)
Occurrence of the vancomycin-resistant genes *vanA*, *vanB*, *vanC1*, *vanC2*
and *vanC3* in *enterococcus* strains isolated from poultry and pork.
Int. J. Food Microbiol. 60, 185-194
- Livermore, D. M. (2000)
Quinupristin/Dalfopristin and linezolid: where, when, which and whether to
use?
J. Antimicrob. Chemother. 46, 347-350
- Lyon, W.J., Olson, D. G. und Murano, E.A. (1995)
Isolation and purification of enterocin EL1, a bacteriocin produced by a
strain of *Enterococcus faecium*.
J. Food Protect. 58, 890-898

- Mac, K., Wichmann-Schauer, H., Peters, J. und Ellerbroek, L. (2002)
 Nachweis von Vancomycin-Resistenzgenen bei Enterokokken-
 Feldstämmen tierischer Herkunft mittels Multiplex-PCR.
 Arch. Lebensmittelhyg. 53, 66-67
- Macedo, A.C., Malcata, F.X. und Hogg, T.A. (1995)
 Microbiological profile in Serra ewes` cheese during ripening.
 J. Appl. Bacteriol. 79, 1-11
- Manero, A. und Blanch, A. R. (1999)
 Identification of *Enterococcus* spp. with a biochemical key.
 Appl. Environ. Microbiol. 65, 4425-4430
- Mann, P. A., Xiong, L., Mankin, A. S., Chau, A. S., Mendrick, C. A., Najarian, D. J.,
 Cramer, C. A., Loebenberg, D., Coates, E., Murgolo, N. J., Aarestrup, F.
 M., Goering, R. V., Black, T. A., Hare, R. S. und McNicholas, P. M. (2001)
 EmtA, a rRNA methyltransferase conferring high-level evernimicin
 resistance.
 Mol. Microbiol. 41, 1349-1356
- Martinez-Murcia, A. J. und Collins, M. D. (1991)
Enterococcus sulfureus, a new yellow-pigmented *Enterococcus* species.
 FEMS Microbiol. Lett. 80, 69-74
- McAlister, T., George, N., Faoagali, J. und Bell, J. (1999)
 Isolation of β -lactamase positive vancomycin resistant *Enterococcus*
faecalis; first case in Australia.
 Communic. Dis. Intellig. 23, 237-239
- McKessar, S. J., Berry, A. M., Bell, J. M., Turnidge, J. D. und Paton, J. C. (2000)
 Genetic characterization of *van G*, a novel vancomycin resistance locus
 of *Enterococcus faecalis*.
 Antimicrob. Agents Chemother. 44, 3224-3228
- McNicholas, P. M., Najarian, D. J., Mann, P. A., Hesk, D., Hare, R. S., Shaw, K. J.
 und Black, T. A. (2000)
 Evernimicin binds exclusively to the 50S ribosomal subunit and inhibits
 translation in cell-free systems derived from both Gram-positive and
 Gram-negative bacteria.
 Antimicrob. Agents Chemother. 44, 1121-1126

- Moellering, R. C., Jr., und Krogstadt, D. J. (1979)
 Antibiotic resistance in enterococci.
 In: Schlessinger, D. (Hrsg.): Microbiology-1979. American Society for
 Microbiology, Washington, D. C., 293-298
- Murray, B. E. (1990)
 The life and times of the *Enterococcus*.
 Clin. Microbiol. Rev. 3, 46-65
- Murray, B. E. (1992)
 β -lactamase-producing enterococci.
 Antimicrob. Agents Chemother. 36, 2355-2359
- Murray, I. A. und Shaw, W. V. (1997)
 O-acetyltransferases for chloramphenicol and other natural products.
 Antimicrob. Agents Chemother. 41, 1-6
- Mutschler, E. (1996)
 Antiinfektiva.
 In: Mutschler, E. und Schäfer-Körting, M.: Arzneimittelwirkungen.
 Wiss. Verl.-Ges. Stuttgart, 651-737
- Najjar, A. und Murray, B. E. (1987)
 Failure to demonstrate a consistent in vitro bactericidal effect of
 trimethoprim-sulfamethoxazole against enterococci.
 Antimicrob. Agents Chemother. 31, 808-810
- Nakanishi, N., Yoshida, S., Wakebe, H., Inoue, M. und Mitsuhashi, S. (1991)
 Mechanisms of clinical resistance to fluoroquinolones in *Enterococcus*
faecalis.
 Antimicrob. Agents Chemother. 35, 1053-1059
- NCCLS-National Committee for Clinical Laboratory Standards (1999)
 Performance standards for antimicrobial disk and dilution susceptibility
 tests for bacteria isolated from animals; Approved Standard.
 NCCLS-Document M31-A, Wayne, Pennsylvania, USA.
- NCCLS-National Committee for Clinical Laboratory Standards (2000)
 Methods for dilution antimicrobial susceptibility tests for bacteria that grow
 aerobically; Approved Standard – Fifth Edition. NCCLS-Document M7-A5,
 Wayne, Pennsylvania, USA.

- NCCLS-National Committee for Clinical Laboratory Standards (2001)
Performance standards for antimicrobial susceptibility testing; eleventh informational supplement. NCCLS-Document M100-S11, Wayne, Pennsylvania, USA.
- Niemi, R.M., Niemelä, S.I., Bamfort, D.H., Hantula, J., Hyvärinen, T., Forsten, T. und Raateland, A. (1993)
Presumptive faecal streptococci in environmental samples characterized by one-dimensional sodium dodecyl sulphate-polyacrylamide gel electrophoresis.
Appl. Environ. Microbiol. 59, 2190-2196
- Nowlan, S. P., und Deibel, R. H. (1967)
Group Q streptococci. I. Ecology, serology, physiology and relationship to established enterococci.
J. Bacteriol. 94, 291-296
- Noble, C.J. (1978)
Carriage of group D streptococci in the human bowel.
J. Clin. Pathol. 4, 1182-1186
- Onodera, Y., Okuda, J., Tanaka, M. und Sato, K. (2002)
Inhibitory activities of quinolones against DNA gyrase and topoisomerase IV of *Enterococcus faecalis*.
Antimicrob. Agents Chemother. 46, 1800-1804
- Ordenez, J.A., Barneto, R. und Ramos, M. (1978)
Studies on Manchego cheese ripened in olive oil.
Milchwiss. 33, 609-613
- Pangon, B., Pina, P., Gazagne, L., Chauvel, M., Godard, T., Courvalin, P. und Allouch, P. (1999)
Antimicrobial susceptibility of 471 enterococcal strains responsible for infections at 101 French hospitals.
Eur. J. Clin. Microbiol. Infect. Dis. 18, 836-838
- Parente, E. und Hill, C. (1992a)
Characterization of Enterocin 1146, a bacteriocin from *Enterococcus faecium* inhibitory to *Listeria monocytogenes*.
J. Food Protect. 55, 497-502

- Parente, E. und Hill, C. (1992b)
Inhibition of *Listeria* in buffer, broth, and milk by Enterocin 1146, a bacteriocin produced by *Enterococcus faecium*.
J. Food Protect. 55, 503-508
- Patel, R., Piper, K., Cockerill III, F.R., Steckelberg, J.M. und Yousten, A.A. (2000)
The biopesticide *Paenibacillus popilliae* has a vancomycin resistance gene cluster homologous to the enterococcal *vanA* vancomycin resistance gene cluster.
Antimicrob. Agents Chemother. 44, 705-709
- Patterson, J. E. und Zervos, M. J. (1989)
Susceptibility and bactericidal activity studies of four β -lactamase-producing enterococci.
Antimicrob. Agents Chemother. 33, 251-253
- Perichon, B., Reynolds, P., Courvalin, P. (1997)
VanD-type glycopeptide-resistant *Enterococcus faecium* BM4339.
Antimicrob. Agents Chemother. 41, 2016-2018
- Pette, J. W. (1955)
De vorming van zwavelwaterstof in Goudse kaas, veroorzaakt door melkzuurbacterien.
Netherlands Milk Dairy J. 10, 291-302
- Phillips, I. (1999)
The use of bacitracin as a growth promoter in animals produces no risk to human health.
J. Antimicrob. Chemother. 44, 725-728
- Pitfall, A. (1984)
In vivo v *in vitro* susceptibility of *Enterococcus* to trimethoprim-sulfamethoxazole.
JAMA 252, 2748-2749
- Poyart, C., Lambert, T., Morand, P., Abassade, P., Quesne, G., Baudouy, Y. und Trieu-Cuot, P. (2002)
Native valve endocarditis due to *Enterococcus hirae*.
J. Clin. Microbiol. 40, 2689-2690
- Rende-Fournier, R., Leclercq, R., Galimand, M., Duval, J. und Courvalin, P. (1993)
Identification of the *satA* gene encoding a streptogramin A

- acetyltransferase in *Enterococcus faecium* BM4145.
Antimicrob. Agents Chemother. 37, 2119-2125
- Rice, L. B., Carias, L. L., Hutton-Thomas, R., Sifaoui, F., Gutmann, L. und Rudin, S. D. (2001)
Penicillin-binding protein 5 and expression of ampicillin resistance in *Enterococcus faecium*.
Antimicrob. Agents Chemother. 45, 1480-1486
- Rice, L. B., Carias, L., Rudin, S., Vael, C., Goossens, H., Konstabel, C., Klare, I., Nallapareddy, S. R., Huang, W. und Murray, B. E. (2003)
A potential virulence gene, *hyl*_{Efm}, predominates in *Enterococcus faecium* of clinical origin.
J. Infect. Dis. 187, 508-512
- Roberts, M. C. (1996)
Tetracycline resistance determinants: mechanisms of action, regulation of expression, genetic mobility, and distribution.
FEMS Microbiol. Rev. 19, 1-24
- Roberts, M. C., Sutcliffe, J., Courvalin, P., Jensen, L. B., Rood, J. und Seppala, H. (1999)
Nomenclature for macrolide and macrolide-lincosamide-streptogramin B resistance determinants.
Antimicrob. Agents Chemother. 43, 2823-2830
- Rodrigues, U. und Collins, M.D. (1990)
Phylogenetic analysis of *Streptococcus saccharolyticus* based on 16S rRNA sequencing.
FEMS Microbiol. Lett. 71, 231-234
- Ruoff, K. L., de la Maza, L., Murtagh, M.J., Spargo, J.D., Ferraro, M. J. (1990)
Species identities of enterococci isolated from clinical specimens.
J. Clin. Microbiol. 28, 435 - 437
- Rybkin, T., Mainardi, J.-L., Sougakoff, W., Collatz, E. und Gutmann, L. (1998)
Penicillin-binding protein 5 sequence alterations in clinical isolates of *Enterococcus faecium* with different levels of β -lactam resistance.
J. Infect. Dis. 178, 159-163
- Sabia, C., Manicardi, G., Messi, P., de Niederhäusern, S. und Bondi, M. (2002)
Enterocin 416K1, an antilisterial bacteriocin produced by *Enterococcus*

- casseliflavus* IM 416K1 isolated from Italian sausages.
Int. J. Food Microbiology 75, 163-170
- Sandoe, J. A. T. , Witherden, I. R. und Settle, C. (2001)
Vertebral osteomyelitis caused by *Enterococcus raffinosus*.
J. Clin. Microbiol. 39, 1678-1679
- Sarantinopoulos, P., Kalantzopoulos, G. und Tsakalidou, E. (2002)
Effect of *Enterococcus faecium* on microbiological, physicochemical and sensory characteristics of Greek Feta cheese.
Int. J. Food Microbiology 76, 93-105
- Schmitz, F.-J., Krey, A., Sadurski, R., Verhoef, J., Milatovic, D. und Fluit, A.C. (2001)
Resistance to tetracycline and distribution of tetracycline resistance genes in European *Staphylococcus aureus* isolates.
J. Antimicrob. Chemother. 47, 239-240
- Shephard, B. D. und Gilmore, M. S. (2002)
Antibiotic-resistant enterococci: the mechanisms and dynamics of drug introduction and resistance.
Microb. Infect. 4, 215-224
- Sherman, J. M. (1937)
The streptococci.
Bacteriol. Rev. 1, 3-97
- Singh, K. V., Qin, X., Weinstock, G.M. und Murray, B. E. (1998)
Generation and testing of mutants of *Enterococcus faecalis* in a mouse peritonitis model.
J. Infect. Dis. 178, 1416-1420
- Singh, K. V., Weinstock, G. M. und Murray, B. E. (2002)
An *Enterococcus faecalis* ABC homologue (Lsa) is required for the resistance of this species to clindamycin and quinupristin-dalfopristin.
Antimicrob. Agents Chemother. 46, 1845-1850
- Stock, I. und Wiedemann, B. (1998)
Die Bestimmung der natürlichen Antibiotika-Empfindlichkeit.
Chemotherapie Journal 7, 127-135
- Svec, P., Devriese, L. A., Sedlacek, I., Baele, M., Vancanneyt, M., Haesebrouck, F., Swings, J. und Doskar, J. (2001)
Enterococcus haemoperoxidus sp. nov. and *Enterococcus moraviensis* sp.

- nov., isolated from water.
 Int. J. Syst. Evol. Microbiol. 51, 1567-1574
- Teixeira, L. M., Carvalho, M. D. G. S., Merquior, V. L. C., Steigerwalt, A. G., Teixeira, M. D. G. M., Brenner, D. J. und Facklam, R. R. (1997)
 Recent approaches on the taxonomy of the enterococci and some related microorganisms.
 Adv. Exp. Med. Biol. 418, 397-400
- Teixeira, L.M., Carvalho, M. G. S., Espinola, M. M. B., Steigerwalt, A. G., Douglas, M. P., Brenner, D.J. und Facklam, R.R. (2001)
Enterococcus porcinus sp. nov. and *Enterococcus ratti* sp. nov., associated with enteric disorders in animals.
 Int. J. Syst. Evol. Microbiol. 51, 1737-1743
- Teuber, M., Meile, L. und Schwarz, F. (1999)
 Acquired antibiotic resistance in lactic acid bacteria from food.
 Antonie van Leeuwenhoek 76, 115-137
- Thal, L.A., Chow, J.W., Mahayni, R., Bonilla, H., Perri, M.B., Donabedian, S.A., Silverman, J., Taber, S. und Zervos, M.J. (1995)
 Characterization of antimicrobial resistance in enterococci of animal origin.
 Antimicrob. Agents Chemother. 39, 2112-2115
- Theuretzbacher, U. (1998)
 Beta-Lactamasen und Beta-Lactamase-Inhibitoren.
 Chemotherapie Journal 7, 136-142
- Trovatelli, L. D. und Schiesser, A. (1987)
 Identification and significance of enterococci in hard cheese made from raw cow and sheep milk.
 Milchwiss. 42, 717-719
- Tyrrell, G. J., Turnbull, L., Teixeira, L. M., Lefebvre, J., Carvalho, M. G. S., Facklam, R. R. und Lovgren, M. (2002)
E. gilvus sp.nov. and *Enterococcus pallens* sp. nov. isolated from human clinical specimens.
 J. Clin. Microbiol. 40, 1140-1145
- Tzipori, S., Hayes, J., Sims, L. und Witters, M. (1984)
Streptococcus durans: an unexpected enteropathogen of foals.
 J. Infect. Dis. 150, 589-593

- Uttley, A.H.C., Collins, C.H., Naidoo, J. und George, R.C. (1988)
Vancomycin-resistant enterococci.
Lancet 335, 57-58
- van den Bogaard, A. E., Hazen, M., Hoyer, M., Oostenbach, P. und Stobberingh, E. E. (2002)
Effects of flavophospholipol on resistance in fecal *Escherichia coli* and enterococci of fattening pigs.
Antimicrob. Agents Chemother. 46, 110-118
- Vancanneyt, M., Snauwaert, C., Cleenwerck, I., Baele, M., Descheemaeker, P., Goossens, H., Pot, B., Vandamme, P., Swings, J., Haesebrouck, F. und Devriese, L. A. (2001)
Enterococcus villorum sp. nov., an enteroadherent bacterium associated with diarrhoea in piglets.
Int. J. Syst. Evol. Microbiol. 51, 393-400
- Villani, F., Salzano, G., Sorrentino, E., Pepe, O., Marino, P. und Coppola, S. (1993)
Enterocin 226NWC, a bacteriocin produced by *Enterococcus faecalis* 226, active against *Listeria monocytogenes*.
J. Appl. Bacteriol. 74, 380-387
- Vlaemynck, G., Herman, L. und Coudijzer, K. (1994)
Isolation and characterization of two bacteriocins produced by *Enterococcus faecium* strains inhibitory to *Listeria monocytogenes*.
Int. J. Food Microbiology 24, 211-225
- Wallrauch, C., Elsner, E., Milatovic, D., Cremer, J. und Braveny, I. (1997)
Antibiotikaresistenz der Enterokokken in Deutschland.
Med. Klin. 92, 464-468
- Watanakunakorn, C. (1990)
Enterococci from blood cultures during 1980-1989: susceptibility to ampicillin, penicillin and vancomycin.
J. Antimicrob. Chemother. 26, 602-604
- Watanakunakorn, C. (1993)
Increasing prevalence of resistance to ampicillin, penicillin and vancomycin of enterococci isolated from blood cultures during 1990-1991.
J. Antimicrob. Chemother. 31, 325-326

- Wegener, H.C., Madsen, M., Nielsen, N. und Aarestrup, F.M. (1997)
Isolation of vancomycin resistant *Enterococcus faecium* from food.
Int. J. Food Microbiol. 35, 57-66
- Wegener, H.C. (1998)
Historical yearly usage of glycopeptides for animals and humans: The
american-european paradox revisited.
Antimicrob. Agents Chemother. 42, 3049
- Weisblum, B. (1995)
Erythromycin resistance by ribosome modification.
Antimicrob. Agents Chemother. 39, 577-585
- Werner, G., Klare, I. und Witte, W. (1998)
Association between quinupristin/dalfopristin resistance in glycopeptide-
resistant *Enterococcus faecium* and the use of additives in animal feed.
Eur. J. Clin. Microbiol. Infect. Dis. 17, 401-402
- Werner, G. und Witte, W. (1999)
Characterization of a new enterococcal gene, *satG*, encoding a putative
acetyltransferase conferring resistance to streptogramin A compounds.
Antimicrob. Agents Chemother. 43, 1813-1814
- Werner, G., Klare, I., Heier, H., Hinz, K.-H., Böhme, G., Wendt, M. und Witte, W.
(2000)
Quinupristin/Dalfopristin-resistant enterococci of the *satA* (*vatD*) and *satG*
(*vatE*) genotypes from different ecological origins in Germany.
Microb. Drug Resist. 6, 37-47
- Wessels, D., Jooste, P. J. und Mostert, J. F. (1990)
Technologically important characteristics of *Enterococcus* isolates from
milk and dairy products.
Int. J. Food Microbiol. 10, 349-352
- Williams, A. M., Farrow, J. A. E. und Collins, M. D. (1989)
Reverse transcriptase sequencing of 16S ribosomal RNA from
Streptococcus cecorum.
Lett. Appl. Microbiol. 8, 185 - 189
- Williams, A. M., Rodrigues, J. A. E. und Collins, M.D. (1991)
Intragenetic relationship of enterococci as determined by reverse
transcriptase sequencing of small – subunit rRNA.

Res. Microb. 142, 67-74

Williamson, R., LeBouguenec, C., Gutmann, L. und Horaud, T. (1985)

One or two low affinity penicillin-binding proteins may be responsible for the range of susceptibility of *Enterococcus faecium* to benzylpenicillin.

J. Gen. Microbiol. 131, 1933-1940

Zervos, M. J. und Schaberg, D. R. (1985)

Reversal of the in vitro susceptibility of enterococci to trimethoprim-sulfamethoxazole by folinic acid.

Antimicrob. Agents Chemother. 28, 446-448

Zorzi, W., Zhou, X. Y., Dardenne, O., Lamotte, J., Raze, D., Pierre, J., Gutmann, L. und Coyette, J. (1996)

Structure of the low-affinity penicillin-binding protein5 PBP5fm in wild-type and highly penicillin-resistant strains of *Enterococcus faecium*.

J. Bacteriol. 178, 4948-4957