

5 REFERENCES

REFERENCES

- Adachi M, Furuichi Y and Miyamoto C (1994a) Identification of specific regions of the human endothelin-B receptor required for high affinity binding with endothelin-3. *Biochim Biophys Acta* 1223(2): 202-208.
- Adachi M, Furuichi Y and Miyamoto C (1994b) Identification of a ligand-binding site of the human endothelin-A receptor and specific regions required for ligand selectivity. *Eur J Biochem* 220(1): 37-43.
- Aimoto S, Hojoh H and Takasaki C (1990) Studies on the disulfide bridges of sarafotoxins. Chemical synthesis of sarafotoxin S6B and its homologue with different disulfide bridges. *Biochem Int* 21(6):1051-1057.
- Aktories K, Schultz G and Jakobs KH (1980) Regulation of adenylate cyclase activity in hamster adipocytes. Inhibition by prostaglandins, alpha-adrenergic agonists and nicotinic acid. *Naunyn Schmiedebergs Arch Pharmacol* 312(2): 167-173.
- Aktories K, Schultz G and Jakobs KH (1982) Inactivation of the guanine nucleotide regulatory site mediating inhibition of the adenylate cyclase in hamster adipocytes. *Naunyn Schmiedebergs Arch Pharmacol* 321(4): 247-252.
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215(3): 403-410.
- Andersen NH, Chen CP, Marschner TM, Krystek SR, Jr. and Bassolino DA (1992) Conformational isomerism of endothelin in acidic aqueous media: a quantitative NOESY analysis. *Biochemistry* 31(5): 1280-1295.
- Aquila E, Whelchel A, Knot HJ, Nelson M and Posada J (1996) Activation of multiple mitogen-activated protein kinase signal transduction pathways by the endothelin B receptor requires the cytoplasmic tail. *J Biol Chem* 271(49): 31572-31579.
- Archer E, Maigret B, Escrieut C, Pradayrol L and Fourmy D (2003) Rhodopsin crystal: new template yielding realistic models of G-protein-coupled receptors? *Trends Pharmacol Sci* 24(1): 36-40.
- Atkins AR, Martin RC and Smith R (1995) 1H NMR studies of sarafotoxin SRTb, a nonselective endothelin receptor agonist, and IRL 1620, an ETB receptor-specific agonist. *Biochemistry* 34(6): 2026-2033.
- Attwood TK and Findlay JB (1994) Fingerprinting G-protein-coupled receptors. *Protein Eng* 7(2): 195-203.
- Azpiroz I and Gautam N (2001) G protein gamma subunit interaction with a receptor regulates receptor-stimulated nucleotide exchange. *J Biol Chem* 276(45): 41742-41747.
- Ballesteros JA, Jensen AD, Liapakis G, Rasmussen SG, Shi L, Gether U and Javitch JA (2001a) Activation of the beta 2-adrenergic receptor involves disruption of an ionic lock between the cytoplasmic ends of transmembrane segments 3 and 6. *J Biol Chem* 276(31): 29171-29177.
- Ballesteros JA, Shi L and Javitch JA (2001b) Structural mimicry in G protein-coupled receptors: implications of the high-resolution structure of rhodopsin for structure-function analysis of rhodopsin-like receptors. *Mol Pharmacol* 60(1): 1-19.
- Balmforth AJ, Lee AJ, Warburton P, Donnelly D and Ball SG (1997) The conformational change responsible for AT1 receptor activation is dependent upon two juxtaposed asparagine residues on transmembrane helices III and VII. *J Biol Chem* 272(7): 4245-4251.
- Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN and Bourne PE (2000) The Protein Data Bank. *Nucleic Acids Res* 28(1): 235-242.
- Bhatnagar S and Rao GS (2000) Molecular modeling of the complex of endothelin-1 (ET-1) with the endothelin type A (ET(A)) receptor and the rational design of a peptide antagonist. *J Biomol Struct Dyn* 17(6): 957-964.
- Blahos J, 2nd, Mary S, Perroy J, de Colle C, Brabec I, Bockaert J and Pin JP (1998) Extreme C terminus of G protein alpha-subunits contains a site that discriminates between Gi-coupled metabotropic glutamate receptors. *J Biol Chem* 273(40): 25765-25769.
- Bockaert J and Pin JP (1999) Molecular tinkering of G protein-coupled receptors: an evolutionary success. *EMBO J* 18(7): 1723-1729.
- Boivin S, Tessier S, Aubin J, Lampron P, Dethieux M and Fournier A (2004) Identification of a binding domain of the endothelin-B receptor using a selective IRL-1620-derived photoprobe. *Biochemistry* 43(36): 11516-11525.

REFERENCES

- Bonini JA, Jones KA, Adham N, Forray C, Artymyshyn R, Durkin MM, Smith KE, Tamm JA, Boteju LW, Lakhlani PP, Raddatz R, Yao WJ, Ogozalek KL, Boyle N, Kouranova EV, Quan Y, Vaysse PJ, Wetzel JM, Branchek TA, Gerald C and Borowsky B (2000) Identification and characterization of two G protein-coupled receptors for neuropeptide FF. *J Biol Chem* 275(50): 39324-39331.
- Breitweg-Lehmann E, Czupalla C, Storm R, Kudlacek O, Schunack W, Freissmuth M and Nurnberg B (2002) Activation and inhibition of G proteins by lipoamines. *Mol Pharmacol* 61(3): 628-636.
- Breu V, Hashido K, Broger C, Miyamoto C, Furuichi Y, Hayes A, Kalina B, Loffler BM, Ramuz H and Clozel M (1995) Separable binding sites for the natural agonist endothelin-1 and the non-peptide antagonist bosentan on human endothelin-A receptors. *Eur J Biochem* 231(1): 266-270.
- Bueb JL, Mousli M, Bronner C, Rouot B and Landry Y (1990) Activation of Gi-like proteins, a receptor-independent effect of kinins in mast cells. *Mol Pharmacol* 38(6): 816-822.
- Cadwallader K, Beltman J, McCormick F and Cook S (1997) Differential regulation of extracellular signal-regulated protein kinase 1 and Jun N-terminal kinase 1 by Ca²⁺ and protein kinase C in endothelin-stimulated Rat-1 cells. *Biochem J* 321 (Pt 3): 795-804.
- Carlson LA and Oro L (1962) The effect of nicotinic acid on the plasma free fatty acid; demonstration of a metabolic type of sympathicolysis. *Acta Med Scand* 172: 641-645.
- Case DA, Pearlman DA, Caldwell JW, Cheatham TE, III, Wang J, Ross WS, Simmerling CL, Darden TA, Merz KM, Stanton RV, Cheng AL, Vincent JJ, Crowley M, Tsui V, Gohlke H, Radmer RJ, Duan Y, Pitera J, Massova I, Seibel GL, Singh UC, Weiner PK and Kollman PA (2002) AMBER7.
- Cheong CG, Escalante-Semerena JC and Rayment I (2001) Structural investigation of the biosynthesis of alternative lower ligands for cobamides by nicotinate mononucleotide: 5,6-dimethylbenzimidazole phosphoribosyltransferase from *Salmonella enterica*. *J Biol Chem* 276(40): 37612-37620.
- Clark M, Cramer RD, III and van Opdenbosch N (1989) Validation of the General Purpose Tripos 5.2 Force Field. *J Comp Chem* 10: 982-1012.
- Clozel M, Breu V, Gray GA, Kalina B, Loffler BM, Burri K, Cassal JM, Hirth G, Muller M and Neidhart W (1994) Pharmacological characterization of bosentan, a new potent orally active nonpeptide endothelin receptor antagonist. *J Pharmacol Exp Ther* 270(1): 228-235.
- Coleman DE, Berghuis AM, Lee E, Linder ME, Gilman AG and Sprang SR (1994) Structures of active conformations of Gi alpha 1 and the mechanism of GTP hydrolysis. *Science* 265(5177): 1405-1412.
- Cornell WD, Cieplak P, Baylay CI, Gould IR, Merz KM, Ferguson DM, Spellmeyer DC, Fox T, Caldwell JW and Kollman PA (1995) A second generation force field for the simulation of proteins and nucleic acids. *J Am Chem Soc* 117: 5179-5197.
- Davenport AP, Mille R, Pelton JT and Huggins J (1997) Distribution of endothelin receptors, in Endothelin in Biology and Medicine pp 45-68, CRC Press, Boca Raton, FL.
- Davenport AP (2002) International Union of Pharmacology. XXIX. Update on endothelin receptor nomenclature. *Pharmacol Rev* 54(2): 219-226.
- Davenport AP and Battistini B (2002) Classification of endothelin receptors and antagonists in clinical development. *Clin Sci (Lond)* 103 Suppl 48: 1S-3S.
- de Nucci G, Thomas R, D'Orleans-Juste P, Antunes E, Walder C, Warner TD and Vane JR (1988) Pressor effects of circulating endothelin are limited by its removal in the pulmonary circulation and by the release of prostacyclin and endothelium-derived relaxing factor. *Proc Natl Acad Sci U S A* 85(24): 9797-9800.
- Doherty AM, Cody WL, Depue PL, He JX, Waite LA, Leonard DM, Leitz NL, Dudley DT, Rapundalo ST, Hingorani GP, Haleen SJ, Ladouceur DM, Hill KE, Flynn MA and Reynolds EE (1993) Structure-Activity-Relationships of C-Terminal Endothelin Hexapeptide Antagonists. *J Med Chem* 36(18):2585-2594.
- Dohlman HG, Caron MG, DeBlasi A, Frielle T and Lefkowitz RJ (1990) Role of extracellular disulfide-bonded cysteines in the ligand binding function of the beta 2-adrenergic receptor. *Biochemistry* 29(9): 2335-2342.

- Eddy SR (1998) Profile hidden Markov models. *Bioinformatics* 14(9): 755-763.
- Ellis PJ, Appleby CA, Guss JM, Hunter WN, Ollis DL and Freeman HC (1997) Structure of ferric soybean leghemoglobin a nicotinate at 2.3 Å resolution. *Acta Crystallogr D Biol Crystallogr* 53(Pt 3): 302-310.
- Ergul A, Tackett RL and Puett D (1995) Identification of receptor binding and activation sites in endothelin-1 by use of site-directed mutagenesis. *Circ Res* 77(6): 1087-1094.
- Feng DF and Doolittle RF (1987) Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *J Mol Evol* 25(4): 351-360.
- Filmore D (2004) It's a GPCR world. *Modern Drug Discovery* 7(11): 24-28.
- Foord SM (2002) Receptor classification: post genome. *Curr Opin Pharmacol* 2(5): 561-566.
- Fotiadis D, Liang Y, Filipek S, Saperstein DA, Engel A and Palczewski K (2003) Atomic-force microscopy: Rhodopsin dimers in native disc membranes. *Nature* 421(6919): 127-128.
- Fraser CM (1989) Site-directed mutagenesis of beta-adrenergic receptors. Identification of conserved cysteine residues that independently affect ligand binding and receptor activation. *J Biol Chem* 264(16): 9266-9270.
- Fredriksson R, Lagerstrom MC, Lundin LG and Schioth HB (2003) The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol* 63(6): 1256-1272.
- Gasteiger J and Marsili M (1978) A New Model for Calculating Atomic Charges in Molecules. *Tetrahedron Lett* 34: 3181-3184.
- Gether U (2000) Uncovering molecular mechanisms involved in activation of G protein-coupled receptors. *Endocr Rev* 21(1): 90-113.
- Gilchrist A, Vanhauwe JF, Li A, Thomas TO, Voyno-Yasenetskaya T and Hamm HE (2001) G alpha minigenes expressing C-terminal peptides serve as specific inhibitors of thrombin-mediated endothelial activation. *J Biol Chem* 276(28): 25672-25679.
- Gohla A, Offermanns S, Wilkie TM and Schultz G (1999) Differential involvement of Galphai2 and Galphai3 in receptor-mediated stress fiber formation. *J Biol Chem* 274(25): 17901-17907.
- Grantcharova E, Ferkert J, Reusch HP, Krell HW, Papsdorf G, Beyermann M, Schulein R, Rosenthal W and Oksche A (2002) The extracellular N terminus of the endothelin B (ETB) receptor is cleaved by a metalloprotease in an agonist-dependent process. *J Biol Chem* 277(46): 43933-43941.
- Gregan B, Jurgensen J, Papsdorf G, Ferkert J, Schaefer M, Beyermann M, Rosenthal W and Oksche A (2004) Ligand-dependent differences in the internalization of endothelin A and endothelin B receptor heterodimers. *J Biol Chem* 279(26): 27679-27687.
- Grigorieff N, Ceska TA, Downing KH, Baldwin JM and Henderson R (1996) Electron-crystallographic refinement of the structure of bacteriorhodopsin. *J Mol Biol* 259(3): 393-421.
- Hamm HE, Deretic D, Arendt A, Hargrave PA, Koenig B and Hofmann KP (1988) Site of G protein binding to rhodopsin mapped with synthetic peptides from the alpha subunit. *Science* 241(4867): 832-835.
- Hamm HE (2001) How activated receptors couple to G proteins. *Proc Natl Acad Sci U S A* 98(9): 4819-4821.
- He W, Miao FJ, Lin DC, Schwandner RT, Wang Z, Gao J, Chen JL, Tian H and Ling L (2004) Citric acid cycle intermediates as ligands for orphan G-protein-coupled receptors. *Nature* 429(6988): 188-193.
- Hechter O and Calek A, Jr. (1974) Principles of hormone action: the problem of molecular linguistics. *Acta Endocrinol Suppl (Copenh)* 191: 39-66.
- Henikoff S and Henikoff JG (1992) Amino acid substitution matrices from protein blocks. *Proc Natl Acad Sci U S A* 89(22): 10915-10919.
- Hermans E (2003) Biochemical and pharmacological control of the multiplicity of coupling at G-protein-coupled receptors. *Pharmacol Ther* 99(1): 25-44.

- Higashijima T, Uzu S, Nakajima T and Ross EM (1988) Mastoparan, a peptide toxin from wasp venom, mimics receptors by activating GTP-binding regulatory proteins (G proteins). *J Biol Chem* 263(14): 6491-6494.
- Hilal-Dandan R, Merck DT, Lujan JP and Brunton LL (1994) Coupling of the type A endothelin receptor to multiple responses in adult rat cardiac myocytes. *Mol Pharmacol* 45(6): 1183-1190.
- Horstmeyer A, Cramer H, Sauer T, Muller-Esterl W and Schroeder C (1996) Palmitoylation of endothelin receptor A. Differential modulation of signal transduction activity by post-translational modification. *J Biol Chem* 271(34): 20811-20819.
- Itoh Y, Yanagisawa M, Ohkubo S, Kimura C, Kosaka T, Inoue A, Ishida N, Mitsui Y, Onda H and Fujino M (1988) Cloning and sequence analysis of cDNA encoding the precursor of a human endothelium-derived vasoconstrictor peptide, endothelin: identity of human and porcine endothelin. *FEBS Lett* 231(2): 440-444.
- Janes RW, Peapus DH and Wallace BA (1994) The crystal structure of human endothelin. *Nat Struct Biol* 1(5): 311-319.
- Janz JM and Farrens DL (2004) Rhodopsin activation exposes a key hydrophobic binding site for the transducin alpha-subunit C terminus. *J Biol Chem* 279(28): 29767-29773.
- Ji TH, Grossmann M and Ji I (1998) G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J Biol Chem* 273(28): 17299-17302.
- Jiang Q, Van Rhee AM, Kim J, Yehle S, Wess J and Jacobson KA (1996) Hydrophilic side chains in the third and seventh transmembrane helical domains of human A2A adenosine receptors are required for ligand recognition. *Mol Pharmacol* 50(3): 512-521.
- Jones LG (1996) Inhibition of cyclic AMP accumulation by endothelin is pertussis toxin sensitive and calcium independent in isolated adult feline cardiac myocytes. *Life Sci* 58(2): 115-123.
- Jones TA and Thirup S (1986) Using known substructures in protein model building and crystallography. *EMBO J* 5(4): 819-822.
- Karle IL and Balaram P (1990) Structural characteristics of alpha-helical peptide molecules containing Aib residues. *Biochemistry* 29(29): 6747-56
- Katahira R, Umemura I, Takai M, Oda K, Okada T and Nosaka AY (1998) Structural studies on endothelin receptor subtype B specific agonist IRL 1620 [suc-[Glu9, Ala11,15]ET-1(8-21)] and its analogs with dipalmitoyl phosphatidylcholine vesicles by NMR spectroscopy. *J Pept Res* 51(2): 155-164.
- Kelley LA, MacCallum RM and Sternberg MJ (2000) Enhanced genome annotation using structural profiles in the program 3D-PSSM. *J Mol Biol* 299(2): 499-520.
- Kim J, Jiang Q, Glashofer M, Yehle S, Wess J and Jacobson KA (1996) Glutamate residues in the second extracellular loop of the human A2a adenosine receptor are required for ligand recognition. *Mol Pharmacol* 49(4): 683-691.
- Kisselev OG, Kao J, Ponder JW, Fann YC, Gautam N and Marshall GR (1998) Light-activated rhodopsin induces structural binding motif in G protein alpha subunit. *Proc Natl Acad Sci U S A* 95(8): 4270-4275.
- Knopp RH (1999) Drug treatment of lipid disorders. *N Engl J Med* 341(7): 498-511.
- Kolakowski LF, Jr. (1994) GCRDb: a G-protein-coupled receptor database. *Receptors Channels* 2(1): 1-7.
- Kostenis E, Zeng FY and Wess J (1998) Functional characterization of a series of mutant G protein alpha_q subunits displaying promiscuous receptor coupling properties. *J Biol Chem* 273(28): 17886-17892.
- Kristiansen K (2004) Molecular mechanisms of ligand binding, signaling, and regulation within the superfamily of G-protein-coupled receptors: molecular modeling and mutagenesis approaches to receptor structure and function. *Pharmacol Ther* 103(1): 21-80.
- Krystek SR, Jr., Patel PS, Rose PM, Fisher SM, Kienzle BK, Lach DA, Liu EC, Lynch JS, Novotny J and Webb ML (1994) Mutation of peptide binding site in transmembrane region of a G protein-coupled receptor accounts for endothelin receptor subtype selectivity. *J Biol Chem* 269(17): 12383-12386.

REFERENCES

- Kusunoki H, Wakamatsu K, Sato K, Miyazawa T and Kohno T (1998) G protein-bound conformation of mastoparan-X: heteronuclear multidimensional transferred nuclear overhauser effect analysis of peptide uniformly enriched with 13C and 15N. *Biochemistry* 37(14): 4782-4790.
- Lambright DG, Sondek J, Bohm A, Skiba NP, Hamm HE and Sigler PB (1996) The 2.0 Å crystal structure of a heterotrimeric G protein. *Nature* 379(6563): 311-319.
- Laskowski RA, Rullmann JA, MacArthur MW, Kaptein R and Thornton JM (1996) AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. *J Biol NMR* 8(4): 477-486.
- Lee JA, Brinkmann JA, Longton ED, Peishoff CE, Lago MA, Leber JD, Cousins RD, Gao A, Stadel JM and Kumar CS (1994a) Lysine 182 of endothelin B receptor modulates agonist selectivity and antagonist affinity: evidence for the overlap of peptide and non-peptide ligand binding sites. *Biochemistry* 33(48): 14543-14549.
- Lee JA, Elliott JD, Sutiphong JA, Friesen WJ, Ohlstein EH, Stadel JM, Gleason JG and Peishoff CE (1994b) Tyr-129 is important to the peptide ligand affinity and selectivity of human endothelin type A receptor. *Proc Natl Acad Sci U S A* 91(15): 7164-7168.
- Leschke C, Storm R, Breitweg-Lehmann E, Exner T, Nurnberg B and Schunack W (1997) Alkyl-substituted amino acid amides and analogous di- and triamines: new non-peptide G protein activators. *J Med Chem* 40(19): 3130-3139.
- Liu B and Wu D (2003) The first inner loop of endothelin receptor type B is necessary for specific coupling to Galphai3. *J Biol Chem* 278(4): 2384-2387.
- Lovering AL, Hyde EI, Searle PF and White SA (2001) The structure of Escherichia coli nitroreductase complexed with nicotinic acid: three crystal forms at 1.7 Å, 1.8 Å and 2.4 Å resolution. *J Mol Biol* 309(1): 203-213.
- Luecke H, Schobert B, Lanyi JK, Spudich EN and Spudich JL (2001) Crystal structure of sensory rhodopsin II at 2.4 angstroms: insights into color tuning and transducer interaction. *Science* 293(5534): 1499-1503.
- Maguire JJ and Davenport AP (1995) ETA receptor-mediated constrictor responses to endothelin peptides in human blood vessels in vitro. *Br J Pharmacol* 115(1): 191-197.
- Menziani M, Cocchi M, Fanelli F, De Benedetti PG, Sanz F, Giraldo J and Manaut F (1995) Theoretical QSAR Analysis on Three Dimensional Models of the Complexes Between Peptide and Non-peptide Antagonists With the EtA and EtB Receptors, in *QSAR and Molecular Modelling: Concepts, Computational Tools and Biological Applications* pp 519-525, Prous Science Publishers.
- Miedlich SU, Gama L, Seuwen K, Wolf RM and Breitwieser GE (2004) Homology modeling of the transmembrane domain of the human calcium sensing receptor and localization of an allosteric binding site. *J Biol Chem* 279(8): 7254-7263.
- Morris AL, MacArthur MW, Hutchinson EG and Thornton JM (1992) Stereochemical quality of protein structure coordinates. *Proteins* 12(4): 345-364.
- Mousli M, Bueb JL, Bronner C, Rouot B and Landry Y (1990) G protein activation: a receptor-independent mode of action for cationic amphiphilic neuropeptides and venom peptides. *Trends Pharmacol Sci* 11(9): 358-362.
- Muller G (2000) Towards 3D structures of G protein-coupled receptors: a multidisciplinary approach. *Curr Med Chem* 7(9): 861-888.
- Muradov KG and Artemyev NO (2000) Coupling between the N- and C-terminal domains influences transducin-alpha intrinsic GDP/GTP exchange. *Biochemistry* 39(14): 3937-3942.
- Murzin AG, Breuer SE, Hubbard T and Chothia C (1995) SCOP: a structural classification of proteins database for the investigation of sequences and structures. *J Mol Biol* 247: 536-540.
- Nakajima K, Kubo S, Kumagaye S, Nishio H, Tsunemi M, Inui T, Kuroda H, Chino N, Watanabe TX and Kimura T (1989a) Structure-activity relationship of endothelin: importance of charged groups. *Biochem Biophys Res Commun* 163(1): 424-429.

REFERENCES

- Nakajima K, Kumagaye S, Nishio H, Kuroda H, Watanabe TX, Kobayashi Y, Tamaoki H, Kimura T and Sakakibara S (1989b) Synthesis of endothelin-1 analogues, endothelin-3, and sarafotoxin S6b: structure-activity relationships. *J Cardiovasc Pharmacol* 13 Suppl 5: S8-12.
- Natochin M, Muradov KG, McEntaffer RL and Artemyev NO (2000) Rhodopsin recognition by mutant G(s)alpha containing C-terminal residues of transducin. *J Biol Chem* 275(4): 2669-2675.
- Nurnberg B, Togel W, Krause G, Storm R, Breitweg-Lehmann E and Schunack W (1999) Non-peptide G-protein activators as promising tools in cell biology and potential drug leads. *Eur J Med Chem* 34: 5-30.
- Okamoto Y, Ninomiya H, Tanioka M, Sakamoto A, Miwa S and Masaki T (1997) Palmitoylation of human endothelinB. Its critical role in G protein coupling and a differential requirement for the cytoplasmic tail by G protein subtypes. *J Biol Chem* 272(34): 21589-21596.
- Oksche A, Boese G, Horstmeyer A, Furkert J, Beyermann M, Bienert M and Rosenthal W (2000) Late endosomal/lysosomal targeting and lack of recycling of the ligand-occupied endothelin B receptor. *Mol Pharmacol* 57(6): 1104-1113.
- Olsson AG (1994) Nicotinic acid and derivatives. Springer-Verlag, Heidelberg.
- Onrust R, Herzmark P, Chi P, Garcia PD, Lichtarge O, Kingsley C and Bourne HR (1997) Receptor and betagamma binding sites in the alpha subunit of the retinal G protein transducin. *Science* 275(5298): 381-384.
- Orry AJW and Wallace BA (2000) Modeling and docking the endothelin G-protein-coupled receptor. *Biophysical Journal* 79(6): 3083-3094.
- Palczewski K, Kumasaka T, Hori T, Behnke CA, Motoshima H, Fox BA, Le TI, Teller DC, Okada T, Stenkamp RE, Yamamoto M and Miyano M (2000) Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* 289(5480): 739-745.
- Patny A, Desai PV and Avery MA (2006) Homology modeling of G-protein-coupled receptors and implications in drug design. *Curr Med Chem* 13(14): 1667-1691.
- Pearson WR and Lipman DJ (1988) Improved tools for biological sequence comparison. *Proc Natl Acad Sci U S A* 85(8): 2444-2448.
- Pellegrini M, Mammi S, Peggion E and Mierke DF (1997) Threonine6-bradykinin: structural characterization in the presence of micelles by nuclear magnetic resonance and distance geometry. *J Med Chem* 40(1): 92-98.
- Pellegrini M and Mierke DF (1997) Threonine6-bradykinin: molecular dynamics simulations in a biphasic membrane mimetic. *J Med Chem* 40(1): 99-104.
- Perlman JH, Wang W, Nussenzveig DR and Gershengorn MC (1995) A disulfide bond between conserved extracellular cysteines in the thyrotropin-releasing hormone receptor is critical for binding. *J Biol Chem* 270(42): 24682-24685.
- Peter MG and Davenport AP (1996) Characterization of the endothelin receptor selective agonist, BQ3020 and antagonists BQ123, FR139317, BQ788, 50235, Ro462005 and bosentan in the heart. *Br J Pharmacol* 117(3):455-462.
- Randall MD, Douglas SA and Hiley CR (1989) Vascular activities of endothelin-1 and some alanyl substituted analogues in resistance beds of the rat. *Br J Pharmacol* 98(2):685-699.
- Reddy SG, Scapin G and Blanchard JS (1996) Interaction of pyridine nucleotide substrates with Escherichia coli dihydrodipicolinate reductase: thermodynamic and structural analysis of binary complexes. *Biochemistry* 35(41): 13294-13302.
- Rose PM, Krystek SR, Jr., Patel PS, Liu EC, Lynch JS, Lach DA, Fisher SM and Webb ML (1995) Aspartate mutation distinguishes ETA but not ETB receptor subtype-selective ligand binding while abolishing phospholipase C activation in both receptors. *FEBS Lett* 361(2-3): 243-249.
- Rovero P, Galloppini C, Laricchia-Robbio L, Mazzoni MR and Revoltella RP (1998) Structure-activity analysis of C-terminal endothelin analogues. *J Cardiovasc Pharmacol* 31 Suppl 1: S251-S254.

- Saeki T, Ihara M, Fukuroda T, Yamagiwa M and Yano M (1991) [Ala1,3,11,15]endothelin-1 analogs with ETB agonistic activity. *Biochem Biophys Res Commun* 179(1):286-292.
- Sakamoto A, Yanagisawa M, Sawamura T, Enoki T, Ohtani T, Sakurai T, Nakao K, Toyo-oka T and Masaki T (1993) Distinct subdomains of human endothelin receptors determine their selectivity to endothelinA-selective antagonist and endothelinB-selective agonists. *J Biol Chem* 268(12): 8547-8553.
- Sakmar TP (1998) Rhodopsin: a prototypical G protein-coupled receptor. *Prog Nucleic Acid Res Mol Biol* 59: 1-34.
- Sansom MS and Weinstein H (2000) Hinges, swivels and switches: the role of prolines in signalling via transmembrane alpha-helices. *Trends Pharmacol Sci* 21(11): 445-451.
- Savarese TM, Wang CD and Fraser CM (1992) Site-directed mutagenesis of the rat m1 muscarinic acetylcholine receptor. Role of conserved cysteines in receptor function. *J Biol Chem* 267(16): 11439-11448.
- Semple G, Skinner PJ, Cherrier MC, Webb PJ, Sage CR, Tamura SY, Chen R, Richman JG and Connolly DT (2006) 1-Alkyl-benzotriazole-5-carboxylic acids are highly selective agonists of the human orphan G-protein-coupled receptor GPR109b. *J Med Chem* 49(4): 1227-1230.
- Sgourakis NG, Bagos PG, Papasaikas PG and Hamodrakas SJ (2005) A method for the prediction of GPCRs coupling specificity to G-proteins using refined profile Hidden Markov Models. *BMC Bioinformatics* 6: 104.
- Shacham S, Topf M, Avisar N, Glaser F, Marantz Y, Bar-Haim S, Noiman S, Naor Z and Becker OM (2001) Modeling the 3D structure of GPCRs from sequence. *Med Res Rev* 21(5): 472-483.
- Shi L and Javitch JA (2002) The binding site of aminergic G protein-coupled receptors: the transmembrane segments and second extracellular loop. *Annu Rev Pharmacol Toxicol* 42: 437-467.
- Shi L and Javitch JA (2006) A role for information collection, management, and integration in structure-function studies of G-protein coupled receptors. *Curr Pharm Des* 12(14): 1771-1783.
- Singh R, Hurst DP, Barnett-Norris J, Lynch DL, Reggio PH and Guarneri F (2002) Activation of the cannabinoid CB1 receptor may involve a W6.48/F3.36 rotamer toggle switch. *J Pept Res* 60(6): 357-370.
- Soga T, Kamohara M, Takasaki J, Matsumoto S, Saito T, Ohishi T, Hiyama H, Matsuo A, Matsushime H and Furuchi K (2003) Molecular identification of nicotinic acid receptor. *Biochem Biophys Res Commun* 303(1): 364-369.
- Srinivasan N, Guruprasad K and Blundell TL (1996) Comparative Modelling of Proteins. IRL Press, Oxford, England.
- Stenkamp RE, Teller DC and Palczewski K (2002) Crystal structure of rhodopsin: a G-protein-coupled receptor. *ChemBioChem* 3(10): 963-967.
- Stitham J, Stojanovic A, Merenick BL, O'Hara KA and Hwa J (2003) The unique ligand-binding pocket for the human prostacyclin receptor. Site-directed mutagenesis and molecular modeling. *J Biol Chem* 278(6): 4250-4257.
- Strader CD, Candelore MR, Hill WS, Sigal IS and Dixon RA (1989) Identification of two serine residues involved in agonist activation of the beta-adrenergic receptor. *J Biol Chem* 264(23): 13572-13578.
- Strader CD, Fong TM, Tota MR, Underwood D and Dixon RA (1994) Structure and function of G protein-coupled receptors. *Annu Rev Biochem* 63: 101-132.
- Strader CD, Gaffney T, Sugg EE, Candelore MR, Keys R, Patchett AA and Dixon RA (1991) Allele-specific activation of genetically engineered receptors. *J Biol Chem* 266(1): 5-8.
- Sukumar M and Higashijima T (1992) G protein-bound conformation of mastoparan-X, a receptor-mimetic peptide. *J Biol Chem* 267(30): 21421-21424.
- Sukumar M, Ross EM and Higashijima T (1997) A Gs-selective analog of the receptor-mimetic peptide mastoparan binds to Gs alpha in a kinked helical conformation. *Biochemistry* 36(12): 3632-3639.

REFERENCES

- Sunahara RK, Tesmer JJ, Gilman AG and Sprang SR (1997) Crystal structure of the adenylyl cyclase activator Gsalpha. *Science* 278(5345): 1943-1947.
- Suryanarayana S, Daunt DA, von Zastrow M and Kobilka BK (1991) A point mutation in the seventh hydrophobic domain of the alpha 2 adrenergic receptor increases its affinity for a family of beta receptor antagonists. *J Biol Chem* 266(23): 15488-15492.
- Swaminath G, Deupi X, Lee TW, Zhu W, Thian FS, Kobilka TS and Kobilka B (2005) Probing the beta2 adrenoceptor binding site with catechol reveals differences in binding and activation by agonists and partial agonists. *J Biol Chem* 280(23): 22165-22171.
- Takagi Y, Ninomiya H, Sakamoto A, Miwa S and Masaki T (1995) Structural basis of G protein specificity of human endothelin receptors. A study with endothelinA/B chimeras. *J Biol Chem* 270(17): 10072-10078.
- Takai M, Umemura I, Yamasaki K, Watakabe T, Fujitani Y, Oda K, Urade Y, Inui T, Yamamura T and Okada T (1992) A potent and specific agonist, Suc-[Glu9,Ala11,15]-endothelin-1(8-21), IRL 1620, for the ETB receptor. *Biochem Biophys Res Commun* 184(2):953-959.
- Takashima H, Mimura N, Ohkubo T, Yoshida T, Tamaoki H and Kobayashi Y (2004a) Distributed computing and NMR constraint-based high-resolution structure determination: applied for bioactive Peptide endothelin-1 to determine C-terminal folding. *J Am Chem Soc* 126(14): 4504-4505.
- Takashima H, Tamaoki H, Teno N, Nishi Y, Uchiyama S, Fukui K and Kobayashi Y (2004b) Hydrophobic core around tyrosine for human endothelin-1 investigated by photochemically induced dynamic nuclear polarization nuclear magnetic resonance and matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Biochemistry* 43(44): 13932-13936.
- Takasuka T, Sakurai T, Goto K, Furuichi Y and Watanabe T (1994) Human endothelin receptor ETB. Amino acid sequence requirements for super stable complex formation with its ligand. *J Biol Chem* 269(10): 7509-7513.
- Tam JP, Liu W, Zhang JW, Galantino M, Bertolero F, Cristiani C, Vaghi F and de Castiglione R (1994) Alanine scan of endothelin: importance of aromatic residues. *Peptides* 15(4): 703-708.
- Teller DC, Okada T, Behnke CA, Palczewski K and Stenkamp RE (2001) Advances in determination of a high-resolution three-dimensional structure of rhodopsin, a model of G-protein-coupled receptors (GPCRs). *Biochemistry* 40(26): 7761-7772.
- ter Laak AM and Kuhne R (1999) Bacteriorhodopsin in a periodic boundary water-vacuum-water box as an example towards stable molecular dynamics simulations of G-protein coupled receptors. *Receptors Channels* 6(4): 295-308.
- Tesmer JJ, Sunahara RK, Gilman AG and Sprang SR (1997) Crystal structure of the catalytic domains of adenylyl cyclase in a complex with Gsalpha.GTPgammaS. *Science* 278(5345): 1907-1916.
- Tesmer JJ, Sunahara RK, Johnson RA, Gosselin G, Gilman AG and Sprang SR (1999) Two-metal-ion catalysis in adenylyl cyclase. *Science* 285(5428): 756-760.
- Thompson JD, Higgins DG and Gibson TJ (1994a) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22(22): 4673-4680.
- Thompson JD, Higgins DG and Gibson TJ (1994b) Improved sensitivity of profile searches through the use of sequence weights and gap excision. *Comput Appl Biosci* 10(1): 19-29.
- Tornvall P, Hamsten A, Johansson J and Carlson LA (1990) Normalisation of the composition of very low density lipoprotein in hypertriglyceridemia by nicotinic acid. *Atherosclerosis* 84(2-3): 219-227.
- Tota RT, Candelore MR, Dixon RAF and Strader CD (1990) Biophysical and genetic analysis of the ligand-binding site of the beta-adrenoceptor. *Trends Pharmacol Sci* 12: 4-6.
- Tunaru S, Kero J, Schaub A, Wufka C, Blaukat A, Pfeffer K and Offermanns S (2003) PUMA-G and HM74 are receptors for nicotinic acid and mediate its anti-lipolytic effect. *Nat Med* 9(3): 352-355.

REFERENCES

- Tunaru S, Lattig J, Kero J, Krause G and Offermanns S (2005) Characterization of determinants of ligand binding to the nicotinic acid receptor GPR109A (HM74A/PUMA-G). *Mol Pharmacol* 68(5): 1271-1280.
- Urade Y, Fujitani Y, Oda K, Watakabe T, Umemura I, Takai M, Okada T, Sakata K and Karaki H (1992) An endothelin B receptor-selective antagonist: IRL 1038, [Cys11-Cys15]-endothelin-1(11-21). *FEBS Lett* 311(1):12-16.
- Vassilatis DK, Hohmann JG, Zeng H, Li F, Ranchalis JE, Mortrud MT, Brown A, Rodriguez SS, Weller JR, Wright AC, Bergmann JE and Gaitanaris GA (2003) The G protein-coupled receptor repertoires of human and mouse. *Proc Natl Acad Sci U S A* 100(8): 4903-4908.
- Vichi P, Whelchel A and Posada J (1999) Transmembrane helix 7 of the endothelin B receptor regulates downstream signaling. *J Biol Chem* 274(15): 10331-10338.
- Wakamatsu K, Okada A, Miyazawa T, Ohya M and Higashijima T (1992) Membrane-bound conformation of mastoparan-X, a G-protein-activating peptide. *Biochemistry* 31(24): 5654-5660.
- Wall MA, Coleman DE, Lee E, Iniguez-Lluhi JA, Posner BA, Gilman AG and Sprang SR (1995) The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2. *Cell* 83(6): 1047-1058.
- Warner TD, de Nucci G and Vane JR (1989) Rat endothelin is a vasodilator in the isolated perfused mesentery of the rat. *Eur J Pharmacol* 159(3): 325-326.
- Watanabe TX, Itahara Y, Nakajima K, Kumagaye S, Kimura T and Sakakibara S (1991) The biological activity of endothelin-1 analogues in three different assay systems. *J Cardiovasc Pharmacol* 17 Suppl 7: S5-S9.
- Webb ML, Patel PS, Rose PM, Liu EC, Stein PD, Barrish J, Lach DA, Stouch T, Fisher SM, Hadjilambris O, Lee H, Skwish S, Dickinson KE and Krystek SR, Jr. (1996) Mutational analysis of the endothelin type A receptor (ETA): interactions and model of selective ETA antagonist BMS-182874 with putative ETA receptor binding cavity. *Biochemistry* 35(8): 2548-2556.
- Wess J (1998) Molecular basis of receptor/G-protein-coupling selectivity. *Pharmacol Ther* 80(3): 231-264.
- Wieland K, Zuurmond HM, Krasel C, Ijzerman AP and Lohse MJ (1996) Involvement of Asn-293 in stereospecific agonist recognition and in activation of the beta 2-adrenergic receptor. *Proc Natl Acad Sci U S A* 93(17): 9276-9281.
- Williams DL, Jr., Jones KL, Pettibone DJ, Lis EV and Clineschmidt BV (1991) Sarafotoxin S6c: an agonist which distinguishes between endothelin receptor subtypes. *Biochem Biophys Res Commun* 175(2):556-561.
- Wise A, Foord SM, Fraser NJ, Barnes AA, Elshourbagy N, Eilert M, Ignar DM, Murdock PR, Steplewski K, Green A, Brown AJ, Dowell SJ, Szekeres PG, Hassall DG, Marshall FH, Wilson S and Pike NB (2003) Molecular identification of high and low affinity receptors for nicotinic acid. *J Biol Chem* 278(11): 9869-9874.
- Withers-Ward ES, Mueller TD, Chen IS and Feigon J (2000) Biochemical and structural analysis of the interaction between the UBA(2) domain of the DNA repair protein HHR23A and HIV-1 Vpr. *Biochemistry* 39(46): 14103-14112.
- Wong SK (2003) G protein selectivity is regulated by multiple intracellular regions of GPCRs. *Neurosignals* 12(1): 1-12.
- Wuller S, Wiesner B, Loffler A, Ferkert J, Krause G, Hermosilla R, Schaefer M, Schulein R, Rosenthal W and Oksche A (2004) Pharmacochaperones post-translationally enhance cell surface expression by increasing conformational stability of wild-type and mutant vasopressin V2 receptors. *J Biol Chem* 279(45): 47254-47263.
- Yanagisawa M and Masaki T (1989) Molecular biology and biochemistry of the endothelins. *Trends Pharmacol Sci* 10(9):374-378.
- Yanagisawa M, Kurihara H, Kimura S, Tomobe Y, Kobayashi M, Mitsui Y, Yazaki Y, Goto K and Masaki T (1988) A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 332(6163):411-415.

REFERENCES

- Zhao MM, Hwa J and Perez DM (1996) Identification of critical extracellular loop residues involved in alpha 1-adrenergic receptor subtype-selective antagonist binding. *Mol Pharmacol* 50(5): 1118-1126.