

Bibliography

- [1] C. Cohen-Tannoudji, B. Liu, and F. Laloe. *Quantum Mechanics*. Wiley, New York, 1977.
- [2] M. Reed and B. Simon. *Functional Analysis*. Methods of Modern Mathematical Physics. Academic Press, New York, 1972.
- [3] P.W. Atkins. *Physical Chemistry*. Oxford University Press, Oxford, fifth edition, 1994.
- [4] O. Bratteli and D. Robinson. *Operator Algebras and Quantum Statistical Mechanics*, volume I and II. Springer, New York, 1979, 1981.
- [5] J. Manz. Molecular wavepacket dynamics: Theory for experiments 1926-1996. In V. Sundström, editor, *Femtochemistry and Femtobiology: Ultrafast Reaction Dynamics at Atomic-Scale Resolution*, pages 80–318. Imperial College Press, London, 1997.
- [6] K. Christmann. *Introduction to Surface Physical Chemistry*. Steinkopff, Darmstadt, 1991.
- [7] J.C. Tully. Theories of the dynamics of inelastic and reactive processes at surfaces. *Ann. Rev. Phys. Chem.*, 31:319–343, 1980.
- [8] B. Jackson. Quantum and semiclassical calculations of gas–surface energy transfer and sticking. *Comp. Phys. Commun.*, 80:119–1944, 1994.
- [9] M.R. Hand and J. Harris. Recoil effects in surface dissociation. *J. Chem. Phys.*, 92:7610–7617, 1990.

- [10] J.W. Gadzuk. Hot-electron femtochemistry at surfaces. In J. Manz and L. Wöste, editors, *Femtosecond Chemistry*, chapter 20, pages 603–624. VCH publishers, Weinheim, 1995.
- [11] R. Gomer. Diffusion of adsorbates on metal surfaces. *Rep. Prog. Phys.*, 53:917–1002, 1990.
- [12] R. Kosloff. Propagation methods for quantum molecular dynamics. *Annu. Rev. Phys. Chem.*, 45:145–178, 1994.
- [13] N. Balakrishnan, C. Kalyanaraman, and N. Sathyamurthy. Time-dependent quantum mechanical approach to reactive scattering and related processes. *Phys. Rep.*, 280:80–144, 1997.
- [14] P. Saalfrank and W.H. Miller. Quantum–mechanical rates for gas–surface processes. *Surf. Sci.*, 303:206–230, 1994.
- [15] R. Baer and R. Kosloff. Quantum dissipative dynamics of adsorbates near metal surfaces: A surrogate Hamiltonian theory applied to hydrogen on nickel. *J. Chem. Phys.*, 106:8862–8875, 1997.
- [16] M. Dohle and P. Saalfrank. Surface oscillator models for dissociative sticking of molecular hydrogen at non–rigid surfaces. *Surf. Sci.*, 373:95–108, 1996.
- [17] E.J. Heller. Time-dependent approach to semiclassical dynamics. *J. Chem. Phys.*, 62:1544–1555, 1975.
- [18] P.A.M. Dirac. Note on exchange phenomena in the Thomas atom. *Proc. Cambridge Phil. Soc.*, 26:376–385, 1930.
- [19] A.D. McLachlan. A variational solution of the time–dependent Schrödinger equation. *Mol. Phys.*, 8:39–44, 1963.
- [20] N. Makri and W.H. Miller. Time–Dependent Self–Consistent Field (TDSCF) approximation for a reaction coordinate coupled to a harmonic bath: Single and multiple configuration treatments. *J. Chem. Phys.*, 87:5781–5787, 1987.
- [21] Z. Kotler, A. Nitzan, and R. Kosloff. Multiconfiguration time-dependent self-consistent field approximation for curve crossing in presence of a bath. A Fast Fourier Transform study. *Chem. Phys. Lett.*, 153:483–498, 1988.

- [22] J. Campos-Martinez, J.R. Waldeck, and R.D. Coalson. Beyond the time-dependent Hartree grid approximation for curve-crossing problems. *J. Chem. Phys.*, 96:3613–3623, 1992.
- [23] H.-D. Meyer, U. Manthe, J. Kucar, and L.S. Cederbaum. Approximate methods for time-evolution of wave packets. In J. Broeckhove and L. Lathouwers, editors, *Time-Dependent Quantum Molecular Dynamics*, pages 233–246, New York, 1992. NATO Advanced Research Workshop, Plenum.
- [24] U. Manthe, H.-D. Meyer, and L.S. Cederbaum. Multiconfigurational time-dependent Hartree study of complex dynamics: Photodissociation of NO₂. *J. Chem. Phys.*, 97:9062, 1992.
- [25] M. Ehara, H.-D. Meyer, and L.S. Cederbaum. Multiconfiguration time-dependent Hartree (MCTDH) study on rotational and diffractive inelastic molecule-surface scattering. *J. Chem. Phys.*, 105:8865–8877, 1996.
- [26] G.A. Worth, H.-D. Meyer, and L.S. Cederbaum. The effect of a model environment on the S₂ absorption spectrum of pyrazine: A wave packet study treating all 24 vibrational modes. *J. Chem. Phys.*, 105:4412–4426, 1996.
- [27] H.-D. Meyer. Multiconfiguration time-dependent Hartree method. In P.v.R. Schleyer et. al., editor, *The Encyclopedia of Computational Chemistry*. John Wiley and Sons, Chichester, 1998. in press.
- [28] R.B. Gerber, V. Buch, and M.A. Ratner. Time-dependent self-consistent field approximation for intramolecular energy transfer. I. Formulation and application to dissociation of van der Waals molecules. *J. Chem. Phys.*, 77:3022–3030, 1982.
- [29] R. Alimi, R.B. Gerber, A.D. Hammerich, R. Kosloff, and M.A. Ratner. Validity of time-dependent self-consistent-field (TDSCF) approximations for unimolecular dynamics: A test for photodissociation of the Xe-HI cluster. *J. Chem. Phys.*, 93:6484–6490, 1990.
- [30] F.A. Bornemann, P. Nettesheim, and Ch. Schütte. Quantum-classical molecular dynamics as an approximation to full quantum dynamics. *J. Chem. Phys.*, 105:1074–1083, 1996.

- [31] P. Nettesheim, F.A. Bornemann, B. Schmidt, and Ch. Schütte. An explicit and symplectic integrator for quantum-classical molecular dynamics. *Chem. Phys. Lett.*, 256:581–588, 1996.
- [32] U. Müller and G. Stock. Consistent treatment of quantum-mechanical and classical degrees of freedom in mixed quantum-classical simulations. *J. Chem. Phys.*, 108:7516–7526, 1998.
- [33] P. Jungwirth and R.B. Gerber. Quantum dynamics of large polyatomic systems using a classically-based separable potential method. *J. Chem. Phys.*, 102:6046–6056, 1995.
- [34] P. Jungwirth, E. Fredj, and R.B. Gerber. Quantum molecular dynamics of large systems beyond separable approximation: The configuration interaction classical separable potential method. *J. Chem. Phys.*, 107:8963–8974, 1997.
- [35] P. Jungwirth and B. Schmidt. Quantum dynamics following electron photodetachment in the I^-Ar_2 complex: How good are new separable and non-separable simulation methods? *Chem. Phys. Lett.*, 275:127–136, 1997.
- [36] M.P. Allen and D.J. Tildesley. *Computer Simulations of Liquids*. Clarendon, Oxford, 1987.
- [37] K. Blum. *Density matrix theory and applications*. Plenum press, New York, 1981.
- [38] J. Harris. On the adsorption and desorption of H_2 at metal surfaces. *Appl. Phys. A*, 47:63–71, 1988.
- [39] M. Gostein, H. Parhikteh, and G.O. Sitz. Survival probability of H_2 ($v = 1, j = 1$) scattered from Cu(110). *Phys. Rev. Lett.*, 75:342–345, 1995.
- [40] M. Gostein, E. Watts, and G.O. Sitz. Vibrational relaxation of H_2 ($v = 1, j = 1$) on Pd(111). *Phys. Rev. Lett.*, 79:2891–2894, 1997.
- [41] J.W. Gadzuk, L.J. Richter, S.A. Buntin, D.S. King, and R.R. Cavanagh. Laser-excited hot-electron induced desorption: A theoretical model applied to NO/Pt(111). *Surf. Sci.*, 235:317–333, 1990.

- [42] S.A. Buntin, L.J. Richter, R.R. Cavanagh, and D.S. King. Optically driven surface reactions: Evidence for the role of hot electrons. *Phys. Rev. Lett.*, 61:1321–1324, 1988.
- [43] Ch. Scheurer and P. Saalfrank. Hydrogen transfer in vibrationally relaxing benzoic acid dimers: Time-dependent density matrix dynamics and infrared spectra. *J. Chem. Phys.*, 104:2869–2882, 1996.
- [44] W.E. Moerner and T. Basche. Optical spectroscopy of single impurity molecules in solids. *Angew. Chem.*, 105:537–557, 1993.
- [45] U.P. Wild, F. Gütter, M. Pirotta, and A. Renn. Single molecule spectroscopy: Stark effect of pentacene in p-terphenyl. *Chem. Phys. Lett.*, 193:451–455, 1992.
- [46] R. Alicki and K. Lendi. *Quantum Dynamical Semigroups and Applications*, volume 286 of *Lecture Notes in Physics*. Springer, Berlin, 1987.
- [47] R. Zwanzig. Ensemble method in the theory of irreversibility. *J. Chem. Phys.*, 33:1338–1341, 1960.
- [48] G. Lindblad. On the existence of quantum subdynamics. *J. Phys. A*, 29:4197–4207, 1996.
- [49] M.V. Korolkov and G.K. Paramonov. State-selective vibrational excitation of diatomic molecules coupled to a quasiresonant environment: Markov approximation and non-Markov approach. *Phys. Rev. A*, 55:589–604, 1997.
- [50] A. Suarez, R. Silbey, and I. Oppenheim. Memory effects in the relaxation of quantum open systems. *J. Chem. Phys.*, 97:5101–5107, 1992.
- [51] P. Talkner. The failure of the quantum regression hypothesis. *Ann. Phys. (NY)*, 167:390–436, 1986.
- [52] C. Meier and D.J. Tannor. Non-Markovian density matrix dynamics in solutions: I methodology. 1998. (in preparation).
- [53] A.G. Redfield. On the theory of relaxation processes. *IBM J. Res. Dev.*, 1:19–31, 1957.

- [54] A.G. Redfield. The theory of relaxation processes. *Adv. Mag. Reson.*, 1:1–32, 1965.
- [55] G. Lindblad. On the generators of quantum dynamical semigroups. *Commun. Math. Phys.*, 48:119–130, 1976.
- [56] V. Gorini, A. Kossakowski, and E.C. Sudarshan. Completely positive dynamical semigroups of N-level systems. *J. Math. Phys.*, 17:821–825, 1976.
- [57] H. Spohn. Kinetic equations from Hamiltonian dynamics: Markovian limits. *Rev. Mod. Phys.*, 52:569–615, 1980.
- [58] A.M. Walch and R.D. Coalson. Redfield theory is quantitative for coupled harmonic oscillators. *Chem. Phys. Lett.*, 198:293–299, 1992.
- [59] S. Gnutzmann and F. Haake. Positivity violation and initial slips in open systems. *Z. Phys. B*, 101:263–273, 1996.
- [60] S. Gao. Lindblad approach to quantum dynamics of open systems. *Phys. Rev. B*, 57:4509–4517, 1998.
- [61] G. Lindblad. Brownian motion of a quantum harmonic oscillator. *Rep. Math. Phys.*, 10:393–406, 1976.
- [62] F. Haake and R. Reinbold. Strong damping and low-temperature anomalies for the harmonic oscillator. *Phys. Rev. A*, 32:2462–2475, 1985.
- [63] D. Kohen and D.J. Tannor. Classical-quantum correspondence in the Redfield equation and its solutions. *J. Chem. Phys.*, 107:5141–5153, 1997.
- [64] D. Kohen, C.C. Marston, and D.J. Tannor. Phase space approach to the theories of quantum dissipation. *J. Chem. Phys.*, 107:5236–5253, 1997.
- [65] A.O. Caldeira and A.J. Leggett. Path integral approach to quantum Brownian motion. *Physica A*, 121:587–616, 1983.
- [66] I. Oppenheim and V. Romero-Rochin. Quantum Brownian motion. *Physica A*, 147:184–202, 1987.

- [67] R.P. Feynman and F.L. Vernon, Jr. The theory of a general quantum system interacting with a linear dissipative system. *Ann. Phys. (NY)*, 24:118–173, 1963.
- [68] O. Linden and V. May. Application of the time-dependent projection operator technique: The nonlinear quantum master equation. *Physica A*, 254:411–432, 1998.
- [69] B.B. Laird, J. Budimir, and J.L. Skinner. Quantum mechanical derivation of the Bloch equations: Beyond the weak-coupling limit. *J. Chem. Phys.*, 94:4391–4404, 1991.
- [70] B. Hellsing and H. Metiu. An efficient method for solving the quantum Liouville equation: Applications to electronic absorption spectroscopy. *Chem. Phys. Lett.*, 127:45–49, 1986.
- [71] M. Berman and R. Kosloff. Time-dependent solution of the Liouville–von Neumann equation: Non-dissipative evolution. *Comp. Phys. Comm.*, 63:1–20, 1991.
- [72] B.M. Garraway and P.L. Knight. Comparison of quantum-state diffusion and quantum-jump simulations of two-photon processes in a dissipative environment. *Phys. Rev. A*, 49:1266–1274, 1994.
- [73] W.K. Lai, K.-A. Suominen, B.M. Garraway, and S. Stenholm. Dissipation effects on wave packets in level crossings: A comparison between two numerical approaches. *Phys. Rev. A*, 47:4779–4785, 1993.
- [74] P. Saalfrank and R. Kosloff. Quantum dynamics of bond breaking in a dissipative environment: Indirect and direct photodesorption of neutrals from metals. *J. Chem. Phys.*, 105:2441–2455, 1996.
- [75] I. Burghardt. The dynamics of predissociation in the condensed phase: Markovian master equation. *J. Phys. Chem. A*, 102:4192–4206, 1998.
- [76] V. May, O. Kühn, and M. Schreiber. Density matrix description of ultrafast dissipative wave packet dynamics. *J. Phys. Chem.*, 97:12591–12595, 1993.

- [77] J.M. Jean, R.A. Friesner, and G.R. Fleming. Application of a multilevel Redfield theory to electron transfer in condensed phases. *J. Chem. Phys.*, 96:5827–5842, 1992.
- [78] W.T. Pollard and R.A. Friesner. Solution of the Redfield equation for the dissipative quantum dynamics of multilevel systems. *J. Chem. Phys.*, 100:5054–5065, 1994.
- [79] B. Jackson. Reduced density matrix description of gas–solid interactions: Scattering, trapping and desorption. *J. Chem. Phys.*, 108:1131–1139, 1997.
- [80] L. Pesce and P. Saalfrank. The coupled channel density matrix method for open quantum systems: Formulation and application to the vibrational relaxation of molecules scattering from nonrigid surfaces. *J. Chem. Phys.*, 108:3045–3056, 1998.
- [81] M. Berman, R. Kosloff, and H. Tal-Ezer. Solution of the time–dependent Liouville–von Neumann equation: Dissipative evolution. *J. Phys. A*, 25:1283–1307, 1992.
- [82] L. Pesce and P. Saalfrank. “Free” nuclear density propagation in two dimensions: The coupled channel density matrix method and its application to inelastic molecule-surface scattering. *Chem. Phys.*, 219:43–55, 1997.
- [83] C. Leforestier, R.H. Bisseling, C. Cerjan, M.D. Feit, R. Friesner, A. Guldberg, A. Hammerich, G. Jolicard, W. Karrlein, H.-D. Meyer, N. Lipkin, O. Roncero, and R. Kosloff. A comparison of different propagation schemes for the time dependent Schrödinger equation. *J. Comp. Phys.*, 94:59–80, 1991.
- [84] R. Dum, A.S. Parkins, P. Zoller, and C.W. Gardiner. Monte Carlo simulations of master equations in quantum optics for vacuum, thermal, and squeezed reservoirs. *Phys. Rev. A*, 46:4382–4396, 1992.
- [85] K. Mølmer, Y. Castin, and J. Dalibard. Monte Carlo wave–function method in quantum optics. *J. Opt. Soc. Am. B*, 10:524–538, 1993.
- [86] H.-P. Breuer and F. Petruccione. Stochastic dynamics of quantum jumps. *Phys. Rev. E*, 52:428–441, 1995.

- [87] J. Steinbach, B.M. Garraway, and P.L. Knight. High-order unreveling of master equations for dissipative evolution. *Phys. Rev. A*, 51:3302–3308, 1995.
- [88] B. Wolfseder and W. Domcke. Multi-mode vibronic coupling with dissipation. Application of the Monte Carlo wavefunction propagation method. *Chem. Phys. Lett.*, 235:370–376, 1995.
- [89] P. Saalfrank. Stochastic wave packet vs. direct solution of Liouville–von–Neumann equations for photodesorption processes. *Chem. Phys.*, 211:365–276, 1996.
- [90] K. Finger and P. Saalfrank. vibrationally excited products after photodesorption of NO from Pt(111): A two-mode open-system density matrix approach. *Chem. Phys. Lett.*, 268:291–299, 1997.
- [91] N. Gisin and I.C. Percival. The quantum state diffusion model applied to open systems. *J. Phys. A*, 25:5677–5691, 1992.
- [92] R. Dum, P. Zoller, and H. Ritsch. Monte Carlo simulation of the atomic master equation for spontaneous emission. *Phys. Rev. A*, 45:4879–4887, 1992.
- [93] N. Gisin and I.C. Percival. Wave–function approach to dissipative processes: Are there quantum jumps? *Phys. Lett. A*, 167:315–318, 1992.
- [94] N. Gisin and I.C. Percival. The quantum state diffusion picture of physical processes. *J. Phys. A*, 26:2245–2260, 1993.
- [95] T. Gerdts and U. Manthe. A wave packet approach to the Liouville–von Neumann equation for dissipative systems. *J. Chem. Phys.*, 106:3017–3023, 1996.
- [96] L. Pesce, T. Gerdts, U. Manthe, and P. Saalfrank. Variational wavepacket method for dissipative photodesorption problems. *Chem. Phys. Lett.*, 288:383–390, 1998.
- [97] H. Tal-Ezer. Polynomial approximation of functions of matrices and its application to the solution of a general system of linear equations. Contractor Report 87-63, NASA, 1987.

- [98] J.B. Conway. *A Course in Functional Analysis*. Springer, Berlin, 1990.
- [99] S. Lang. *Complex Analysis*. Springer, Berlin, 1993.
- [100] R. Kosloff. Pseudospectral methods for solving the time-dependent Schrödinger equation. In J. Broeckhove and L. Lathouwers, editors, *Time-Dependent Quantum Molecular Dynamics*, New York, 1992. NATO Advanced Research Workshop, Plenum.
- [101] J.C. Light, I.P. Hamilton, and J.V. Lill. Generalized discrete variable approximation in quantum mechanics. *J. Chem. Phys.*, 83:1400–1409, 1985.
- [102] W.T. Pollard, A.K. Felts, and R.A. Friesner. The Redfield equation in condensed-phase quantum dynamics. *Adv. Chem. Phys.*, 93:77, 1996.
- [103] S.W. Ellacott. Computation of Faber series with application to numerical polynomial approximation in the complex plane. *Mathematics of Computation*, 40:575–587, 1983.
- [104] S.W. Ellacott. A survey of Faber methods in numerical approximation. *Comput. Math. Appl.*, 12B:1103–1107, 1986.
- [105] L.N. Trefethen. Near-circularity of the error curve in complex Chebychev approximation. *J. Approx. Theory*, 31:344–367, 1981.
- [106] L. Reichel. Newton interpolation at Lejá points. *BIT*, 30:332–346, 1990.
- [107] Y. Huang, D.J. Kouri, and D.K. Hoffman. A general, energy-separable polynomial representation of the time-independent full Green operator with application to time-independent wavepacket forms of Schrödinger and Lippmann–Schwinger equations. *Chem. Phys. Lett.*, 225:37–45, 1994.
- [108] H. Tal-Ezer and R. Kosloff. An accurate and efficient scheme for propagating the time dependent Schrödinger equation. *J. Chem. Phys.*, 81:3967–3971, 1984.
- [109] D. Gaier. *Vorlesungen über Approximation im Komplexen*. Birkhäuser, Basel, 1980.

- [110] P. Henrici. *Applied and Computational Complex Analysis*, volume 3. John Wiley and Sons, New York, 1986.
- [111] Y. Huang, D.J. Kouri, and D.K. Hoffman. General energy–separable Faber polynomial representation of operator functions: Theory and application in quantum scattering. *J. Chem. Phys.*, 101:10493–10506, 1994.
- [112] W. Huiszinga. Faber–, Newton– und Krylovapproximation zur Integration grosser Differentialgleichungssysteme aus der Quantenmechanik. Diploma Thesis, Free University Berlin. Available via <http://www.zib.de/huiszinga>, 1997.
- [113] V.I. Smirnov and N.A. Lebedev. *Functions of a Complex Variable, Constructive Theory*. M.I.T. Press, Boston, 1968.
- [114] I.N. Bronstein and K.A. Semendjajew. *Taschenbuch der Mathematik*. Verlag Harry Deutsch, Zürich, 1989.
- [115] G. Ashkenazi, R. Kosloff, S. Ruhmann, and H. Tal-Ezer. Newtonian propagation methods applied to the photodissociation dynamics of I_3^- . *J. Chem. Phys.*, 103:10005–10014, 1995.
- [116] H. Tal-Ezer. High degree interpolation polynomial in Newton form. Contractor Report 88-39, NASA, 1988.
- [117] P. Nettesheim, W. Huiszinga, and C. Schütte. Chebyshev-approximation for wavepacket-dynamics: Better than expected. Technical Report SC-96-47, Konrad-Zuse-Zentrum für Informationstechnik Berlin, 1996.
- [118] E. Kreyszig. *Advanced Engineering Mathematics*. John Wiley & Sons, New York, seventh edition, 1993.
- [119] N. Došlić, O. Kühn, and J. Manz. Infrared laser pulse controlled ultrafast H-atom switching in two-dimensional asymmetric double well potentials. *Ber. Bunsenges. Phys. Chem.*, 102:292–297, 1998.
- [120] F. Neugebauer, D. Malzahn, and V. May. Calculation of molecular absorption spectra using a density matrix propagation scheme: An extension of Heller’s formula. *Chem. Phys.*, 201:151–162, 1995.

- [121] R. Kosloff. Time-dependent quantum-mechanical methods for molecular dynamics. *J. Phys. Chem.*, 92:2087–2100, 1988.
- [122] H. Guo. Calculation of resonance Raman amplitudes via polynomial propagation. *J. Chem. Phys.*, 1998. submitted.
- [123] I.S. Gradshteyn and I.M. Ryzhik. *Tables of Integrals, Series, and Products*. Academic, New York, 1980.
- [124] U. Manthe, H.-D. Meyer, and L.S. Cederbaum. Wave-packet dynamics within the multiconfiguration Hartree framework: General aspects and applications to NOCl. *J. Chem. Phys.*, 97:3199–3213, 1992.
- [125] A.P.J. Jansen. A multiconfiguration time-dependent Hartree approximation based on natural single-particle states. *J. Chem. Phys.*, 99:4055–4058, 1993.
- [126] U. Manthe. Comment on "A multiconfigurational time-dependent Hartree approximation based on natural single-particle states "[J.Chem.Phys. 99,4055(1993)]. *J. Chem. Phys.*, 101:2652–2653, 1994.
- [127] R. Kosloff. Quantum molecular dynamics on grids. In R.E. Wyatt and J.Z.H. Zhang, editors, *Dynamics of Molecules and Chemical Reactions*. M. Decker, New York, 1996.
- [128] J.T. Muckerman and T. Uzer. A theoretical investigation of microscopic dynamics and time scales of predesorption processes. *J. Chem. Phys.*, 90:1968–1973, 1989.
- [129] Y. Guan, J.T. Muckerman, and T. Uzer. Desorption of vibrationally excited adsorbates in competition with relaxation: A quantal picture. *J. Chem. Phys.*, 93:4400–4412, 1990.
- [130] L.J. Richter, S.A. Buntin, R.R. Cavanagh, and D.S. King. Non-Boltzmann rotational and inverted spin-orbit state distributions for laser-induced desorption of NO from Pt(111). *J. Chem. Phys.*, 89:5344–5345, 1988.
- [131] S.A. Buntin, L.J. Richter, D.S. King, and R.R. Cavanagh. State-resolved evidence for hot carrier driven surface reactions: Laser-induced desorption of NO from Pt(111). *J. Chem. Phys.*, 91:6429–6446, 1989.

- [132] J. Manz and L. Wöste, editors. *Femtosecond Chemistry*, Weinheim, 1995. VCH publishers.
- [133] P. Saalfrank, R. Baer, and R. Kosloff. Density matrix description of laser-induced hot electron mediated photodesorption of NO from Pt(111). *Chem. Phys. Lett.*, 230:463–472, 1994.
- [134] P. Saalfrank. Photodesorption of neutrals from metal surfaces: A wave packet study. *Chem. Phys.*, 193:119–139, 1995.
- [135] N. Chakrabarti, V. Balasubramanian, N. Sathyamurthy, and J.W. Gadzuk. Photoinduced desorption in NO/Pt: A time-dependent quantum mechanical study. *Chem. Phys. Lett.*, 242:490–498, 1995.
- [136] W. Reimer, T. Fink, and J. Küppers. Inverse photoemission spectroscopy of H, CO and NO adsorbed at Ni(100) and Ni(111) surfaces. *Surf. Sci.*, 193:259–270, 1988.
- [137] J. Dai and J.C. Light. The steric effect in a full dimensional quantum dynamics simulation for the dissociative adsorption of H₂ on Cu(111). *J. Chem. Phys.*, 108:7816–8920, 1998.
- [138] J. Dai and J.C. Light. Six dimensional quantum dynamics study for dissociative adsorption of H₂ on Cu(111) surface. *J. Chem. Phys.*, 107:1676–1679, 1997.
- [139] A. Gross, S. Wilke, and M. Scheffler. Six-dimensional quantum dynamics of adsorption and desorption of H₂ at Pd(100): No need for a molecular precursor adsorption state. *Surf. Sci.*, 357:614–358, 1996.
- [140] A. Gross, S. Wilke, and M. Scheffler. Six-dimensional quantum dynamics of adsorption and desorption of H₂ at Pd(100): Steering and steric effects. *Phys. Rev. Lett.*, 75:2718–2721, 1995.
- [141] G.J. Kroes, E.J. Baerends, and R.C. Mowrey. Six-dimensional quantum dynamics of dissociative chemisorption of ($v = 0, j = 0$) H₂ on Cu(100). *Phys. Rev. Lett.*, 78:3583–3586, 1997.

- [142] G.J. Kroes, E.J. Baerends, and R.C. Mowrey. Six-dimensional quantum dynamics of dissociative chemisorption of H₂ on Cu(100). *J. Chem. Phys.*, 107:3309–3323, 1997.
- [143] G.R. Darling and S. Holloway. The dissociation of diatomic molecules at surfaces. *Rep. Prog. Phys.*, 58:1595–1672, 1996.
- [144] B. Jackson. Time-dependent wave packet approach to quantum reactive scattering. *Annu. Rev. Phys. Chem.*, 46:251–274, 1995.
- [145] C.T. Rettner, D.J. Auerbach, J.C. Tully, and A.W. Kleyn. Chemical dynamics at the gas–surface interface. *J. Phys. Chem.*, 100:13021–13033, 1996.
- [146] H.A. Michelsen, C.T. Rettner, D.J. Auerbach, and R.N. Zare. Effect of rotation on the translational and vibrational energy dependence of the dissociative adsorption of D₂ on Cu(111). *J. Chem. Phys.*, 98:8294–8307, 1993.
- [147] C.T. Rettner, D.J. Auerbach, and H.A. Michelsen. Role of vibrational and translational energy in the activated dissociative adsorption of D₂ on Cu(111). *Phys. Rev. Lett.*, 68:1164–1167, 1992.
- [148] C.T. Rettner, D.J. Auerbach, and H.A. Michelsen. Observation of direct vibrational excitation in collisions of H₂ and D₂ with a Cu(111) surface. *Phys. Rev. Lett.*, 68:2547–2550, 1992.
- [149] H.A. Michelsen, C.T. Rettner, and D.J. Auerbach. On the influence of surface temperature on adsorption and desorption in the D₂/Cu(111) system. *Surf. Sci.*, 272:65–72, 1992.
- [150] D.T. Colbert and W.H. Miller. A novel discrete variable representation for quantum mechanical reactive scattering via the S-matrix Kohn method. *J. Chem. Phys.*, 96:1982–1991, 1992.
- [151] R.C. Mowrey and D.J. Kouri. On a hybrid close-coupling-wave packet approach to molecular scattering. *Chem. Phys. Lett.*, 119:285–289, 1985.
- [152] R.C. Mowrey and D.J. Kouri. Close-coupling wave packet approach to numerically exact molecule-surface scattering calculations. *J. Chem. Phys.*, 84:6466–6473, 1986.

- [153] H.F. Bowen, D.J. Kouri, R.C. Mowrey, A.T. Yinnon, and R.B. Gerber. Molecule–corrugated surface collisions: Converged close coupling wave packet and quasiclassical trajectory calculations for N₂ scattering from corrugated lattices. *J. Chem. Phys.*, 99:704–720, 1993.
- [154] G.-J. Kroes and R.C. Mowrey. Calculations on rotationally and diffractionally inelastic molecule–surface scattering for arbitrary angles of incidence: A new wave packet technique. *J. Chem. Phys.*, 101:805–813, 1994.
- [155] G.-J. Kroes and R.C. Mowrey. Scattering of H₂ by LiF(001) studied using a new model potential. I. Prediction of large differences in diffraction of cold beams of para-H₂ and normal H₂. *J. Chem. Phys.*, 103:2186–2201, 1995.
- [156] G.R. Darling and S. Holloway. Rotational motion and the dissociation of H₂ on Cu(111). *J. Chem. Phys.*, 101:3268–3281, 1994.
- [157] A. Bartana, R. Kosloff, and D. Tannor. Laser cooling of molecular internal degrees of freedom by a series of shaped pulses. *J. Chem. Phys.*, 99:196–210, 1993.
- [158] U. Banin, A. Bartana, S. Ruhman, and R. Kosloff. Impulsive excitation of coherent vibrational motion ground surface dynamics induced by intense short pulses. *J. Chem. Phys.*, 101:8461–8481, 1994.
- [159] B. Jackson and H. Metiu. The dynamics of H₂ dissociation on Ni(100): a quantum mechanical study of a restricted two-dimensional model. *J. Chem. Phys.*, 86:1026–1035, 1987.
- [160] G.A. Gates, G.R. Darling, and S. Holloway. A theoretical study of the vibrational excitation of NO/Ag(111). *J. Chem. Phys.*, 101:6281–6288, 1994.
- [161] A. Hodgson, J. Moryl, P. Traversaro, and H. Zhao. Energy transfer and vibrational effects in the dissociation and scattering of D₂ from Cu(111). *Nature*, 356:501–504, 1992.
- [162] R.B. Gerber. Molecular scattering from surfaces: Theoretical methods and results. *Chem. Rev.*, 87:29–79, 1987.

- [163] M. Morin, N.J. Levinos, and A.L. Harris. Vibrational energy transfer of CO/Cu(100): Nonadiabatic vibration/electron coupling. *J. Chem. Phys.*, 96:3950–3956, 1992.
- [164] M. Head-Gordon and J.C. Tully. Vibrational relaxation on metal surfaces: Molecular-orbital theory and application to CO/Cu(100). *J. Chem. Phys.*, 96:3939–3949, 1992.
- [165] M. Head-Gordon and J.C. Tully. Molecular-orbital calculation of the lifetimes of the vibrational modes of CO on Cu(100). *Phys. Rev. B*, 46:1853–1856, 1992.
- [166] B. Jackson. Reduced density matrix approach for gas–solid scattering. *Chem. Phys. Lett.*, 270:484–490, 1997.
- [167] R.E. Walkup, D.M. Newns, and Ph. Avouris. Vibrational heating and atom transfer with the STM. *J. Electron Spectrosc. Relat. Phenom.*, 64/65:523, 1993.
- [168] S. Gao, B.I. Lundqvist, and W. Ho. Hot-electron-induced vibrational heating at surface: Importance of quantum-mechanical description. *Surf. Sci.*, 341:L1031–L1036, 1995.
- [169] S. Efrima, K.F. Freed, C. Jedrzejek, and H. Metiu. A one-dimensional microscopic model for thermal desorption of an atom. Applications to the case of weak bonding. *Chem. Phys. Lett.*, 74:43–48, 1980.
- [170] R. Kosloff and C. Cerjan. Dynamical atom/surface effects: Quantum mechanical scattering and desorption. *J. Chem. Phys.*, 81:3722–3729, 1984.
- [171] O. Kühn and V. May. Density matrix approach to the hot-electron stimulated photodesorption. *Chem. Phys.*, 208:117–126, 1996.
- [172] E.R. Bittner and J.C. Light. Quantum stochastic approach for molecule/surface scattering. II. Adsorption resonances of He atoms on Xe overlayers. *J. Chem. Phys.*, 101:2446–2454, 1994.
- [173] D. Bekšić and D.A. Micha. Electronically diabatic quantum dynamics of molecular desorption. *J. Chem. Phys.*, 103:3795–3808, 1995.

- [174] G.R. Darling and S. Holloway. Rotational effects in the dissociative adsorption of H₂ on Cu(111). *Faraday Disc. Chem. Soc.*, 96:43–54, 1993.
- [175] B.Hammer and J.Nørskov. Why gold is the noblest of all the metals? *Nature*, 376:238–240, 1995.
- [176] C.T. Rettner, F. Fabre, J. Kimman, and D.J. Auerbach. Observation of direct vibrational excitation in gas surface collisions: NO on Ag(111). *Phys. Rev. Lett.*, 55:1904–1907, 1985.
- [177] E.K. Watts, J.L.W. Siders, and G.O. Sitz. Vibrational excitation of NO scattered from Cu(110). *Surf. Sci.*, 374:191–196, 1997.
- [178] S. Andersson and J. Harris. Trapping of molecular hydrogen at surfaces via translational–rotational energy conversion. *Phys. Rev. B*, 27:9–14, 1983.
- [179] S. Andersson and M. Persson. Sticking in the physisorption well: Influence of surface structure. *Phys. Rev. Lett.*, 70:202–205, 1993.
- [180] M.H. Beck and H.-D. Meyer. An efficient and robust integration scheme for the equations of motion of the multiconfiguration time-dependent Hartree (MCTDH) method. *Z. Phys. D*, 42:113–129, 1997.
- [181] M. Hochbruck and C. Lubich. On Krylov subspace approximations to the matrix exponential operator. *SIAM J. Numer. Anal.*, 34:1911–1925, 1995.
- [182] E. Fattal, R. Baer, and R. Kosloff. A phase space approach for optimizing grid representation: The mapped Fourier method. *Phys. Rev. E*, 53:1217–1227, 1996.