

7 Literatur

1. Julius D, Basbaum AI: Molecular mechanisms of nociception. *Nature* 2001; 413: 203-10
2. Scholz J, Woolf CJ: Can we conquer pain? *Nat Neurosci* 2002; 5 Suppl: 1062-7
3. Merskey H, Bogduk N: Pain terminology. 1994; 209-214
4. Zubieta JK, Smith YR, Bueller JA, et al.: Regional mu opioid receptor regulation of sensory and affective dimensions of pain. *Science* 2001; 293: 311-5
5. Przewlocki R, Hassan AH, Lason W, Epplen C, Herz A, Stein C: Gene expression and localization of opioid peptides in immune cells of inflamed tissue: functional role in antinociception. *Neuroscience* 1992; 48: 491-500
6. Ji RR, Zhang Q, Law PY, Low HH, Elde R, Hokfelt T: Expression of mu-, delta-, and kappa-opioid receptor-like immunoreactivities in rat dorsal root ganglia after carrageenan-induced inflammation. *J Neurosci* 1995; 15: 8156-66
7. Rittner HL, Brack A, Machelska H, et al.: Opioid peptide-expressing leukocytes: identification, recruitment, and simultaneously increasing inhibition of inflammatory pain. *Anesthesiology* 2001; 95: 500-8
8. Rittner HL, Machelska H, Schäfer M, Stein C, Brack A: Comment on "Neutrophils: are they hyperalgesic or anti-hyperalgesic?" *J Leukoc Biol* 2006
9. Rittner HL, Brack A, Mousa SA, et al.: Selective local PMN recruitment by CXCL1 or CXCL2/3 injection does not cause inflammatory pain. *J Leukoc Biol* 2006; 79: 1022-32
10. Stein C, Gramsch C, Herz A: Intrinsic mechanisms of antinociception in inflammation: local opioid receptors and beta-endorphin. *J Neurosci* 1990; 10: 1292-8
11. Stein C, Hassan AH, Przewlocki R, Gramsch C, Peter K, Herz A: Opioids from immunocytes interact with receptors on sensory nerves to inhibit nociception in inflammation. *Proc Natl Acad Sci U S A* 1990; 87: 5935-9
12. Stein C, Schäfer M, Machelska H: Attacking pain at its source: new perspectives on opioids. *Nat Med* 2003; 9: 1003-8
13. Machelska H, Cabot PJ, Mousa SA, Zhang Q, Stein C: Pain control in inflammation governed by selectins. *Nat Med* 1998; 4: 1425-8
14. Brack A, Rittner HL, Machelska H, et al.: Control of inflammatory pain by chemokine-mediated recruitment of opioid-containing polymorphonuclear cells. *Pain* 2004; 112: 229-38
15. Rittner HL, Brack A, Stein C: [Pain and the immune system: friend or foe?]. *Anaesthetist* 2002; 51: 351-8
16. Rittner HL, Brack A, Stein C: Pro-algesic versus analgesic actions of immune cells. *Curr Opin Anaesthesiol* 2003; 16: 527-33
17. Jones AK, al-Janabi MA, Solanki K, et al.: In vivo leukocyte migration in arthritis. *Arthritis Rheum* 1991; 34: 270-5
18. Levine JD, Lau W, Kwiat G, Goetzl EJ: Leukotriene B4 produces hyperalgesia that is dependent on polymorphonuclear leukocytes. *Science* 1984; 225: 743-5
19. Levine JD, Gooding J, Donatoni P, Borden L, Goetzl EJ: The role of the polymorphonuclear leukocyte in hyperalgesia. *J Neurosci* 1985; 5: 3025-9
20. Baggiolini M: Chemokines and leukocyte traffic. *Nature* 1998; 392: 565-8
21. Luster AD: Chemokines--chemotactic cytokines that mediate inflammation. *N Engl J Med* 1998; 338: 436-45

22. Rollins BJ: Chemokines. *Blood* 1997; 90: 909-28
23. Middleton J, Patterson AM, Gardner L, Schmutz C, Ashton BA: Leukocyte extravasation: chemokine transport and presentation by the endothelium. *Blood* 2002; 100: 3853-60
24. Nakagawa H, Komorita N, Shibata F, et al.: Identification of cytokine-induced neutrophil chemoattractants (CINC), rat GRO/CINC-2 alpha and CINC-2 beta, produced by granulation tissue in culture: purification, complete amino acid sequences and characterization. *Biochem J* 1994; 301 (Pt 2): 545-50
25. Shibata F, Konishi K, Kato H, et al.: Recombinant production and biological properties of rat cytokine-induced neutrophil chemoattractants, GRO/CINC-2 alpha, CINC-2 beta and CINC-3. *Eur J Biochem* 1995; 231: 306-11
26. Wagner JG, Roth RA: Neutrophil migration mechanisms, with an emphasis on the pulmonary vasculature. *Pharmacol Rev* 2000; 52: 349-74
27. Chandrasekar B, Smith JB, Freeman GL: Ischemia-reperfusion of rat myocardium activates nuclear factor-KappaB and induces neutrophil infiltration via lipopolysaccharide-induced CXC chemokine. *Circulation* 2001; 103: 2296-302
28. Amano H, Oishi K, Sonoda F, et al.: Role of cytokine-induced neutrophil chemoattractant-2 (CINC-2) alpha in a rat model of chronic bronchopulmonary infections with *Pseudomonas aeruginosa*. *Cytokine* 2000; 12: 1662-8
29. Garcia-Ramallo E, Marques T, Prats N, Beleta J, Kunkel SL, Godessart N: Resident cell chemokine expression serves as the major mechanism for leukocyte recruitment during local inflammation. *J Immunol* 2002; 169: 6467-73
30. Zhang XW, Wang Y, Liu Q, Thorlacius H: Redundant function of macrophage inflammatory protein-2 and KC in tumor necrosis factor-alpha-induced extravasation of neutrophils in vivo. *Eur J Pharmacol* 2001; 427: 277-83
31. Zwijnenburg PJ, Polfliet MM, Florquin S, et al.: CXC-chemokines KC and macrophage inflammatory protein-2 (MIP-2) synergistically induce leukocyte recruitment to the central nervous system in rats. *Immunol Lett* 2003; 85: 1-4
32. Stein C, Millan MJ, Yassouridis A, Herz A: Antinociceptive effects of mu- and kappa-agonists in inflammation are enhanced by a peripheral opioid receptor-specific mechanism. *Eur J Pharmacol* 1988; 155: 255-64
33. Hargreaves K, Dubner R, Brown F, Flores C, Joris J: A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* 1988; 32: 77-88
34. Hunt SP, Pini A, Evan G: Induction of c-fos-like protein in spinal cord neurons following sensory stimulation. *Nature* 1987; 328: 632-4
35. Ma QP, Woolf CJ: Basal and touch-evoked fos-like immunoreactivity during experimental inflammation in the rat. *Pain* 1996; 67: 307-16
36. Presley RW, Menetrey D, Levine JD, Basbaum AI: Systemic morphine suppresses noxious stimulus-evoked Fos protein-like immunoreactivity in the rat spinal cord. *J Neurosci* 1990; 10: 323-35
37. Abbadie C, Besson JM: C-fos expression in rat lumbar spinal cord following peripheral stimulation in adjuvant-induced arthritic and normal rats. *Brain Res* 1993; 607: 195-204
38. Li X, Lighthall G, Liang DY, Clark JD: Alterations in spinal cord gene expression after hindpaw formalin injection. *J Neurosci Res* 2004; 78: 533-41
39. Brack A, Rittner HL, Machelska H, et al.: Mobilization of opioid-containing polymorphonuclear cells by hematopoietic growth factors and influence on inflammatory pain. *Anesthesiology* 2004; 100: 149-57

40. Zimmermann M: Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* 1983; 16: 109-10
41. Brack A, Rittner HL, Younge BR, Kaltschmidt C, Weyand CM, Goronzy JJ: Glucocorticoid-mediated repression of cytokine gene transcription in human arteritis-SCID chimeras. *J Clin Invest* 1997; 99: 2842-50
42. Endlich B, Armstrong D, Brodsky J, Novotny M, Hamilton TA: Distinct temporal patterns of macrophage-inflammatory protein-2 and KC chemokine gene expression in surgical injury. *J Immunol* 2002; 168: 3586-94
43. Rovai LE, Herschman HR, Smith JB: The murine neutrophil-chemoattractant chemokines LIX, KC, and MIP-2 have distinct induction kinetics, tissue distributions, and tissue-specific sensitivities to glucocorticoid regulation in endotoxemia. *J Leukoc Biol* 1998; 64: 494-502
44. Takano K, Nakagawa H: Contribution of cytokine-induced neutrophil chemoattractant CINC-2 and CINC-3 to neutrophil recruitment in lipopolysaccharide-induced inflammation in rats. *Inflamm Res* 2001; 50: 503-8
45. Frendeus B, Godaly G, Hang L, Karpman D, Svanborg C: Interleukin-8 receptor deficiency confers susceptibility to acute pyelonephritis. *J Infect Dis* 2001; 183 Suppl 1: S56-60
46. Hall LR, Diaconu E, Patel R, Pearlman E: CXC chemokine receptor 2 but not C-C chemokine receptor 1 expression is essential for neutrophil recruitment to the cornea in helminth-mediated keratitis (river blindness). *J Immunol* 2001; 166: 4035-41
47. Moore TA, Newstead MW, Strieter RM, Mehrad B, Beaman BL, Standiford TJ: Bacterial clearance and survival are dependent on CXC chemokine receptor-2 ligands in a murine model of pulmonary Nocardia asteroides infection. *J Immunol* 2000; 164: 908-15
48. Blackwell TS, Lancaster LH, Blackwell TR, Venkatakrishnan A, Christman JW: Chemotactic gradients predict neutrophilic alveolitis in endotoxin-treated rats. *Am J Respir Crit Care Med* 1999; 159: 1644-52
49. Schmal H, Shanley TP, Jones ML, Friedl HP, Ward PA: Role for macrophage inflammatory protein-2 in lipopolysaccharide-induced lung injury in rats. *J Immunol* 1996; 156: 1963-72
50. Wiekowski MT, Chen SC, Zalamea P, et al.: Disruption of neutrophil migration in a conditional transgenic model: evidence for CXCR2 desensitization in vivo. *J Immunol* 2001; 167: 7102-10
51. Frevert CW, Farone A, Danaee H, Paulauskis JD, Kobzik L: Functional characterization of rat chemokine macrophage inflammatory protein-2. *Inflammation* 1995; 19: 133-42
52. Mercer-Jones MA, Shrotri MS, Heinzelmann M, Peyton JC, Cheadle WG: Regulation of early peritoneal neutrophil migration by macrophage inflammatory protein-2 and mast cells in experimental peritonitis. *J Leukoc Biol* 1999; 65: 249-55
53. Bennett G, al-Rashed S, Hoult JR, Brain SD: Nerve growth factor induced hyperalgesia in the rat hind paw is dependent on circulating neutrophils. *Pain* 1998; 77: 315-22
54. Foster PA, Wicks S, Foster M, Brain SD: Cellular pathology changes in rat skin following intradermal injection of nerve growth factor: neutrophil-dependent and - independent events. *J Pathol* 2002; 197: 245-55
55. Sacerdote P, Manfredi B, Gaspani L, Panerai AE: The opioid antagonist naloxone induces a shift from type 2 to type 1 cytokine pattern in BALB/cJ mice. *Blood* 2000; 95: 2031-6

56. Soruri A, Riggert J, Schlott T, Kiafard Z, Dettmer C, Zwirner J: Anaphylatoxin C5a induces monocyte recruitment and differentiation into dendritic cells by TNF-alpha and prostaglandin E2-dependent mechanisms. *J Immunol* 2003; 171: 2631-6
57. Susaki Y, Shimizu S, Katakura K, et al.: Functional properties of murine macrophages promoted by nerve growth factor. *Blood* 1996; 88: 4630-7
58. Cunha FQ, Lorenzetti BB, Poole S, Ferreira SH: Interleukin-8 as a mediator of sympathetic pain. *Br J Pharmacol* 1991; 104: 765-7
59. Cunha FQ, Poole S, Lorenzetti BB, Ferreira SH: The pivotal role of tumour necrosis factor alpha in the development of inflammatory hyperalgesia. *Br J Pharmacol* 1992; 107: 660-4
60. Cunha JM, Cunha FQ, Poole S, Ferreira SH: Cytokine-mediated inflammatory hyperalgesia limited by interleukin-1 receptor antagonist. *Br J Pharmacol* 2000; 130: 1418-24
61. Cunha TM, Verri WA, Jr., Silva JS, Poole S, Cunha FQ, Ferreira SH: A cascade of cytokines mediates mechanical inflammatory hypernociception in mice. *Proc Natl Acad Sci U S A* 2005; 102: 1755-60
62. Lorenzetti BB, Poole S, Veiga FH, Cunha FQ, Ferreira SH: Cytokine-mediated inflammatory hyperalgesia limited by interleukin-13. *Eur Cytokine Netw* 2001; 12: 260-7
63. Lorenzetti BB, Veiga FH, Canetti CA, Poole S, Cunha FQ, Ferreira SH: Cytokine-induced neutrophil chemoattractant 1 (CINC-1) mediates the sympathetic component of inflammatory mechanical hypersensitivity in rats. *Eur Cytokine Netw* 2002; 13: 456-61
64. Oh SB, Tran PB, Gillard SE, Hurley RW, Hammond DL, Miller RJ: Chemokines and glycoprotein120 produce pain hypersensitivity by directly exciting primary nociceptive neurons. *J Neurosci* 2001; 21: 5027-35
65. Opree A, Kress M: Involvement of the proinflammatory cytokines tumor necrosis factor-alpha, IL-1 beta, and IL-6 but not IL-8 in the development of heat hyperalgesia: effects on heat-evoked calcitonin gene-related peptide release from rat skin. *J Neurosci* 2000; 20: 6289-93
66. Brack A, Rittner HL, Machelska H, et al.: Endogenous peripheral antinociception in early inflammation is not limited by the number of opioid-containing leukocytes but by opioid receptor expression. *Pain* 2004; 108: 67-75
67. Ribeiro RA, Vale ML, Thomazzi SM, et al.: Involvement of resident macrophages and mast cells in the writhing nociceptive response induced by zymosan and acetic acid in mice. *Eur J Pharmacol* 2000; 387: 111-8