

## **6 LITERATURVERZEICHNIS**

---

- Ahmed, N., Maines-Bandiera, S., Quinn, M.A., Unger, W.G., Dedhar, S. and Auersperg, N. (2006) Molecular pathways regulating EGF-induced epithelio-mesenchymal transition in human ovarian surface epithelium. *Am J Physiol Cell Physiol*, 290, C1532-1542.
- Ahmed, N., Riley, C., Rice, G. and Quinn, M. (2005) Role of integrin receptors for fibronectin, collagen and laminin in the regulation of ovarian carcinoma functions in response to a matrix microenvironment. *Clin Exp Metastasis*, 22, 391-402.
- Akao, Y., Banno, Y., Nakagawa, Y., Hasegawa, N., Kim, T.J., Murate, T., Igarashi, Y. and Nozawa, Y. (2006) High expression of sphingosine kinase 1 and S1P receptors in chemotherapy-resistant prostate cancer PC3 cells and their camptothecin-induced up-regulation. *Biochem Biophys Res Commun*, 342, 1284-1290.
- Akiyoshi, S., Ishii, M., Nemoto, N., Kawabata, M., Aburatani, H. and Miyazono, K. (2001) Targets of transcriptional regulation by transforming growth factor-beta: expression profile analysis using oligonucleotide arrays. *Jpn J Cancer Res*, 92, 257-268.
- Alexandrow, M.G. and Moses, H.L. (1995) Transforming growth factor beta and cell cycle regulation. *Cancer Res*, 55, 1452-1457.
- Allende, M.L., Yamashita, T. and Proia, R.L. (2003) G-protein-coupled receptor S1P1 acts within endothelial cells to regulate vascular maturation. *Blood*, 102, 3665-3667.
- Ancellin, N. and Hla, T. (1999) Differential pharmacological properties and signal transduction of the sphingosine 1-phosphate receptors EDG-1, EDG-3, and EDG-5. *J Biol Chem*, 274, 18997-19002.
- Angst, B.D., Marcozzi, C. and Magee, A.I. (2001) The cadherin superfamily. *J Cell Sci*, 114, 625-626.
- Anliker, B. and Chun, J. (2004a) Cell surface receptors in lysophospholipid signaling. *Semin Cell Dev Biol*, 15, 457-465.
- Anliker, B. and Chun, J. (2004b) Lysophospholipid G protein-coupled receptors. *J Biol Chem*, 279, 20555-20558.
- Ara, T., Deyama, Y., Yoshimura, Y., Higashino, F., Shindoh, M., Matsumoto, A. and Fukuda, H. (2000) Membrane type 1-matrix metalloproteinase expression is regulated by E-cadherin through the suppression of mitogen-activated protein kinase cascade. *Cancer Lett*, 157, 115-121.
- Arikawa, K., Takuwa, N., Yamaguchi, H., Sugimoto, N., Kitayama, J., Nagawa, H., Takehara, K. and Takuwa, Y. (2003) Ligand-dependent inhibition of B16 melanoma cell migration and invasion via endogenous S1P2 G protein-coupled receptor. Requirement of inhibition of cellular RAC activity. *J Biol Chem*, 278, 32841-32851.
- Ashcroft, G.S., Yang, X., Glick, A.B., Weinstein, M., Letterio, J.L., Mizel, D.E., Anzano, M., Greenwell-Wild, T., Wahl, S.M., Deng, C. and Roberts, A.B. (1999) Mice lacking Smad3 show accelerated wound healing and an impaired local inflammatory response. *Nat Cell Biol*, 1, 260-266.
- Azuma, H., Takahara, S., Ichimaru, N., Wang, J.D., Itoh, Y., Otsuki, Y., Morimoto, J., Fukui, R., Hoshiga, M., Ishihara, T., Nonomura, N., Suzuki, S., Okuyama, A. and Katsuoaka, Y. (2002) Marked prevention of tumor growth and metastasis by a novel immunosuppressive agent, FTY720, in mouse breast cancer models. *Cancer Res*, 62, 1410-1419.

- Bajjalieh, S.M., Martin, T.F. and Floor, E. (1989) Synaptic vesicle ceramide kinase. A calcium-stimulated lipid kinase that co-purifies with brain synaptic vesicles. *J Biol Chem*, 264, 14354-14360.
- Bajou, K., Masson, V., Gerard, R.D., Schmitt, P.M., Albert, V., Praus, M., Lund, L.R., Frandsen, T.L., Brunner, N., Dano, K., Fusenig, N.E., Weidle, U., Carmeliet, G., Loskutoff, D., Collen, D., Carmeliet, P., Foidart, J.M. and Noel, A. (2001) The plasminogen activator inhibitor PAI-1 controls in vivo tumor vascularization by interaction with proteases, not vitronectin. Implications for antiangiogenic strategies. *J Cell Biol*, 152, 777-784.
- Bakin, A.V., Tomlinson, A.K., Bhowmick, N.A., Moses, H.L. and Arteaga, C.L. (2000) Phosphatidylinositol 3-kinase function is required for transforming growth factor beta-mediated epithelial to mesenchymal transition and cell migration. *J Biol Chem*, 275, 36803-36810.
- Bates, R.C. and Mercurio, A.M. (2003) Tumor necrosis factor-alpha stimulates the epithelial-to-mesenchymal transition of human colonic organoids. *Mol Biol Cell*, 14, 1790-1800.
- Batlle, E., Sancho, E., Franci, C., Dominguez, D., Monfar, M., Baulida, J. and Garcia De Herreros, A. (2000) The transcription factor snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat Cell Biol*, 2, 84-89.
- Baudhuin, L.M., Cristina, K.L., Lu, J. and Xu, Y. (2002) Akt activation induced by lysophosphatidic acid and sphingosine-1-phosphate requires both mitogen-activated protein kinase kinase and p38 mitogen-activated protein kinase and is cell-line specific. *Mol Pharmacol*, 62, 660-671.
- Benezra, R., Davis, R.L., Lockshon, D., Turner, D.L. and Weintraub, H. (1990) The protein Id: a negative regulator of helix-loop-helix DNA binding proteins. *Cell*, 61, 49-59.
- Benitah, S.A., Valeron, P.F., Rui, H. and Lacal, J.C. (2003) STAT5a activation mediates the epithelial to mesenchymal transition induced by oncogenic RhoA. *Mol Biol Cell*, 14, 40-53.
- Bernardo, K., Hurwitz, R., Zenk, T., Desnick, R.J., Ferlinz, K., Schuchman, E.H. and Sandhoff, K. (1995) Purification, characterization, and biosynthesis of human acid ceramidase. *J Biol Chem*, 270, 11098-11102.
- Bhowmick, N.A., Ghiassi, M., Bakin, A., Aakre, M., Lundquist, C.A., Engel, M.E., Arteaga, C.L. and Moses, H.L. (2001) Transforming growth factor-beta1 mediates epithelial to mesenchymal transdifferentiation through a RhoA-dependent mechanism. *Mol Biol Cell*, 12, 27-36.
- Billich, A., Bornancin, F., Devay, P., Mechtcheriakova, D., Urtz, N. and Baumruker, T. (2003) Phosphorylation of the immunomodulatory drug FTY720 by sphingosine kinases. *J Biol Chem*, 278, 47408-47415.
- Bindels, S., Mestdagt, M., Vandewalle, C., Jacobs, N., Volders, L., Noel, A., Roy, F.V., Berx, G., Foidart, J.M. and Gilles, C. (2006) Regulation of vimentin by SIP1 in human epithelial breast tumor cells. *Oncogene*.
- Bishop, A.L. and Hall, A. (2000) Rho GTPases and their effector proteins. *Biochem J*, 348 Pt 2, 241-255.
- Blobe, G.C., Liu, X., Fang, S.J., How, T. and Lodish, H.F. (2001) A novel mechanism for regulating transforming growth factor beta (TGF-beta) signaling. Functional modulation of type III TGF-beta receptor expression through interaction with the PDZ domain protein, GIPC. *J Biol Chem*, 276, 39608-39617.

- Blobe, G.C., Schieman, W.P. and Lodish, H.F. (2000) Role of transforming growth factor beta in human disease. *N Engl J Med*, 342, 1350-1358.
- Bolos, V., Peinado, H., Perez-Moreno, M.A., Fraga, M.F., Esteller, M. and Cano, A. (2003) The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors. *J Cell Sci*, 116, 499-511.
- Botti, C., Seregini, E., Ferrari, L., Martinetti, A. and Bombardieri, E. (1998) Immunosuppressive factors: role in cancer development and progression. *Int J Biol Markers*, 13, 51-69.
- Brew, K., Dinakarandian, D. and Nagase, H. (2000) Tissue inhibitors of metalloproteinases: evolution, structure and function. *Biochim Biophys Acta*, 1477, 267-283.
- Brinkmann, V. (2004) FTY720: mechanism of action and potential benefit in organ transplantation. *Yonsei Med J*, 45, 991-997.
- Brinkmann, V., Chen, S., Feng, L., Pinschewer, D., Nikolova, Z. and Hof, R. (2001) FTY720 alters lymphocyte homing and protects allografts without inducing general immunosuppression. *Transplant Proc*, 33, 530-531.
- Brinkmann, V., Davis, M.D., Heise, C.E., Albert, R., Cottens, S., Hof, R., Bruns, C., Prieschl, E., Baumruker, T., Hiestand, P., Foster, C.A., Zollinger, M. and Lynch, K.R. (2002) The immune modulator FTY720 targets sphingosine 1-phosphate receptors. *J Biol Chem*, 277, 21453-21457.
- Brinkmann, V., Pinschewer, D., Chiba, K. and Feng, L. (2000) FTY720: a novel transplantation drug that modulates lymphocyte traffic rather than activation. *Trends Pharmacol Sci*, 21, 49-52.
- Buisson-Legendre, N., Bernard, P., Bobichon, H., Emonard, H., Schneider, C., Maquart, F.X., Hays, B. and Hornebeck, W. (1999) Involvement of the 92-kDa gelatinase (matrix metalloproteinase-9) in the ceramide-mediated inhibition of human keratinocyte growth. *Biochem Biophys Res Commun*, 260, 634-640.
- Burridge, K., Fath, K., Kelly, T., Nuckolls, G. and Turner, C. (1988) Focal adhesions: transmembrane junctions between the extracellular matrix and the cytoskeleton. *Annu Rev Cell Biol*, 4, 487-525.
- Chacko, B.M., Qin, B., Correia, J.J., Lam, S.S., de Caestecker, M.P. and Lin, K. (2001) The L3 loop and C-terminal phosphorylation jointly define Smad protein trimerization. *Nat Struct Biol*, 8, 248-253.
- Chakraborti, S., Mandal, M., Das, S., Mandal, A. and Chakraborti, T. (2003) Regulation of matrix metalloproteinases: an overview. *Mol Cell Biochem*, 253, 269-285.
- Chen, L., Klass, C. and Woods, A. (2004) Syndecan-2 regulates transforming growth factor-beta signaling. *J Biol Chem*, 279, 15715-15718.
- Chua, C.W., Lee, D.T., Ling, M.T., Zhou, C., Man, K., Ho, J., Chan, F.L., Wang, X. and Wong, Y.C. (2005) FTY720, a fungus metabolite, inhibits in vivo growth of androgen-independent prostate cancer. *Int J Cancer*, 117, 1039-1048.
- Coffey, R.J., Jr., Bascom, C.C., Sipes, N.J., Graves-Deal, R., Weissman, B.E. and Moses, H.L. (1988) Selective inhibition of growth-related gene expression in murine keratinocytes by transforming growth factor beta. *Mol Cell Biol*, 8, 3088-3093.

- Comijn, J., Berx, G., Vermassen, P., Verschuere, K., van Grunsven, L., Bruyneel, E., Mareel, M., Huylebroeck, D. and van Roy, F. (2001) The two-handed E box binding zinc finger protein SIP1 downregulates E-cadherin and induces invasion. *Mol Cell*, 7, 1267-1278.
- Cui, W., Fowles, D.J., Bryson, S., Duffie, E., Ireland, H., Balmain, A. and Akhurst, R.J. (1996) TGFbeta1 inhibits the formation of benign skin tumors, but enhances progression to invasive spindle carcinomas in transgenic mice. *Cell*, 86, 531-542.
- Curran, S. and Murray, G.I. (1999) Matrix metalloproteinases in tumour invasion and metastasis. *J Pathol*, 189, 300-308.
- Datto, M.B., Frederick, J.P., Pan, L., Borton, A.J., Zhuang, Y. and Wang, X.F. (1999) Targeted disruption of Smad3 reveals an essential role in transforming growth factor beta-mediated signal transduction. *Mol Cell Biol*, 19, 2495-2504.
- Datto, M.B., Li, Y., Panus, J.F., Howe, D.J., Xiong, Y. and Wang, X.F. (1995) Transforming growth factor beta induces the cyclin-dependent kinase inhibitor p21 through a p53-independent mechanism. *Proc Natl Acad Sci U S A*, 92, 5545-5549.
- Davaille, J., Li, L., Mallat, A. and Lotersztajn, S. (2002) Sphingosine 1-phosphate triggers both apoptotic and survival signals for human hepatic myofibroblasts. *J Biol Chem*, 277, 37323-37330.
- Davies, G., Jiang, W.G. and Mason, M.D. (2001) Matrilysin mediates extracellular cleavage of E-cadherin from prostate cancer cells: a key mechanism in hepatocyte growth factor/scatter factor-induced cell-cell dissociation and in vitro invasion. *Clin Cancer Res*, 7, 3289-3297.
- Davies, M., Robinson, M., Smith, E., Huntley, S., Prime, S. and Paterson, I. (2005) Induction of an epithelial to mesenchymal transition in human immortal and malignant keratinocytes by TGF-beta1 involves MAPK, Smad and AP-1 signalling pathways. *J Cell Biochem*, 95, 918-931.
- de Caestecker, M.P., Parks, W.T., Frank, C.J., Castagnino, P., Bottaro, D.P., Roberts, A.B. and Lechleider, R.J. (1998) Smad2 transduces common signals from receptor serine-threonine and tyrosine kinases. *Genes Dev*, 12, 1587-1592.
- Derynck, R. and Feng, X.H. (1997) TGF-beta receptor signaling. *Biochim Biophys Acta*, 1333, F105-150.
- Derynck, R. and Zhang, Y.E. (2003) Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature*, 425, 577-584.
- Di Guglielmo, G.M., Le Roy, C., Goodfellow, A.F. and Wrana, J.L. (2003) Distinct endocytic pathways regulate TGF-beta receptor signalling and turnover. *Nat Cell Biol*, 5, 410-421.
- Doll, F., Pfeilschifter, J. and Huwiler, A. (2005) The epidermal growth factor stimulates sphingosine kinase-1 expression and activity in the human mammary carcinoma cell line MCF7. *Biochim Biophys Acta*, 1738, 72-81.
- Dorsett, Y. and Tuschl, T. (2004) siRNAs: applications in functional genomics and potential as therapeutics. *Nat Rev Drug Discov*, 3, 318-329.
- Downward, J. (1998) Ras signalling and apoptosis. *Curr Opin Genet Dev*, 8, 49-54.
- Dudley, D.T., Pang, L., Decker, S.J., Bridges, A.J. and Saltiel, A.R. (1995) A synthetic inhibitor of the mitogen-activated protein kinase cascade. *Proc Natl Acad Sci U S A*, 92, 7686-7689.

- Edlund, S., Landstrom, M., Heldin, C.H. and Aspenstrom, P. (2002) Transforming growth factor-beta-induced mobilization of actin cytoskeleton requires signaling by small GTPases Cdc42 and RhoA. *Mol Biol Cell*, 13, 902-914.
- Edsall, L.C., Van Brocklyn, J.R., Cuvillier, O., Kleuser, B. and Spiegel, S. (1998) N,N-Dimethylsphingosine is a potent competitive inhibitor of sphingosine kinase but not of protein kinase C: modulation of cellular levels of sphingosine 1-phosphate and ceramide. *Biochemistry*, 37, 12892-12898.
- Edwards, D.R., Beaudry, P.P., Laing, T.D., Kowal, V., Leco, K.J., Leco, P.A. and Lim, M.S. (1996) The roles of tissue inhibitors of metalloproteinases in tissue remodelling and cell growth. *Int J Obes Relat Metab Disord*, 20 Suppl 3, S9-15.
- Eger, A., Aigner, K., Sonderegger, S., Dampier, B., Oehler, S., Schreiber, M., Berx, G., Cano, A., Beug, H. and Foisner, R. (2005) DeltaEF1 is a transcriptional repressor of E-cadherin and regulates epithelial plasticity in breast cancer cells. *Oncogene*, 24, 2375-2385.
- Eger, A., Stockinger, A., Park, J., Langkopf, E., Mikula, M., Gotzmann, J., Mikulits, W., Beug, H. and Foisner, R. (2004) beta-Catenin and TGFbeta signalling cooperate to maintain a mesenchymal phenotype after FosER-induced epithelial to mesenchymal transition. *Oncogene*, 23, 2672-2680.
- El Bawab, S., Roddy, P., Qian, T., Bielawska, A., Lemasters, J.J. and Hannun, Y.A. (2000) Molecular cloning and characterization of a human mitochondrial ceramidase. *J Biol Chem*, 275, 21508-21513.
- Ellenrieder, V., Hendler, S.F., Boeck, W., Seufferlein, T., Menke, A., Ruhland, C., Adler, G. and Gress, T.M. (2001) Transforming growth factor beta1 treatment leads to an epithelial-mesenchymal transdifferentiation of pancreatic cancer cells requiring extracellular signal-regulated kinase 2 activation. *Cancer Res*, 61, 4222-4228.
- Endo, A., Nagashima, K., Kurose, H., Mochizuki, S., Matsuda, M. and Mochizuki, N. (2002) Sphingosine 1-phosphate induces membrane ruffling and increases motility of human umbilical vein endothelial cells via vascular endothelial growth factor receptor and CrkII. *J Biol Chem*, 277, 23747-23754.
- Engel, M.E., McDonnell, M.A., Law, B.K. and Moses, H.L. (1999) Interdependent SMAD and JNK signaling in transforming growth factor-beta-mediated transcription. *J Biol Chem*, 274, 37413-37420.
- Feng, X.H. and Derynck, R. (2005) Specificity and versatility in tgf-beta signaling through Smads. *Annu Rev Cell Dev Biol*, 21, 659-693.
- Flanders, K.C. (2004) Smad3 as a mediator of the fibrotic response. *Int J Exp Pathol*, 85, 47-64.
- Fowlkes, J.L., Enghild, J.J., Suzuki, K. and Nagase, H. (1994) Matrix metalloproteinases degrade insulin-like growth factor-binding protein-3 in dermal fibroblast cultures. *J Biol Chem*, 269, 25742-25746.
- Franz, T.J. (1975) Percutaneous absorption on the relevance of in vitro data. *J Invest Dermatol*, 64, 190-195.
- French, K.J., Upson, J.J., Keller, S.N., Zhuang, Y., Yun, J.K. and Smith, C.D. (2006) Antitumor Activity of Sphingosine Kinase Inhibitors. *J Pharmacol Exp Ther*.

- Funaba, M., Zimmerman, C.M. and Mathews, L.S. (2002) Modulation of Smad2-mediated signaling by extracellular signal-regulated kinase. *J Biol Chem*, 277, 41361-41368.
- Gilles, C., Polette, M., Piette, J., Birembaut, P. and Foidart, J.M. (1994) Epithelial-to-mesenchymal transition in HPV-33-transfected cervical keratinocytes is associated with increased invasiveness and expression of gelatinase A. *Int J Cancer*, 59, 661-666.
- Gon, Y., Wood, M.R., Kiosses, W.B., Jo, E., Sanna, M.G., Chun, J. and Rosen, H. (2005) S1P3 receptor-induced reorganization of epithelial tight junctions compromises lung barrier integrity and is potentiated by TNF. *Proc Natl Acad Sci U S A*, 102, 9270-9275.
- Goodman, L.V. and Majack, R.A. (1989) Vascular smooth muscle cells express distinct transforming growth factor-beta receptor phenotypes as a function of cell density in culture. *J Biol Chem*, 264, 5241-5244.
- Gotzmann, J., Huber, H., Thallinger, C., Wolschek, M., Jansen, B., Schulte-Hermann, R., Beug, H. and Mikulits, W. (2002) Hepatocytes convert to a fibroblastoid phenotype through the cooperation of TGF-beta1 and Ha-Ras: steps towards invasiveness. *J Cell Sci*, 115, 1189-1202.
- Goumans, M.J., Valdimarsdottir, G., Itoh, S., Rosendahl, A., Sideras, P. and ten Dijke, P. (2002) Balancing the activation state of the endothelium via two distinct TGF-beta type I receptors. *Embo J*, 21, 1743-1753.
- Graler, M.H. and Goetzl, E.J. (2004) The immunosuppressant FTY720 down-regulates sphingosine 1-phosphate G-protein-coupled receptors. *Faseb J*, 18, 551-553.
- Greenburg, G. and Hay, E.D. (1982) Epithelia suspended in collagen gels can lose polarity and express characteristics of migrating mesenchymal cells. *J Cell Biol*, 95, 333-339.
- Greene, R.M., Nugent, P., Mukhopadhyay, P., Warner, D.R. and Pisano, M.M. (2003) Intracellular dynamics of Smad-mediated TGFbeta signaling. *J Cell Physiol*, 197, 261-271.
- Grunert, S., Jechlinger, M. and Beug, H. (2003) Diverse cellular and molecular mechanisms contribute to epithelial plasticity and metastasis. *Nat Rev Mol Cell Biol*, 4, 657-665.
- Hait, N.C., Sarkar, S., Le Stunff, H., Mikami, A., Maceyka, M., Milstien, S. and Spiegel, S. (2005) Role of sphingosine kinase 2 in cell migration toward epidermal growth factor. *J Biol Chem*, 280, 29462-29469.
- Hammer, S., Sauer, B., Spika, I., Schraut, C., Kleuser, B. and Schafer-Korting, M. (2004) Glucocorticoids mediate differential anti-apoptotic effects in human fibroblasts and keratinocytes via sphingosine-1-phosphate formation. *J Cell Biochem*, 91, 840-851.
- Han, G., Lu, S.L., Li, A.G., He, W., Corless, C.L., Kulesz-Martin, M. and Wang, X.J. (2005) Distinct mechanisms of TGF-beta1-mediated epithelial-to-mesenchymal transition and metastasis during skin carcinogenesis. *J Clin Invest*, 115, 1714-1723.
- Hanahan, D. and Weinberg, R.A. (2000) The hallmarks of cancer. *Cell*, 100, 57-70.
- Hannon, G.J. and Beach, D. (1994) p15INK4B is a potential effector of TGF-beta-induced cell cycle arrest. *Nature*, 371, 257-261.
- Hanyu, A., Ishidou, Y., Ebisawa, T., Shimanuki, T., Imamura, T. and Miyazono, K. (2001) The N domain of Smad7 is essential for specific inhibition of transforming growth factor-beta signaling. *J Cell Biol*, 155, 1017-1027.

- Hata, A., Shi, Y. and Massague, J. (1998) TGF-beta signaling and cancer: structural and functional consequences of mutations in Smads. *Mol Med Today*, 4, 257-262.
- Hennings, H., Michael, D., Cheng, C., Steinert, P., Holbrook, K. and Yuspa, S.H. (1980) Calcium regulation of growth and differentiation of mouse epidermal cells in culture. *Cell*, 19, 245-254.
- Hla, T., Lee, M.J., Ancellin, N., Paik, J.H. and Kluk, M.J. (2001) Lysophospholipids--receptor revelations. *Science*, 294, 1875-1878.
- Hla, T. and Maciag, T. (1990) An abundant transcript induced in differentiating human endothelial cells encodes a polypeptide with structural similarities to G-protein-coupled receptors. *J Biol Chem*, 265, 9308-9313.
- Huber, M.A., Azoitei, N., Baumann, B., Grunert, S., Sommer, A., Pehamberger, H., Kraut, N., Beug, H. and Wirth, T. (2004) NF-kappaB is essential for epithelial-mesenchymal transition and metastasis in a model of breast cancer progression. *J Clin Invest*, 114, 569-581.
- Hung, R.J., Hsu Ia, W., Dreiling, J.L., Lee, M.J., Williams, C.A., Oberst, M.D., Dickson, R.B. and Lin, C.Y. (2004) Assembly of adherens junctions is required for sphingosine 1-phosphate-induced matriptase accumulation and activation at mammary epithelial cell-cell contacts. *Am J Physiol Cell Physiol*, 286, C1159-1169.
- Huntley, S.P., Davies, M., Matthews, J.B., Thomas, G., Marshall, J., Robinson, C.M., Eveson, J.W., Paterson, I.C. and Prime, S.S. (2004) Attenuated type II TGF-beta receptor signalling in human malignant oral keratinocytes induces a less differentiated and more aggressive phenotype that is associated with metastatic dissemination. *Int J Cancer*, 110, 170-176.
- Idzko, M., Panther, E., Corinti, S., Morelli, A., Ferrari, D., Herouy, Y., Dichmann, S., Mockenhaupt, M., Gebicke-Haerter, P., Di Virgilio, F., Girolomoni, G. and Norgauer, J. (2002) Sphingosine 1-phosphate induces chemotaxis of immature and modulates cytokine-release in mature human dendritic cells for emergence of Th2 immune responses. *Faseb J*, 16, 625-627.
- Ignatov, A., Lintzel, J., Hermans-Borgmeyer, I., Kreienkamp, H.J., Joost, P., Thomsen, S., Methner, A. and Schaller, H.C. (2003a) Role of the G-protein-coupled receptor GPR12 as high-affinity receptor for sphingosylphosphorylcholine and its expression and function in brain development. *J Neurosci*, 23, 907-914.
- Ignatov, A., Lintzel, J., Kreienkamp, H.J. and Schaller, H.C. (2003b) Sphingosine-1-phosphate is a high-affinity ligand for the G protein-coupled receptor GPR6 from mouse and induces intracellular Ca<sup>2+</sup> release by activating the sphingosine-kinase pathway. *Biochem Biophys Res Commun*, 311, 329-336.
- Imai, K., Hiramatsu, A., Fukushima, D., Pierschbacher, M.D. and Okada, Y. (1997) Degradation of decorin by matrix metalloproteinases: identification of the cleavage sites, kinetic analyses and transforming growth factor-beta1 release. *Biochem J*, 322 (Pt 3), 809-814.
- Inman, G.J., Nicolas, F.J., Callahan, J.F., Harling, J.D., Gaster, L.M., Reith, A.D., Laping, N.J. and Hill, C.S. (2002) SB-431542 is a potent and specific inhibitor of transforming growth factor-beta superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5, and ALK7. *Mol Pharmacol*, 62, 65-74.
- Inohara, S., Kitano, Y. and Kitagawa, K. (1995) Cell cycle regulators in the keratinocyte (cyclin-cdk). *Exp Dermatol*, 4, 1-8.



- Itagaki, K. and Hauser, C.J. (2003) Sphingosine 1-phosphate, a diffusible calcium influx factor mediating store-operated calcium entry. *J Biol Chem*, 278, 27540-27547.
- Itoh, F., Asao, H., Sugamura, K., Heldin, C.H., ten Dijke, P. and Itoh, S. (2001) Promoting bone morphogenetic protein signaling through negative regulation of inhibitory Smads. *Embo J*, 20, 4132-4142.
- Itoh, S., Itoh, F., Goumans, M.J. and Ten Dijke, P. (2000) Signaling of transforming growth factor-beta family members through Smad proteins. *Eur J Biochem*, 267, 6954-6967.
- Itoh, S., Thorikay, M., Kowanetz, M., Moustakas, A., Itoh, F., Heldin, C.H. and ten Dijke, P. (2003) Elucidation of Smad requirement in transforming growth factor-beta type I receptor-induced responses. *J Biol Chem*, 278, 3751-3761.
- Janda, E., Lehmann, K., Killisch, I., Jechlinger, M., Herzig, M., Downward, J., Beug, H. and Grunert, S. (2002) Ras and TGF[beta] cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. *J Cell Biol*, 156, 299-313.
- Jechlinger, M., Grunert, S., Tamir, I.H., Janda, E., Ludemann, S., Waerner, T., Seither, P., Weith, A., Beug, H. and Kraut, N. (2003) Expression profiling of epithelial plasticity in tumor progression. *Oncogene*, 22, 7155-7169.
- Jeckel, D., Karrenbauer, A., Burger, K.N., van Meer, G. and Wieland, F. (1992) Glucosylceramide is synthesized at the cytosolic surface of various Golgi subfractions. *J Cell Biol*, 117, 259-267.
- Jo, E., Sanna, M.G., Gonzalez-Cabrera, P.J., Thangada, S., Tigyi, G., Osborne, D.A., Hla, T., Parrill, A.L. and Rosen, H. (2005) S1P1-selective in vivo-active agonists from high-throughput screening: off-the-shelf chemical probes of receptor interactions, signaling, and fate. *Chem Biol*, 12, 703-715.
- Jorda, M., Olmeda, D., Vinyals, A., Valero, E., Cubillo, E., Llorens, A., Cano, A. and Fabra, A. (2005) Upregulation of MMP-9 in MDCK epithelial cell line in response to expression of the Snail transcription factor. *J Cell Sci*, 118, 3371-3385.
- Kaartinen, V., Voncken, J.W., Shuler, C., Warburton, D., Bu, D., Heisterkamp, N. and Groffen, J. (1995) Abnormal lung development and cleft palate in mice lacking TGF-beta 3 indicates defects of epithelial-mesenchymal interaction. *Nat Genet*, 11, 415-421.
- Kalluri, R. and Neilson, E.G. (2003) Epithelial-mesenchymal transition and its implications for fibrosis. *J Clin Invest*, 112, 1776-1784.
- Kawamori, T., Osta, W., Johnson, K.R., Pettus, B.J., Bielawski, J., Tanaka, T., Wargovich, M.J., Reddy, B.S., Hannun, Y.A., Obeid, L.M. and Zhou, D. (2006) Sphingosine kinase 1 is up-regulated in colon carcinogenesis. *Faseb J*, 20, 386-388.
- Kim, D.S. (1999) Wirkungen von Wachstumsfaktoren und Sphingolipiden auf humane Keratinozyten und Fibroblasten in vitro (Freie Universität Berlin).
- Kim, E.S., Kim, M.S. and Moon, A. (2004) TGF-beta-induced upregulation of MMP-2 and MMP-9 depends on p38 MAPK, but not ERK signaling in MCF10A human breast epithelial cells. *Int J Oncol*, 25, 1375-1382.
- Kim, J.W., Kim, Y.W., Inagaki, Y., Hwang, Y.A., Mitsutake, S., Ryu, Y.W., Lee, W.K., Ha, H.J., Park, C.S. and Igarashi, Y. (2005) Synthesis and evaluation of sphingoid analogs as inhibitors of sphingosine kinases. *Bioorg Med Chem*, 13, 3475-3485.

- Kingsley, D.M. (1994) The TGF-beta superfamily: new members, new receptors, and new genetic tests of function in different organisms. *Genes Dev*, 8, 133-146.
- Kleuser, B., Cuvillier, O. and Spiegel, S. (1998) 1Alpha,25-dihydroxyvitamin D3 inhibits programmed cell death in HL-60 cells by activation of sphingosine kinase. *Cancer Res*, 58, 1817-1824.
- Kobayashi, T., Hattori, S., Nagai, Y., Tajima, S. and Nishikawa, T. (1998) Differential regulation of MMP-2 and MMP-9 gelatinases in cultured human keratinocytes. *Dermatology*, 197, 1-5.
- Kolesnick, R.N. and Hemer, M.R. (1990) Characterization of a ceramide kinase activity from human leukemia (HL-60) cells. Separation from diacylglycerol kinase activity. *J Biol Chem*, 265, 18803-18808.
- Kondo, M., Cubillo, E., Tobiume, K., Shirakihara, T., Fukuda, N., Suzuki, H., Shimizu, K., Takehara, K., Cano, A., Saitoh, M. and Miyazono, K. (2004) A role for Id in the regulation of TGF-beta-induced epithelial-mesenchymal transdifferentiation. *Cell Death Differ*, 11, 1092-1101.
- Kowanetz, M., Valcourt, U., Bergstrom, R., Heldin, C.H. and Moustakas, A. (2004) Id2 and Id3 define the potency of cell proliferation and differentiation responses to transforming growth factor beta and bone morphogenetic protein. *Mol Cell Biol*, 24, 4241-4254.
- Kretschmar, M., Doody, J., Timokhina, I. and Massague, J. (1999) A mechanism of repression of TGFbeta/ Smad signaling by oncogenic Ras. *Genes Dev*, 13, 804-816.
- Kuphal, S., Palm, H.G., Poser, I. and Bosserhoff, A.K. (2005) Snail-regulated genes in malignant melanoma. *Melanoma Res*, 15, 305-313.
- Kupperman, E., An, S., Osborne, N., Waldron, S. and Stainier, D.Y. (2000) A sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature*, 406, 192-195.
- Kurahara, S., Shinohara, M., Ikebe, T., Nakamura, S., Beppu, M., Hiraki, A., Takeuchi, H. and Shirasuna, K. (1999) Expression of MMPS, MT-MMP, and TIMPs in squamous cell carcinoma of the oral cavity: correlations with tumor invasion and metastasis. *Head Neck*, 21, 627-638.
- Kwon, Y.G., Min, J.K., Kim, K.M., Lee, D.J., Billiar, T.R. and Kim, Y.M. (2001) Sphingosine 1-phosphate protects human umbilical vein endothelial cells from serum-deprived apoptosis by nitric oxide production. *J Biol Chem*, 276, 10627-10633.
- LaGamba, D., Nawshad, A. and Hay, E.D. (2005) Microarray analysis of gene expression during epithelial-mesenchymal transformation. *Dev Dyn*, 234, 132-142.
- Lai, Y.T., Beason, K.B., Brames, G.P., Desgrosellier, J.S., Cleggett, M.C., Shaw, M.V., Brown, C.B. and Barnett, J.V. (2000) Activin receptor-like kinase 2 can mediate atrioventricular cushion transformation. *Dev Biol*, 222, 1-11.
- LaMontagne, K., Littlewood-Evans, A., Schnell, C., O'Reilly, T., Wyder, L., Sanchez, T., Probst, B., Butler, J., Wood, A., Liau, G., Billy, E., Theuer, A., Hla, T. and Wood, J. (2006) Antagonism of sphingosine-1-phosphate receptors by FTY720 inhibits angiogenesis and tumor vascularization. *Cancer Res*, 66, 221-231.
- Laping, N.J., Grygielko, E., Mathur, A., Butter, S., Bomberger, J., Tweed, C., Martin, W., Fornwald, J., Lehr, R., Harling, J., Gaster, L., Callahan, J.F. and Olson, B.A. (2002) Inhibition of transforming growth factor (TGF)-beta1-induced extracellular matrix with a novel inhibitor of the TGF-beta type I receptor kinase activity: SB-431542. *Mol Pharmacol*, 62, 58-64.

- Lauffenburger, D.A. and Horwitz, A.F. (1996) Cell migration: a physically integrated molecular process. *Cell*, 84, 359-369.
- Le Stunff, H., Milstien, S. and Spiegel, S. (2004) Generation and metabolism of bioactive sphingosine-1-phosphate. *J Cell Biochem*, 92, 882-899.
- Lebrin, F., Goumans, M.J., Jonker, L., Carvalho, R.L., Valdimarsdottir, G., Thorikay, M., Mummery, C., Arthur, H.M. and ten Dijke, P. (2004) Endoglin promotes endothelial cell proliferation and TGF-beta/ALK1 signal transduction. *Embo J*, 23, 4018-4028.
- Lee, C.W., Rivera, R., Gardell, S., Dubin, A.E. and Chun, J. (2006) GPR92 as a new G12/13 and Gq coupled lysophosphatidic acid receptor that increases cAMP: LPA5. *J Biol Chem*.
- Lee, M.J., Thangada, S., Claffey, K.P., Ancellin, N., Liu, C.H., Kluk, M., Volpi, M., Sha'afi, R.I. and Hla, T. (1999) Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell*, 99, 301-312.
- Lee, M.J., Thangada, S., Paik, J.H., Sapkota, G.P., Ancellin, N., Chae, S.S., Wu, M., Morales-Ruiz, M., Sessa, W.C., Alessi, D.R. and Hla, T. (2001) Akt-mediated phosphorylation of the G protein-coupled receptor EDG-1 is required for endothelial cell chemotaxis. *Mol Cell*, 8, 693-704.
- Lee, M.J., Van Brocklyn, J.R., Thangada, S., Liu, C.H., Hand, A.R., Menzeleev, R., Spiegel, S. and Hla, T. (1998) Sphingosine-1-phosphate as a ligand for the G protein-coupled receptor EDG-1. *Science*, 279, 1552-1555.
- Lehmann, K., Janda, E., Pierreux, C.E., Rytomaa, M., Schulze, A., McMahon, M., Hill, C.S., Beug, H. and Downward, J. (2000) Raf induces TGFbeta production while blocking its apoptotic but not invasive responses: a mechanism leading to increased malignancy in epithelial cells. *Genes Dev*, 14, 2610-2622.
- Lemaitre, V. and D'Armiento, J. (2006) Matrix metalloproteinases in development and disease. *Birth Defects Res C Embryo Today*, 78, 1-10.
- Letamendia, A., Lastres, P., Botella, L.M., Raab, U., Langa, C., Velasco, B., Attisano, L. and Bernabeu, C. (1998) Role of endoglin in cellular responses to transforming growth factor-beta. A comparative study with betaglycan. *J Biol Chem*, 273, 33011-33019.
- Letterio, J.J. and Roberts, A.B. (1998) Regulation of immune responses by TGF-beta. *Annu Rev Immunol*, 16, 137-161.
- Levade, T. and Jaffrezou, J.P. (1999) Signalling sphingomyelinases: which, where, how and why? *Biochim Biophys Acta*, 1438, 1-17.
- Li, J.H., Zhu, H.J., Huang, X.R., Lai, K.N., Johnson, R.J. and Lan, H.Y. (2002) Smad7 inhibits fibrotic effect of TGF-Beta on renal tubular epithelial cells by blocking Smad2 activation. *J Am Soc Nephrol*, 13, 1464-1472.
- Lillie, F.R. (1908) *The Development of the Chick*, Henry Holt and Co, New York.
- Liu, B., Andrieu-Abadie, N., Levade, T., Zhang, P., Obeid, L.M. and Hannun, Y.A. (1998) Glutathione regulation of neutral sphingomyelinase in tumor necrosis factor-alpha-induced cell death. *J Biol Chem*, 273, 11313-11320.
- Liu, B. and Hannun, Y.A. (1997) Inhibition of the neutral magnesium-dependent sphingomyelinase by glutathione. *J Biol Chem*, 272, 16281-16287.

- Liu, G., Zhang, X., Lin, H., Wang, H., Li, Q., Ni, J. and Zhu, C. (2006) Effects of E-cadherin on mouse embryo implantation and expression of matrix metalloproteinase-2 and -9. *Biochem Biophys Res Commun*, 343, 832-838.
- Liu, Y., Wada, R., Yamashita, T., Mi, Y., Deng, C.X., Hobson, J.P., Rosenfeldt, H.M., Nava, V.E., Chae, S.S., Lee, M.J., Liu, C.H., Hla, T., Spiegel, S. and Proia, R.L. (2000) Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation. *J Clin Invest*, 106, 951-961.
- Lobmann, R., Ambrosch, A., Schultz, G., Waldmann, K., Schiweck, S. and Lehnert, H. (2002) Expression of matrix-metalloproteinases and their inhibitors in the wounds of diabetic and non-diabetic patients. *Diabetologia*, 45, 1011-1016.
- Lopez-Casillas, F., Wrana, J.L. and Massague, J. (1993) Betaglycan presents ligand to the TGF beta signaling receptor. *Cell*, 73, 1435-1444.
- Maddala, R., Reddy, V.N., Epstein, D.L. and Rao, V. (2003) Growth factor induced activation of Rho and Rac GTPases and actin cytoskeletal reorganization in human lens epithelial cells. *Mol Vis*, 9, 329-336.
- Magnusson, M.K. and Mosher, D.F. (1998) Fibronectin: structure, assembly, and cardiovascular implications. *Arterioscler Thromb Vasc Biol*, 18, 1363-1370.
- Majno, G., Gabbiani, G., Hirschel, B.J., Ryan, G.B. and Statkov, P.R. (1971) Contraction of granulation tissue in vitro: similarity to smooth muscle. *Science*, 173, 548-550.
- Makela, M., Larjava, H., Pirila, E., Maisi, P., Salo, T., Sorsa, T. and Uitto, V.J. (1999) Matrix metalloproteinase 2 (gelatinase A) is related to migration of keratinocytes. *Exp Cell Res*, 251, 67-78.
- Mandala, S.M., Thornton, R., Galve-Roperh, I., Poulton, S., Peterson, C., Olivera, A., Bergstrom, J., Kurtz, M.B. and Spiegel, S. (2000) Molecular cloning and characterization of a lipid phosphohydrolase that degrades sphingosine-1-phosphate and induces cell death. *Proc Natl Acad Sci U S A*, 97, 7859-7864.
- Manggau, M. (2002) Untersuchungen zu Signalwegen von SPP in humanen Keratinozyten (Freie Universität Berlin).
- Manggau, M., Kim, D.S., Ruwisch, L., Vogler, R., Korting, H.C., Schafer-Korting, M. and Kleuser, B. (2001) 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> protects human keratinocytes from apoptosis by the formation of sphingosine-1-phosphate. *J Invest Dermatol*, 117, 1241-1249.
- Maschler, S., Wirl, G., Spring, H., Bredow, D.V., Sordat, I., Beug, H. and Reichmann, E. (2005) Tumor cell invasiveness correlates with changes in integrin expression and localization. *Oncogene*, 24, 2032-2041.
- Massague, J. (1990) The transforming growth factor-beta family. *Annu Rev Cell Biol*, 6, 597-641.
- Massague, J., Blain, S.W. and Lo, R.S. (2000) TGFbeta signaling in growth control, cancer, and heritable disorders. *Cell*, 103, 295-309.
- Masson, V., de la Ballina, L.R., Munaut, C., Wielockx, B., Jost, M., Maillard, C., Blacher, S., Bajou, K., Itoh, T., Itohara, S., Werb, Z., Libert, C., Foidart, J.M. and Noel, A. (2005) Contribution of host MMP-2 and MMP-9 to promote tumor vascularization and invasion of malignant keratinocytes. *Faseb J*, 19, 234-236.

- Masszi, A., Fan, L., Rosivall, L., McCulloch, C.A., Rotstein, O.D., Mucsi, I. and Kapus, A. (2004) Integrity of cell-cell contacts is a critical regulator of TGF-beta 1-induced epithelial-to-myofibroblast transition: role for beta-catenin. *Am J Pathol*, 165, 1955-1967.
- Matloubian, M., Lo, C.G., Cinamon, G., Lesneski, M.J., Xu, Y., Brinkmann, V., Allende, M.L., Proia, R.L. and Cyster, J.G. (2004) Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature*, 427, 355-360.
- Mazars, A., Lallemand, F., Prunier, C., Marais, J., Ferrand, N., Pessah, M., Cherqui, G. and Atfi, A. (2001) Evidence for a role of the JNK cascade in Smad7-mediated apoptosis. *J Biol Chem*, 276, 36797-36803.
- McCawley, L.J., O'Brien, P. and Hudson, L.G. (1998) Epidermal growth factor (EGF)- and scatter factor/hepatocyte growth factor (SF/HGF)- mediated keratinocyte migration is coincident with induction of matrix metalloproteinase (MMP)-9. *J Cell Physiol*, 176, 255-265.
- Mehnert, W. and Mader, K. (2001) Solid lipid nanoparticles: production, characterization and applications. *Adv Drug Deliv Rev*, 47, 165-196.
- Mei, J.M., Borchert, G.L., Donald, S.P. and Phang, J.M. (2002) Matrix metalloproteinase(s) mediate(s) NO-induced dissociation of beta-catenin from membrane bound E-cadherin and formation of nuclear beta-catenin/LEF-1 complex. *Carcinogenesis*, 23, 2119-2122.
- Meyer Zu Heringdorf, D. (2004) Lysophospholipid receptor-dependent and -independent calcium signaling. *J Cell Biochem*, 92, 937-948.
- Miettinen, P.J., Ebner, R., Lopez, A.R. and Derynck, R. (1994) TGF-beta induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J Cell Biol*, 127, 2021-2036.
- Mignatti, P. and Rifkin, D.B. (1993) Biology and biochemistry of proteinases in tumor invasion. *Physiol Rev*, 73, 161-195.
- Mitchison, T.J. and Cramer, L.P. (1996) Actin-based cell motility and cell locomotion. *Cell*, 84, 371-379.
- Morales-Ruiz, M., Lee, M.J., Zollner, S., Gratton, J.P., Scotland, R., Shiojima, I., Walsh, K., Hla, T. and Sessa, W.C. (2001) Sphingosine 1-phosphate activates Akt, nitric oxide production, and chemotaxis through a Gi protein/phosphoinositide 3-kinase pathway in endothelial cells. *J Biol Chem*, 276, 19672-19677.
- Morali, O.G., Jouneau, A., McLaughlin, K.J., Thiery, J.P. and Larue, L. (2000) IGF-II promotes mesoderm formation. *Dev Biol*, 227, 133-145.
- Muller, N., Reinacher-Schick, A., Baldus, S., van Hengel, J., Berx, G., Baar, A., van Roy, F., Schmiegel, W. and Schwarte-Waldhoff, I. (2002) Smad4 induces the tumor suppressor E-cadherin and P-cadherin in colon carcinoma cells. *Oncogene*, 21, 6049-6058.
- Müller, R.H., Mehnert, W., Lucks, J.S., Schwarz, C., Mühlen, A.z., H.Weyers, Freytas, C. and Rühl, D. (1995) Solid lipid nanoparticles (SLN)- an alternative colloidal drug carrier system for controlled drug delivery. *Eur J Biopharm*, 41, 62-69.
- Müller, R.H., Rühl, D., Runge, S., Schulze-Forster, K. and Mehnert, W. (1997) Cytotoxicity of solid lipid nanoparticles as a function of the lipid matrix and the surfactant. *Pharm Res*, 14, 458-462.

- Munster, U., Nakamura, C., Haberland, A., Jores, K., Mehnert, W., Rummel, S., Schaller, M., Korting, H.C., Zouboulis Ch, C., Blume-Peytavi, U. and Schafer-Korting, M. (2005) RU 58841-myristate--prodrug development for topical treatment of acne and androgenetic alopecia. *Pharmazie*, 60, 8-12.
- Murayama, T. and Ui, M. (1983) Loss of the inhibitory function of the guanine nucleotide regulatory component of adenylate cyclase due to its ADP ribosylation by islet-activating protein, pertussis toxin, in adipocyte membranes. *J Biol Chem*, 258, 3319-3326.
- Myeroff, L.L., Parsons, R., Kim, S.J., Hedrick, L., Cho, K.R., Orth, K., Mathis, M., Kinzler, K.W., Lutterbaugh, J., Park, K. and et al. (1995) A transforming growth factor beta receptor type II gene mutation common in colon and gastric but rare in endometrial cancers with microsatellite instability. *Cancer Res*, 55, 5545-5547.
- Nakao, A., Afrakhte, M., Moren, A., Nakayama, T., Christian, J.L., Heuchel, R., Itoh, S., Kawabata, M., Heldin, N.E., Heldin, C.H. and ten Dijke, P. (1997) Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. *Nature*, 389, 631-635.
- Naldini, L., Weidner, K.M., Vigna, E., Gaudino, G., Bardelli, A., Ponzetto, C., Narsimhan, R.P., Hartmann, G., Zarnegar, R., Michalopoulos, G.K. and et al. (1991) Scatter factor and hepatocyte growth factor are indistinguishable ligands for the MET receptor. *Embo J*, 10, 2867-2878.
- Nawrocki-Raby, B., Gilles, C., Polette, M., Bruyneel, E., Laronze, J.Y., Bonnet, N., Foidart, J.M., Mareel, M. and Birembaut, P. (2003a) Upregulation of MMPs by soluble E-cadherin in human lung tumor cells. *Int J Cancer*, 105, 790-795.
- Nawrocki-Raby, B., Gilles, C., Polette, M., Martinella-Catusse, C., Bonnet, N., Puchelle, E., Foidart, J.M., Van Roy, F. and Birembaut, P. (2003b) E-Cadherin mediates MMP down-regulation in highly invasive bronchial tumor cells. *Am J Pathol*, 163, 653-661.
- Nawshad, A., Lagamba, D., Polad, A. and Hay, E.D. (2005) Transforming growth factor-beta signaling during epithelial-mesenchymal transformation: implications for embryogenesis and tumor metastasis. *Cells Tissues Organs*, 179, 11-23.
- Nobes, C.D. and Hall, A. (1995) Rho, rac, and cdc42 GTPases regulate the assembly of multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell*, 81, 53-62.
- Noe, V., Fingleton, B., Jacobs, K., Crawford, H.C., Vermeulen, S., Steelant, W., Bruyneel, E., Matrisian, L.M. and Mareel, M. (2001) Release of an invasion promoter E-cadherin fragment by matrilysin and stromelysin-1. *J Cell Sci*, 114, 111-118.
- Noguchi, K., Ishii, S. and Shimizu, T. (2003) Identification of p2y9/GPR23 as a novel G protein-coupled receptor for lysophosphatidic acid, structurally distant from the Edg family. *J Biol Chem*, 278, 25600-25606.
- O'Kane, S. and Ferguson, M.W. (1997) Transforming growth factor beta s and wound healing. *Int J Biochem Cell Biol*, 29, 63-78.
- O'Toole, E.A. (2001) Extracellular matrix and keratinocyte migration. *Clin Exp Dermatol*, 26, 525-530.

- O'Toole, T.E., Katagiri, Y., Faull, R.J., Peter, K., Tamura, R., Quaranta, V., Loftus, J.C., Shattil, S.J. and Ginsberg, M.H. (1994) Integrin cytoplasmic domains mediate inside-out signal transduction. *J Cell Biol*, 124, 1047-1059.
- Oft, M., Akhurst, R.J. and Balmain, A. (2002) Metastasis is driven by sequential elevation of H-ras and Smad2 levels. *Nat Cell Biol*, 4, 487-494.
- Oft, M., Heider, K.H. and Beug, H. (1998) TGFbeta signaling is necessary for carcinoma cell invasiveness and metastasis. *Curr Biol*, 8, 1243-1252.
- Okamoto, H., Takuwa, N., Yokomizo, T., Sugimoto, N., Sakurada, S., Shigematsu, H. and Takuwa, Y. (2000) Inhibitory regulation of Rac activation, membrane ruffling, and cell migration by the G protein-coupled sphingosine-1-phosphate receptor EDG5 but not EDG1 or EDG3. *Mol Cell Biol*, 20, 9247-9261.
- Olivera, A., Rosenfeldt, H.M., Bektas, M., Wang, F., Ishii, I., Chun, J., Milstien, S. and Spiegel, S. (2003) Sphingosine kinase type 1 induces G12/13-mediated stress fiber formation, yet promotes growth and survival independent of G protein-coupled receptors. *J Biol Chem*, 278, 46452-46460.
- Olivera, A. and Spiegel, S. (1993) Sphingosine-1-phosphate as second messenger in cell proliferation induced by PDGF and FCS mitogens. *Nature*, 365, 557-560.
- Orlati, S., Porcelli, A.M., Hrelia, S. and Rugolo, M. (1998) Sphingosylphosphorylcholine and sphingosine-1-phosphate mobilize cytosolic calcium through different mechanisms in human airway epithelial cells. *Cell Calcium*, 23, 387-394.
- Osada, M., Yatomi, Y., Ohmori, T., Ikeda, H. and Ozaki, Y. (2002) Enhancement of sphingosine 1-phosphate-induced migration of vascular endothelial cells and smooth muscle cells by an EDG-5 antagonist. *Biochem Biophys Res Commun*, 299, 483-487.
- Paik, J.H., Chae, S., Lee, M.J., Thangada, S. and Hla, T. (2001) Sphingosine 1-phosphate-induced endothelial cell migration requires the expression of EDG-1 and EDG-3 receptors and Rho-dependent activation of alpha vbeta3- and beta1-containing integrins. *J Biol Chem*, 276, 11830-11837.
- Paik, J.H., Skoura, A., Chae, S.S., Cowan, A.E., Han, D.K., Proia, R.L. and Hla, T. (2004) Sphingosine 1-phosphate receptor regulation of N-cadherin mediates vascular stabilization. *Genes Dev*, 18, 2392-2403.
- Papathoma, A.S., Zoumpourlis, V., Balmain, A. and Pintzas, A. (2001) Role of matrix metalloproteinase-9 in progression of mouse skin carcinogenesis. *Mol Carcinog*, 31, 74-82.
- Paterson, I.C., Davies, M., Stone, A., Huntley, S., Smith, E., Pring, M., Eveson, J.W., Robinson, C.M., Parkinson, E.K. and Prime, S.S. (2002) TGF-beta1 acts as a tumor suppressor of human malignant keratinocytes independently of Smad 4 expression and ligand-induced G(1) arrest. *Oncogene*, 21, 1616-1624.
- Peinado, H., Portillo, F. and Cano, A. (2004) Transcriptional regulation of cadherins during development and carcinogenesis. *Int J Dev Biol*, 48, 365-375.
- Peinado, H., Quintanilla, M. and Cano, A. (2003) Transforming growth factor beta-1 induces snail transcription factor in epithelial cell lines: mechanisms for epithelial mesenchymal transitions. *J Biol Chem*, 278, 21113-21123.

- Pessah, M., Marais, J., Prunier, C., Ferrand, N., Lallemand, F., Mauviel, A. and Atfi, A. (2002) c-Jun associates with the oncoprotein Ski and suppresses Smad2 transcriptional activity. *J Biol Chem*, 277, 29094-29100.
- Phanish, M.K., Wahab, N.A., Colville-Nash, P., Hendry, B.M. and Dockrell, M.E. (2006) The differential role of Smad2 and Smad3 in the regulation of pro-fibrotic TGFbeta1 responses in human proximal-tubule epithelial cells. *Biochem J*, 393, 601-607.
- Piek, E., Ju, W.J., Heyer, J., Escalante-Alcalde, D., Stewart, C.L., Weinstein, M., Deng, C., Kucherlapati, R., Bottinger, E.P. and Roberts, A.B. (2001) Functional characterization of transforming growth factor beta signaling in Smad2- and Smad3-deficient fibroblasts. *J Biol Chem*, 276, 19945-19953.
- Piek, E., Moustakas, A., Kurisaki, A., Heldin, C.H. and ten Dijke, P. (1999) TGF-(beta) type I receptor/ALK-5 and Smad proteins mediate epithelial to mesenchymal transdifferentiation in NMuMG breast epithelial cells. *J Cell Sci*, 112 (Pt 24), 4557-4568.
- Pinschewer, D.D., Ochsenbein, A.F., Odermatt, B., Brinkmann, V., Hengartner, H. and Zinkernagel, R.M. (2000) FTY720 immunosuppression impairs effector T cell peripheral homing without affecting induction, expansion, and memory. *J Immunol*, 164, 5761-5770.
- Postigo, A.A., Depp, J.L., Taylor, J.J. and Kroll, K.L. (2003) Regulation of Smad signaling through a differential recruitment of coactivators and corepressors by ZEB proteins. *Embo J*, 22, 2453-2462.
- Pozzi, A., Moberg, P.E., Miles, L.A., Wagner, S., Soloway, P. and Gardner, H.A. (2000) Elevated matrix metalloprotease and angiostatin levels in integrin alpha 1 knockout mice cause reduced tumor vascularization. *Proc Natl Acad Sci U S A*, 97, 2202-2207.
- Radef-Huang, J., Seasholtz, T.M., Matteo, R.G. and Brown, J.H. (2004) G protein mediated signaling pathways in lysophospholipid induced cell proliferation and survival. *J Cell Biochem*, 92, 949-966.
- Reichmann, E., Schwarz, H., Deiner, E.M., Leitner, I., Eilers, M., Berger, J., Busslinger, M. and Beug, H. (1992) Activation of an inducible c-FosER fusion protein causes loss of epithelial polarity and triggers epithelial-fibroblastoid cell conversion. *Cell*, 71, 1103-1116.
- Rhyu, D.Y., Yang, Y., Ha, H., Lee, G.T., Song, J.S., Uh, S.T. and Lee, H.B. (2005) Role of reactive oxygen species in TGF-beta1-induced mitogen-activated protein kinase activation and epithelial-mesenchymal transition in renal tubular epithelial cells. *J Am Soc Nephrol*, 16, 667-675.
- Roberts, A.B. and Sporn, M.B. (1987) Transforming growth factor-beta: potential common mechanisms mediating its effects on embryogenesis, inflammation-repair, and carcinogenesis. *Int J Rad Appl Instrum B*, 14, 435-439.
- Roberts, A.B. and Sporn, M.B. (1993) Physiological actions and clinical applications of transforming growth factor-beta (TGF-beta). *Growth Factors*, 8, 1-9.
- Robinson, C.M., Stone, A.M., Shields, J.D., Huntley, S., Paterson, I.C. and Prime, S.S. (2003) Functional significance of MMP-2 and MMP-9 expression by human malignant oral keratinocyte cell lines. *Arch Oral Biol*, 48, 779-786.
- Ruoslahti, E. (1991) Integrins. *J Clin Invest*, 87, 1-5.



- Ruoslahti, E. (1996) RGD and other recognition sequences for integrins. *Annu Rev Cell Dev Biol*, 12, 697-715.
- Ruwisch, L. (2002) Sphingolipide: Potentielle Arzneistoffe zur topischen Therapie hyperproliferativer Hauterkrankungen. Analytik und in-vitro Untersuchungen zu Apoptose, Proliferation und Penetration (Freie Universität Berlin).
- Saika, S., Kono-Saika, S., Ohnishi, Y., Sato, M., Muragaki, Y., Ooshima, A., Flanders, K.C., Yoo, J., Anzano, M., Liu, C.Y., Kao, W.W. and Roberts, A.B. (2004) Smad3 signaling is required for epithelial-mesenchymal transition of lens epithelium after injury. *Am J Pathol*, 164, 651-663.
- Sanceau, J., Truchet, S. and Bauvois, B. (2003) Matrix metalloproteinase-9 silencing by RNA interference triggers the migratory-adhesive switch in Ewing's sarcoma cells. *J Biol Chem*, 278, 36537-36546.
- Sanchez, T., Estrada-Hernandez, T., Paik, J.H., Wu, M.T., Venkataraman, K., Brinkmann, V., Claffey, K. and Hla, T. (2003) Phosphorylation and action of the immunomodulator FTY720 inhibits vascular endothelial cell growth factor-induced vascular permeability. *J Biol Chem*, 278, 47281-47290.
- Sanchez, T. and Hla, T. (2004) Structural and functional characteristics of S1P receptors. *J Cell Biochem*, 92, 913-922.
- Sanford, L.P., Ormsby, I., Gittenberger-de Groot, A.C., Sariola, H., Friedman, R., Boivin, G.P., Cardell, E.L. and Doetschman, T. (1997) TGFbeta2 knockout mice have multiple developmental defects that are non-overlapping with other TGFbeta knockout phenotypes. *Development*, 124, 2659-2670.
- Sanna, M.G., Liao, J., Jo, E., Alfonso, C., Ahn, M.Y., Peterson, M.S., Webb, B., Lefebvre, S., Chun, J., Gray, N. and Rosen, H. (2004) Sphingosine 1-phosphate (S1P) receptor subtypes S1P1 and S1P3, respectively, regulate lymphocyte recirculation and heart rate. *J Biol Chem*, 279, 13839-13848.
- Santibanez, J.F., Guerrero, J., Quintanilla, M., Fabra, A. and Martinez, J. (2002) Transforming growth factor-beta1 modulates matrix metalloproteinase-9 production through the Ras/MAPK signaling pathway in transformed keratinocytes. *Biochem Biophys Res Commun*, 296, 267-273.
- Santibanez, J.F., Olivares, D., Guerrero, J. and Martinez, J. (2003) Cyclic AMP inhibits TGFbeta1-induced cell-scattering and invasiveness in murine-transformed keratinocytes. *Int J Cancer*, 107, 715-720.
- Santos Maia, C., Mehnert, W., Schaller, M., Korting, H.C., Gysler, A., Haberland, A. and Schafer-Korting, M. (2002) Drug targeting by solid lipid nanoparticles for dermal use. *J Drug Target*, 10, 489-495.
- Sarkar, S., Maceyka, M., Hait, N.C., Paugh, S.W., Sankala, H., Milstien, S. and Spiegel, S. (2005) Sphingosine kinase 1 is required for migration, proliferation and survival of MCF-7 human breast cancer cells. *FEBS Lett*, 579, 5313-5317.
- Sauer, B., Ruwisch, L. and Kleuser, B. (2003) Antiapoptotic action of 1alpha,25-dihydroxyvitamin D3 in primary human melanocytes. *Melanoma Res*, 13, 339-347.

- Sauer, B., Vogler, R., von Wenckstern, H., Fujii, M., Anzano, M.B., Glick, A.B., Schafer-Korting, M., Roberts, A.B. and Kleuser, B. (2004a) Involvement of Smad signaling in sphingosine 1-phosphate-mediated biological responses of keratinocytes. *J Biol Chem*, 279, 38471-38479.
- Sauer, B., Vogler, R., Zimmermann, K., Fujii, M., Anzano, M.B., Schafer-Korting, M., Roberts, A.B. and Kleuser, B. (2004b) Lysophosphatidic acid interacts with transforming growth factor-beta signaling to mediate keratinocyte growth arrest and chemotaxis. *J Invest Dermatol*, 123, 840-849.
- Savagner, P., Kusewitt, D.F., Carver, E.A., Magnino, F., Choi, C., Gridley, T. and Hudson, L.G. (2005) Developmental transcription factor slug is required for effective re-epithelialization by adult keratinocytes. *J Cell Physiol*, 202, 858-866.
- Savagner, P., Yamada, K.M. and Thiery, J.P. (1997) The zinc-finger protein slug causes desmosome dissociation, an initial and necessary step for growth factor-induced epithelial-mesenchymal transition. *J Cell Biol*, 137, 1403-1419.
- Schonbeck, U., Mach, F. and Libby, P. (1998) Generation of biologically active IL-1 beta by matrix metalloproteinases: a novel caspase-1-independent pathway of IL-1 beta processing. *J Immunol*, 161, 3340-3346.
- Shi, Y. and Massague, J. (2003) Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell*, 113, 685-700.
- Shu, X., Wu, W., Mosteller, R.D. and Broek, D. (2002) Sphingosine kinase mediates vascular endothelial growth factor-induced activation of ras and mitogen-activated protein kinases. *Mol Cell Biol*, 22, 7758-7768.
- Shull, M.M., Ormsby, I., Kier, A.B., Pawlowski, S., Diebold, R.J., Yin, M., Allen, R., Sidman, C., Proetzel, G., Calvin, D. and et al. (1992) Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease. *Nature*, 359, 693-699.
- Sirard, C., Kim, S., Mirtsos, C., Tadich, P., Hoodless, P.A., Itie, A., Maxson, R., Wrana, J.L. and Mak, T.W. (2000) Targeted disruption in murine cells reveals variable requirement for Smad4 in transforming growth factor beta-related signaling. *J Biol Chem*, 275, 2063-2070.
- Spiegel, S. and Kolesnick, R. (2002) Sphingosine 1-phosphate as a therapeutic agent. *Leukemia*, 16, 1596-1602.
- Spiegel, S. and Milstien, S. (2002) Sphingosine 1-phosphate, a key cell signaling molecule. *J Biol Chem*, 277, 25851-25854.
- Spiegel, S. and Milstien, S. (2003a) Exogenous and intracellularly generated sphingosine 1-phosphate can regulate cellular processes by divergent pathways. *Biochem Soc Trans*, 31, 1216-1219.
- Spiegel, S. and Milstien, S. (2003b) Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat Rev Mol Cell Biol*, 4, 397-407.
- Steffensen, B., Hakkinen, L. and Larjava, H. (2001) Proteolytic events of wound-healing--coordinated interactions among matrix metalloproteinases (MMPs), integrins, and extracellular matrix molecules. *Crit Rev Oral Biol Med*, 12, 373-398.
- Stetler-Stevenson, W.G., Aznavoorian, S. and Liotta, L.A. (1993) Tumor cell interactions with the extracellular matrix during invasion and metastasis. *Annu Rev Cell Biol*, 9, 541-573.

- Stoker, M., Gherardi, E., Perryman, M. and Gray, J. (1987) Scatter factor is a fibroblast-derived modulator of epithelial cell mobility. *Nature*, 327, 239-242.
- Stoker, M. and Perryman, M. (1985) An epithelial scatter factor released by embryo fibroblasts. *J Cell Sci*, 77, 209-223.
- Sutphen, R., Xu, Y., Wilbanks, G.D., Fiorica, J., Grendys, E.C., Jr., LaPolla, J.P., Arango, H., Hoffman, M.S., Martino, M., Wakeley, K., Griffin, D., Blanco, R.W., Cantor, A.B., Xiao, Y.J. and Krischer, J.P. (2004) Lysophospholipids are potential biomarkers of ovarian cancer. *Cancer Epidemiol Biomarkers Prev*, 13, 1185-1191.
- Taki, M., Verschueren, K., Yokoyama, K., Nagayama, M. and Kamata, N. (2006) Involvement of Ets-1 transcription factor in inducing matrix metalloproteinase-2 expression by epithelial-mesenchymal transition in human squamous carcinoma cells. *Int J Oncol*, 28, 487-496.
- Tanimoto, T., Jin, Z.G. and Berk, B.C. (2002) Transactivation of vascular endothelial growth factor (VEGF) receptor Flk-1/KDR is involved in sphingosine 1-phosphate-stimulated phosphorylation of Akt and endothelial nitric-oxide synthase (eNOS). *J Biol Chem*, 277, 42997-43001.
- Tanimoto, T., Lungu, A.O. and Berk, B.C. (2004) Sphingosine 1-phosphate transactivates the platelet-derived growth factor beta receptor and epidermal growth factor receptor in vascular smooth muscle cells. *Circ Res*, 94, 1050-1058.
- Tester, A.M., Ruangpanit, N., Anderson, R.L. and Thompson, E.W. (2000) MMP-9 secretion and MMP-2 activation distinguish invasive and metastatic sublines of a mouse mammary carcinoma system showing epithelial-mesenchymal transition traits. *Clin Exp Metastasis*, 18, 553-560.
- Thiery, J.P. (2002) Epithelial-mesenchymal transitions in tumour progression. *Nat Rev Cancer*, 2, 442-454.
- Thiery, J.P. and Chopin, D. (1999) Epithelial cell plasticity in development and tumor progression. *Cancer Metastasis Rev*, 18, 31-42.
- Thomas, G.J., Poomsawat, S., Lewis, M.P., Hart, I.R., Speight, P.M. and Marshall, J.F. (2001) alpha v beta 6 Integrin upregulates matrix metalloproteinase 9 and promotes migration of normal oral keratinocytes. *J Invest Dermatol*, 116, 898-904.
- Tian, F., DaCosta Byfield, S., Parks, W.T., Yoo, S., Felici, A., Tang, B., Piek, E., Wakefield, L.M. and Roberts, A.B. (2003) Reduction in Smad2/3 signaling enhances tumorigenesis but suppresses metastasis of breast cancer cell lines. *Cancer Res*, 63, 8284-8292.
- Tolle, M., Levkau, B., Keul, P., Brinkmann, V., Giebing, G., Schonfelder, G., Schafers, M., von Wnuck Lipinski, K., Jankowski, J., Jankowski, V., Chun, J., Zidek, W. and Van der Giet, M. (2005) Immunomodulator FTY720 Induces eNOS-dependent arterial vasodilatation via the lysophospholipid receptor S1P3. *Circ Res*, 96, 913-920.
- Tsai, C.H., Hsieh, Y.S., Yang, S.F., Chou, M.Y. and Chang, Y.C. (2003) Matrix metalloproteinase 2 and matrix metalloproteinase 9 expression in human oral squamous cell carcinoma and the effect of protein kinase C inhibitors: preliminary observations. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 95, 710-716.

- Tsuchida, K., Zhu, Y., Siva, S., Dunn, S.R. and Sharma, K. (2003) Role of Smad4 on TGF-beta-induced extracellular matrix stimulation in mesangial cells. *Kidney Int*, 63, 2000-2009.
- Uhlenbrock, K., Gassenhuber, H. and Kostenis, E. (2002) Sphingosine 1-phosphate is a ligand of the human gpr3, gpr6 and gpr12 family of constitutively active G protein-coupled receptors. *Cell Signal*, 14, 941-953.
- Uhlenbrock, K., Huber, J., Ardati, A., Busch, A.E. and Kostenis, E. (2003) Fluid shear stress differentially regulates gpr3, gpr6, and gpr12 expression in human umbilical vein endothelial cells. *Cell Physiol Biochem*, 13, 75-84.
- Valcourt, U., Kowanetz, M., Niimi, H., Heldin, C.H. and Moustakas, A. (2005) TGF-beta and the Smad signaling pathway support transcriptomic reprogramming during epithelial-mesenchymal cell transition. *Mol Biol Cell*, 16, 1987-2002.
- Valles, A.M., Boyer, B., Badet, J., Tucker, G.C., Barritault, D. and Thiery, J.P. (1990) Acidic fibroblast growth factor is a modulator of epithelial plasticity in a rat bladder carcinoma cell line. *Proc Natl Acad Sci U S A*, 87, 1124-1128.
- Van Brocklyn, J.R., Lee, M.J., Menzeleev, R., Olivera, A., Edsall, L., Cuvillier, O., Thomas, D.M., Coopman, P.J., Thangada, S., Liu, C.H., Hla, T. and Spiegel, S. (1998) Dual actions of sphingosine-1-phosphate: extracellular through the Gi-coupled receptor Edg-1 and intracellular to regulate proliferation and survival. *J Cell Biol*, 142, 229-240.
- Van Veldhoven, P.P. and Mannaerts, G.P. (1994) Sphinganine 1-phosphate metabolism in cultured skin fibroblasts: evidence for the existence of a sphingosine phosphatase. *Biochem J*, 299 (Pt 3), 597-601.
- Visentin, B., Vekich, J.A., Sibbald, B.J., Cavalli, A.L., Moreno, K.M., Matteo, R.G., Garland, W.A., Lu, Y., Yu, S., Hall, H.S., Kundra, V., Mills, G.B. and Sabbadini, R.A. (2006) Validation of an anti-sphingosine-1-phosphate antibody as a potential therapeutic in reducing growth, invasion, and angiogenesis in multiple tumor lineages. *Cancer Cell*, 9, 225-238.
- Vogler, R., Sauer, B., Kim, D.S., Schafer-Korting, M. and Kleuser, B. (2003) Sphingosine-1-phosphate and its potentially paradoxical effects on critical parameters of cutaneous wound healing. *J Invest Dermatol*, 120, 693-700.
- Von Wenckstern, H. (2006) Signalwege von Sphingosin-1-Phosphat und deren Bedeutung in der Migration von Langerhanszellen (Freie Universität Berlin).
- Vu, T.H. and Werb, Z. (2000) Matrix metalloproteinases: effectors of development and normal physiology. *Genes Dev*, 14, 2123-2133.
- Walters, K.A. (1989) Penetration Enhancers and their use in Transdermal Therapeutic Systems, *Transdermal Drug Delivery*, ed. J. Hadgraft, Guy, R.H. (Hrsg.), Marcel Dekker, New York.
- Wang, F., Van Brocklyn, J.R., Edsall, L., Nava, V.E. and Spiegel, S. (1999) Sphingosine-1-phosphate inhibits motility of human breast cancer cells independently of cell surface receptors. *Cancer Res*, 59, 6185-6191.
- Wang, J.F., Olson, M.E., Reno, C.R., Kulyk, W., Wright, J.B. and Hart, D.A. (2000) Molecular and cell biology of skin wound healing in a pig model. *Connect Tissue Res*, 41, 195-211.
- Weeks, B.H., He, W., Olson, K.L. and Wang, X.J. (2001) Inducible expression of transforming growth factor beta1 in papillomas causes rapid metastasis. *Cancer Res*, 61, 7435-7443.

- Weidner, K.M., Sachs, M. and Birchmeier, W. (1993) The Met receptor tyrosine kinase transduces motility, proliferation, and morphogenic signals of scatter factor/hepatocyte growth factor in epithelial cells. *J Cell Biol*, 121, 145-154.
- Wrana, J.L. and Attisano, L. (2000) The Smad pathway. *Cytokine Growth Factor Rev*, 11, 5-13.
- Wu, J.W., Hu, M., Chai, J., Seoane, J., Huse, M., Li, C., Rigotti, D.J., Kyin, S., Muir, T.W., Fairman, R., Massague, J. and Shi, Y. (2001) Crystal structure of a phosphorylated Smad2. Recognition of phosphoserine by the MH2 domain and insights on Smad function in TGF-beta signaling. *Mol Cell*, 8, 1277-1289.
- Xia, P., Gamble, J.R., Wang, L., Pitson, S.M., Moretti, P.A., Wattenberg, B.W., D'Andrea, R.J. and Vadas, M.A. (2000) An oncogenic role of sphingosine kinase. *Curr Biol*, 10, 1527-1530.
- Xie, L., Law, B.K., Chytil, A.M., Brown, K.A., Aakre, M.E. and Moses, H.L. (2004) Activation of the Erk pathway is required for TGF-beta1-induced EMT in vitro. *Neoplasia*, 6, 603-610.
- Xin, C., Ren, S., Eberhardt, W., Pfeilschifter, J. and Huwiler, A. (2006) The immunomodulator FTY720 and its phosphorylated derivative activate the Smad signalling cascade and upregulate connective tissue growth factor and collagen type IV expression in renal mesangial cells. *Br J Pharmacol*, 147, 164-174.
- Xin, C., Ren, S., Kleuser, B., Shabahang, S., Eberhardt, W., Radeke, H., Schafer-Korting, M., Pfeilschifter, J. and Huwiler, A. (2004) Sphingosine 1-phosphate cross-activates the Smad signaling cascade and mimics transforming growth factor-beta-induced cell responses. *J Biol Chem*, 279, 35255-35262.
- Yamaguchi, K., Nagai, S., Ninomiya-Tsuji, J., Nishita, M., Tamai, K., Irie, K., Ueno, N., Nishida, E., Shibuya, H. and Matsumoto, K. (1999) XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAB1-TAK1 in the BMP signaling pathway. *Embo J*, 18, 179-187.
- Yamashita, H., Kitayama, J., Shida, D., Yamaguchi, H., Mori, K., Osada, M., Aoki, S., Yatomi, Y., Takuwa, Y. and Nagawa, H. (2006) Sphingosine 1-phosphate receptor expression profile in human gastric cancer cells: differential regulation on the migration and proliferation. *J Surg Res*, 130, 80-87.
- Yang, J., Mani, S.A., Donaher, J.L., Ramaswamy, S., Itzykson, R.A., Come, C., Savagner, P., Gitelman, I., Richardson, A. and Weinberg, R.A. (2004) Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. *Cell*, 117, 927-939.
- Yatomi, Y., Ruan, F., Hakomori, S. and Igarashi, Y. (1995) Sphingosine-1-phosphate: a platelet-activating sphingolipid released from agonist-stimulated human platelets. *Blood*, 86, 193-202.
- Ye, X., Ishii, I., Kingsbury, M.A. and Chun, J. (2002) Lysophosphatidic acid as a novel cell survival/apoptotic factor. *Biochim Biophys Acta*, 1585, 108-113.
- Yin, F. and Watsky, M.A. (2005) LPA and S1P increase corneal epithelial and endothelial cell transcellular resistance. *Invest Ophthalmol Vis Sci*, 46, 1927-1933.
- Yin, J.J., Selander, K., Chirgwin, J.M., Dallas, M., Grubbs, B.G., Wieser, R., Massague, J., Mundy, G.R. and Guise, T.A. (1999) TGF-beta signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. *J Clin Invest*, 103, 197-206.

- Yokoyama, K., Kamata, N., Fujimoto, R., Tsutsumi, S., Tomonari, M., Taki, M., Hosokawa, H. and Nagayama, M. (2003) Increased invasion and matrix metalloproteinase-2 expression by Snail-induced mesenchymal transition in squamous cell carcinomas. *Int J Oncol*, 22, 891-898.
- Yu, L., Hebert, M.C. and Zhang, Y.E. (2002) TGF-beta receptor-activated p38 MAP kinase mediates Smad-independent TGF-beta responses. *Embo J*, 21, 3749-3759.
- Yu, Q. and Stamenkovic, I. (2000) Cell surface-localized matrix metalloproteinase-9 proteolytically activates TGF-beta and promotes tumor invasion and angiogenesis. *Genes Dev*, 14, 163-176.
- Yue, J. and Mulder, K.M. (2000) Activation of the mitogen-activated protein kinase pathway by transforming growth factor-beta. *Methods Mol Biol*, 142, 125-131.
- Zavadil, J., Bitzer, M., Liang, D., Yang, Y.C., Massimi, A., Kneitz, S., Piek, E. and Bottinger, E.P. (2001) Genetic programs of epithelial cell plasticity directed by transforming growth factor-beta. *Proc Natl Acad Sci U S A*, 98, 6686-6691.
- Zavadil, J. and Bottinger, E.P. (2005) TGF-beta and epithelial-to-mesenchymal transitions. *Oncogene*, 24, 5764-5774.
- Zavadil, J., Cermak, L., Soto-Nieves, N. and Bottinger, E.P. (2004) Integration of TGF-beta/Smad and Jagged1/Notch signalling in epithelial-to-mesenchymal transition. *Embo J*, 23, 1155-1165.
- Zeisberg, M., Hanai, J., Sugimoto, H., Mammoto, T., Charytan, D., Strutz, F. and Kalluri, R. (2003) BMP-7 counteracts TGF-beta1-induced epithelial-to-mesenchymal transition and reverses chronic renal injury. *Nat Med*, 9, 964-968.
- Zhang, H., Desai, N.N., Olivera, A., Seki, T., Brooker, G. and Spiegel, S. (1991) Sphingosine-1-phosphate, a novel lipid, involved in cellular proliferation. *J Cell Biol*, 114, 155-167.
- Zhao, Y., Usatyuk, P.V., Cummings, R., Saatian, B., He, D., Watkins, T., Morris, A., Spannhake, E.W., Brindley, D.N. and Natarajan, V. (2005) Lipid phosphate phosphatase-1 regulates lysophosphatidic acid-induced calcium release, NF-kappaB activation and interleukin-8 secretion in human bronchial epithelial cells. *Biochem J*, 385, 493-502.