

6. Literaturverzeichnis

Achenbach S, Moshage W, Gohl K, Abraham-Fuchs K, Schneider S, Backmann K. Magnetocardiographic investigation of the origin and propagation of cardiac arrhythmias. In: Hoke M, Ern  SN, Okada YC, Romani GL, eds. Biomagnetism: clinical aspects. Amsterdam: Elsevier, 1992: 453-7.

Achenbach S. Investigation of magnetocardiographic parameters for the detection of graft rejection after heart transplantation. In: Baumgartner C, Deecke L, Stroink G, Williamson S, eds. Biomagnetism: fundamentals, research and clinical applications. Amsterdam: IOS Press, 1995: 619-23.

Akers WS, Cross A, Speth R, Dwoskin LP, Cassis LA. Renin-angiotensin system and sympathetic nervous system in cardiac pressure-overload hypertrophy. *Am J Physiol Heart Circ Physiol* 2000; 279 (6): H2797-H2806.

Antzelevitch C, Shimizu W, Yan GX, Sicouri S, Weissenburger J, Nesterenko VV, Burashnikov A, Di Diego J, Saffitz J, Thomas GP. The M cell: its contribution to the ECG and to normal and abnormal electrical function of the heart. *J Cardiovasc Electrophysiol*. 1999 Aug; 10 (8): 1124-52.

Armoundas AA, Wu R, Juang G, Marb n E, Tomaselli GF. Electrical and structural remodeling of the failing ventricle. *Pharmacology & Therapeutics* 2001; 92: 213-230.

Baule GM, McFee R. Detection of the Magnetic Field of the Heart. *Am Heart J* 1963; 66: 95-96.

Brisinda D, Caristo ME, Fenici R. Contactless magnetocardiographic mapping in anaesthetized Wistar Rats: Evidence of age-related changes of cardiac electrical activity. *Am J Physiol Heart Circ Physiol*. 2005 Dec 22.

Brisinda D, Meloni AM, Fenici R. Contactless magnetocardiographic study of ventricular repolarization in intact Wistar rats: Evidence of gender-related differences. *Basic Res Cardiol* 2004; 99: 193-203.

Brisinda D, Meloni AM, Fenici R. Magnetocardiographic differences of ventricular repolarization parameters between Wistar rats and guinea pigs. *Neuro and Clin Neurophysiol* 2004a; 9: 1-4.

Brockmeier K, Casciardi S, Comani S, Del Gratta C, Di Donato L, Di Luzio S, Erne SN, Pasquarelli A, Peresson M, Romani GL. Dynamic magneto-cardiography, *Biomagnetism: clinical aspects*, Excerpta Medica, 1992.

Brockmeier K, Schmitz L, Bobadilla Chavez JDJ, Burghoff M, Koch H, Zimmermann R, Trahms L. Magnetocardiography and 32-lead potential mapping: repolarization in normal subjects during pharmacologically induced stress. *J Cardiovasc Electrophysiol* 1997; 18: 615-626.

Chaikovsky I, Hohler J, Hecker T, et al. Detection of coronary artery disease in patients with normal or unspecifically changed ECG on the basis of magnetocardiography. In: *Biomag 2000 Proceedings of the 12th International Conference on Biomagnetism*. Helsinki: University of Technology, Espoo, 2001: 565-8.

Chen JH. Two components of delayed rectifier current in heart: molecular basis functional diversity and contribution to repolarization. *Acta Pharmacol Sin* 2004; 25 (2): 137-145.

Cleutjens JP, Kandala JC, Guarda E, Guntaka RV, Weber KT. Regulation of collagen degradation in the rat myocardium after infarction. *J Mol Cell Cardiol*. 1995; 27: 1281–1292.

Cohen D, Edelsack EA, Zimmermann JE. Magnetocardiograms taken inside a shielded room with a superconducting point-contact magnetometer. *Appl Phys Lett* 1970; 16: 278-80.

Cohen D, Kaufman LA. Magnetic Determination of the relationship between the S-T segment shift and the injury current produced by coronary artery occlusion. *Circ Res* 1975; 36: 414-424.

Cohen D, Savard P, Rifkin RD, Lepeschkin E, Strauss WE. Magnetic measurement of S-T and T-Q segment shifts in humans. Part II: exercise-induced S-T segment depression. *Circ Res* 1983; 53: 274-279.

Cohen D. Steady fields of the heart: Coronary Artery Occlusion in Dogs. In : Biomagnetism an interdisciplinary approach, Plenum Press. New York 1983a; 267-270.

De Angelis N, Fiordaliso F, Latini R, Calvillo L, Funicello M, Gobbi M, Mennini T, Masson S. Appraisal of the role of angiotensin II and aldosterone in ventricular myocyte apoptosis in adult normotensive rat. *J Mo. Cell Cardiol* 2002; 34: pp. 1655–1665.

Delpón E, Caballero R, Gómez R, Núñez L, Tamargo J. Angiotensin II, angiotensin II antagonists and spironolactone and their modulation of cardiac repolarization. *Trends in Pharmacological Sciences* 2005; 26 (3): 155-161.

Dietz R, Osterziel KJ, Willenbrock R, Gulba DC, von Harsdorf R. Ventricular remodeling after acute myocardial infarction. *Thromb Haemost.* 1999 Sep; 82 (1):73-5.

Erlebacher JA, Weiss JL, Weisfeldt ML, Bulkley BH. Early dilation of the infarcted segment in acute transmural myocardial infarction: role of infarct expansion in acute left ventricular enlargement. *J Am Coll Cardiol.* 1984; 4: 201–208.

Fenici RR, Melillo G, Cappelli A, De Luca C, Masselli M. Magnetocardiographic localization of Kent bundles. In: Williamson S, Hoke M, Stroink G, Kotani M, eds. *Advances in biomagnetism.* New York, NY: Plenum Press, 1989: 365-8.

Fenici RR, Melillo M. Biomagnetic study of cardiac arrhythmias. *Clin Phys Physiol Meas* 1991; 12: A5-A10.

Fiebeler A, Schmidt F, Muller DN, Park JK, Dechend R, Bieringer M, Shagdarsuren E, Breu V, Haller H, Luft FC. Mineralocorticoid receptor affects AP-1 and nuclear factor-kappaB activation in angiotensin II-induced cardiac injury. *Hypertension* 2001; 37: (2 Part 2), pp. 787–793.

Fozzard HA, Makielski JC. The electrophysiology of acute myocardial ischemia. *Ann Rev Med* 1985; 36: 275-284.

Grande P, Christiansen C, Pedersen A, Christiansen MS. Optimal diagnosis in acute myocardial infarction. A cost-effectiveness study. *Circulation* 1980; 61:723.

Gudden F, Hoenig E, Reichenberger H, Schittenhelm R, Schneider A. A multichannel system for use in biomagnetic diagnosis. *Electromedica* 1989; 57:2-7.

Hailer B, van Leeuwen P, Donker D, Grönemeyer D, Seibel R, Wehr M. Die Anwendung des Biomagnetismus in der Kardiologie. MZ Verlag 1995; 15:90-103.

Hänninen H, Takala P, Mäkijärvi M, Montonen J, Korhonen P, Oikarinen L, Simelius K, Nenonen J, Katila T, Toivonen L. Recording locations in multichannel magnetocardiography and body surface potential mapping sensitive for regional exercise-induced myocardial ischemia. *Basic Res Cardiol* 2001; 96: 405-414.

Heger JJ, Weyman AE, Wann LS, Dillon JC, Feigenbaum H. Cross-sectional echocardiographic analysis of the extent of left ventricular asynergy in acute myocardial infarction. *Circulation* 1980; 61: 1113.

Holland RP, Brooks H. TQ-ST segment mapping: critical review and analysis of current concepts. *Am J Cardiol* 1977; 40: 110-129.

Janse MJ, Wit AL. Electrophysiological mechanisms of ventricular arrhythmias resulting from myocardial ischemia and infarction. *Electrophysiological effects of acute myocardial ischemia. Physiol Rex* 1989; 69: 1058-1076.

Johns TNP & Olsen JB. Experimentel myocardial infarction: a method of coronary occlusion in small animals. *Ann. Surg* 1954; 140: 675-682.

Kariniemi V, Ahopelto J, Karp PJ, Katila TE. The fetal magnetocariogram. *J Perinat Med* 1974; 2: 214-6.

Kirchhof P, Eckardt L, Arslan O, Reinhardt L, Monnig G, Fetsch T, Breithardt G, Borggrefe M, Haverkamp W. Prolonged QRS duration increases QT dispersion but does not relate to arrhythmias in survivors of acute myocardial infarction. *Pacing Clin Electrophysiol.* 2001 May; 24 (5): 789-95.

Kléber AG, Fleischhauer J, Cascio WE. Ischemia-induced propagation failure in the heart. In: Zipes DP, Jalife J, eds. *Cardiac Electrophysiology: From cell to bedside*. 2nd edition, W.B. Saunders Company, Philadelphia 1995: 175-182.

Koumi S, Backer CL, Arentzen CE. Characterization of a $[Ca^{2+}]$ -independent current in human atrial and ventricular cardiomyocytes in the absence of Na^+ and K^+ . *Circulation* 1995; 92: 164-174.

Lange C. Myokardiale Vitalitätsdiagnostik bei Patienten mit koronarer Herzkrankheit mittels Magnetokardiographie. Dissertation 2004.

Leder U, Frankenstein L. Temporal properties of high frequency intra-QRS signals in myocardial infarction and healthy hearts; *Biomed Tech (Berl)*. 2000 Sep; 45 (9): 243-7.

Lüderitz B, Jung W, Wolpert C, Lewalter T, Schumacher B, Lickfett L, Vahlhaus C. Neue Methoden in der kardialen Funktionsdiagnostik – QT-Dispersion. *Dt. Ärzteblatt* 1999; 96 (27): A-1835.

Luft FC. Cardiac angiotensin is upregulated in the hearts of unstable angina patients. *Circ Res* 2004; 94: 1530.

Lukas A, Antzelevitch C. Differences in the electrophysiological response of canine ventricular epicardium and endocardium to ischemia. *Circ* 1993 Dez; 88 (6): 2903-15.

Mäkijärvi M, Montonen J, Toivonen L, Siltanen P, Nieminen MS, Leinio M, Katila T. Identification of patients with ventricular tachycardia after myocardial infarction by high-resolution magnetocardiography and electrocardiography. *J Electrocardiol* 1993; 26: 117-24.

Martin G. *Clinical Cardiology: New Frontiers*. Left Ventricular Remodeling After Myocardial Infarction. *Pathophysiology and Therapy*. *Circulation* 2000; 101: 2981.

Martinez EA, Nass CM, Jermyn RM, Rosenbaum SH, Akhtar S, Chan DW, Malkus H, Weiss JL, Fleisher LA. Intermittent cardiac troponin-I screening in an effective means of surveillance for a perioperative myocardial infarction. *J Cardiothorac Vasc Anesth*. 2005; 19 (5): 577-82.

Mittmann C, Eschenhagen T and Scholtz H. Cellular and molecular aspects of contractile dysfunction in heart failure. *Cardiovasc Res* 1998; 39: 267–275.

Monteiro EC, Della Penna S, Di Donato L, Di Luzio S, Pasquarelli A, Erne SN. The study of steady magnetic fields associated with primary and secondary ST shift in ischemic rabbit hearts. *Phyiol Meas* 1997; 18: 191-200.

Moshage W, Achenbach S, Gohl K, Bachmann K. Evaluation of the noninvasive localization accuracy of cardiac arrhythmias attainable by multichannel magnetocardiography. *Int J Card Imaging* 1996; 12: 47-59.

Nenonen J. Solving the inverse problem in magnetocardiography. *IEEE Engineering in Medicine and Biology* 1994; 13: 487-496.

Nikitin IP, Shabalin AV, Ermakova EN, Kytmanov AV, Golyshev NV, Rogachevskii BM, Motorin SV. Magneto-, electro- and echocardiography in detecting symptoms of the „hypertensive heart“. *Clin Med* 1996; 74: 29-31.

Oikarinen L, Paavola M, Montonen J, Viitasalo M, Makijarvi M, Toivonen L, Katila T. Magnetocardiographic QT interval dispersion in postmyocardial infarction patients with sustained ventricular tachycardia: validation of automated QT measurements. *Pacing Clin Electrophysiol* 1998; 21 (10): 1934-42.

Perrier E, Kerfant BG, Lalevee N, Bideaux P, Rossier MF, Richard S, Gomez AM, Benitah JP. Mineralocorticoid Receptor Antagonism Prevents the Electrical Remodeling That Precedes Cellular Hypertrophy After Myocardial Infarction. *Circulation* 2004; 110 (7): 777-783.

Peters NS, Wit AL. Myocardial Architecture and Ventricular Arrhythmogenesis. *Circulation* 1998; 97: 1746-1754.

Peuhkurinen KJ. Ischemic Heart Disease at the Cellular Level. *International Journal of Bioelectromagnetism* 2000; Number 1, Vol 2.

Pfeffer JM, Pfeffer MA, Fishbein MC, Fletcher PJ, Braunwald E. Progressive ventricular remodeling in rat with myocardial infarction. *Am J Physiol Heart Circ Physiol* 1991; 260: H1406-H1414.

Pfeffer MA, Pfeffer JM, Fishbein MC, Fletcher PJ, Spadaro J, Kloner RA, Braunwald E. Myocardial infarct size and ventricular function in rats. *Circ Res* 1979; Vol 44, 503-512.

Pfeffer MA. Left ventricular remodeling after acute myocardial infarction. *Annu Rev Med* 1995; 46: 455-466.

Plonsey R. Comparative capabilities of electrocardiography and magnetocardiography. *Am J Cardiol* 1972; 29: 735-736.

Rengli J, Schweizer W. Diagnostische Schwierigkeiten. *Schweiz med Wochenschr* 1977; 107 (Suppl 6): 37.

Saarinen M, Siltanen P, Karp PJ, Katila TE. The normal magnetocardiogram: I Morphology. *Ann Clin Res* 1978; 10 (suppl 21): 1-22.

Sadoshima J, Izumo S. Mechanical stretch rapidly activates multiple signal transduction pathways in cardiac myocytes: potential involvement of an autocrine/paracrine mechanism. *EMBO Journal*. 1993a; 12: 1681-92.

Sadoshima J, Izumo S. Molecular characterization of angiotensin II-induced hypertrophy of cardiac myocytes and hyperplasia of cardiac fibroblasts. Critical role of the AT1 receptor subtype. *Circ Res* 1993; 73: 413-423.

Sadoshima J, Xu Y, Slayter HS and Izumo S. Autocrine release of angiotensin II mediates stretch-induced hypertrophy of cardiac myocytes in vitro. *Cell* 1993b; 75: 977-984.

Sato M, Terada Y, Mitsui T, Miyashita T, Kandori A, Tsukada K. Detection of myocardial ischemia by magnetocardiogram using 64-channel SQUID system. In: *Biomag 2000 Proceedings of the 12th International Conference on Biomagnetism*. Helsinki: University of Technology, Espoo, 2001: 523-6.

Savard P, Cohen D, Lepeschkin E, Cuffin BN, Madias JE. Magnetic measurement of S-T and T-Q segment shifts in humans. Part I: early repolarization and left bundle branch block. *Circ Res* 1983; 53: 264-273.

Schmitz L, Koch H, Brockmeier K. Magnetocardiographic diagnosis of graft rejection after heart transplantation. In: Hoke M, Erné S, Okada YC, Romani GL, eds. *Biomagnetism: clinical aspects*. Amsterdam: Elsevier, 1992: 555-61.

Schneider S, Hoenig E, Reichenberger H, Abraham-Fucks, Moshage W, Oppelt W, Stefan A, Weikl A, Wirth A. Multichannel biomagnetic system for study of electrical activity of the brain and the heart. *Radiology* 1990; 176: 825-830.

Schunkert H, Sadoshima JI, Cornelius T, Kagaya Y, Weinberg EO, Izumo S, Riegger G, Lorell BH. Angiotensin II-induced growth responses in isolated adult rat hearts: evidence for load-independent induction of cardiac protein synthesis by angiotensin II. *Circ Res*. 1995; 76 (3): 489-497.

Selker HP. Coronary care unit triage decision aids: how do we know when they work? *Am J Med* 1989; 87: 491.

Siltanen P. Magnetocardiography. In: Marcfarlane PW, Lawrie TDV, eds. *Comprehensive Electrocardiography Volume II*, Pergamon Press, New York Oxford Beijing Frankfurt Sao Paolo Sidney Tokyo Toronto: 1989: 1408-1438.

Solomon SD. Remodeling and recovery following myocardial infarction. *Ital Heart J* 2003; 4 (6): 379-382.

Statistisches Bundesamt Deutschland – Pressestelle, Wiesbaden “Krankheitskosten 2002“, Impressum.

Steinhoff U. MCG as a tool for drug safety testing and knock-out model animal studies. *Electrocardiol*. 2004; 37 Suppl: 187-92.

Stroink G, Lant J, Elliott P, Charlebois P, Gardner MJ. Discrimination between myocardial infarction and ventricular tachycardia patients using magnetocardiographic trajectory plots and iso-integral maps. *J Electrocardiol* 1992; 25: 129-142.

Surawicz B, Saito S. Exercise testing for detection of myocardial ischemia in patients with abnormal electrocardiograms at rest. *Am J Cardiol* 1978; 41: 943-951.

Surawicz B. Electrophysiology of myocardial ischemia. *Electrophysiological basis of ECG and cardiac arrhythmias*. Williams and Wilkins, Philadelphia 1995a: 378-401.

Uchida S, Goto K, Tachikawa A, Iramina K. Magnetocardiographic Imaging for ischemic myocardial muscles on rats. *IEICE Trans. Inf. & Syst.* 2002; Vol E85, No.1: 30-34.

Tunstall-Pedoe H, Kuulasmaa K, Mahonen M, Tolonen H, Ruokokoski E, Amouvel P. Contribution of trends in survival and coronary-event rates to changes in coronary heart disease mortality: 10-year results from 37 WHO MONICA project populations. *Monitoring trends and determinants in cardiovascular disease*. *Lancet* 1999; 353 (9164):1547-57.

Van de Werf F, Ardissino D, Betriu A, Cokkinos DV, Falk E, Fox KA, Julian D, Lengyel M, Neumann FJ, Ruzyllo W, Thygesen C, Underwood SR, Vahanian A, Verheugt FW, Wijns W; Task Force on the Management of Acute Myocardial Infarction of the European Society of Cardiology. Management of acute myocardial infarction in patients presenting with ST-segment elevation. *European Heart J* 2003; 24: 28-66.

Van Leeuwen P, Hailer B, Lange S, Donker D, Grönemeyer D. Spatial and temporal changes during QT-interval in the magnetic field of patients with coronary artery disease. *Biomed Tech (Berl)* 1999c; 44: 139-142.

Van Leeuwen P, Hailer B, Wehr M. Spatial distribution of QT intervals: an alternative approach to QT dispersion. *Pacing Clin Elctrophysiol* 1996; 19: 1894-9.

Vatner SF, Baig H, Manders WT, Maroko PR. Effects of coronary artery reperfusion on myocardial infarct size calculated from creatine kinase. *J Clin Invest* 1978; 61: 1048-56.

Vincent GM, Abildskov JA, Burgess MJ. Mechanisms of ischemic ST-segment displacement. Evaluation by direct current recordings. *Circulation* 1977; 56 (4): 559-566.

Visser Ca, Kan G, Meltzer RS. Incidence, timing and prognostic value of left ventricular aneurysm formation after myocardial infarction. *Am J Cardiol* 1986; 57: 729-32.

Voller H, Dietz R. Left ventricular remodeling: pathophysiological mechanisms and therapeutic recommendations. *Z Kardiol* 1999 Dez; 88 (12): 982-90.

Watanabe S, Yamaguchi I. Magnetocardiographic diagnosis for myocardial ischemia and arrhythmias. *Rinsho Byori* 2006; 54 (5): 466-76.

Weiss JL, Bulkley BH, Mason SJ. Two-dimensional echocardiographic quantification of myocardial injury in man: Comparison with post mortem studies (abstract). *Circulation* 1978 a; 57/58 (Suppl II): 153.

Weissmüller P, Abraham-Fuchs K, Schneider S, Richter P, Kochs M, Hombach V. Magnetocardiographic non invasive localizations of accessory pathways in the Wolff-Parkinson-White syndrome by a multichannel system. *Eur Heart J* 1992; 13: 616-22.

Weissmüller P, Richter P, Abraham-Fuchs K. Spatial differences of the duration of ventricular late fields in the signal-averaged magnetocardiogram in patients with ventricular late potentials. *Pacing Clin Electrophysiol* 1993; 16: 70-9.

Willenheimer R. Left ventricular remodelling and dysfunction. Can the process be prevented? *Int Journal of Cardiology* 2000; 72 (2): 143-150.

Yamazaki T, Komuro I, Kudoh S. Endothelin-1 is involved in mechanical stress-induced cardiomyocyte hypertrophy. *J Biol Chem* 1996; 271: 3221–3227.

Yan GX, Yamada JK, Kléber AG, McHowat J, Corr PB. Dissociation between cellular K^+ loss, reduction in repolarisation time, and tissue ATP levels during myocardial hypoxia and ischemia. *Circ Res* 1993; 72: 560-570.