

Contents

1	Introduction	1
2	Objects of Investigation	5
2.1	C ₆₀ Fullerene and its Properties	5
2.2	Model Peptides and Methods of Investigations	12
3	Interaction of Strong Laser Fields with Matter	17
3.1	General Aspects	17
3.2	Multiphoton Ionisation and Fragmentation	19
3.3	Energy Redistribution	27
4	Experimental Apparatus	31
4.1	Vacuum Chamber	31
4.2	Molecular Beam Source	32
4.3	Time of Flight Mass Spectrometry	35
4.4	Photoelectron Spectroscopy	40
4.5	Ion and Electron Detection	41
5	Femtosecond Laser Pulses	51
5.1	Mathematical Description of Laser Pulses	51
5.2	Generation of Femtosecond Laser Pulses	57
5.2.1	Femtosecond Oscillators	57
5.2.2	Chirped Pulse Amplification	60
5.2.3	Multipass Amplification Laser System	62
5.3	Generation of sub-10 fs Pulses	64
5.4	Laser Intensity and Polarisation Control	66
5.5	Average Intensity of Elliptically Polarised Light	70
5.6	Pulse Shaping	71
6	Pulse Characterisation Techniques	79
6.1	Temporal Pulse Characterisation	79
6.1.1	Methods of Temporal Pulse Characterisation	79
6.1.2	Autocorrelation Measurement	80
6.1.3	FROG Technique	83
6.1.4	XFROG Method	86
6.1.5	SPIDER Technique	87
6.2	Spatial Laser Beam Characterisation	90

6.3	Pulse Intensity and Fluence	93
7	Optimisation	95
7.1	Introduction to Mathematical Optimisation	95
7.2	Methods of Stochastic Optimisation	97
7.2.1	Simulated Annealing	97
7.2.2	Evolutionary Algorithms	100
7.3	Practical Implementation of Optimisation	104
8	Excitation of C₆₀	113
8.1	Multiphoton Ionisation of C ₆₀	113
8.1.1	Different Ionisation Mechanisms	113
8.1.2	Saturation Intensity	115
8.1.3	Ionisation Energy and Model Potential	117
8.1.4	Calculation of Ionisation Rates for C ₆₀ ^{q+}	119
8.2	Pulse Duration Dependent Study	120
8.2.1	Experimental Observations	120
8.2.2	Sequential Ionisation	124
8.2.3	Saturation Intensities	125
8.3	Polarisation Dependent Study	128
8.3.1	Photoion Spectroscopy	128
8.3.2	Photoelectron Spectroscopy	140
9	Dynamics of Ultrafast Energy Redistribution in C₆₀	143
9.1	Energy Coupling	143
9.2	Single Pulse Study	145
9.3	One Colour Pump-Probe Study	148
9.4	Two Colour Pump-Probe Study	149
9.5	Pulse Shaping Study	158
10	Mass Spectrometry of Model Peptides	165
10.1	Experimental Results	165
11	Summary and Outlook	179
	Bibliography	183
	Curriculum Vitae	223
	Acknowledgements	225

List of Figures

2.1	The structure of C ₆₀ fullerene	6
2.2	Single electron energy diagram and electronic absorption spectra of C ₆₀ fullerene	10
2.3	Illustration of the energetics for plasmon enhanced multiphoton process in the generation of C ₆₀ ^{q+} with different final charge state q (adopted from [HSV92]) .	11
2.4	Model peptides used in present work	14
3.1	Different photoionisation regimes as a function of the laser intensity and photon energy	20
3.2	Perturbative nonlinear regime of ionisation ($\gamma > 1$)	21
3.3	Strong field regime of ionisation ($\gamma \ll 1$)	23
3.4	Illustration of energy relaxation processes in large but finite systems	29
4.1	Photograph of the vacuum apparatus	32
4.2	Photograph of the oven	34
4.3	Photoelectron and reflectron TOF mass spectrometer	36
4.4	Experimentally measured shape and Gaussian fit of Xenon peak	38
4.5	Charged particles detection schemes	43
4.6	Pulse height distribution	45
4.7	The standard deviation of the integrated ion yield	47
4.8	TOF mass and photoelectron spectra of C ₆₀ fullerene	48
5.1	Pulses of different shapes (Lorentzian, Gaussian, and sech^2 with 100 fs FWHM) and their autocorrelation functions	55
5.2	Optical layout of the multipass amplification laser system	63
5.3	Laser system for the generation of sub-10 fs pulses and an example of the pulse temporal profile produced by this system	65
5.4	The action of $\lambda/2$ - and $\lambda/4$ -plates on linearly polarised light	69
5.5	Scheme of the pulse shaping setup	73
5.6	The layout of SLM-S 640/12	74
5.7	Calibration of LCM	76
6.1	Schematic diagram of an autocorrelator/FROG setup	81
6.2	Pulse duration measurement by the autocorrelation technique for the pulse produced by the Multipass laser system	82
6.3	Temporal characterisation of the pulses produced by the Multipass laser system	84
6.4	Schematic diagram of a cross-correlator/XFROG setup	87
6.5	Temporal characterisation of the pulses passed through the shaper setup	88
6.6	Principle of SPIDER apparatus	89
6.7	Spatial beam characterisation	92

7.1	Diagram of the simulated annealing algorithm	98
7.2	Basic scheme of the genetic algorithm	101
7.3	Scheme of the adaptive closed loop setup	105
7.4	Results of the test optimisations	109
8.1	Typical mass spectra of C_{60} produced by laser pulses of 795 nm wavelength with a pulse duration of 25 fs (top) and 5 ps (bottom)	114
8.2	Ionisation potentials for different charge states of C_{60}^{q+}	118
8.3	Comparison between experimental data of C_{60} photoionisation and the theoretical ionisation rates derived from S-matrix theory by A. Becker and F. H. M. Faisal	119
8.4	Mass spectra obtained from photoionisation of C_{60} with 9 fs laser pulses centred at 765 nm in intensity range between $4 \times 10^{13} \text{ W/cm}^2$ and $4.0 \times 10^{14} \text{ W/cm}^2$	121
8.5	Experimental yields of C_{60}^{q+} ions measured as a function of intensity for 27 fs and 9 fs laser pulses and their fits	122
8.6	Yields of C_{60}^+ and C_{60}^{2+} calculated by solving the coupled differential equations	125
8.7	Saturation intensities for C_{60}^{q+} ions as a function of the final charge state q	126
8.8	Mass spectra of C_{60} after excitation with linearly and circularly polarised laser radiation of 27 fs pulses at 797 nm	129
8.9	Integrated yield of Xe^+ measured at 797 nm normalised to the yield with linear light polarisation plotted as a function of ellipticity angle β and laser intensity	131
8.10	Integrated yield of C_{60}^+ and C_{60}^{3+} measured at 797 nm normalised to the yield with linear light polarisation plotted as a function of ellipticity angle and laser intensity	133
8.11	Integrated yield of C_{60}^+ and C_{60}^{3+} ions normalised to the yield with linear light polarisation plotted as a function of ellipticity angle and laser intensity after interaction of C_{60} with 399 nm laser pulses	134
8.12	Integrated yield of fragments ΣC_{60-2n}^+ and ΣC_{60-2n}^{3+} measured at 797 nm normalised to the yield with linear light polarisation plotted as a function of ellipticity angle and laser intensity	135
8.13	Examples of classical electron trajectories in the combined field of C_{60}^+ and linearly or circularly polarised light	136
8.14	Probability to find different recollision energies for electrons emitted from the inner ($0.65a$) and outer radius ($1.00a$) of the C_{60} molecule in linearly or circularly polarised light at $I = 4.3 \times 10^{14} \text{ W/cm}^2$, 797 nm	139
8.15	Photoelectron spectra of C_{60} obtained with laser intensity of $1.4 \times 10^{14} \text{ W/cm}^2$ for linear and circular light polarisations	141
9.1	Illustration of energy redistribution processes in laser excited C_{60} molecule monitored in the present work with two colour pump-probe spectroscopy	144
9.2	Comparison between different parts of the C_{60} mass spectrum obtained with 9 fs laser pulses and 27 fs laser pulses for equivalent intensities	146
9.3	ΣC_{60-2n}^+ , ΣC_{60-2n}^{2+} , ΣC_{60-2n}^{3+} fragment ion yield obtained with 27 fs and 9 fs laser pulses plotted as a function of the laser intensity on log-log scale	147
9.4	Results of one colour pump-probe experiment with 9 fs laser pulses	149
9.5	Mass spectra with blue (399 nm, $3.4 \times 10^{12} \text{ W/cm}^2$) and/or red (797 nm, $5.1 \times 10^{13} \text{ W/cm}^2$) laser pulses	150

9.6	Total ion yield of different charge states of C_{60}^{q+} , $q = 1 - 4$ as a function of the time delay between 399 nm pump pulse ($3.4 \times 10^{12} \text{ W/cm}^2$) and 797 nm probe pulse ($5.1 \times 10^{13} \text{ W/cm}^2$)	152
9.7	Total ion yield of different masses of C_{60-4n}^{3+} , $n = 0 - 3$ as a function of the time delay between 399 nm pump pulse ($3.4 \times 10^{12} \text{ W/cm}^2$) and 797 nm probe pulse ($5.1 \times 10^{13} \text{ W/cm}^2$)	153
9.8	Relaxation times τ_{el} for highly excited electrons due to electron-electron and electron-vibrational coupling	155
9.9	Ratios H_{br}/O_r , H_{br}/O_r , and H_{br}/M_{br}	156
9.10	Result of C_{50}^+ optimisation	159
9.11	Correlation between C_{50}^+ and C_{48}^+ , C_{52}^+ , C_{60}^+ , C_{60}^{2+} , and C_{50}^{2+} ions during the optimisation run	160
9.12	Comparison between two-colour pump-probe experiment and pulse shaping optimisation	161
9.13	Period of the $a_g(1)$ breathing mode as a function of the number of excited electrons derived from NA-QMD simulations and as a function of charge state derived from hybrid B3LYP level of the DFT method	163
10.1	Mass spectra of the Ac-Phe-NHMe molecular system recorded with laser pulses of $3.7 \times 10^{13} \text{ W/cm}^2$ intensity and 32 fs pulse duration centred at 797 nm	166
10.2	Different possible fragmentation channels of the Ac-Phe-NHMe molecular system	167
10.3	Fragment ion yields of mass 43 u, mass 162 u, and the parent ion (mass 220 u) of the Ac-Phe-NHMe molecular system recorded with 34 fs laser pulses plotted as a function of the laser intensity on the log-log scale	168
10.4	Fitness f characterising the predominant formation of fragment mass 43 u as a function of the generation in the adaptive feedback loop and SH-XFROG map of the optimal pulse shape	169
10.5	Temporal envelope and corresponding mass spectra recorded for the unshaped, stretched to 246 fs (FWHM), and optimal pulses	170
10.6	Distribution of individual optimal pulses grouped according to the similarities in pulses structure	171
10.7	Calculated triple pulse sequence in the time domain illustrating the effect of different phase parameters Φ on the pulse sequence	172
10.8	Ion yields plotted as a function of the relative phase shift in the parameterised triple pulses and measured temporal envelope of the pulse sequence applied for three different relative phase shifts $\Phi = 0, \pi/2$, and $3\pi/2$	173
10.9	Comparison between power spectra of the triple pulses assuming a third order process and resonant two photon ionisation spectrum of Ac-Phe-NHMe	174
10.10	Fitness f characterising the enhanced formation of fragment mass 77 u as a function of the generation in the adaptive feedback loop	175
10.11	Temporal structure and corresponding mass spectra of Ac-Ala-NHMe are recorded with unshaped and optimal pulse	177

List of Tables

2.1	Optically active vibrational modes of C ₆₀ fullerene	8
4.1	Distances and voltage settings for the reflectron TOF mass spectrometer	37
4.2	Distances and voltage settings for the photoelectron spectrometer	41
5.1	The mathematical description of some pulse shapes in the time and frequency domains	54
5.2	Deconvolution factors D_{AC} and time-bandwidth products K_{TB} for the various shapes of laser pulses	57
5.3	Output parameters of oscillator and amplifier of the multipass amplification laser system	64
7.1	Standard parameters for a free optimisation with the genetic algorithm	110

