

## **8. Literaturverzeichnis**

- Adolphs J., Schmidt D.K., Korsukewitz I., Kamin B., Habazettl H., Schafer M., and Welte M. (2004):  
Effects of thoracic epidural anaesthesia on intestinal microvascular perfusion in a rodent model of normotensive endotoxaemia.  
Intensive Care Med 30 (11), 2094-2101.
- Andlauer W., Kolb J., and Furst P. (2000):  
Isoflavones from tofu are absorbed and metabolized in the isolated rat small intestine.  
J Nutr 130 (12), 3021-3027.
- Andlauer W., Kolb J., and Furst P. (2004):  
Phloridzin improves absorption of genistin in isolated rat small intestine.  
Clin Nutr 23 (5), 989-995.
- Angus D.C., Linde-Zwirble W.T., Lidicker J., Clermont G., Carcillo J., and Pinsky M.R. (2001):  
Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care.  
Crit Care Med 29 (7), 1303-1310.
- Baue A.E., Durham R., and Faist E. (1998):  
Systemic inflammatory response syndrome (SIRS), multiple organ dysfunction syndrome (MODS), multiple organ failure (MOF): are we winning the battle?  
Shock 10 (2), 79-89.
- Bauer A.J., Schwarz N.T., Moore B.A., Turler A., and Kalff J.C. (2002):  
Ileus in critical illness: mechanisms and management.  
Curr Opin Crit Care 8 (2), 152-157.
- Bauldry S.A., Elsey K.L., and Bass D.A. (1992):  
Activation of NADPH oxidase and phospholipase D in permeabilized human neutrophils. Correlation between oxidase activation and phosphatidic acid production.  
J Biol Chem 267 (35), 25141-25152.
- Baumgartner J.D. and Calandra T. (1999):  
Treatment of sepsis: past and future avenues.  
Drugs 57 (2), 127-132.
- Bertuglia S. and Giusti A. (2003):  
Microvascular oxygenation, oxidative stress, NO suppression and superoxide dismutase during postischemic reperfusion.  
Am J Physiol Heart Circ Physiol 285 (3), H1064-H1071.
- Bhakdi S., Muhly M., Korom S., and Schmidt G. (1990):  
Effects of *Escherichia coli* hemolysin on human monocytes. Cytocidal action and stimulation of interleukin 1 release.  
J Clin Invest 85 (6), 1746-1753.

Bhakdi S. and Tranum-Jensen J. (1991):  
Alpha-toxin of *Staphylococcus aureus*.  
Microbiol Rev 55 (4), 733-751.

Bohlen H.G. (1998):  
Integration of intestinal structure, function, and microvascular regulation.  
Microcirculation 5 (1), 27-37.

Bone R.C. (1993):  
Why new definitions of sepsis and organ failure are needed.  
Am J Med 95 (4), 348-350.

Bone R.C., Balk R.A., Cerra F.B., Dellinger R.P., Fein A.M., Knaus W.A., Schein R.M., and Sibbald W.J. (1992):  
Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine.  
Chest 101 (6), 1644-1655.

Bonten M.J., Gaillard C.A., de Leeuw P.W., and Stobberingh E.E. (1997):  
Role of colonization of the upper intestinal tract in the pathogenesis of ventilator-associated pneumonia.  
Clin Infect Dis 24 (3), 309-319.

Brell B. (2004):  
Untersuchung zu Störung der mucosalen Mikrozirkulation durch *Staphylococcus aureus* alpha-Toxin an einem isoliert perfundierten Kaninchenileum-Modell.  
Berlin, Humboldt-Universität, Dissertation 2004.

Brell B., Temmesfeld-Wollbrück B., Altzschner I., Frisch E., Schmeck B., Hocke A.C., Suttorp N., and Hippenstiel S. (2005a):  
Adrenomedullin reduces *Staphylococcus aureus* alpha-toxin-induced rat ileum microcirculatory damage.  
Crit Care Med 33 (4), 819-826.

Brell B., Temmesfeld-Wollbrück B., Altzschner I., Frisch E., Schmeck B., Hocke A.C., Suttorp N., and Hippenstiel S. (2005b):  
Adrenomedullin reduces *Staphylococcus aureus* alpha-toxin-induced rat ileum microcirculatory damage.  
Crit Care Med 33 (4), 819-826.

Brun-Buisson C. (2000):  
The epidemiology of the systemic inflammatory response.  
Intensive Care Med 26 Suppl 1 , S64-S74.

Brun-Buisson C., Doyon F., Carlet J., Dellamonica P., Gouin F., Lepoutre A., Mercier J.C., Ofenstadt G., and Regnier B. (1995):  
Incidence, risk factors, and outcome of severe sepsis and septic shock in adults. A multicenter prospective study in intensive care units. French ICU Group for Severe Sepsis.

JAMA 274 (12), 968-974.

Caron K.M. and Smithies O. (2001):

Extreme hydrops fetalis and cardiovascular abnormalities in mice lacking a functional

Adrenomedullin gene.

Proc Natl Acad Sci U S A 98 (2), 615-619.

Ceppa E.P., Fuh K.C., and Bulkley G.B. (2003):

Mesenteric hemodynamic response to circulatory shock.

Curr Opin Crit Care 9 (2), 127-132.

Chiu C.J., McArdle A.H., Brown R., Scott H.J., and Gurd F.N. (1970):

Intestinal mucosal lesion in low-flow states. I. A morphological, hemodynamic, and metabolic reappraisal.

Arch Surg 101 (4), 478-483.

De Vriese A.S., Verbeuren T.J., Vallez M.O., Lameire N.H., De Buyzere M., and Vanhoutte

P.M. (2000):

Off-line analysis of red blood cell velocity in renal arterioles.

J Vasc Res 37 (1), 26-31.

Dregelid E., Haukaas S., Amundsen S., Eide G.E., Soreide O., Lekven J., and Svanes K. (1986):

Microsphere method in measurement of blood flow to wall layers of small intestine.

Am J Physiol 250 (5 Pt 1), G670-G678.

Evaldson G., Heimdahl A., Kager L., and Nord C.E. (1982):

The normal human anaerobic microflora.

Scand J Infect Dis Suppl 35 , 9-15.

Fink M.P. (1993):

Adequacy of gut oxygenation in endotoxemia and sepsis.

Crit Care Med 21 (2 Suppl), S4-S8.

Fink M.P. (2003):

Intestinal epithelial hyperpermeability: update on the pathogenesis of gut mucosal barrier dysfunction in critical illness.

Curr Opin Crit Care 9 (2), 143-151.

Fowler D.E. and Wang P. (2002):

The cardiovascular response in sepsis: proposed mechanisms of the beneficial effect of adrenomedullin and its binding protein (review).

Int J Mol Med 9 (5), 443-449.

Germann R., Hasibeder W., Haisjackl M., Sparr H., Luz G., Pernthaler H., Friesenecker B., Bonatti J., Gruber E., and Schwarz B. (1995):

Dopamine-1-receptor stimulation and mucosal tissue oxygenation in the porcine jejunum.

Crit Care Med 23 (9), 1560-1566.

- Grimminger F., Rose F., Sibelius U., Meinhardt M., Potzsch B., Spriestersbach R., Bhakdi S., Suttorp N., and Seeger W. (1997):  
 Human endothelial cell activation and mediator release in response to the bacterial exotoxins *Escherichia coli* hemolysin and staphylococcal alpha-toxin.  
*J Immunol* 159 (4), 1909-1916.
- Grover R., Zaccardelli D., Colice G., Guntupalli K., Watson D., and Vincent J.L. (1999a):  
 An open-label dose escalation study of the nitric oxide synthase inhibitor, N(G)-methyl-L-arginine hydrochloride (546C88), in patients with septic shock. Glaxo Wellcome International Septic Shock Study Group.  
*Crit Care Med* 27 (5), 913-922.
- Grover R., Lopez A., Lorente J., Steingrub J., Bakker J., Willatts S., McLuckie A., and Takala J. (1999b):  
 Multi-center, randomized, placebo-controlled, double blind study of the nitric oxide synthase inhibitor 546c88: Effect on survival in patients with septic shock. [Report].  
*Critical Care Medicine* 27 (1S), 33A.
- Haglund U., Abe T., Ahren C., Braide I., and Lundgren O. (1976):  
 The intestinal mucosal lesions in shock. II. The relationship between the mucosal lesions and the cardiovascular derangement following regional shock.  
*Eur Surg Res* 8 (5), 448-460.
- Harshman S., Lefferts P.L., and Snapper J.R. (1992):  
 Staphylococcal alpha toxin: a study with chronically instrumented awake sheep.  
*Infect Immun* 60 (9), 3489-3496.
- Hatoum O.A., Binion D.G., Otterson M.F., and Guterman D.D. (2003):  
 Acquired microvascular dysfunction in inflammatory bowel disease: Loss of nitric oxide-mediated vasodilation.  
*Gastroenterology* 125 (1), 58-69.
- Hildebrand A., Pohl M., and Bhakdi S. (1991):  
*Staphylococcus aureus* alpha-toxin. Dual mechanism of binding to target cells.  
*J Biol Chem* 266 (26), 17195-17200.
- Hinshaw L.B. (1996):  
 Sepsis/septic shock: participation of the microcirculation: an abbreviated review.  
*Crit Care Med* 24 (6), 1072-1078.
- Hippenstiel S., Witzenrath M., Schmeck B., Hocke A., Krisp M., Krüll M., Seybold J., Seeger W., Rascher W., Schütte H., and Suttorp N. (2002):  
 Adrenomedullin reduces endothelial hyperpermeability.  
*Circ Res* 91 (7), 618-625.
- Hirata Y., Mitaka C., Sato K., Nagura T., Tsunoda Y., Amaha K., and Marumo F. (1996):  
 Increased circulating adrenomedullin, a novel vasodilatory peptide, in sepsis.  
*J Clin Endocrinol Metab* 81 (4), 1449-1453.

Horton J.W. (1992):  
Alterations in intestinal permeability and blood flow in a new model of mesenteric ischemia.  
*Circ Shock* 36 (2), 134-139.

Ince C. (2005):  
The microcirculation is the motor of sepsis.  
*Crit Care* 9 Suppl 4 , S13-S19.

Ince C. and Sinaasappel M. (1999):  
Microcirculatory oxygenation and shunting in sepsis and shock.  
*Crit Care Med* 27 (7), 1369-1377.

Kato H., Shichiri M., Marumo F., and Hirata Y. (1997):  
Adrenomedullin as an autocrine/paracrine apoptosis survival factor for rat endothelial cells.  
*Endocrinology* 138 (6), 2615-2620.

Kernodle D.S., Voladri R.K., Menzies B.E., Hager C.C., and Edwards K.M. (1997):  
Expression of an antisense hla fragment in *Staphylococcus aureus* reduces alpha-toxin production  
in vitro and attenuates lethal activity in a murine model.  
*Infect Immun* 65 (1), 179-184.

King C.J., Tytgat S., Delude R.L., and Fink M.P. (1999):  
Ileal mucosal oxygen consumption is decreased in endotoxemic rats but is restored toward normal by treatment with aminoguanidine.  
*Crit Care Med* 27 (11), 2518-2524.

Kitamura K., Kangawa K., Kawamoto M., Ichiki Y., Nakamura S., Matsuo H., and Eto T. (1993):  
Adrenomedullin: a novel hypotensive peptide isolated from human pheochromocytoma.  
*Biochem Biophys Res Commun* 192 (2), 553-560.

Kiyomizu A., Kitamura K., Kawamoto M., and Eto T. (2001):  
Distribution and molecular forms of adrenomedullin and proadrenomedullin N-terminal 20 peptide in the porcine gastrointestinal tract.  
*J Gastroenterol* 36 (1), 18-23.

Kohno M., Hanehira T., Kano H., Horio T., Yokokawa K., Ikeda M., Minami M., Yasunari K., and Yoshikawa J. (1996):  
Plasma adrenomedullin concentrations in essential hypertension.  
*Hypertension* 27 (1), 102-107.

Kullak K. (1997):  
Bedeutung der Darmflora für den Menschen.  
*Medizin und Ernährung Supplement* (6), 56-59.

Laroux F.S. and Grisham M.B. (2001):  
Immunological basis of inflammatory bowel disease: role of the microcirculation.  
*Microcirculation* 8 (5), 283-301.

Laupland K.B., Gregson D.B., Zygun D.A., Doig C.J., Mortis G., and Church D.L. (2004):

Severe bloodstream infections: a population-based assessment.  
Crit Care Med 32 (4), 992-997.

Le Gall J.R., Brun-Buisson C., Trunet P., Latournerie J., Chantereau S., and Rapin M. (1982):  
Influence of age, previous health status, and severity of acute illness on outcome from intensive  
care.  
Crit Care Med 10 (9), 575-577.

Lee J.S. (1994):  
Microvascular volume changes induced by exercise, heat exposure, or endotoxin injection.  
Am J Physiol 267 (3 Pt 2), H1142-H1150.

Leung F.W., Morishita T., Livingston E.H., Reedy T., and Guth P.H. (1987):  
Reflectance spectrophotometry for the assessment of gastroduodenal mucosal perfusion.  
Am J Physiol 252 (6 Pt 1), G797-G804.

Li S., Arvidson S., and Mollby R. (1997):  
Variation in the agr-dependent expression of alpha-toxin and protein A among clinical isolates of  
*Staphylococcus aureus* from patients with septicaemia.  
FEMS Microbiol Lett 152 (1), 155-161.

Lindert J., Werner J., Redlin M., Kuppe H., Habazettl H., and Pries A.R. (2002):  
OPS imaging of human microcirculation: a short technical report.  
J Vasc Res 39 (4), 368-372.

Lowy F.D. (1998):  
*Staphylococcus aureus* infections.  
N Engl J Med 339 (8), 520-532.

Madorin W.S., Martin C.M., and Sibbald W.J. (1999):  
Dopexamine attenuates flow motion in ileal mucosal arterioles in normotensive sepsis.  
Crit Care Med 27 (2), 394-400.

Martin G.S., Mannino D.M., Eaton S., and Moss M. (2003):  
The epidemiology of sepsis in the United States from 1979 through 2000.  
N Engl J Med 348 (16), 1546-1554.

Mayer K., Temmesfeld-Wollbrück B., Friedland A., Olschewski H., Reich M., Seeger W., and  
Grimminger A.F. (1999):  
Severe microcirculatory abnormalities elicited by *E. coli* hemolysin in the rabbit ileum mucosa.  
Am J Respir Crit Care Med 160 (4), 1171-1178.

Minamino N., Kikumoto K., and Isumi Y. (2002):  
Regulation of adrenomedullin expression and release.  
Microsc Res Tech 57 (1), 28-39.

Nagaya N., Miyatake K., Kyotani S., Nishikimi T., Nakanishi N., and Kangawa K. (2003):  
Pulmonary vasodilator response to adrenomedullin in patients with pulmonary hypertension.  
Hypertens Res 26 Suppl , S141-S146.

Nakajima Y., Baudry N., Duranteau J., and Vicaut E. (2001):  
Microcirculation in intestinal villi: a comparison between hemorrhagic and endotoxin shock.  
Am J Respir Crit Care Med 164 (8 Pt 1), 1526-1530.

Nakamura K., Toda H., Terasako K., Kakuyama M., Hatano Y., Mori K., and Kangawa K. (1995):  
Vasodilative effect of adrenomedullin in isolated arteries of the dog.  
Jpn J Pharmacol 67 (3), 259-262.

Nishikimi T., Yoshihara F., Mori Y., Kangawa K., and Matsuoka H. (2003):  
Cardioprotective effect of adrenomedullin in heart failure.  
Hypertens Res 26 Suppl , S121-S127.

Nuki C., Kawasaki H., Kitamura K., Takenaga M., Kangawa K., Eto T., and Wada A. (1993):  
Vasodilator effect of adrenomedullin and calcitonin gene-related peptide receptors in rat mesenteric vascular beds.  
Biochem Biophys Res Commun 196 (1), 245-251.

Nystrom P.O. (1998):  
The systemic inflammatory response syndrome: definitions and aetiology.  
J Antimicrob Chemother 41 Suppl A , 1-7.

Opal S.M. and Cross A.S. (1999):  
Clinical trials for severe sepsis. Past failures, and future hopes.  
Infect Dis Clin North Am 13 (2), 285-97, vii.

Ornan D.A., Chaudry I.H., and Wang P. (1999):  
Pulmonary clearance of adrenomedullin is reduced during the late stage of sepsis.  
Biochim Biophys Acta 1427 (2), 315-321.

Pajk W., Schwarz B., Knotzer H., Friesenecker B., Mayr A., Dunser M., and Hasibeder W. (2002):  
Jejunal tissue oxygenation and microvascular flow motion during hemorrhage and resuscitation.  
Am J Physiol Heart Circ Physiol 283 (6), H2511-H2517.

Patel A.H., Nowlan P., Weavers E.D., and Foster T. (1987):  
Virulence of protein A-deficient and alpha-toxin-deficient mutants of *Staphylococcus aureus* isolated by allele replacement.  
Infect Immun 55 (12), 3103-3110.

Pittet D., Thievent B., Wenzel R.P., Li N., Gurman G., and Suter P.M. (1993):  
Importance of pre-existing co-morbidities for prognosis of septicemia in critically ill patients.  
Intensive Care Med 19 (5), 265-272.

Pittner A., Nalos M., Asfar P., Yang Y., Ince C., Georgieff M., Bruckner U.B., Radermacher P., and Froba G. (2003):  
Mechanisms of inducible nitric oxide synthase (iNOS) inhibition-related improvement of gut mucosal acidosis during hyperdynamic porcine endotoxemia.  
Intensive Care Med 29 (2), 312-316.

Pries A.R. (1988):

A versatile video image analysis system for microcirculatory research.  
Int J Microcirc Clin Exp 7 (4), 327-345.

Pullamsetti S.S., Maring D., Ghofrani H.A., Mayer K., Weissmann N., Rosengarten B., Lehner M., Schudt C., Boer R., Grimminger F., Seeger W., and Schermuly R.T. (2006):  
Effect of nitric oxide synthase (NOS) inhibition on macro- and microcirculation in a model of rat endotoxic shock.  
Thromb Haemost 95 (4), 720-727.

Rigalli A., Beinlich A., and Puche R.C. (2001):

Intestinal absorption of fluoride at high luminal concentration of fluoride.  
Arzneimittelforschung 51 (2), 151-155.

Russell D.H., Barreto J.C., Klemm K., and Miller T.A. (1995):

Hemorrhagic shock increases gut macromolecular permeability in the rat.  
Shock 4 (1), 50-55.

Sakata J., Asada Y., Shimokubo T., Kitani M., Inatsu H., Kitamura K., Kangawa K., Matsuo H., Sumiyoshi A., and Eto T. (1998):

Adrenomedullin in the gastrointestinal tract. Distribution and gene expression in rat and augmented gastric adrenomedullin after fasting.

J Gastroenterol 33 (6), 828-834.

Salzman A.L., Wang H., Wollert P.S., VanderMeer T.J., Compton C.C., Denenberg A.G., and Fink M.P. (1994):

Endotoxin-induced ileal mucosal hyperpermeability in pigs: role of tissue acidosis.  
Am J Physiol 266 (4 Pt 1), G633-G646.

Sato N., Kamada T., Shichiri M., Kawano S., Abe H., and Hagihara B. (1979):

Measurement of hemoperfusion and oxygen sufficiency in gastric mucosa in vivo. Evidence of mucosal hypoxia as the cause of hemorrhagic shock- induced gastric mucosal lesion in rats.  
Gastroenterology 76 (4), 814-819.

Schmidt H., Secchi A., Wellmann R., Bach A., Bohrer H., Gebhard M.M., and Martin E. (1996a):  
Effect of endotoxemia on intestinal villus microcirculation in rats.

J Surg Res 61 (2), 521-526.

Schmidt H., Secchi A., Wellmann R., Bach A., Bohrer H., Gebhard M.M., and Martin E. (1996b):

Effect of endotoxemia on intestinal villus microcirculation in rats.  
J Surg Res 61 (2), 521-526.

Schmidt W., Tinelli M., Secchi A., Gebhard M.M., Martin E., and Schmidt H. (2000):

Influence of amrinone on intestinal villus blood flow during endotoxemia.  
J Crit Care 15 (3), 97-102.

Schoenberg M.H., Weiss M., and Radermacher P. (1998):

Outcome of patients with sepsis and septic shock after ICU treatment.

Langenbecks Arch Surg 383 (1), 44-48.

Schottmueller H. (1914):

Wesen und Behandlung der Sepsis.

Verh Dtsch Ges Inn Med , 31 , 257-280.

Secchi A., Wellmann R., Martin E., and Schmidt H. (1997):

Dobutamine maintains intestinal villus blood flow during normotensive endotoxemia: an intravital microscopic study in the rat.

J Crit Care 12 (3), 137-141.

Seeger W., Bauer M., and Bhakdi S. (1984):

Staphylococcal alpha-toxin elicits hypertension in isolated rabbit lungs. Evidence for thromboxane formation and the role of extracellular calcium.

J Clin Invest 74 (3), 849-858.

Seeger W., Birkemeyer R.G., Ermert L., Suttorp N., Bhakdi S., and Duncker H.R. (1990):

Staphylococcal alpha-toxin-induced vascular leakage in isolated perfused rabbit lungs.

Lab Invest 63 (3), 341-349.

Shepherd A.P. and Kiel J.W. (1992):

A model of countercurrent shunting of oxygen in the intestinal villus.

Am J Physiol 262 (4 Pt 2), H1136-H1142.

Shindo T., Kurihara H., Maemura K., Kurihara Y., Kuwaki T., Izumida T., Minamino N., Ju K.H., Morita H., Oh-hashi Y., Kumada M., Kangawa K., Nagai R., and Yazaki Y. (2000): Hypotension and resistance to lipopolysaccharide-induced shock in transgenic mice overexpressing adrenomedullin in their vasculature.

Circulation 101 (19), 2309-2316.

Sibelius U., Grandel U., Buerke M., Mueller D., Kiss L., Kraemer H.J., Braun-Dullaeus R., Haberbosch W., Seeger W., and Grimminger F. (2000):

Staphylococcal alpha-toxin provokes coronary vasoconstriction and loss in myocardial contractility in perfused rat hearts: role of thromboxane generation.

Circulation 101 (1), 78-85.

Sielenkamper A.W., Eicker K., and Van Aken H. (2000):

Thoracic epidural anesthesia increases mucosal perfusion in ileum of rats.

Anesthesiology 93 (3), 844-851.

Soderquist B., Colgue-Navarro P., Blomquist L., Olcén P., Holmberg H., and Möllby R. (1993): Staphylococcal alpha-toxin in septicaemic patients; detection in serum, antibody response and production in isolated strain.

Serodiagn Immunother Infect Dis 5 , 139-144.

Soderquist B., Kallman J., Holmberg H., Vikerfors T., and Kihlstrom E. (1998):

Secretion of IL-6, IL-8 and G-CSF by human endothelial cells in vitro in response to *Staphylococcus aureus* and staphylococcal exotoxins.

APMIS 106 (12), 1157-1164.

Sorrells D.L., Friend C., Koltuksuz U., Courcoulas A., Boyle P., Garrett M., Watkins S., Rowe M.I., and Ford H.R. (1996):

Inhibition of nitric oxide with aminoguanidine reduces bacterial translocation after endotoxin challenge in vivo.

Arch Surg 131 (11), 1155-1163.

Sriskandan S. and Cohen J. (1999):

Gram-positive sepsis. Mechanisms and differences from gram-negative sepsis.

Infect Dis Clin North Am 13 (2), 397-412.

Stonestreet B.S., Ocampo S.S., and Oh W. (1998):

Reductions in cardiac output in hypoxic young pigs: systemic and regional perfusion and oxygen metabolism.

J Appl Physiol 85 (3), 874-882.

Suttorp N., Buerke M., and Tannert-Otto S. (1992):

Stimulation of PAF-synthesis in pulmonary artery endothelial cells by *Staphylococcus aureus* alpha-toxin.

Thromb Res 67 (2), 243-252.

Suttorp N., Fuhrmann M., Tannert-Otto S., Grimminger F., and Bhakdi S. (1993):

Pore-forming bacterial toxins potently induce release of nitric oxide in porcine endothelial cells.

J Exp Med 178 (1), 337-341.

Suttorp N., Hessz T., Seeger W., Wilke A., Koob R., Lutz F., and Drenckhahn D. (1988):

Bacterial exotoxins and endothelial permeability for water and albumin in vitro.

Am J Physiol 255 (3 Pt 1), C368-C376.

Suttorp N., Seeger W., Dewein E., Bhakdi S., and Roka L. (1985):

Staphylococcal alpha-toxin-induced PGI<sub>2</sub> production in endothelial cells: role of calcium.

Am J Physiol 248 (1 Pt 1), C127-C134.

Suttorp N., Seeger W., Zucker-Reimann J., Roka L., and Bhakdi S. (1987):

Mechanism of leukotriene generation in polymorphonuclear leukocytes by staphylococcal alpha-toxin.

Infect Immun 55 (1), 104-110.

Temmesfeld-Wollbrück B., Szalay A., Mayer K., Olschewski H., Seeger W., and Grimminger F. (1998):

Abnormalities of gastric mucosal oxygenation in septic shock: partial responsiveness to dopexamine.

Am J Respir Crit Care Med 157 (5 Pt 1), 1586-1592.

Theuer C.J., Wilson M.A., Steeb G.D., and Garrison R.N. (1993):

Microvascular vasoconstriction and mucosal hypoperfusion of the rat small intestine during bacteremia.

Circ Shock 40 (1), 61-68.

Trager K., Radermacher P., Rieger K.M., Grover R., Vlatten A., Iber T., Adler J., Georgieff M., and Santak B. (2000):

Norepinephrine and N(G)-monomethyl-L-arginine in hyperdynamic septic shock in pigs: effects on intestinal oxygen exchange and energy balance.

Crit Care Med 28 (6), 2007-2014.

Tugtekin I.F., Radermacher P., Theisen M., Matejovic M., Stehr A., Ploner F., Matura K., Ince C., Georgieff M., and Trager K. (2001):

Increased ileal-mucosal-arterial PCO<sub>2</sub> gap is associated with impaired villus microcirculation in endotoxic pigs.

Intensive Care Med 27 (4), 757-766.

Ueda S., Nishio K., Minamino N., Kubo A., Akai Y., Kangawa K., Matsuo H., Fujimura Y., Yoshioka A., Masui K., Doi N., Murao Y., and Miyamoto S. (1999):

Increased plasma levels of adrenomedullin in patients with systemic inflammatory response syndrome.

Am J Respir Crit Care Med 160 (1), 132-136.

Valeva A., Walev I., Pinkernell M., Walker B., Bayley H., Palmer M., and Bhakdi S. (1997): Transmembrane beta-barrel of staphylococcal alpha-toxin forms in sensitive but not in resistant cells.

Proc Natl Acad Sci U S A 94 (21), 11607-11611.

Vallet B. (2002):

Endothelial cell dysfunction and abnormal tissue perfusion.

Crit Care Med 30 (5 Suppl), S229-S234.

Vallet B., Lund N., Curtis S.E., Kelly D., and Cain S.M. (1994):

Gut and muscle tissue PO<sub>2</sub> in endotoxemic dogs during shock and resuscitation.

J Appl Physiol 76 (2), 793-800.

van Bommel J., Siegemund M., Henny C.P., van den Heuvel D.A., Trouwborst A., and Ince C. (2000):

Preservation of intestinal microvascular Po<sub>2</sub> during normovolemic hemodilution in a rat model.

J Lab Clin Med 135 (6), 476-483.

VanderMeer T.J., Wang H., and Fink M.P. (1995):

Endotoxemia causes ileal mucosal acidosis in the absence of mucosal hypoxia in a normodynamic porcine model of septic shock.

Crit Care Med 23 (7), 1217-1226.

Vanner S., Jiang M.M., Brooks V.L., and Surprenant A. (1990):

Characterization of vasopressin actions in isolated submucosal arterioles of the intestinal microcirculation.

Circ Res 67 (4), 1017-1026.

Venkataraman R and Fink MP. 2002. Mechanisms responsible for intestinal barrier dysfunction in critical illness. In: Vincent JL, Evans TW, and Fink MP, editors. Mechanism of organ dysfunction in critical illness. Springer.

- Vijay P. (2000):  
Adrenomedullin in the treatment of pulmonary hypertension.  
Heart 84 (6), 575-576.
- Vincent J.L., Sakr Y., Sprung C.L., Ranieri V.M., Reinhart K., Gerlach H., Moreno R., Carlet J., Le Gall J.R., and Payen D. (2006):  
Sepsis in European intensive care units: results of the SOAP study.  
Crit Care Med 34 (2), 344-353.
- Volk T. and Kox W.J. (2000):  
Endothelium function in sepsis.  
Inflamm Res 49 (5), 185-198.
- Walley K.R. (1996):  
Heterogeneity of oxygen delivery impairs oxygen extraction by peripheral tissues: theory.  
J Appl Physiol 81 (2), 885-894.
- Walmrath D., Scharmann M., Konig R., Pilch J., Grimminger F., and Seeger W. (1993):  
Staphylococcal alpha-toxin induced ventilation-perfusion mismatch in isolated blood-free perfused rabbit lungs.  
J Appl Physiol 74 (4), 1972-1980.
- Wang P. (2001):  
Adrenomedullin and cardiovascular responses in sepsis.  
Peptides 22 (11), 1835-1840.
- Wang P., Yoo P., Zhou M., Cioffi W.G., Ba Z.F., and Chaudry I.H. (1999):  
Reduction in vascular responsiveness to adrenomedullin during sepsis.  
J Surg Res 85 (1), 59-65.
- Westphal M., Freise H., Kehrel B.E., Bone H.G., Van Aken H., and Sielenkamper A.W. (2004):  
Arginine vasopressin compromises gut mucosal microcirculation in septic rats.  
Crit Care Med 32 (1), 194-200.
- Yang S., Zhou M., Chaudry I.H., and Wang P. (2002a):  
Novel approach to prevent the transition from the hyperdynamic phase to the hypodynamic phase of sepsis: role of adrenomedullin and adrenomedullin binding protein-1.  
Ann Surg 236 (5), 625-633.
- Yang S., Zhou M., Fowler D.E., and Wang P. (2002b):  
Mechanisms of the beneficial effect of adrenomedullin and adrenomedullin-binding protein-1 in sepsis: down-regulation of proinflammatory cytokines.  
Crit Care Med 30 (12), 2729-2735.
- Yoshimoto R., Mitsui-Saito M., Ozaki H., and Karaki H. (1998):  
Effects of adrenomedullin and calcitonin gene-related peptide on contractions of the rat aorta and porcine coronary artery.  
Br J Pharmacol 123 (8), 1645-1654.

Zhou M., Chaudry I.H., and Wang P. (2001):  
The small intestine is an important source of adrenomedullin release during polymicrobial sepsis.  
*Am J Physiol Regul Integr Comp Physiol* 281 (2), R654-R660.