

6 Literaturverzeichnis:

- Aarnio M, Mecklin JP, Aaltonen LA, Nystrom-Lahti M, Jarvinen HJ. Life-time risk of different cancers in hereditary non-polyposis colorectal cancer (HNPCC) syndrome. *Int J Cancer* 1995;64:430-3
- Al-Chalabi, Leigh PN. Recent advances in amyotrophic lateral sclerosis. *Curr Opin Neurol* 2000;13:397-405
- Almoguera C, Shibata D, Forrester K, Martin J, Arnheim N, Perucho M. Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes. *Cell* 1988;53:549-54
- Ammann RW, Bühler H, Münch R, Freiburghaus AW, Siegenthaler W. Differences in the natural history of idiopathic (nonalcoholic) and alcoholic chronic pancreatitis. A comparative long-term study of 287 patients. *Pancreas* 1987;2:368-77
- Antonarakis SE. Recommendations for a nomenclature system for human gene mutations. Nomenclature Working Group. *Hum Mutat* 1998;11:1-3
- Audrézet MP, Chen JM, Le Marèchal C, Ruszniewski P, Robaszkiewicz M, Raguénès O, Quéré I, Scotet V, Férec C. Determination of the relative contribution of three genes – the cystic fibrosis transmembrane conductance regulator gene, the cationic trypsinogen gene, and the pancreatic secretory trypsin inhibitor gene – to the etiology of idiopathic chronic pancreatitis. *Eur J Hum Gen* 2002;10:100-06
- Bank S, Marks IN, Novis B. Sweat electrolytes in chronic pancreatitis. *Am J Dig Dis* 1978;23:178-81
- Baribault H, Price J, Miyai K, Oshima RG. Mid-gestational lethality in mice lacking keratin 8. *Genes Dev* 1993;7:1191-202
- Baribault H, Penner J, Iozzo RV, Wilson-Heiner M. Colorectal hyperplasia and inflammation in keratin 8-deficient FVB/N mice. *Genes Dev* 1994;15:2964-73
- Bartelt DC, Shapanka R, Greene LJ. The primary structure of the human pancreatic secretory trypsin inhibitor. Amino acid sequence of the reduced S-aminoethylated protein. *Arch Biochem Biophys* 1977;179:189-99
- Bhatia E, Choudhuri G, Sikora SS, Landt O, Kage A, Becker M, Witt H. Tropical calcific

- pancreatitis: strong association with SPINK1 trypsin inhibitor mutations. *Gastroenterology* 2002;123:1020-25
- Blessing M, Rüther U, Franke WW. Ectopic synthesis of epidermal cytokeratins in pancreatic islet cells of transgenic mice interferes with cytoskeletal order and insulin production. *J Cell Biol* 1993;120:743-66.
- Bornman PC, Beckingham IJ. Chronic Pancreatitis *BMJ* 2001;322:660-3
- Bradley EL 3rd. A clinically based classification system for acute pancreatitis. Summary of the International Symposium on Acute Pancreatitis, Atlanta, Ga, September 11 through 13, 1992. *Arch Surg* 1993; 128:586-90.
- Buning C, Halangk J, Dignass A, Ockenga J, Deindl P, Nickel R, Genschel J, Landt O, Lochs H, Schmidt H, Witt H. Keratin 8 Y54H and G62C mutations are not associated with inflammatory bowel disease. *Dig Liver Dis.* 2004;36:388-91
- Caldas C, Kern SE. K-ras mutation and pancreatic adenocarcinoma. *Int J Pancreatol* 1995;18:1-6
- Casanova ML, Bravo A, Ramirez A, Morreale de Escobar G, Were F, Merlino G, Vidal M, Jorcano JL. Exocrine pancreatic disorders in transgenic mice expressing human keratin 8. *J Clin Invest* 1999;103:1587-95
- Cavestro GM, Frulloni L, Nouvenne A, Neri TM, Calore B, Ferri B, Bovo P, Okolicsanyi L, Di Mario F, Cavallini G. Keratin 8 mutations in patients with chronic pancreatitis. *Dig Liver Dis* 2003;35:416-20
- Chandak GR, Idris MM, Reddy DN, Bhaskar S, Sriram PV, Singh L. Mutations in the pancreatic secretory trypsin inhibitor gene (PSTI/SPINK1) rather than the cationic trypsinogen gene (PRSS1) are significantly associated with tropical calcific pancreatitis. *J Med Genet* 2002;39:347-51
- Cheng JQ, Ruggeri B, Klein WM, Sonoda G, Altomare DA, Watson DK, Testa JR. Amplification of AKT2 in human pancreatic cells and inhibition of AKT2 expression and tumorigenicity by antisense RNA. *Proc Natl Acad Sci U S A* 1996;93:3636-41
- Chou CF, Smith AJ, Omary MB. Characterization and Dynamics of O-Linked Glycosylation of Human Cytokeratin 8 and 18. *J Biol Chem* 1992;267:3901-6
- Chou CF, Omary MB. Mitotic arrest with anti-microtubule agents or okadaic acid is

- associated with increased glycoprotein terminal GlcNAc's. *J Cell Sci* 1994;*107*:1833-43
- Cohn JA, Friedman KJ, Noone PG, Knowles MR, Silverman LM, Jowell PS. Relation between mutations of the cystic fibrosis gene and idiopathic pancreatitis. *N Engl J Med* 1998;*339*:653-8
- Colomb E, Guy O, Deprez P, Michel R, Figarella C. The two human trypsinogens: catalytic properties of the corresponding trypsins. *Biochim Biophys Acta* 1978;*525*:186-93
- Colomb E, Figarella C. Comparative studies on the mechanism of activation of the two human trypsinogens. *Biochim Biophys Acta* 1979;*571*:343-51
- Criddle DN, Murphy J, Fistetto G, Barrow S, Tepikin AV, Neoptolemos JP, Sutton R, Petersen OH. Fatty acid ethyl esters cause pancreatic calcium toxicity via inositol trisphosphate receptors and loss of ATP synthesis. *Gastroenterology* 2006;*130*:781-93
- De Angelis C, Valente G, Spaccapietra M, Angonese C, Del Favero G, Naccarato R, Andriulli A. Histological study of alcoholic, nonalcoholic, and obstructive chronic pancreatitis. *Pancreas* 1992;*7*:193-6
- Durbec JP, Sarles H. Multicenter survey of the etiology of pancreatic diseases. Relationship between the relative risk of developing chronic pancreatitis and alcohol, protein and lipid consumption. *Digestion* 1978;*18*:337-50
- Freedman SD, Blanco P, Shea JC, Alvarez JG. Mechanisms to explain pancreatic dysfunction in cystic fibrosis. *Med Clin North Am* 2000;*84*:657-64
- Fuchs E, Weber K. Intermediate filaments: structure, dynamics, function and disease. *Annu Rev Biochem* 1994;*63*:345-82
- Fuchs E, Cleveland DW. A Structural Scaffolding of Intermediate Filaments in Health and Disease. *Science* 1998;*279*:514-19
- Gilbert S, Loranger A, Daigle N, Marceau N. Simple epithelium keratins 8 and 18 provide resistance to FAS-mediated apoptosis. The protection occurs through a receptor-targeting modulation. *J Cell Biol* 2001;*154*:763-73
- Goggins M, Hruban RH, Kern SE. BRCA2 is inactivated late in the development of pancreatic intraepithelial neoplasia: evidence and implications. *Am J Pathol*

2000;156:1767-71

- Goldstein AM, Fraser MC, Struewing JP, Hussussian CJ, Ranade K, Zametkin DP, Fontaine LS, Organic SM, Dracopoli NC, Clark WH Jr. Increased risk of pancreatic cancer in melanoma-prone kindreds with p16INK4 mutations. *N Engl J Med* 1995;333:970-4
- Gorry MC, Ghabbaizedeh D, Furey W, Gates LK Jr, Preston RA, Aston CE, Zhang Y, Ulrich C, Ehrlich GD, Whitcomb DC. Mutations in the cationic trypsinogen gene are associated with recurrent acute and chronic pancreatitis. *Gastroenterology* 1997;113:1063-8
- Gruis NA, van der Velden PA, Sandkuijl LA, Prins DE, Weaver-Feldhaus J, Kamb A, Bergman W, Frants RR. Homozygotes for CDKN2 (p16) germline mutation in Dutch familial melanoma kindreds. *Nat Genet* 1995;10:351-3
- Hahn SA, Schutte M, Hoque AT, Moskaluk CA, da Costa LT, Rozenblum E, Weinstein CL, Fischer A, Yeo CJ, Hruban RH, Kern SE. DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1. *Science* 1996;271:350-3
- Halangk J, Berg T, Puhl G, Müller T, Nickel R, Kage A, Landt O, Luck W, Wiedenmann B, Neuhaus P, Witt H. Keratin 8 Y54H and G62C mutations are not associated with liver disease. *J Med Gen* 2004;41:e92
- Hassan Z, Mohan V, Ali L, Allotey R, Barakat K, Faruque MO, Deepa R, McDermott MF, Jackson AE, Cassell P, Curtis D, Gelding SV, Vijayaravaghan S, Gyr N, Whitcomb DC, Khan AK, Hitman GA. SPINK1 is a susceptibility gene for fibrocalculous pancreatic diabetes in subjects from the Indian subcontinent. *Am J Hum Genet* 2002;71:964-8
- Hesse M, Berg T, Wiedenmann B, Spengler U, Woitas RP, Magin TM. A frequent keratin 8 p.L227L polymorphism, but no point mutations in keratin 8 and 18 genes, in patients with various liver disorders. *J Med Genet* 2004;41:e42
- Hruban RH, Wilentz RE, Kern SE. Genetic progression in the pancreatic ducts. *Am J Pathol* 2000;156:1821-5
- Hruban RH, Adsay NV, Albores-Saavedra J, Compton C, Garrett ES, Goodman SN, Kern SE, Klimstra DS, Kloppel G, Longnecker DS, Luttges J, Offerhaus GJ. Pancreatic intraepithelial neoplasia. A new nomenclature and classification

- system for pancreatic duct lesions. *Am J Surg Pathol* 2001;25:579-86
- Hutchinson CJ, Alvarez-Reyes M, Vaughan OA. Lamins in disease: why do ubiquitously expressed nuclear envelope proteins give rise to tissue-specific disease phenotypes? *J Cell Sci* 2001;114:9-19
- Irvine AD, McLean WH. Human keratin diseases: the increasing spectrum of disease and subtlety of the phenotype-genotype correlation. *Br J Dermatol* 1999;140:815-28
- Janmey PA, Shah JV, Janssen KP, Schliwa M. Viscoelasticity of intermediate filament networks. *Subcell Biochem* 1998;31:381-97
- Jemal A, Thomas A, Murray T, Thun M. Cancer statistics, 2002. *CA Cancer J Clin* 2002;52:23-47
- Kazal LA, Spicer DS, Brahinsky RA. Isolation of a crystalline trypsininhibitor-anticoagulant protein from the pancreas. *J Am Chem Soc* 1948;70:304-40
- Kiraly O, Guan L, Szepessy E, Toth M, Kukor Z, Sahin-Toth M. Expression of human cationic trypsinogen with an authentic N terminus using intein-mediated splicing in aminopeptidase P deficient *Escherichia coli*. *Protein Expr Purif.* 2006;48:104-11
- Krauss S, Franke WW. Organization and sequence of the human gene encoding cytokeratin 8. *Gene* 1990;86:241-9
- Ku NO, Omary MB. Identification of the major physiologic phosphorylation site of human keratin 18: potential kinases and a role in filament reorganization. *J Cell Biol* 1994;127:161-71
- Ku NO, Liao J, Chou CF, Omary MB. Implications of intermediate filament protein phosphorylation. *Cancer Metastasis Rev* 1996;15:429-44
- Ku NO, Michie SA, Soetikno RM, Resurreccion EZ, Broome RL, Oshima RG, Omary MB. Susceptibility to hepatotoxicity in transgenic mice that express a dominant-negative human keratin 18 mutant. *J Clin Invest* 1996;98:1034-46
- Ku NO, Omary MB. Phosphorylation of human keratin 8 in vivo at conserved head domain serine 23 and at epidermal growth factor-stimulated tail domain serine 431. *J Biol Chem* 1997;272:7556-64
- Ku NO, Zhou X, Toivola DM, Omary MB. The cytoskeleton of digestive epithelia in

- health and disease. *Am J Physiol Gastrointest Liver Physiol* 1999;277:G1108-G1137
- Ku NO, Gish R, Wright TL, Omary MB. Keratin 8 mutations in patients with cryptogenic liver disease. *N Engl J Med* 2001;344:1580-7
- Ku NO, Darling JM, Krams SM, Esquivel CO, Keeffe EB, Sibley RK, Lee YM, Wright TL, Omary MB. Keratin 8 and 18 mutations are risk factors for developing liver disease of multiple etiologies. *Proc Natl Acad Sci USA* 2003;100:6063-8
- Ku NO, Lim JK, Krams SM, Esquivel CO, Keeffe EB, Wright TL, Parry DA, Omary MB. Keratins as susceptibility genes for end-stage liver disease. *Gastroenterology* 2005;129:885-93
- Lal G, Liu L, Hogg D, Lassam NJ, Redston MS, Gallinger S. Patients with both pancreatic adenocarcinoma and melanoma may harbor germline CDKN2A mutations. *Genes Chromosomes Cancer* 2000;27:358-61
- Lankisch PG, Assmus C, Maisonneuve P, Lowenfels AB. Epidemiology of pancreatic diseases in Luneburg County. A study in a defined German population. *Pancreatology* 2001;1:129-99
- Laskowski M, Wu FC. Temporary inhibition of trypsin. *J Biol Chem* 1953;204:797-805
- Lerch MM, Hoppe-Seyler P, Gerok W. Origin and development of exocrine pancreatic insufficiency in experimental renal failure. *Gut* 1994;35:401-7
- Lerch MM, Gorelick FS. Early trypsinogen activation in acute pancreatitis. *Med Clin North Am* 2000;84:549-63
- Levine AJ. p53, the cellular gatekeeper for growth and division. *Cell* 1997;88:323-31
- Li FP, Fraumeni JF Jr, Mulvihill JJ, Blattner WA, Dreyfus MG, Tucker MA, Miller RW. A cancer family syndrome in twenty-four kindreds. *Cancer Res* 1988;48:5358-62
- Liao J, Lowthert LA, Ku NO, Fernandez R, Omary MB. Dynamics of human keratin 18 phosphorylation: polarized distribution of phosphorylated keratins in simple epithelial tissues. *J Cell Biol* 1995;131:1291-301 (a)
- Liao J, Lowthert LA, Omary MB. Heat stress or rotavirus infection of human epithelial cells generates a distinct hyperphosphorylated form of keratin 8. *Exp Cell Res* 1995;219:348-57 (b)
- Liao J, Omary MB. 14-3-3 proteins associate with phosphorylated simple epithelial

- keratins during cell cycle progression and act as a solubility cofactor. *J Cell Biol* 1996;133:345-57
- Liao J, Ku NO, Omary MB. Stress, apoptosis, and mitosis induce phosphorylation of human keratin 8 at Ser-73 in tissues and cultured cells. *J Biol Chem* 1997;272:17565-73
- Loranger A, Duclos S, Grenier A, Price J, Wilson-Heiner M, Baribault H, Marceau N. Simple epithelium keratins are required for maintenance of hepatocyte integrity. *Am J Pathol* 1997;151:1673-83
- Ma L, Xu J, Coulombe PA, Wirtz D. Keratin filament suspensions show unique micromechanical properties. *J Biol Chem* 1999;274:19145-51
- Malka D, Hammel P, Maire F, Rufat P, Madeira I, Pessione F, Levy P, Ruzniewski P. Risk of pancreatic adenocarcinoma in chronic pancreatitis. *Gut* 2002;51:849-52
- Maniotis AJ, Chen CS, Ingber DE. Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure. *Proc Natl Acad Sci U S A* 1997;94:849-54
- McMillan DE, Geevarghese PJ. Dietary cyanide and tropical malnutrition diabetes. *Diabetes Care* 1979;2:202-8
- Mohan V, Nagalotimath SJ, Yajnik CS, Tripathy BB. Fibrocalculous pancreatic diabetes. *Diabetes Metab Rev* 1998;14:153-70
- Moskaluk CA, Hruban RH, Kern SE. p16 and K-ras gene mutations in the intraductal precursors of human pancreatic adenocarcinoma. *Cancer Res* 1997;57:2140-3
- Mullis KB, Faloona FA. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol* 1987;155:335-50
- Nakamura K, Sarles H, Payan H. Three-dimensional reconstruction of the pancreatic ducts in chronic pancreatitis. *Gastroenterology* 1972;62:942-9
- Nemoda Z, Sahin-Toth M. Chymotrypsin C (caldecrin) stimulates autoactivation of human cationic trypsinogen. *J Biol Chem* 2006;281:11879-86
- Noone PG, Zhou Z, Silverman LM, Jowell PS, Knowles MR, Cohn JA. Cystic fibrosis gene mutations and pancreatitis risk: relation to epithelial ion transport and trypsin inhibitor gene mutations. *Gastroenterology* 2001;121:1310-19
- Otsuki M. Chronic pancreatitis in Japan: epidemiology, prognosis, diagnostic criteria, and

- future problems. *J Gastroenterol* 2003;38:315-26
- Pfützer RH, Barmada MM, Brunskill AP, Finch R, Hart PS, Neoptolemos J, Furey WF, Whitcomb DC. SPINK1/PSTI polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. *Gastroenterology* 2000;119:615-23
- Rozenblum E, Schutte M, Goggins M, Hahn SA, Panzer S, Zahurak M, Goodman SN, Sohn TA, Hruban RH, Yeo CJ, Kern SE. Tumor-suppressive pathways in pancreatic carcinoma. *Cancer Res* 1997;57:1731-4
- Sahin-Tóth M, Tóth M. Gain-of function mutations associated with hereditary pancreatitis enhance autoactivation of human cationic trypsinogen. *Biochem Biophys Res Comm* 2000;278:286-9
- Sarles H, Adler G, Dani R, Frey C, Gullo L, Harada H, Martin E, Norohna M, Scuro LA. The pancreatitis classification of Marseilles-Rome 1988. *Scand J Gastroenterol* 1989;24:641-2
- Sarner M, Cotton PB. Definitions of acute and chronic pancreatitis. *Clin Gastroenterol* 1984;13:865-70
- Scheele G, Bartelt D, Bieger W. Characterization of human exocrine pancreatic proteins by two-dimensional isoelectric focusing/sodium dodecyl sulfate gel electrophoresis. *Gastroenterology* 1981;80:461-73
- Schneider A, Whitcomb DC, Singer MV. Animal Models in Chronic Pancreatitis – What Can We Learn? *Pancreatology* 2002;2:189-203
- Schneider A, Suman A, Rossi L, Barmada MM, Beglinger C, Parvin S, Sattar S, Ali L, Khan AK, Gyr N, Whitcomb DC. SPINK1/PSTI mutations are associated with tropical pancreatitis and type II diabetes mellitus in Bangladesh. *Gastroenterology* 2002;123:1026-30
- Schneider A, Lamb J, Barmada MM, Cuneo A, Money ME, Whitcomb DC. Keratin 8 Mutations Are Not Associated with Familial, Sporadic and Alcoholic Pancreatitis in a Population from the United States. *Pancreatology* 2006;6:103-8
- Schneider G, Lersch C, Schmid RM. Genetic alterations in pancreatic carcinoma. *Mol Cancer* 2003;2:15
- Shaper AG. Chronic pancreatic disease and protein malnutrition. *Lancet* 1960;1:1223-4
- Sharer N, Schwarz M, Malone G, Howarth A, Painter J, Super M, Braganza J.

- Mutations of the cystic fibrosis gene in patients with chronic pancreatitis. *N Engl J Med* 1998;339:645-52
- Shields JM, Pruitt K, McFall A, Shaub A, Der CJ. Understanding Ras: 'it ain't over 'til it's over'. *Trends Cell Biol* 2000;10:147-54
- Shwachman H, Lebenthal E, Khaw KT. Recurrent acute pancreatitis in patients with cystic fibrosis with normal pancreatic enzymes. *Pediatrics* 1975;55:86-95
- Silverman DT. Risk factors for pancreatic cancer: a case control study based on direct interviews. *Teratog Carcinog Mutagen* 2001;21:7-15
- Simon B, Printz H. Epidemiological trends in pancreatic neoplasias. *Dig Dis* 2001;19:6-14
- Steer ML, Meldolesi J. The cell biology of experimental pancreatitis. *N Engl J Med* 1987;316:144-50
- Strelkov SV, Herrmann H, Aebi U. Molecular architecture of intermediate filaments. *BioEssays* 2003;25:243-51
- Strnad P, Lienau TC, Tao GZ, Ku NO, Magin TM, Omary MB. Denaturing temperature selection may underestimate keratin mutation detection by DHPLC. *Hum Mutat* 2006;27:444-52
- Strong LC, Stine M, Norsted TL. Cancer in survivors of childhood soft tissue sarcoma and their relatives. *J Natl Cancer Inst* 1987;79:1213-20
- Su GH, Hruban RH, Bansal RK, Bova GS, Tang DJ, Shekher MC, Westerman AM, Entius MM, Goggins M, Yeo CJ, Kern SE. Germline and somatic mutations of the STK11/LKB1 Peutz-Jeghers gene in pancreatic and biliary cancers. *Am J Pathol* 1999;154:1835-40
- Tada M, Omata M, Kawai S, Saisho H, Ohto M, Saiki RK, Sninsky JJ. Detection of ras gene mutations in pancreatic juice and peripheral blood of patients with pancreatic adenocarcinoma. *Cancer Res* 1993;53:2472-4
- Tada M, Ohashi M, Shiratori Y, Okudaira T, Komatsu Y, Kawabe T, Yoshida H, Machinami R, Kishi K, Omata M. Analysis of K-ras gene mutation in hyperplastic duct cells of the pancreas without pancreatic disease. *Gastroenterology* 1996;110:227-31
- Tandon RK, Sato N, Garg PK. Chronic pancreatitis: Asia-Pacific consensus report. *J*

Gastroenterol Hep 2002;17:508-18

Terhune PG, Phifer DM, Tosteson TD, Longnecker DS. K-ras mutation in focal proliferative lesions of human pancreas. *Cancer Epidemiol Biomarkers Prev* 1998;7:515-21

Toivola DM, Omary MB, Ku NO, Peltola O, Baribault H, Eriksson JE. Protein phosphatase inhibition in normal and keratin 8/18 assembly-incompetent mouse strains supports a functional role of keratin intermediate filaments in preserving hepatocyte integrity. *Hepatology* 1998;28:116-28

Toivola DM, Baribault H, Magin T, Michie SA, Omary MB. Simple epithelial keratins are dispensable for cytoprotection in two pancreatitis models. *Am J Physiol Gastrointest Liver Physiol* 2000;279:G1343-G1354

Weizman Z, Durie PR. Acute pancreatitis in childhood. *J Pediatr* 1988;113:24-9

Werner J, Laposata M, Fernandez-del Castillo C, Saghir M, Iozzo RV, Lewandrowski KB, Warshaw AL. Pancreatic injury in rats induced by fatty acid ethyl ester, a nonoxidative metabolite of alcohol. *Gastroenterology* 1997;113:286-94

Whitcomb DC, Gorry MC, Preston RA, Furey W, Sossenheimer MJ, Ulrich CD, Martin SP, Gates LK Jr, Amann ST, Toskes PP, Liddle R, McGrath K, Uomo G, Post JC, Ehrlich GD. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nat Genet* 1996;14:141-5

Whitcomb DC. Early trypsinogen activation in acute pancreatitis. *Gastroenterology* 1999;116:770-2

Wilentz RE, Geradts J, Maynard R, Offerhaus GJ, Kang M, Goggins M, Yeo CJ, Kern SE, Hruban RH. Inactivation of the p16 (INK4A) tumor-suppressor gene in pancreatic duct lesions: loss of intranuclear expression. *Cancer Res* 1998;58:4740-4

Wilentz RE, Iacobuzio-Donahue CA, Argani P, McCarthy DM, Parsons JL, Yeo CJ, Kern SE, Hruban RH. Loss of expression of Dpc4 in pancreatic intraepithelial neoplasia: evidence that DPC4 inactivation occurs late in neoplastic progression. *Cancer Res* 2000;60:2002-6

Witt H, Luck W, Becker M. A signal peptide cleavage site mutation in the cationic trypsinogen gene is strongly associated with chronic pancreatitis.

Gastroenterology 1999;117:7-10

- Witt H, Luck W, Hennies HC, Claßen M, Kage A, Laß U, Landt O, Becker M. Mutations in the gene encoding the serine protease inhibitor, Kazal type 1 are associated with chronic pancreatitis. *Nat Genet* 2000;25:213-6
- Witt H, Hennies HC, Becker M. SPINK1 Mutations in Chronic Pancreatitis. *Gastroenterology* 2001;120:1060-1
- Witt H, Luck W, Becker M, Böhmig M, Kage A, Truninger K, Ammann RW, O'Reilly D, Kingsnorth A, Schulz HU, Halangk W, Kielstein V, Knoefel WT, Teich N, Keim V. Mutation in the SPINK1 trypsin inhibitor gene, alcohol use, and chronic pancreatitis. *JAMA* 2001;285:2716-17
- Witt H. Chronic pancreatitis and cystic fibrosis. *Gut* 2003;52(Suppl II):ii31-41
- Witt H, Sahin-Toth M, Landt O, Chen JM, Kahne T, Drenth JP, Kukor Z, Szepessy E, Halangk W, Dahm S, Rohde K, Schulz HU, Le Marechal C, Akar N, Ammann RW, Truninger K, Bargetzi M, Bhatia E, Castellani C, Cavestro GM, Cerny M, Destro-Bisol G, Spedini G, Eiberg H, Jansen JB, Koudova M, Rausova E, Macek M Jr, Malats N, Real FX, Menzel HJ, Moral P, Galavotti R, Pignatti PF, Rickards O, Spicak J, Zarnescu NO, Bock W, Gress TM, Friess H, Ockenga J, Schmidt H, Pfutzer R, Lohr M, Simon P, Weiss FU, Lerch MM, Teich N, Keim V, Berg T, Wiedenmann B, Luck W, Groneberg DA, Becker M, Keil T, Kage A, Bernardova J, Braun M, Guldner C, Halangk J, Rosendahl J, Witt U, Treiber M, Nickel R, Ferec C. A degradation-sensitive anionic trypsinogen (PRSS2) variant protects against chronic pancreatitis. *Nat Genet* 2006;38:668-673
- Yanagisawa A, Ohtake K, Ohashi K, Hori M, Kitagawa T, Sugano H, Kato Y. Frequent c-Ki-ras oncogene activation in mucous cell hyperplasias of pancreas suffering from chronic inflammation. *Cancer Res* 1993;53:953-6
- Zatloukal K, Stumptner C, Lehner M, Denk H, Baribault H, Eshkind LG, Franke WW. Cytokeratin 8 protects from hepatotoxicity, and its ratio to cytokeratin 18 determines the ability of hepatocytes to form Mallory bodies. *Am J Pathol* 2000;156:1263-74