The Role of Transcription Factors Klf4 and Icsbp in the Development of Myeloid Cells

Dissertation

zur Erlangung des Doktorgrades der Naturwissenschaften

eingereicht am Fachbereich Biologie, Chemie, Pharmazie der Freien Universität Berlin

vorgelegt von

Maja Milanovic

aus Zrenjanin, Serbien

Berlin, 2006

This work was executed at the Leibniz Institute for Molecular Pharmacology, Berlin, in the Department of Molecular Genetics under the supervision of Dr. Dirk Carstanjen and Prof. Dr. Ivan Horak.

Referees:

Prof. Dr. I. Horak, Department of Molecular Genetics, Leibniz Institute for Molecular Pharmacology, Berlin

Prof. Dr. H. Oschkinat, Department of NMR-supported Structural Biology, Leibniz Institute for Molecular Pharmacology, Berlin

Date of defense: 9.10.2007

Statement

Herewith, I declare that I prepared the presented dissertation myself and without the use illegitimate aids. I used no other but the indicated sources and accessories. Further, I insure that this dissertation has not before been submitted to any other faculty for examination.

Berlin, _____

Maja Milanovic

Table of Contents

1. INTRODUCTION	1
1.1. Hematopoiesis, general terms	1
1.2. Hierarchy of the hematopoietic process	1
1.2.1. Hematopoietic stem cells	1
1.2.2. Maturation of hematopoietic progenitor cells	1
1.2.3. The mature cells of the myeloid lineage	3
1.2.3.1. Granulocytes	3
1.2.3.2. Monocytes/macrophages	4
1.3. Regulation of hematopoiesis	5
1.3.1 Molecular basis of hematopoietic lineage commitment	5
1.3.2 The role of cytokines in controlling hematopoiesis1.3.3. Transcriptional control of hematopoiesis	6 9
1.4 The study of transcription factors in the myeloid development	11
1.4.1. Icsbp	11
1.4.1.1. Mice with a targeted deletion of Icsbp	14
1.4.1.2. Pursuing the mechanism of Icsbp action in myeloid development –	
global gene expression analysis	15
1.5. Klf4	15
1.6. The expression of eosinophil specific genes in the Icsbp ^{-/-} GMP	17
2. PROBLEM STATEMENT	18
	10
3. MATERIALS	19
3.1. Chemicals and reagents	19
3.2. Buffers	19
3.3. Antibodies	21
3.3.1. Antibodies used for Fluorescence Activated Cell Sorting (FACS) and	21
Magnetic Cell Sorting (MACS) 3.3.2. Antibodies used for Western blotting	21 21
3.4. Recombinant cytokines and proteins	21
3.5. Enzymes	21
3.6. Kits	22
3.7. Cell culture media	22
3.7.1. Basic components and supplements	22
3.7.2. Media	22
3.8. Bacterial media and supplements	23
3.9. Oligonucleotides	23
3.9.1. Primers for RT-PCR and Northern blotting probes	23
3.9.2. Primers used for real-time PCR	23
3.9.3. Primers for PCR fragments used for cloning	24
3.10. Equipment	25
3.11. Mice	25
4. METHODS	26
4.1. Cell culture	26
4.1.1. Isolation and cultivation of primary cells	26
4.1.2. Cultivation of cell lines	27
4.2. Molecular biology methods	27
4.2.1. Genomic DNA isolation from cells and organs	27
4.2.2. Plasmid DNA isolation from bacteria	27 27
4.2.3. Electrophoretic separation of DNA fragments 4.2.4. Total RNA isolation from cells	27
4.2.4. Potar KNA Isolation from cens 4.2.5. Photometric determination of the DNA and RNA concentration	28 28
4.2.6. Reverse transcription from the RNA	28
4.2.7. Polymerase Chain Reaction (PCR)	29
4.2.7.1. Qualitative and semiquantitative PCR amplification	29

4.2.7.2. Real-time quantitative PCR	29
4.2.8. Northern blotting	30
4.2.8.1. The preparation of the gel and the probes	30
4.2.8.2. The electrophoresis and blotting procedure	30
4.2.8.3. The preparation of the radioactive labeled probe	30
4.2.8.4. The hybridization, washing and exposing procedure	31
4.2.8.5. The stripping and reprobing procedure	31
4.3. The retroviral constructs	31
4.3.1. Vector	31
4.3.2. Cloning of the retroviral constructs	32
4.3.2.1. PCR amplification of desired cDNA sequences (<i>Icsbp</i> , <i>Klf4</i> , $p21^{Waf1}$)	32
4.3.2.2. Digestion	33
4.3.2.3. Ligation	33
4.3.2.4. Retransformation	34
4.3.2.5. The colony screening	34
4.3.3. Production of retrovirus containing supernatants	35
4.3.4. Determining the titer of the retrovirus containing supernatans	35
4.3.5. Infection of bone marrow cells by retrovirus constructs	36
4.4. Progenitor enrichment of bone marrow cells	37
4.4.1. 5-fluorouracil (5-FU) treatment of mice	37
4.4.2. Depletion of mature cells by MACS	37
4.4.3. Mononuclear cell enrichment by Biocoll gradient centrifugation	38
4.5. The Colony Forming Unit (CFU) assay	38
4.6. Fluorescence Assisted Cell Sorting (FACS)	38
4.6.1. Cell labeling for FACS	38
4.6.2. Propidium iodide staining for cell cycle and apoptosis analysis	39
4.6.3. The cell sorting	39
4.7. Analysis of hematopoietic parameters	40
4.7.1. Automated blood cell counting	40
4.7.2. Analysis of cell distribution in organs	40
4.8. The mouse inflammation models	40
4.8.1. Thioglycolate induced sterile peritoneal inflammation	40
4.8.2. Nippostrogylus brasiliensis infection	40
	40
4.9. Histopathology	
4.10. Protein biochemistry methods	41
4.10.1. Total protein extraction from cells	41
4.10.2. Determining the protein concentration by Bradford method	41
4.10.3. The denaturing discontinuous polyacrylamide gel electrophoresis	
(SDS-PAGE) of proteins	42
4.10.3.1. The gel preparation	42
4.10.3.2. The electrophoresis and blotting procedure	42
4.10.3.3. The specific protein detection	43
4.10.4. The cytokine concentration determination by ELISA	43
4.11. Microscopy	43
4.11.1. Preparation of slides	43
4.11.2. May-Grünwald-Giemsa staining	43
4.11.3. Fluorescent microscopy	44
4.11.5. Fuorescent meroscopy	
5. RESULTS	45
5.1. Klf4 over-expression in bone marrow progenitors results in increased percent	
of macrophages	45
5.1.1. Constitutive Klf4 over-expression	45
5.1.2. Inducible Klf4 over-expression	48
5.1.2.1. Klf4 over-expression reduces proliferative capacity of cells	49
5.1.2.2. Klf4 over-expression increases the percentage of M-CFUs	50
5.1.2.2. Klif4 is expressed in primary macrophages, but not in granulocytes	54
5.1.2.5. Kn4 is expressed in primary macrophages, but not in granulocytes 5.2 . Analysis of the hematopoietic system in Klf4 ^{-/-} mice	55
5.2. Analysis of the hematopoletic system in K114 mice $5.2.1$. Conditional Klf4 deletion in Klf4 ^{fl/fl} mice	
	56
5.2.2. Analysis of the immune cell distribution in hematopoietic tissues shows no	50
overt deregulation in hematopoiesis of Klf4 ^{-/-} mice	58

5.2.3. Clonogenic capacity and differentiation pattern of bone marrow progenitors	
is not altered by Klf4 deletion	59
5.3. p21 ^{Waf1} is a downstream target gene of K1f4	60
5.3.1. Expression of p21 in mature immune cells is limited to monocytes/macrophage lineage 5.3.2. Klf4 up regulates $p21^{Waf1}$ expression in bone marrow macrophages	61
5.3.2. Klf4 up regulates p21 ^{Waf1} expression in bone marrow macrophages	62
5.3.3. Over-expression of $p21^{war1}$ in the bone marrow progenitors enhances	
macrophage differentiation	63
5.3.3.1 Constitutive p21 ^{Waf1} over-expression 5.3.3.2. p21 ^{Waf1} -ER ^{T2} over-expression	63
5.3.3.2. $p21^{Waf1}$ -ER ^{T2} over-expression	65
5.4. Icsbp and Klf4 induce macrophage differentiation through independent pathways	69
5.4.1. Icsbp over-expression leads to macrophage development in Klf4 ^{-/-} progenitors	69
5.5. Analysis of the molecular mechanism of Klf4 activity in hematopoietic cells	71
5.5.1. The over-expression of Klf4-ER ^{$T2$} actively induces the macrophage	
differentiation program in uncommitted myeloid progenitors	71
5.5.2. The p21-ER ^{T_2} over-expression leads to the same gene expression changes as	
Klf4-ER ^{T2} over-expression	72
5.5.3. The analysis of the molecular mechanism of Klf4 activity by using Klf4 mutants	74
5.6 Analysis of the Icsbp role in another aspect of myelopoiesis – the generation	
of eosinophils	80
5. 6.1. The number of eosinophils is reduced in Icsbp ^{-/-} mice	81
5. 6.1.1. The local eosinophilia in response to thioglycolate elicited inflammation	
is reduced in Icsbp ^{-/-} mice	81
5.6.1.2. The systemic eosinophilia in response to parasite infection is reduced	
in Icsbp ^{-/-} mice	82
5.6.2. The generation of eosinophils from the bone marrow progenitors in response	- -
to II-5 is impaired in Icsbp ^{-/-} mice	85
5.6.2.1. The number of eosinophil committed progenitors is reduced in Icsbp ^{-/-} mice	86
5.6.2.2. The signal transduction through β_c -unit of the II-5 receptor is not disabled,	00
but the expression of the cytokine-binding α -unit is reduced in Icsbp ^{-/-} eosinophils 5.6.2.3. Icsbp ^{-/-} eosinophils are more susceptible to starvation-induced apoptosis	88 89
5.6.3. The differentiation of Icsbp ^{-/-} eosinophil progenitors is shifted towards other	09
myeloid lineages	90
5.6.4. The expression of <i>Gata1</i> is reduced in Icsbp ^{-/-} eosinophils and their precursors	90 92
5.0.4. The expression of <i>Guar</i> is reduced in resolve cosmophils and then precursors)2
6. DISCUSSION	95
))
7. CONCLUSIONS	105
7. CONCLUSIONS	103
8. SUMMARY	106
6. SOMINIAR I	100
8. ZUSAMMENFASSUNG	108
REFERENCES	111
APPENDIX	127